


Prairie Wetland Carbon Research Meeting



Northern Prairie Wildlife
Research Center
Jamestown, North Dakota
May 15-16, 2002



Prairie Wetlands of North America Important for Carbon Storage

Ned H. Euliss, Jr.,¹ R. A. Gleason,¹ A. Olness,² R. L. McDougal,³ H. R. Murkin,³ R. D. Robarts,⁴ R. A. Bourbonniere,⁵ B. G. Warner⁶

¹U.S. Geological Survey, Northern Prairie Wildlife Research Center, Jamestown, ND 58401-7317 USA. ²U.S. Department of Agriculture, North Central Soil Conservation Research Laboratory, Morris, MN 56267-1065 USA. ³Ducks Unlimited Canada, Institute for Wetland and Waterfowl Research, Oak Hammock, MB R0C 2Z0 Canada. ⁴Environment Canada, National Water Research Institute, Saskatoon, SK S7N 35H Canada. ⁵Environment Canada, National Water Research Institute, Burlington, ON L7R 4A6 Canada. ⁶Wetlands Research Centre, University of Waterloo, Waterloo, ON N2L 3G1 Canada.

Wetlands in the prairie region of North America are generally assumed to be inconsequential to the total wetland carbon pool. In an effort to determine what these wetland stocks might be and if they represent a significant pool, we evaluated the potential of prairie pothole wetlands in North America to sequester carbon. Previous research suggests that prairie wetlands traditionally functioned as net sinks for atmospheric carbon, but cultivation, the current principal land use, has shifted their function from net sinks to net sources of atmospheric carbon. Land use changes that lead to protection of existing wetland carbon stocks, restoration of new wetland stocks, and cessation of activities that remove carbon stocks such as conventional agriculture may be expected to have a significant impact on total wetland carbon stocks in this region. Of the wetland area in the prairie pothole region, over 50% in the United States and 46% in Canada has been converted to cropland. Our analysis suggests that greater amounts of atmospheric carbon could be stored in wetlands in the prairie pothole region through restoration programs than on no-tillage cropland even though the acreage of wetlands is much smaller. However, both land use activities should be encouraged given they lead to net gain in carbon sinks.