

ISOMERIC EFFECTS ON THIOSULFATE TRANSFORMATION AND
DETOXIFICATION OF 1,3-DICHLOROPROPENE

QIQUANG WANG, JIANYING GAN,* SHARON K. PAPIERNIK, and SCOTT R. YATES

U.S. Department of Agriculture Agricultural Research Service, Soil Physics and Pesticides Research Unit, 450 West Big Springs Road,
Riverside, California 92507

(Received 20 June 2000; Accepted 26 September 2000)

Abstract—The fumigant 1,3-dichloropropene (1,3-D) is one of the most heavily used pesticides but also a suspected carcinogen. Previous research has shown that 1,3-D was rapidly transformed and detoxified by ammonium thiosulfate (ATS), a sulfur and nitrogen fertilizer. As common formulations contain *cis* and *trans* isomers at roughly equivalent ratios, this study was conducted to understand isomeric differences in thiosulfate transformation and detoxification of 1,3-D. Under the same conditions, reaction of *cis*-1,3-D with thiosulfate was more than three times faster than *trans*-1,3-D, which was correlated with a lower reaction activation energy for the *cis* isomer. The *trans* isomer was considerably more toxic to the luminescent bacteria *Vibrio fischeri* than the *cis* isomer, but the toxicity was reduced by 14 times after thiosulfate transformation. Mutagenic activity to strains of *Salmonella typhimurium* was observed for *trans*-1,3-D but was not detected after thiosulfate transformation. These results suggest that thiosulfate transformation detoxifies 1,3-D primarily by deactivating the *trans* isomer, and the reaction is toxicologically beneficial, as it negates the potential harmful effects of 1,3-D to the environment and human health.

Keywords—1,3-Dichloropropene Ammonium thiosulfate Soil fumigants *Vibrio fischeri* Ames test

INTRODUCTION

The fumigant 1,3-dichloropropene (1,3-D) is widely used to fumigate soil for controlling soilborne nematodes and other pests. The annual use of 1,3-D during 1987–1997 was 13 to 20 million kg, which ranked at fourth to sixth among all pesticides used in the United States [1]. This use may increase further as the other major soil fumigant, methyl bromide (MeBr), is scheduled for a complete phaseout by 2005 [2]. Formulations of 1,3-D are typically injected into soils at depths of 30 to 45 cm by tractor-pulled chisels. Because of its high vapor pressure, 1,3-D has been shown to rapidly volatilize from treated fields, losing 20 to 77% of the applied material to the atmosphere [3–7]. Because 1,3-D is a B2 carcinogen and a skin and eye sensitizer, atmospheric emission of 1,3-D is a source of air pollution [8,9]. The use of 1,3-D was suspended for four years in California during 1990–1994 after detection of high 1,3-D concentrations in the air near fumigation sites. Therefore, risk-mitigation practices are needed to prevent excessive 1,3-D emissions during its use.

In a previous study, 1,3-D was found to rapidly undergo a nucleophilic substitution reaction with thiosulfate in water and soil, producing nonvolatile ionic products [10]. It was further shown that when the soil surface was sprayed with ammonium thiosulfate (ATS) solution, 1,3-D volatilization was greatly minimized while its ability to control nematodes was not affected [11]. Thiosulfate salts such as ATS and potassium thiosulfate are fertilizers, which make this approach highly feasible for application. Because commonly used 1,3-D formulations (e.g., Telone II®, DowAgro Sciences, Indianapolis, IN, USA) are roughly 50/50 mixtures of the *cis* and *trans* isomers, it is

important to understand whether isomeric effects occur during 1,3-D's transformation by thiosulfate. Isomers of 1,3-D were found to differ significantly in their degradation rates in adapted soils [12,13]. In this study, we measured individual reaction rates of 1,3-D isomers with thiosulfate in aqueous solutions and the changes in acute toxicity and mutagenicity caused by the transformation for each isomer. The observed differences were further correlated with the structural characteristics of the isomers, and the implication of the isomeric differences for application was discussed.

MATERIALS AND METHODS

Chemicals and test organisms

The *cis* and *trans* isomers of 1,3-D were donated by Dow AgroSciences (Indianapolis, IN, USA) (purity >97.6%). Ammonium thiosulfate was purchased from Fluka (Buchs, Switzerland) (purity >99.5%). All other reagents used in this study were of analytical grade.

The test organism used in the acute toxicity assay was luminescent bacteria *Vibrio fischeri* purchased from AZUR Environmental (Carlsbad, CA, USA). The *Salmonella typhimurium* strains TA97a, TA98, TA100, and TA102 used in the mutagenicity test were obtained from the Division of Biochemistry and Cell Molecular Biology, University of California, Berkeley (Berkeley, CA, USA). Rat hepatic fractions (S9) were purchased from BioReliance-Tox Labs (Rockville, MD, USA).

Kinetic experiments in aqueous solution

In the first aqueous-phase experiment, disappearance of *cis*- and *trans*-1,3-D was determined in aqueous solutions of different thiosulfate concentrations and was used to calculate the reaction rate constant for each isomer. Solutions of ATS were prepared by dissolving the salt in deionized water at 0, 1.0, 2.0, and 4.0 mM, and 100 ml of each solution were then

* To whom correspondence may be addressed
(jgan@ussl.ars.usda.gov).

Reference to a company name or product does not imply an approval or recommendation by the U.S. Department of Agriculture to the exclusion of the others that may be suitable.

transferred into 125-ml glass serum bottles. To initiate the reaction, 0.10 ml of *cis*- or *trans*-1,3-D stock solution (10^3 mM in acetone) were added to the ATS solutions, and the initial concentration of 1,3-D isomers was thus 1.0 mM. The spiked containers were crimp sealed with aluminum caps and Teflon[®]-lined butyl rubber septa and equilibrated at $20 \pm 0.5^\circ\text{C}$ in the dark.

At different times, 0.5 ml ($\times 3$) of aliquot were withdrawn from each reacted solution using a syringe, and the sample was transferred into a 10-ml glass vial containing 5.0 ml ethyl acetate and 3.0 g anhydrous sodium sulfate. The sample vials were immediately crimp sealed and mechanically mixed on a vortexer for 2 min. A portion of the ethyl acetate phase was then transferred into a gas chromatography (GC) vial and analyzed using GC. The GC system was a Hewlett Packard HP6890 GC (Hewlett Packard, Avondale, PA, USA) with an electron capture detector. Analytical conditions were as follows: Rtx-624 column ($30 \text{ m} \times 250 \mu\text{m} \times 1.4 \mu\text{m}$, Restek, Bellefonte, PA, USA), 1.2 ml/min column flow rate, 220°C inlet temperature, and 300°C detector temperature. The oven was held at 70°C for 0.5 min and then increased at $20^\circ\text{C}/\text{min}$ to 140°C and kept at 140°C for 2.5 min. The injection volume was $2.0 \mu\text{l}$, and the split ratio was 10:1. External standards were used for calibration. The decline of fumigant concentration with time was fitted to first-order decay kinetics to obtain the degradation rate constant k_f (1/h) for each fumigant isomer. Values of k_f were further correlated with the initial thiosulfate concentration by linear regression to estimate the second-order rate constant k (1/Ms) for the $\text{S}_{\text{N}}2$ reaction of each isomer with thiosulfate.

Temperature dependence was studied in a separate aqueous-phase experiment to understand the energy dynamics of the transformation of 1,3-D isomers by thiosulfate. The reaction solutions initially contained 1.0 mM *cis*- or *trans*-1,3-D, and 2.0 mM ATS. The reaction solutions were equilibrated at 6, 20, 30, and $40 \pm 0.5^\circ\text{C}$ in the dark, and the decline of fumigant concentration was measured over time by using the same procedures as given previously. The first-order degradation rate constants were correlated to the different temperatures using the Arrhenius equation to derive the activation energy E_a (kJ/mol) for the reaction of each isomer with thiosulfate.

Toxicological assessments

The change in acute toxicity of 1,3-D isomers caused by thiosulfate transformation was evaluated using luminescent bacteria *V. fischeri*. This method has been frequently used for testing acute toxicity of both organic and inorganic toxicants in environmental samples [14–16]. Briefly, prior to toxicity assay, 1,3-D isomers (2.0 mM) and ATS (20 mM) were mixed, and the reaction solutions were equilibrated at room temperature. Analysis by GC showed that 1,3-D isomers were completely transformed by ATS in 3 d. For toxicity assay, the reacted samples were serially diluted with 2% NaCl and transferred into cuvettes. An aliquot of the revived bacteria solution was added into the cuvettes, and after 5.0 min of contact at 20°C , the light intensity at 490 nm was measured on an SLM8000 Spectrofluorometer (SLM Instruments, Urbana, IL, USA). Solutions of 1,3-D isomers of the same concentrations were simultaneously measured under the same conditions. Light intensity of ATS solutions without 1,3-D was also measured and was used as the background for calculating the relative light intensity. The effective concentration at which 50% suppression occurred, or EC50, was obtained by plotting the




Fig. 1. Enhanced transformation of *cis*- and *trans*-1,3-dichloropropene (1,3-D) in the aqueous phase by ammonium thiosulfate at 20°C . Symbols are measured data, and lines are first-order regressions. ●, *cis*-1,3-D in water; ○, *trans*-1,3-D in water; ▼, *cis*-1,3-D in 2.0 mM ATS solution; and ▽, *trans*-1,3-D in 2.0 mM ATS solution.

relative light intensity against the logarithmic sample concentration and solving for the concentration at which 50% reduction in luminescence occurred.

The mutagenic activity of 1,3-D isomers before and after thiosulfate transformation was assessed using *S. typhimurium* strains TA97a, TA98, TA100, and TA102 following the Ames test method [17]. Prior to the test, 1,3-D isomers (100 mM) and ATS (500 mM) were allowed to react in water:acetone solution (4:1) at 20°C . Analysis by GC showed that complete transformation of 1,3-D was achieved in 4 d. The reacted solutions were then used to preincubate the test strains, with or without the addition of S9, for 20 min at 20°C in test tubes and then mixed with top agar. The rat hepatic fraction S9 was used as an exogenous metabolic activation system. The samples were poured into petri dishes containing minimal nutrition media and incubated at 37°C for 2 to 3 d [18]. The amounts of transformed products added to each plate were equivalent to 0.11 to 1,100 μg fumigant before transformation. Values of mutagenicity ratio (MR) were obtained by dividing number of revertant colonies of sample plates by that of the blank control. In principle, an $\text{MR} \geq 2$ with a clear dose response suggests mutagenic activity, while an $\text{MR} < 2$ indicates a lack of activity [19]. Samples of 1,3-D isomers that did not undergo transformation by thiosulfate were also subject to the same test under similar conditions.

RESULTS AND DISCUSSION

Reaction kinetics in aqueous phase

Hydrolysis in deionized water was relatively slow for both isomers of 1,3-D, with only a limited loss occurring during the 96-h equilibration period (Fig. 1). Addition of ATS to the solution significantly accelerated the disappearance of 1,3-D isomers, and the rate of fumigant disappearance was proportional to the initial thiosulfate concentration (Fig. 1 and Table 1). Regression of fumigant concentration with time showed a first-order relationship, and the goodness of fit (R) improved as the initial thiosulfate concentration increased (Table 1), which is characteristic of a second-order reaction. Linear regression of the k_f values from Table 1 to initial thiosulfate

Table 1. First-order transformation rate constant k_f (1/h) and half-life $t_{1/2}$ (h) of *cis*- and *trans*-1,3-dichloropropene in blank and ammonium thiosulfate (ATS) solutions at 20°C

Thiosulfate concn. (mM)	<i>cis</i> -1,3-D			<i>trans</i> -1,3-D		
	k_f	$t_{1/2}$	R^a	k_f	$t_{1/2}$	R
0 (no ATS)	0.0021	330	0.80	0.0032	216	0.86
1.0	0.0216	32.1	0.97	0.0137	50.6	0.99
2.0	0.0788	8.8	0.99	0.0262	26.5	0.99
4.0	0.2076	3.3	1.00	0.0664	10.4	1.00

^a R = correlation coefficient.

concentrations validated that the overall reaction closely followed the second-order kinetics that is characteristic of S_N2 nucleophilic substitution reactions, with the correlation coefficient $R > 0.99$ for either isomer. In ATS solutions, pH (6.20) was not greatly different from that in the control solution, suggesting that pH did not play a role in thiosulfate-initiated transformation of 1,3-D isomers.

In solutions with the same initial thiosulfate concentration, transformation of *cis*-1,3-D was consistently faster than that of *trans*-1,3-D (Fig. 1 and Table 1). From the k_f values given in Table 1 for each isomer, the second-order reaction rate constant was calculated to be $1.73 \times 10^{-2}/\text{Ms}$ for *cis*-1,3-D but only $5.0 \times 10^{-3}/\text{Ms}$ for *trans*-1,3-D. This suggests that thiosulfate was about 3.5 times more efficient in reacting with *cis*-1,3-D than with *trans*-1,3-D. An examination of the stereochemistry of 1,3-D suggests that the C=C bond and the four connected single bonds (C₁-Cl, C₁-H, C₂-C₃, and C₂-H) are in the same plane (Fig. 2). Therefore, the space hindrance for thiosulfate ion to approach C₃ should be very similar for both isomers. On the other hand, it is likely that in the transition state, charge density distribution for *cis*-1,3-D is poorly balanced, with the charge center tilting to one side of the C=C bond (Fig. 2). The more polarized state may favor its interaction with the polar solvent (water), thus lowering the energy level of *cis*-1,3-D and facilitating its reaction with thiosulfate ion [20].

The transformation rate of 1,3-D isomers increased with increasing temperature (Table 2). Temperature dependence of fumigant transformation rate was found to fit well to the Arrhenius relationship, with $R \geq 0.99$. The activation energy E_a was subsequently calculated using the Arrhenius equation (Table 2). The E_a value (72.9 kJ/mol) for the reaction of *cis*-1,3-D with thiosulfate was smaller than that for *trans*-1,3-D (78.3 kJ/mol). The difference was tested to be significant at $P = 0.01$ using t test. This agrees with the observation that under the same conditions, reaction of *cis*-1,3-D with ATS was ap-




Fig. 2. Proposed steric structures and charge distribution patterns for *cis*- and *trans*-1,3-dichloropropene during their nucleophilic substitution reaction with thiosulfate in water.

preciably faster than that of *trans*-1,3-D. This also provides support to the previous postulation that during reaction with thiosulfate, the energy level of the transition state for the *cis* isomer should be lower than that for the *trans* isomer.

Reduction in toxicity of 1,3-D isomers by thiosulfate transformation

Acute toxicity of 1,3-D isomers to the luminescent bacteria was determined before and after transformation by ATS. As indicated by the bacterial EC₅₀ summarized in Table 3, the *trans* isomer was about seven times more toxic than the *cis* isomer to the test organism. Thiosulfate transformation did not significantly change the EC₅₀ of *cis*-1,3-D, which may be attributed to the fact that the nontransformed *cis* isomer (EC₅₀ >1 mM) was relatively innocuous to the test organism. However, the EC₅₀ for *trans*-1,3-D increased 14-fold after transformation by ATS, indicating that thiosulfate transformation was indeed a detoxification process for the *trans* isomer. In our previous study, where a mixture of 1,3-D isomers (*cis/trans*: 47/51) was used, an overall increase in EC₅₀ of 7.4 times was found after ATS transformation [21]. The present study showed that the bacterial toxicity of 1,3-D isomer mix-

Table 2. Effect of temperature on transformation kinetics of *cis*- and *trans*-1,3-dichloropropene (1,3-D) by ammonium thiosulfate (ATS) in aqueous solution^a

Temperature (°C)	<i>cis</i> -1,3-D			<i>trans</i> -1,3-D		
	k_f	$t_{1/2}$	R	k_f	$t_{1/2}$	R
6.0	0.0154	4.50	0.99	0.0053	132	0.95
20.0	0.0788	8.8	0.99	0.0262	26.5	0.99
30.0	0.2025	3.4	1.00	0.0760	9.1	0.99
40.0	0.4643	1.5	1.00	0.2081	3.3	1.00
E_a (kJ/mol) ^b		72.9 ± 2.1			78.3 ± 0.3	

^a k_f = first-order transformation rate constant in 1/h; $t_{1/2}$ = first-order transformation half-life in h; R = correlation coefficient.

^b E_a is activation energy that was calculated from the Arrhenius equation: $k = A \cdot e^{-E_a/RT}$, where k is the reaction rate constant at a given temperature T (°K), R is the gas molar constant, and A is a fitted constant.

Table 3. Changes in bacterial EC50 against *Vibrio fischeri* for 1,3-dichloropropene isomers after transformation by ammonium thiosulfate (ATS) in aqueous solution

1,3-D isomer	ATS transformation	EC50 (mM)	Difference
<i>cis</i>	No	1.17	No difference
	Yes	1.09	
<i>trans</i>	No	0.15	14.2 times
	Yes	2.27	

ture in the previous study could be mainly attributed to the *trans* isomer and that the effect of thiosulfate detoxification was caused primarily by the deactivation of *trans*-1,3-D.

Reduction in mutagenicity of 1,3-D isomers by thiosulfate transformation

The Ames test method has been extensively used for screening genotoxicity of environmental samples because of its high sensitivity and simplicity [19,22,23]. In this study, mutagenicity of *cis*- and *trans*-1,3-D and their thiosulfate transformation products were evaluated using the revised Ames test method [17]. For the *cis* isomer and its transformation products, all MR values were <2.0 for the test concentration range of 0.01 to 100 mM, with or without activation by S9 mixture. This suggests that *cis*-1,3-D itself was not mutagenic to the test systems under the experimental conditions, and transformation by thiosulfate did not alter this status. On the other hand, the same test showed that at the highest concentration used (100 mM), *trans*-1,3-D was mutagenic to two (TA97a and TA100) of the four test strains without S9 activation and three (TA97a, TA98, and TA100) of the four strains with S9 activation (Table 4). The highest MR values were observed with the TA100 strain, which was >4.0 with or without S9 activation. A dose-response effect for TA100 was apparent, for both systems with and without the treatment of the S9 mixture. The S9 activation system appeared to enhance the expression of mutagenicity of *trans*-1,3-D. These observations together suggest that the *trans* isomer of 1,3-D possessed mutagenic activity to some strains of *S. typhimurium*. However, after transformation by thiosulfate, all MR values fell below 2.0 for the tested concentrations, and no clear dose response (shown in Table 5 for S9-activated samples) was seen. Therefore, it may be concluded that thiosulfate transformation reduced the mutagenicity of *trans*-1,3-D below the detection limit of the method used.

Table 4. Values of mutagenicity ratio of *trans*-1,3-dichloropropene (mean \pm standard deviation)

Strains	1,110 $\mu\text{g}/\text{plate}^a$	55.5 $\mu\text{g}/\text{plate}$	2.22 $\mu\text{g}/\text{plate}$	0.11 $\mu\text{g}/\text{plate}$
Without S9 activation				
TA97a	2.10 \pm 0.05 ^b	1.02 \pm 0.08	0.51 \pm 0.08	1.14 \pm 0.12
TA98	1.06 \pm 0.10	1.14 \pm 0.18	0.92 \pm 0.01	1.10 \pm 0.05
TA100	4.26 \pm 0.28 ^b	1.26 \pm 0.00	1.04 \pm 0.14	1.00 \pm 0.04
TA102	1.38 \pm 0.16	1.13 \pm 0.03	1.10 \pm 0.04	1.15 \pm 0.10
With S9 activation				
TA97a	2.36 \pm 0.01 ^b	1.16 \pm 0.14	0.96 \pm 0.09	0.96 \pm 0.23
TA98	2.10 \pm 0.06 ^b	0.82 \pm 0.07	0.92 \pm 0.06	1.18 \pm 0.02
TA100	5.73 \pm 0.03 ^b	1.60 \pm 0.10	1.31 \pm 0.17	1.16 \pm 0.04
TA102	1.55 \pm 0.14	1.12 \pm 0.02	1.03 \pm 0.09	1.03 \pm 0.13

^a 0.11 to 1,110 $\mu\text{g}/\text{plate}$ was equivalent to 0.01 to 100 mM fumigant concentration range used for preincubation.

^b Values are >2 and defined as positive.

Table 5. Values of mutagenicity ratio for *trans*-1,3-dichloropropene after transformation by ammonium thiosulfate (with S9 activation) (mean \pm standard deviation)

Strains	1,110 $\mu\text{g}/\text{plate}^a$	55.5 $\mu\text{g}/\text{plate}$	2.22 $\mu\text{g}/\text{plate}$	0.11 $\mu\text{g}/\text{plate}$
TA97a	0.94 \pm 0.21	0.74 \pm 0.09	0.63 \pm 0.03	0.82 \pm 0.12
TA98	1.08 \pm 0.15	1.05 \pm 0.14	1.41 \pm 0.21	1.26 \pm 0.03
TA100	1.25 \pm 0.06	0.75 \pm 0.02	0.77 \pm 0.22	0.98 \pm 0.09
TA102	0.98 \pm 0.02	0.88 \pm 0.11	1.00 \pm 0.03	0.92 \pm 0.05

^a 0.11 to 1,110 $\mu\text{g}/\text{plate}$ was equivalent to 0.01 to 100 mM fumigant concentration range used for preincubation.

Bacterial mutagenicity of 1,3-D isomers was previously demonstrated using other *S. typhimurium* strains [24,25]. The reactivity was attributed to the electrophilicity of the methylene carbon (C₃ in Fig. 2), which stimulates nucleophilic substitution reactions of 1,3-D with nucleophilic centers in DNA. In mammals, glutathione-initiated biotransformation was shown to deactivate 1,3-D by causing concomitant loss of its alkylating reactivity. The reaction with thiosulfate may have deactivated 1,3-D in a similar manner by eliminating its reactivity with essential cellular macromolecules.

CONCLUSIONS

Many of the currently used chlorinated pesticides, including the soil fumigant 1,3-D, have suspected carcinogenicity and other undesirable toxicological properties, making their potential to contaminate air and water resources a concern. Common formulations of 1,3-D are mixtures of its *cis* and *trans* isomers at roughly equivalent ratios. Our study showed that the *trans* isomer was much more active than the *cis* isomer against the indicator organism *V. fischeri*. At high concentrations, *trans*-1,3-D also caused positive mutagenic reaction to strains of *S. typhimurium*. These results suggest that the *trans* isomer of 1,3-D is toxicologically more active than the *cis* isomer, and formulations that have a greater ratio of *cis*-1,3-D may be less harmful to the environment or human health and therefore should be recommended. Such formulations should not compromise the effectiveness of 1,3-D, as early studies showed that *cis*-1,3-D was considerably more effective against soil nematodes [26,27]. In fact, formulations consisting mainly of the *cis* isomer were recently introduced in countries such as the Netherlands. Because of the higher potency of *cis*-1,3-D, the overall dosage has been reduced to only a half of what was used before [28].

Emission of 1,3-D into the atmosphere is of concern because of its toxicity and carcinogenicity. Previous studies showed that 1,3-D was rapidly transformed by thiosulfate salts, and surface spray of thiosulfate salt solution substantially reduced 1,3-D emission. As ammonium and potassium thiosulfates are fertilizers, this reaction may be used as a preventive or remedial measure to mitigate the risks involved in 1,3-D uses. This study showed that the *cis* and *trans* isomers of 1,3-D differed significantly in their reaction rate with thiosulfate, with the *cis* isomer being transformed about 3.5 times faster than the *trans* isomer. This difference correlates well with the differences in steric characteristics of these isomers. Reaction with thiosulfate decreased *trans*-1,3-D's activity against *V. fischeri* by 14 times and also eliminated its mutagenicity to strains of *S. typhimurium*. These preliminary assays imply that thiosulfate transformation detoxifies 1,3-D and therefore is a toxicologically advantageous process. Further evaluation

should be conducted to develop this approach into practices for reducing 1,3-D emissions from fumigated fields or leaching into groundwater and for safer disposal of 1,3-D wastes.

Acknowledgement—The authors thank the Department of Molecular and Cell Biology, University of California, Berkeley, for providing *Salmonella* tester strains and test protocols and J. Knuteson for donating *cis*- and *trans*-1,3-D.

REFERENCES

- U.S. Environmental Protection Agency. 2000. Pesticides industry sales and usage: 1996 and 1997 market estimates. Report 733-R-99-001. Office of Pesticide Programs, Washington DC.
- U.S. Department of Agriculture. 1999. Administration extends deadline on methyl bromide ban to 2005. *Methyl Bromide Alternatives* 5:1.
- Basile M, Senesi N, Lamberti F. 1986. A study of some factors affecting volatilization losses of 1,3-dichloropropene (1,3-D) from soil. *Agric Ecosyst Environ* 17:269–279.
- Chen C, Green RE, Thomas DM, Knuteson JA. 1995. Modeling 1,3-dichloropropene fumigant volatilization with vapor-phase advection in the soil profile. *Environ Sci Technol* 29:1816–1821.
- Chen C, Green RE, Thomas DM, Knuteson JA. 1996. 1995. Modeling 1,3-dichloropropene fumigant volatilization with vapor-phase advection in the soil profile. *Environ Sci Technol* 30:359 (Correction/addition).
- Gan J, Yates SR, Wang D, Ernst FF. 1998. Effect of application methods on 1,3-dichloropropene volatilization from soil under controlled conditions. *J Environ Qual* 27:432–438.
- Wang D, Knuteson JA, Yates SR. 2000. Two-dimensional model simulation of 1,3-dichloropropene volatilization and transport in a field soil. *J Environ Qual* 29:639–644.
- Baker LW, et al. 1996. Ambient air concentrations of pesticides in California. *Environ Sci Technol* 30:1365–1368.
- United Nations Environment Programmes. 1995. The Montreal protocol on substances that deplete the ozone layer: 1994 report of the methyl bromide technical option committee. UNEP Secretariat, Nairobi, Kenya.
- Gan J, Yates SR, Becker JO, Knuteson JA. 2000. Transformation of 1,3-dichloropropene by thiosulfate salts in soil. *J Environ Qual* 29:1476–1481.
- Gan J, Becker JO, Ernst FF, Hutchinson C, Knuteson JA, Yates SR. 2000. Surface application of ammonium thiosulfate fertilizer to reduce volatilization of 1,3-dichloropropene from soil. *Pest Manage Sci* 56:264–270.
- Ou LT. 1998. Enhanced degradation of the volatile fumigant-nematicides 1,3-D and methyl bromide in soil. *J Nematol* 30:56–64.
- Chung KY, Dickson DW, Ou LT. 1999. Differential enhanced degradation of *cis* and *trans*-1,3-D in soil with a history of repeated field applications of 1,3-D. *J Environ Sci Health B* 34:749–768.
- Johnson BT. 1992. An evaluation of a genotoxicity assay with liver S9 for activation and luminescent bacteria for detection. *Environ Toxicol Chem* 11:473–480.
- Liu F, Chen JW, Kong LR, Wang LS, Wei LP, Zhang Z. 1996. Application of *Photobacterium phosphoreum* to toxicity assessment of phenylthio-carboxylates. *Chemosphere* 32:2077–2082.
- Kahru A, Tomson K, Pall T, Kulm I. 1996. Study of toxicity of pesticides using luminescent bacteria, *Photobacterium phosphoreum*. *Water Sci Technol* 33:147–154.
- Maron DM, Ames BN. 1983. Revised methods for the *Salmonella* mutagenicity test. *Mutat Res* 113:173–215.
- Schneider M, Quistad GB, Casida JE. 1999. Glutathione activation of chloropicrin in the *Salmonella* mutagenicity test. *Mutat Res* 439:233–238.
- Ames BN, McCann J, Yamasaki E. 1975. Methods for detecting carcinogens and mutagens with the *Salmonella*/mammalian-microsome mutagenicity test. *Mutat Res* 31:347–364.
- McMurray J. 1992. *Organic Chemistry*. Brooks/Cole, Pacific Grove, CA, USA, pp 364–374.
- Wang Q, Gan J, Papiernik SK, Yates SR. 2000. Transformation and detoxification of halogenated fumigants by ammonium thiosulfate. *Environ Sci Technol* 34:3717–3721.
- Schenck KM, Meier JR, Ringhand HP, Kopfler FC. 1990. Recovery of 3-chloro-4-(dichloromethyl)-5-hydroxy-2(5H)-furanone from water samples on XAD resins and the effect of chlorine on its mutagenicity. *Environ Sci Technol* 24:863–867.
- Blakey DH, Maus KL, Bell R, Bayley J, Douglas GR, Nestmann ER. 1994. Mutagenic activity of 3 industrial chemicals in a battery of in vitro and in vivo tests. *Mutat Res* 320:273–283.
- Neudecker T, Stefani A, Henschler D. 1977. In vitro mutagenicity of the soil nematicide 1,3-dichloropropene. *Experientia* 33:1084–1085.
- Creedy CL, Brooks TM, Dean BJ, Hutson DH, Wright AS. 1984. The protective action of glutathione on the microbial mutagenicity of the *Z*- and *E*-isomers of 1,3-dichloropropene. *Chem Biol Interact* 50:39–48.
- Moje W, Martin JP, Baines RC. 1957. Structural effect of some organic compounds on soil organisms and citrus seedlings grown in an old citrus soil. *J Agric Food Chem* 5:32–36.
- Youngson CR, Goring CAI. 1970. Nematicidal activity of 1,3-dichloropropene and 1,2-dichloropropane to three types of plant-parasitic nematodes. *Plant Dis Rep* 54:196–199.
- Schomaker CH, Been T. 1999. Compound models describing the relationship between dosage of (*Z*)- or (*E*)-isomers of 1,3-dichloropropene and hatching behavior of *Globodera rostochiensis*. *Nematology* 1:19–29.