

Article

Inheritance and Genetic Mapping of the Reduced Height (*Rht18*) Gene in Wheat

 Nathan P. Grant ¹, Amita Mohan ¹, Devinder Sandhu ^{2,*}  and Kulvinder S. Gill ^{1,*}
¹ Department of Crop and Soil Sciences, Washington State University—Pullman, 277 Johnson Hall, PO Box 646420, Pullman, WA 99164-6420, USA; ngrant@wsu.edu (N.P.G.); amitamohan@wsu.edu (A.M.)

² USDA-ARS Salinity Lab., 450 W. Big Springs Rd., Riverside, CA 92507, USA

* Correspondence: devinder.sandhu@ars.usda.gov (D.S.); ksgill@wsu.edu (K.S.G.); Tel.: +951-369-4832 (D.S.); +509-335-4666 (K.S.G.)

Received: 12 June 2018; Accepted: 11 July 2018; Published: 15 July 2018



Abstract: Short-statured plants revolutionized agriculture during the 1960s due to their ability to resist lodging, increased their response to fertilizers, and improved partitioning of assimilates which led to yield gains. Of more than 21 reduced-height (*Rht*) genes reported in wheat, only three—*Rht-B1b*, *Rht-D1b*, and *Rht8*—were extensively used in wheat breeding programs. The remaining reduced height mutants have not been utilized in breeding programs due to the lack of characterization. In the present study, we determined the inheritance of *Rht18* and developed a genetic linkage map of the region containing *Rht18*. The height distribution of the F₂ population was skewed towards the mutant parent, indicating that the dwarf allele (*Rht18*) is semi-dominant over the tall allele (*rht18*). *Rht18* was mapped on chromosome 6A between markers *barc146* and *cf190* with a genetic distance of 26.2 and 17.3 cM, respectively. In addition to plant height, agronomically important traits, like awns and tiller numbers, were also studied in the bi-parental population. Although the average tiller number was very similar in both parents, the F₂ population displayed a normal distribution for tiller number with the majority of plants having phenotype similar to the parents. Transgressive segregation was observed for plant height and tiller number in F₂ population. This study enabled us to select a semi-dwarf line with superior agronomic characteristics that could be utilized in a breeding program. The identification of SSRs associated with *Rht18* may improve breeders' effectiveness in selecting desired semi-dwarf lines for developing new wheat cultivars.

Keywords: *Rht18*; reduced height; wheat; semi-dwarf; linkage map

1. Introduction

The Green Revolution, in the mid-twentieth century, brought about advancements in agriculture that are still in practice to date. The introduction of semi-dwarf varieties that are more responsive to changing agriculture practices like response to fertilizers was pivotal in bringing the green revolution by increasing cereal production to meet the population demands particularly in developing countries like China, India, Brazil, and Egypt [1]. Two genotypes, Norin10 (*Rht1* (*Rht-B1b*) and *Rht2* (*Rht-D1b*)) and Akakomugi (*Rht8*), were first incorporated into breeding programs to introduce the semi-dwarf genes in wheat cultivars in the United States and Italy [2,3].

The development of semi-dwarf cultivars can be attributed to a shorter yet stronger culm that accommodates high yields and prevents lodging [4,5]. Of the 21 wheat mutants reported to be associated with height reduction, only *Rht-B1b* and/or *Rht-D1b*, *Rht8* and *Rht12* have been characterized in detail [6,7]. *Rht-B1* and *Rht-D1* are two homoeologous genes present on B and D genomes in hexaploid wheat that code for DELLA proteins, which suppress gibberellin (GA)-responsive growth [8]. Normally, GA regulates binding of the GA insensitive dwarf 1 (GID1) receptor protein with DELLA

proteins and promotes their degradation. Mutant alleles, *Rht-B1b* and *Rht-D1b*, produce DELLA proteins that do not bind GID1 resulting in growth inhibition due to insensitivity to GA [8]. Similarly, modulation in GA synthesis or signaling is known to be involved in reducing plant height in different species. Studies in *Arabidopsis*, maize [6], rice [9,10], and barley [11], suggest that GA affects the inter-nodal elongation and thus alters plant height.

Height reduction in present day cultivars of wheat is achieved mainly by *Rht-B1b* and/or *Rht-D1b*, accounting for ~95% of the cultivated wheat around the world [2]. Of the other 19 height mutants reported in wheat, only *Rht8* has been used in some European wheat cultivars. The rest have not been utilized either because of the lack of genetic characterization or mapping information. The limited genetic variability in semi-dwarf lines used in breeding programs is becoming a bottleneck for further wheat improvement, due to the association of some negative effects with the *Rht-B1b* and *Rht-D1b* genes, particularly under abiotic stresses or changing environmental conditions [12]. Currently used semi-dwarf wheat lines are defective from the perspective of GA, which plays an important role in the growth and development of the plant. These genotypes display a significant effect on early seedling growth. Specifically, coleoptile length, first leaf elongation, seedling emergence, and plant height reduction have been reported in the genotypes carrying *Rht-B1b* and *Rht-D1b* compared to tall parents [6,13]. The GA-responsive *Rht12* and *Rht13* were reported to reduce plant height with no adverse effect on the coleoptile and root trait during the seedling stage [14,15]. *Rht12* delayed ear emergence, reducing flag leaf length and grain size, while *Rht13* adversely affected the 1000 kernel weight and flag leaf length. Initially classified as GA-responsive, *Rht8* was reported to be involved in reduced sensitivity to brassinosteroids that resulted in reducing plant height [16]. The 17 other reduced-height mutants have not been fully characterized. *Rht18* was found to be GA-sensitive and was identified as a possible reduced height mutant candidate for future breeding programs [4,17]. In durum wheat, *Rht18* was previously mapped to the short arm of chromosome 6A at the same locus as *Rht14* and *Rht16* [17,18]. Applications of exogenous gibberellins (GA₃) restored plant height and other agronomic traits of *Rht18* dwarf lines to the wild-type levels, indicating that *Rht18* dwarf mutants are impaired in GA biosynthesis [19]. In this investigation, we have mapped *Rht18* to chromosome 6A using a cross between a pre-green revolution tall line (Indian) devoid of any known height reducing genes and *Rht18* mutant Icaro. The transfer of the *Rht18* allele into bread wheat and the selection of potential semi-dwarf lines with good agronomic characteristics can be useful for wheat breeding programs.

2. Results and Discussion

2.1. Plant Height of F₂ and F_{2,3} Progenies

The plant height of the F₂ population was recorded under controlled environmental conditions in a greenhouse along with parental lines Indian and Icaro. The height of the tall parent Indian and dwarf mutant parent Icaro averaged 86 ± 2.82 cm (Mean \pm S.E.) and 44 ± 1.02 cm (Mean \pm S.E.), respectively (Figure 1). Of the 94 F₂ plants, approximately 55 were within 10 cm of the Icaro height range. Only four of the plants in the F₂ population had a phenotype similar to Indian (86 ± 10 cm). Three of the originally-sown plants did not grow to maturity. This is expected as sterility is often associated with the incompatibility among the tetraploid and hexaploid crosses [20,21]. The F₂ population had a height distribution skewed towards the parent Icaro (Figure 1). The skewed distribution towards reduced height parents was also reported in the *Rht3* F₂ mapping population [22]. This distribution suggests that the mutant phenotype is dominant, as only a few plants had the tall phenotype. Interestingly, a few F₂ plants were taller than the tall parent and many were shorter than the dwarf parent. The height distribution pattern suggests that additional modifier genes might be involved in regulating plant height. Plant height is known to be a complex trait regulated by interaction and interplay among major and minor genes [23]. The transgressive segregation observed for plant height might be due to epistatic gene actions [23]. Transgressive segregation was reported earlier in wheat for several

agronomic traits, including plant height [24], grain yield and its components [25], heading date, and vernalization requirement [26]. In a previous study involving *Rht8*, transgressive segregants were observed for longer peduncles and grains per spike with no significant change in spike length, spikelet number, or number of fertile tillers [27]. Additionally, no significant effect was observed on roots, while a slight decrease in coleoptile length occurred. Partitioning of dry matter to ears was increased at anthesis, however, dry weight of stems and above-ground biomass, including ears, decreased [27].




Figure 1. Height distribution in the F₂ population. Plant height was grouped into 10 cm series. The star represents plant height of tall or dwarf mutant parent. The average plant height of Indian is recorded 86 cm (from eight plants) and Icaro as 44 cm (from six plants).

Forty seeds from each individual F₂ plant representing the F_{2.3} progenies were sown in the field the following summer to evaluate the genotypes of the F₂ plants, as it was difficult to classify plants into distinct categories in F₂. The F_{2.3} population showed segregation for plant height (Figure 2), with 14 progenies classified as homozygous short, one as homozygous tall, and 54 were classified as heterozygous. Highly significant effects were found for the plant height (Table 1). For the F_{2.3} population the height was found to be on average taller than the F₂, possibly due to the photoperiod effect in the field. As seen in the F₂ generation, we observed some very dwarf and some very tall plants in F_{2.3} progenies (Figure 2), indicating the role of additional modifier genes in transgressive segregation.

Table 1. Analysis of variance (ANOVA) of plant height for the F_{2.3} population.

Source	DF	SS	MS	F Value	Pr > F
Model	76	155,138.4	2041.29	14.49	<0.0001
Error	648	9131.879	140.91		
Corrected total	724	246,452.3			




Figure 2. Plant height distribution among $F_{2.3}$ families. (A) Indian and Icaro; and (B–H) different $F_{2.3}$ families.

2.2. Spike Morphology

Along with the plant height, the F_2 population also segregated for awn-less/short or long, black awns. Among the parents, Indian spikes were awn-less and Icaro spikes bear long black awns (Figure 3). Among the F_2 plants, 55 plants had awns and 36 plants were awn-less. We have also observed a difference in spike morphology among the F_2 and $F_{2.3}$ plants (Figure 3). The Indian spike is long and had loose spikelets, while the Icaro head is small with compact spikelets (Figure 3). We have observed plants with Indian-type heads with awns and Icaro-type heads without awns (Figure 3).



Figure 3. Spike morphology of parents and $F_{2.3}$ families in the mapping population. (A) Indian; (B) Icaro; and (C–H) different $F_{2.3}$ progenies.

2.3. Tiller Number

The F_2 population displayed a range for the number of tillers per plant ranging from three tillers per plant to 28 tillers per plant (Figure 4). Fifty-five percent of plants have tiller numbers ranging from 9 to 15 per plant, resembling the average for both Indian and Icaro, which were approximately 11 and 12 tillers per plant, respectively. The highest tillering plants were usually dwarf and sterile or contained only a few seeds in a spike. This might be due to incompatibility between the two genotypes. The higher or lower number of tillers compared to the average of both the parents might be due to multigenic nature of the trait. Extreme dwarf plants were sterile and did not set seeds. Further, of the 91 F_2 lines used for $F_{2,3}$ field evaluation, only 75 plants produced seed. This is expected for a hexaploid and tetraploid cross due to pollen viability issues restricting the seed set [28]. Among the plants that set seeds, some had good seed sets while others only contributed a few per plant.




Figure 4. Tiller number distribution in the F_2 population. The average number of tillers for parents Indian and Icaro were 11 and 12, respectively.

The variation observed in spike morphology was not associated with the height phenotype each plant displayed (data not shown). The seed weight did not correlate with plant height. Tiller number in F_2 plants did not associate with the seed weight or number of seeds harvested at maturity (data not shown). The 100 seed weight for Indian and Icaro were 3.43 g and 3.69 g, respectively. Among the $F_{2,3}$ families, the short families had an average 100 seed weight of 2.8 g while the tall families had 3.1 g. The height mutation in wheat is reported to have affected the seed weight compared to the tall counterpart. Of the studied reduced height mutants, *Rht12* reduces the grain weight more compared to *Rht-B1b*, *Rht-B1c*, and *Rht8* [29]. The reduction in grain weight might be due to the adverse effect of *Rht18* on grain size [27,30]. In fields conditions, the tiller number per plant was difficult to measure, hence, was not recorded for the $F_{2,3}$ plants. The $F_{2,3}$ families were also evaluated in the field for their agronomic characteristics to identify the agronomically important plant to be utilized in hexaploid wheat breeding. We have selected one line (line 29) based on plant height, stem strength, and spike morphology. More detailed agronomic and molecular analysis will be performed on the selected line to determine its suitability for utilization in a breeding program.

2.4. Genetic Mapping of *Rht18*

In order to map the gene on a wheat chromosome, over 700 SSR markers [31] were used to screen parents Indian and Icaro. Of these, 154 markers showed polymorphism between the parents and were used to genotype the population. The *Rht18* gene was mapped to the short arm of chromosome 6A and was flanked by *barc146* and *cfid190* (Figure 5). The SSR marker *cfid190* was placed at a

distance of 17.3 cM away from *Rht18*. Previously, *barc003*, a marker from the short arm of the chromosome 6A, was mapped 25.1 cM away from *Rht18* in durum wheat [17,32]. Earlier, *Rht18* was mapped on chromosome 6A between *barc118* and *IWA4371* using recombinant inbred lines (RILs) in durum wheat [18]. The mapping location of *Rht18* in our study is consistent with the previous map position [17,18]. Recently, several independent single nucleotide variants in the *GA2oxA9* gene located on chromosome 6A were associated with the *Rht18* mutant phenotype [33]. *GS2oxA9* is predicted to encode GA 2-oxidase, which reduces the amount of bioactive GA (GA1).




Figure 5. Genetic linkage map showing the position of *Rht18* on chromosome 6A. Genetic distances are shown in centiMorgans (cM).

Reduced-height genes in wheat have been imperative to the agronomic success of the crop. The resulting yield increases have been credited to the improved structure of the plant that responded better to the agronomic practices in use today. The semi-dwarf phenotype increases resistance to lodging along with increasing the number of grains per plants. Incorporating additional reduced-height genes into breeding programs could help contribute to the diversification of the genotype. Considering climate change and the demand for food security, incorporating additional dwarfing genes into the germplasm and evaluating their agronomic worth might help to address the wheat productivity under a changing climate. As the photoperiod and the background of a genotype affect height, a marker close to the gene may assist in easy and precise selection of the locus. Thus, identification of SSR markers closer to the *Rht18* locus may assist breeders in early identification of dwarfing lines for breeding populations. Further, conducting the genomic and agronomic characterization of this mutant gene may become instrumental in developing a better dwarfing system in wheat. Additionally, we have identified a semi-dwarf line from $F_{2:3}$ families with superior agronomic characters that might have potential to be used in wheat breeding to incorporate the gene into the hexaploid background of Pacific Northwest region.

3. Materials and Methods

3.1. Plant Materials

The dwarf parent, *Icaro* (tetraploid; $4\times$) (*Rht18*; PI 503555), was originally derived in 1987 in Italy from fast-neutron treatment of cv. *Anhinga* (PI 428455). The tall line *Indian* (hexaploid; $6\times$) (CI 4489), was developed at the University of Idaho, Idaho before 1915. As the tall parent is released before the introduction of reduced height genes, we presumed that cv. *Indian* would be devoid of the *Rht18* allele in the background. Both the germplasms were procured from GRIN [34].

3.2. *F₁* and Plant Growing Conditions

The *F₁* produced by crossing Indian as the female parent and Icaro as the male parent was self-pollinated, and 120 *F₂* seeds were collected. The *F₂* mapping population was grown at the plant growth facility, Washington State University, under controlled conditions of 16 h days (22 °C) and 8 h (18 °C) nights. For ease of genotyping, 94 randomly-selected *F₂* plants were selected for further analysis. Forty seeds of each *F₂* plant were grown in three-foot rows at the Spillman Agronomy Farm, Pullman, WA for phenotypic screening. Four rows were planted in each plot with a row-to-row and plot-to-plot spacing of one foot. Each row represented progeny of a single *F₂* plant. The seeds were planted mechanically using four planter drills and the plants were grown until maturity using the standard regional agricultural practices with no irrigation.

3.3. Phenotypic Screening

The phenotypic data for height, awns, tiller number, and seed weight was collected for the *F₂* and *F_{2:3}* populations. The plant height was recorded at maturity to the nearest cm excluding the awns. The population was characterized into tall, intermediate, and dwarf based on the plant height at maturity. Tiller numbers were counted manually per plant and seed weight was measured for each individual plant.

3.4. DNA Isolation and Genotyping

Young leaf tissue of *F₂* plants was collected in 96-well DNA extraction plates. Four, 2-cm long leaf segments were clipped and lyophilized for three days. The lyophilized tissue was used for DNA isolation using a modified SDS extraction method [35]. The DNA was diluted to a final concentration of 25 ng/μL. Primer sequence information for simple sequence repeat (SSR) markers were obtained from GrainGenes website [36].

Over 700 SSR markers were first screened for polymorphism between the parental genotypes. The PCR was performed in 12 μL reaction volume containing 1× NEB reaction buffer, 200 μM of dNTPs, 2.5 mM MgCl₂, 0.05 μM forward primer, 0.25 μM reverse primer, 0.2 μM M13 forward-labeled primer, and 1U homemade Taq polymerase. For multiplexing, the M13 sequence was fluorescently labeled separately with FAM, HEX, NED, and PET dyes. The amplification of SSR loci was performed using the protocol consisted of 94 °C/4 min for initial denaturation, followed by 37 cycles (94 °C/30 s, 60 °C/45 s, 72 °C/60 s), with final extension at 72 °C/10 min. The amplification products were separated using ABI DNA analyzer 3100 (Applied Biosystems Inc., Carlsbad, CA, USA). Alleles were sized relative to internal size standard (cassual445 labeled with Dy630) using GeneMarker software (Softgenetics, State College, PA, USA). MapMaker 2.0 was used to construct the genetic linkage map using the Kosambi mapping function [37,38].

Author Contributions: N.P.G., A.M. and D.S. carried out the research study, analyzed and interpreted the data, and drafted the manuscript. N.P.G. and A.M. collected the phenotypic data. K.S.G., D.S. and A.M. conceived and designed the study. Authors read and approved the final manuscript.

Funding: Financial assistance from NSF BREAD-0965533 to support Nathan Grant's undergraduate research and for conducting the experiment is acknowledged.

Acknowledgments: We thank Deven R. See and the Western Regional Small Grains Genotyping Laboratory (WRSGGL), Pullman, WA, for providing the genotyping facility.

Conflicts of Interest: The authors declare no conflict of interest. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

References

1. Germplasm Resources Information Network. Available online: <https://www.ars-grin.gov/> (accessed on 12 July 2018).
2. Hedden, P. The genes of the Green Revolution. *Trends Genet.* **2003**, *19*, 5–9. [[CrossRef](#)]
3. Borojevic, K.; Borojevic, K. The transfer and history of “reduced height genes” (Rht) in wheat from Japan to Europe. *J. Hered.* **2005**, *96*, 455–459. [[CrossRef](#)] [[PubMed](#)]
4. Ellis, M.H.; Rebetzke, G.J.; Chandler, P.; Bonnett, D.; Spielmeyer, W.; Richards, R.A. The effect of different height reducing genes on the early growth of wheat. *Funct. Plant Biol.* **2004**, *31*, 583–589. [[CrossRef](#)]
5. Chapman, S.C.; Mathews, K.L.; Trethowan, R.M.; Singh, R.P. Relationships between height and yield in near-isogenic spring wheats that contrast for major reduced height genes. *Euphytica* **2007**, *157*, 391–397. [[CrossRef](#)]
6. Peng, J.; Richards, D.E.; Hartley, N.M.; Murphy, G.P.; Devos, K.M.; Flintham, J.E.; Beales, J.; Fish, L.J.; Worland, A.J.; Pelica, F.; et al. ‘Green revolution’ genes encode mutant gibberellin response modulators. *Nature* **1999**, *400*, 256–261. [[CrossRef](#)] [[PubMed](#)]
7. Ellis, M.H.; Rebetzke, G.J.; Azanza, F.; Richards, R.A.; Spielmeyer, W. Molecular mapping of gibberellin-responsive dwarfing genes in bread wheat. *Theor. Appl. Genet.* **2005**, *111*, 423–430. [[CrossRef](#)] [[PubMed](#)]
8. Pearce, S.; Saville, R.; Vaughan, S.P.; Chandler, P.M.; Wilhelm, E.P.; Sparks, C.A.; Al-Kaff, N.; Korolev, A.; Boulton, M.I.; Phillips, A.L.; et al. Molecular characterization of *Rht-1* dwarfing genes in hexaploid wheat. *Plant Physiol.* **2011**, *157*, 1820–1831. [[CrossRef](#)] [[PubMed](#)]
9. Ogawa, M.; Kusano, T.; Katsumi, M.; Sano, H. Rice gibberellin-insensitive gene homolog, *OsGAI*, encodes a nuclear-localized protein capable of gene activation at transcriptional level. *Gene* **2000**, *245*, 21–29. [[CrossRef](#)]
10. Ikeda, A.; Ueguchi-Tanaka, M.; Sonoda, Y.; Kitano, H.; Koshioka, M.; Futsuhara, Y.; Matsuoka, M.; Yamaguchi, J. *Slender* rice, a constitutive gibberellin response mutant, is caused by a null mutation of the *SLR1* gene, an ortholog of the height-regulating gene *GAI/RGA/RHT/D8*. *Plant Cell* **2001**, *13*, 999–1010. [[CrossRef](#)] [[PubMed](#)]
11. Chandler, P.M.; Marion-Poll, A.; Ellis, M.; Gubler, F. Mutants at the *Slender1* locus of barley cv Himalaya. Molecular and physiological characterization. *Plant Physiol.* **2002**, *129*, 181–190. [[CrossRef](#)] [[PubMed](#)]
12. Tolmay, V.L. Resistance to biotic and abiotic stress in the Triticeae. *Hereditas* **2001**, *135*, 239–242. [[CrossRef](#)] [[PubMed](#)]
13. Schillinger, W.F.; Donaldson, E.; Allan, R.E.; Jones, S.S. Winter wheat seedling emergence from deep sowing depths. *Agron. J.* **1998**, *90*, 582–586. [[CrossRef](#)]
14. Chen, L.; Phillips, A.L.; Condon, A.G.; Parry, M.A.; Hu, Y.G. GA-responsive dwarfing gene *Rht12* affects the developmental and agronomic traits in common bread wheat. *PLoS ONE* **2013**, *8*, e62285. [[CrossRef](#)] [[PubMed](#)]
15. Wang, Y.; Chen, L.; Du, Y.; Yang, Z.; Condon, A.G.; Hu, Y.-G. Genetic effect of dwarfing gene *Rht13* compared with *Rht-D1b* on plant height and some agronomic traits in common wheat (*Triticum aestivum* L.). *Field Crops Res.* **2014**, *162*, 39–47. [[CrossRef](#)]
16. Gasperini, D.; Greenland, A.; Hedden, P.; Dreos, R.; Harwood, W.; Griffiths, S. Genetic and physiological analysis of *Rht8* in bread wheat: An alternative source of semi-dwarfism with a reduced sensitivity to brassinosteroids. *J. Exp. Bot.* **2012**, *63*, 4419–4436. [[CrossRef](#)] [[PubMed](#)]
17. Watanabe, N. Genetic collection and development of near-isogenic lines in durum wheat. *Vavilov J. Genet. Breed.* **2008**, *12*, 636–643.
18. Vikhe, P.; Patil, R.; Chavan, A.; Oak, M.; Tamhankar, S. Mapping gibberellin-sensitive dwarfing locus *Rht18* in durum wheat and development of SSR and SNP markers for selection in breeding. *Mol. Breed.* **2017**, *37*, 28. [[CrossRef](#)]
19. Yang, Z.Y.; Liu, C.Y.; Du, Y.Y.; Chen, L.; Chen, Y.F.; Hu, Y.G. Dwarfing gene *Rht18* from tetraploid wheat responds to exogenous GA3 in hexaploid wheat. *Cereal Res. Commun.* **2017**, *45*, 23–34. [[CrossRef](#)]
20. Padmanaban, S.; Zhang, P.; Hare, R.A.; Sutherland, M.W.; Martin, A. Pentaploid wheat hybrids: Applications, characterisation, and challenges. *Front. Plant Sci.* **2017**, *8*, 358. [[CrossRef](#)] [[PubMed](#)]
21. Martin, A.; Simpfendorfer, S.; Hare, R.A.; Eberhard, F.S.; Sutherland, M.W. Retention of D genome chromosomes in pentaploid wheat crosses. *Heredity* **2011**, *107*, 315–319. [[CrossRef](#)] [[PubMed](#)]

22. Navarro, C.; Yang, Y.; Mohan, A.; Grant, N.; Gill, K.S.; Sandhu, D. Microsatellites based genetic linkage map of the *Rht3* locus in bread wheat. *Mol. Plant Breed.* **2014**, *5*, 43–46. [[CrossRef](#)]
23. Rieseberg, L.H.; Archer, M.A.; Wayne, R.K. Transgressive segregation, adaptation and speciation. *Heredity* **1999**, *83*, 363–372. [[CrossRef](#)] [[PubMed](#)]
24. Bhatt, G.M. Inheritance of heading date, plant height, and kernel weight in two spring wheat Crosses. *Crop Sci.* **1972**, *12*, 95–98. [[CrossRef](#)]
25. El-Hennawy, M.A.; Abdalla, A.F.; Shafey, S.A.; Al-Ashkar, I.M. Production of doubled haploid wheat lines (*Triticum aestivum* L.) using anther culture technique. *Ann. Agric. Sci.* **2011**, *56*, 63–72. [[CrossRef](#)]
26. Shindo, C.; Tsujimoto, H.; Sasakuma, T. Segregation analysis of heading traits in hexaploid wheat utilizing recombinant inbred lines. *Heredity* **2003**, *90*, 56–63. [[CrossRef](#)] [[PubMed](#)]
27. Yang, Z.; Zheng, J.; Liu, C.; Wang, Y.; Condon, A.G.; Chen, Y.; Hu, Y.-G. Effects of the GA-responsive dwarfing gene *Rht18* from tetraploid wheat on agronomic traits of common wheat. *Field Crops Res.* **2015**, *183*, 92–101. [[CrossRef](#)]
28. Bhagyalakshmi, K.; Vinod, K.K.; Kumar, M.; Arumugachamy, S.; Prabhakaran, A.; Raveendran, T.S. Interspecific hybrids from wild x cultivated *Triticum* crosses—a study on the cytological behavior and molecular relations. *J. Crop Sci. Biotechnol.* **2008**, *11*, 257–262.
29. Casebow, R.; Hadley, C.; Uppal, R.; Addisu, M.; Loddo, S.; Kowalski, A.; Griffiths, S.; Gooding, M. Reduced height (*Rht*) alleles affect wheat grain quality. *PLoS ONE* **2016**, *11*, e0156056. [[CrossRef](#)] [[PubMed](#)]
30. Tang, T. *Physiological and Genetic Studies of an Alternative Semi-Dwarfing Gene Rht18 in Wheat*; University of Tasmania: Tasmania, Australia, 2015.
31. Somers, D.J.; Isaac, P.; Edwards, K. A high-density microsatellite consensus map for bread wheat (*Triticum aestivum* L.). *Theor. Appl. Genet.* **2004**, *109*, 1105–1114. [[CrossRef](#)] [[PubMed](#)]
32. Haque, M.A.; Martinek, P.; Watanabe, N.; Kuboyama, T. Genetic mapping of gibberellic acid-sensitive genes for semi-dwarfism in durum wheat. *Cereal Res. Commun.* **2011**, *39*, 171–178. [[CrossRef](#)]
33. Ford, B.; Foo, E.; Sharwood, R.E.; Karafiatova, M.; Vrána, J.; MacMillan, C.; Nichols, D.S.; Steuernagel, B.; Uauy, C.; Doležel, J.; et al. *Rht18* semi-dwarfism in wheat is due to increased expression of GA 2-oxidaseA9 and lower GA Content. *Plant Physiol.* **2018**. [[CrossRef](#)] [[PubMed](#)]
34. Gerritsen, V.B. All things dwarfed and beautiful. Available online: <https://web.expasy.org/spotlight/pdf/sptlt070.pdf> (accessed on 12 July 2018).
35. Sandhu, D.; Champoux, J.A.; Bondareva, S.N.; Gill, K.S. Identification and physical localization of useful genes and markers to a major gene-rich region on wheat group 1S chromosomes. *Genetics* **2001**, *157*, 1735–1747. [[PubMed](#)]
36. GrainGenes. Available online: <https://wheat.pw.usda.gov/cgi-bin/GG3/browse.cgi?class=marker;begin=8251> (accessed on 12 July 2018).
37. Lander, E.S.; Green, P.; Abrahamson, J.; Barlow, A.; Daly, M.J.; Lincoln, S.E.; Newburg, L. MAPMAKER: An interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. *Genomics* **1987**, *1*, 174–181. [[CrossRef](#)]
38. Kosambi, D.D. The estimation of map distance from recombination values. *Ann. Eugen.* **1944**, *12*, 172–175. [[CrossRef](#)]

