

Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at ScienceDirect

Journal of Invertebrate Pathology

journal homepage: www.elsevier.com/locate/yjipa

Infection characteristics of *Solenopsis invicta* virus 2 in the red imported fire ant, *Solenopsis invicta*

Yoshifumi Hashimoto¹, Steven M. Valles^{*}

Center for Medical, Agricultural and Veterinary Entomology, USDA-ARS, 1600 SW 23rd Drive, Gainesville, FL 32608, USA

ARTICLE INFO

Article history:

Received 25 January 2008

Accepted 11 June 2008

Available online 19 June 2008

Keywords:

Solenopsis invicta virus 2

Positive-strand RNA virus

Prevalence

Tissue tropism

Transmission

qPCR

ABSTRACT

Solenopsis invicta virus 2 (SINV-2) is the second virus identified from the fire ant, *S. invicta* Buren. SINV-2 is unique among positive-strand RNA viruses from insects by possessing four cistrons in a monopartite genome. Fire ant colonies testing positive for SINV-2 by RT-PCR did not exhibit any discernable symptoms. RT-PCR-based surveys for SINV-2 among 688 fire ant mounds in Alachua County, Florida, sampled during the period January, 2006 through December, 2007 showed that the prevalence of SINV-2 among nests ranged from 1.6% to 16.4%. Unlike *S. invicta* virus 1, no seasonal-associated prevalence was observed with regard to SINV-2 infection among fire ant colonies. No social form specificity was evident; SINV-2 was found in both monogyne and polygyne *S. invicta* ants. Real-time quantitative PCR experiments showed that SINV-2 genome equivalents per individual ant ranged from 1.9×10^7 in pupae to 4.3×10^{11} in inseminated queens. The SINV-2 infection was detected in all ant stages examined (eggs, larvae, pupae, workers, and queens). Tissue tropism studies indicated that the alimentary canal (specifically the midgut) is most likely the susceptible tissue. SINV-2 was successfully transmitted to uninfected *S. invicta* ants by feeding a partially purified homogenate of SINV-2-infected ants. The SINV-2 transmission rate ranged from 30% to 80%, and both positive (genomic) and negative (replicative) SINV-2 RNA strands accumulated in recipient ants over the course of the experiment. These results indicated that SINV-2 replicates within *S. invicta*.

© 2008 Elsevier Inc. All rights reserved.

1. Introduction

Red imported fire ant, *Solenopsis invicta* Buren, was introduced into the United States from South America in the early 1900s (Creighton, 1950). Damage attributed to this aggressive, territorial ant species is wide-ranging and includes physical damage to agricultural commodities and equipment, roads, electrical equipment, and livestock, decreased biological diversity, and human health concerns associated with stings (Williams et al., 2001). Nearly all of the natural enemies of *S. invicta* identified in South America are absent from the US population which is likely the reason *S. invicta* has become such a major pest in the US (Porter et al., 1992, 1997). Thus, identification and utilization of natural enemies specific to *S. invicta* are considered to be crucial in achieving sustainable control (Williams et al., 2003).

We reported recently the discovery of a new virus (*S. invicta* virus 2 [SINV-2]) from *S. invicta* (Hashimoto and Valles, 2008; Valles et al., 2007a). SINV-2 is only the second virus known to infect this ant species. Molecular characterization of SINV-2 revealed a single-stranded positive sense RNA genome comprised of 10,303

nucleotides and a polyadenylated tail at the 3' terminus. The SINV-2 genome is unique among other insect-infecting RNA viruses in that it possesses four open reading frames (ORFs) in a monopartite genome. Protein domains consistent with positive-strand RNA viruses (i.e., an RNA-dependent RNA polymerase [RdRp], helicase, and protease) were present in the ORF 4 of the SINV-2 genome. Phylogenetic analysis of SINV-2 RdRp and helicase amino acid sequences indicated unique placement of SINV-2 exclusive from the *Dicistroviridae* (including the first identified fire ant virus, SINV-1), iflaviruses, and plant small RNA viruses. Although an initial molecular characterization of SINV-2 has been completed (Valles et al., 2007a), information concerning the infection characteristics of this virus in the ant host is lacking. Therefore, descriptive studies focused on the infection characteristics (tissue tropism, stage-specificity, transmission, replication, and prevalence) of SINV-2 in the fire ant were conducted.

2. Materials and methods

2.1. Field surveys and social form specificity

A field survey was conducted to examine the extent of SINV-2 infection among *S. invicta* mounds in Alachua County, Florida. Samples of workers were retrieved from the field and RNA

^{*} Corresponding author. Fax: +1 352 374 5818.

E-mail address: steven.valles@ars.usda.gov (S.M. Valles).

¹ Present address: Boyce Thompson Institute for Plant Research at Cornell University, Tower Road, Ithaca, NY 14850, USA.

was extracted from the ants with Trizol reagent (Invitrogen, Carlsbad, CA). SINV-2-specific oligonucleotide primers (p64, 5'-ATTTGTTTGGCCACGGTCAAC, genome position 10,758→10,780, and p65, 5'-GATGATACAAAGCATTAGCGTAGGTAACG, genome position 11,047→11,076) were used in an RT-PCR reaction to determine the presence of infection. In an effort to determine possible ant social form specificity of SINV-2, each mound that tested positive for SINV-2 was also genotyped at the *Gp-9* locus to determine the social form of the nest (Valles and Porter, 2003).

2.2. Stage specificity and tissue tropism

Experiments were conducted to examine the developmental stages of *S. invicta* for SINV-2 infection. Samples of queens ($n = 4$ individuals), workers ($n = 5$ groups of 10 individuals), larvae ($n = 5$ groups of 10 individuals), sexual larvae ($n = 12$ individuals), eggs ($n = 2$ masses of an unknown number of eggs), and pupae ($n = 5$ groups of 10 individuals) were sampled directly from the field from SINV-2-positive colonies. RNA was extracted from the ants with Trizol reagent. The RNA concentration was determined spectrophotometrically.

Tissue tropism of SINV-2 was examined by dissecting different tissues and quantifying the number of SINV-2 genome equivalents in each respective preparation. Ice-chilled fire ant workers from a SINV-2-positive colony were placed on a glass microscope slide, and head, thorax, and abdomen were separated with a surgical blade under a Leica MS5 dissecting microscope. The head and thorax were placed directly in Trizol and the abdomen was immersed in 10 mM Tris-HCl, pH 8.0, for further dissection. Tissues were isolated and removed from the abdomen with micro-dissecting forceps in the following order: crop, poison sac, alimentary canal (comprised of the midgut, hindgut, and Malpighian tubules), and the remaining abdominal carcass (comprised of the Dufour's gland, ovary, fat body, muscle, and cuticle). The alimentary canal was later examined further by separating the midgut, hindgut, and Malpighian tubules. Larvae from SINV-2 positive colonies were dissected to the following groups: Malpighian tubules, alimentary canal, and remaining carcass. Five replicates were conducted. RNA was extracted with Trizol.

The number of SINV-2 genome equivalents in different developmental stages and tissues were quantified by QPCR. cDNA was synthesized from the SINV-2 genome region corresponding to an area between the helicase and RdRp with total RNA isolated from dissected tissues and stages using SuperScript III Reverse Transcriptase (SsRT; Invitrogen) and a gene-specific primer (p514, 5'-TACACTTGGGTCTCAGGAACC, genome position 8816→8836). In a 0.5 ml PCR tube, 2 μ l of primer p514 (1 μ M), 1 μ l of a dNTP mix (10 mM), and 10 μ l of total RNA (50 ng) were mixed and heated to 65 °C for 5 min in a PTC 100 thermal cycler, followed by incubation on ice for 1 min. Then, 4 μ l of first-strand buffer (250 mM Tris-HCl, pH 8.3; 375 mM KCl, 15 mM MgCl₂), 2.75 μ l of DEPC-treated water, and 0.25 μ l of SsRT (200 U/ μ l) were added. The mixture was incubated at 55 °C for 30 min, followed by inactivation of SsRT by heating to 70 °C for 15 min.

QPCR was performed on an ABI PRISM 7000 Sequence Detection System interfaced to the ABI prism 7000 SDS software (Applied Biosystems, Foster City, CA) in a 25 μ l reaction volume. The reaction contained 12.5 μ l of SYBR Green SuperMix (with UDG and ROX, Invitrogen), 0.4 μ l each of 10 μ M SINV-2-specific primers (p511, 5'-CGGAGACTGAGCCTTTCTGGACTCCATAG, genome position 8677→8707 and p515, 5'-TGTATCGCGGAAATTACCAACATCAAC, genome position 8584→8613), 3 mM MgCl₂, 1 μ l of the cDNA synthesis reaction, and 10.7 μ l of DEPC-water. QPCR conditions consisted of one cycle at 50 °C for 2 min and 95 °C for 2 min, followed by 40 cycles at 95 °C for 15 s, 64 °C for 15 s, 72 °C for 1 min. The non-template control for QPCR included a

complete cDNA synthesis reaction devoid of RNA template. A standard curve was constructed from a plasmid clone of the corresponding SINV-2 genome region using a copy number range of 5–5 $\times 10^7$ copies. Reaction efficiencies were determined by regressing C_T values against the template copy number (log) and calculated according to the formula [$E = (10^{-1/\text{slope}}) - 1$] (Klein et al., 1999). Reaction efficiencies routinely exceeded 96%.

QPCR was also conducted to separately quantify the plus (genomic) and minus (replicative) RNA strands of SINV-2. cDNA was synthesized from the SINV-2 plus strand with oligonucleotide primer p511 and minus strand with oligonucleotide primer p515 as described above. After cDNA synthesis, the RNA templates were digested with RNase A and RNase H at 37 °C for 30 min. After RNA digestion, QPCR was conducted as described above with oligonucleotide primers p511 and p515.

2.3. SINV-2 transmission to uninfected fire ants

SINV-2-uninfected laboratory-reared monogyne colonies were identified by RT-PCR and divided into four equivalent fragment colonies comprised of 0.5 g of brood and 4 ml of workers. Colonies were infected by a modified method described by Ackey and Beck (1972). Workers and brood (0.15 g) from a SINV-2-infected colony were homogenized in an equal volume of 10% sucrose with a Potter-Elvehjem Teflon pestle and glass mortar. The homogenate was filtered through four layers of cheesecloth. Approximately 4 ml of the homogenate/sucrose solution was placed into a cotton-stopped test tube and presented to three of the four fragment colonies; one fragment colony was provided a homogenate of uninfected ants and served as control. After 2 days, the homogenate was removed and replaced with unadulterated 10% sucrose, water, frozen crickets (*Acheta domesticus*), and egg yolk (hard-boiled). Three experiments were conducted comprised of three colonies that had been fragmented into four sub-colonies (one control and three treated for a total of nine experimental units experiencing treatment). Fragment colonies were examined for the presence of SINV-2 by extracting total RNA from 20 worker ants and conducting RT-PCR at 0, 7, 14, 21, and 28 days after exposure to the homogenate. Percent infection, by day, among recipient fragment colonies was analyzed by analysis of variance (SAS Institute, 1988). Samples testing positive for SINV-2 by RT-PCR were analyzed by strand-specific QPCR to quantify the genome equivalents (plus and minus strand).

3. Results and discussion

3.1. Field prevalence and fire ant social form

Among 688 *S. invicta* nests sampled from January, 2006 through December, 2007, 38 (5.5%) were found to be infected with SINV-2 by RT-PCR (Table 1). Sampling during the first quarter of 2006 yielded the lowest SINV-2 prevalence (1.6%) and the fourth quarter of 2006 the highest (16.4%). The nest infection rate for SINV-2 was lower than either *S. invicta* virus 1 (SINV-1) genotype; SINV-1 and SINV-1A were found previously in 22.9% and 55% of *S. invicta* nests sampled, respectively (Valles et al., 2004; Valles and Strong, 2005). The prevalence of the SINV-2 infection was independent of collection time. In other words, the presence of the SINV-2 infection in *S. invicta* nests did not exhibit any apparent pattern with respect to season. SINV-1 and SINV-1A infections of fire ants were found in significantly greater frequency during the late spring, summer, and early fall in Alachua County, Florida (Valles et al., 2007b). Further examination revealed that the SINV-1/SINV-1A infection prevalence exhibited a significant correlation with temperature (Valles et al., 2007b). A number of other insect-infecting, positive-strand RNA viruses have been shown to exhibit either seasonal fluctuations (Bailey et al., 1981, 1983; Tentcheva et al., 2004) or greater

Table 1
Field prevalence of SINV-2 infection in Alachua County, Florida, and corresponding Gp-9 genotype of *S. invicta* workers from SINV-2-infected nests

Period of collection ^a	Nests sampled	Nests infected with SINV-2 (%)	Gp-9 genotype of SINV-2-infected nests	
			BB (monogyne)	Bb (polygyne)
Quarter 1, 2006	64	1 (1.6)		1
Quarter 2, 2006	35	1 (2.9)		1
Quarter 3, 2006	85	3 (3.5)	1	2
Quarter 4, 2006	55	9 (16.4)	5	4
Quarter 1, 2007	213	9 (4.2)	3	6
Quarter 2, 2007	108	12 (11.1)	1	11
Quarter 3, 2007	108	2 (1.9)		2
Quarter 4, 2007	20	1 (5)	1	
Grand total	688	38 (5.5)	11 (28.9)	27 (71.1)

^a Quarter 1 corresponds to January, February, and March; Quarter 2 to April, May, and June; Quarter 3 to July, August, and September; Quarter 4 to October, November, and December. Percentage nests (grand mean) infected with SINV-2 are expressed as a proportion of the total samples (688) and percentage Gp-9 genotypes are expressed as a proportion of the total nests infected (38).

abundance in warmer areas (Plus et al., 1975). These associations are hypothesized to be linked to host behavior patterns.

Solenopsis invicta social form, as determined by Gp-9 allele composition (Valles and Porter, 2003), had no bearing on the presence of SINV-2 infection; monogyne (28.9%) and polygyne (71.1%) ants were found to harbor the virus (Table 1). Alachua County, Florida, is populated by an overwhelming majority of the polygyne social form (Porter, 1992), so the higher percentage of SINV-2 observed in polygyne ants is likely from a sampling bias. Regardless, the data clearly show that both social forms are capable of being infected with the virus.

3.2. Stage specificity

QPCR revealed that the SINV-2 genome was present in all developmental stages of infected *S. invicta* colonies (Fig. 1). Whether dissemination was transmitted horizontally or (and) vertically was not determined. However, vertical transmission was implicated by detection of SINV-2 in eggs. Whether SINV-2 was inside the eggs (transovarial) or on the surface of the eggs (transovum) was not determined. Among the four queens from SINV-2 positive col-




Fig. 1. SINV-2 genome equivalents determined for individual ants in different developmental stages. Genome equivalents were interpolated from a standard curve generated simultaneously from a plasmid construct. Mean and standard deviation values are indicated by the bar and error bars, respectively. Eggs were counted and prepared as a pooled group.

onies, all possessed significant numbers of SINV-2 genome equivalents (1.7×10^8 to 1.7×10^{12}). Vertical transmission has been shown for a number of positive-strand RNA viruses that infect the honey bee, *Apis mellifera* (Chen et al., 2006; Chen and Siede, 2007). However, the *Plautia stali* intestine virus was not transovarially transmitted in its host, rather, the virus was passed to progeny by surface contamination of eggs and from feces of infected aphids (Nakashima et al., 1998).

Based on SINV-2 genome equivalents, the infection was quite large ($>10^8$ copies per ant) in all fire ant stages, except eggs, ranging from 1.9×10^8 copies (pupae) to 4.3×10^{11} copies (queens) (Fig. 1). These genome copy values were comparable to those reported for SINV-1-infection of *S. invicta* (Hashimoto et al., 2007). None of these ants (or colonies) exhibited any obvious signs of distress. Thus, SINV-2 appears to cause an asymptomatic, chronic infection in *S. invicta*. Whether the virus is capable of causing disease or mortality among the ants under certain conditions is unknown. Again, using the honey bee as an example, among the 18 known positive-strand RNA viruses infecting honey bees, most are present in the population as an asymptomatic infection. It is postulated that under certain environmental conditions (e.g. stress), the virus(es) begin to replicate actively resulting in the exhibition of disease symptoms and sometimes death (Chen and Siede, 2007).

3.3. Tissue tropism

QPCR examination of tissues for the SINV-2 genome showed that the virus was most prevalent in the alimentary canal of larvae and workers (Table 2). In workers, 99% of the SINV-2 genome equivalents were found in the alimentary canal. When the alimentary canal was further dissected to midgut, hindgut, and Malpighian tubules, the majority of the virus (based on detection of genome equivalents) was found in the midgut. Larvae exhibited a similar tissue tropism; 99.5% of the SINV-2 genome equivalents were found in the alimentary canal. The gut contents of the infected ants also possessed large quantities of SINV-2 genome equivalents (data not shown). Therefore, it is highly possible that detection of SINV-2 in tissues other than midgut is from virus release and subsequent contamination during the dissection process. The alimentary tract is a common target of insect-infecting positive-strand RNA viruses (Gildow and D'Arcy, 1990; Nakashima et al., 1998; van den Heuvel et al., 1997). SINV-1 also has been shown to exhibit tropism toward the alimentary canal of fire ants (Hashimoto and Valles, 2007). It is likely that dissemination of SINV-2 is similar to the mode of dispersal postulated for SINV-1. That is, virus appears to replicate in

Table 2
Presence of SINV-2 genome equivalents (based on RdRp amplification) among different tissues and body parts of workers and larvae of *S. invicta*

Developmental stage	Tissue/body parts	Distribution of genome equivalents by QPCR (% of total ^a)
Adult	Head	0.1 ± 0.01
	Thorax	0.6 ± 1.4
	Crop	0.1 ± 0.2
	Poison sac	0.1 ± 0.2
	Abdominal carcass	0.1 ± 0.1
	Alimentary canal	99.0 ± 1.4
	Midgut	95.8 ± 8.4
Larvae	Hindgut	1.3 ± 2.8
	Malpighian tubules	2.9 ± 4.8
	Malpighian tubules	0.1 ± 0.1
	Carcass	0.4 ± 1.3
	Alimentary canal	0.4 ± 1.3

^a Mean (±SD) total genome equivalents; $9.9 \times 10^8 \pm 1.5 \times 10^9$ (workers) and $4.1 \times 10^7 \pm 7.8 \times 10^7$ (larvae).




Fig. 2. SINV-2 transmission to uninfected *S. invicta* fragment colonies. Worker ants were sampled from the week specified from each colony and tested for the presence of SINV-2 by RT-PCR and QPCR. Bars indicate the percentage of fragment colonies ($n = 9$) testing positive for SINV-2 by RT-PCR. Open circles and squares indicate negative and positive-strand copy numbers from samples testing positive for SINV-2 by RT-PCR among the SINV-2-treated fragment colonies. Filled circles and squares represent samples taken from fragment colonies fed with homogenate prepared from SINV-2 uninfected colonies (i.e., control).

the midgut epithelial cells and shed viral particles into the midgut lumen where they are ultimately dispersed by trophalaxis or contamination of surfaces by fecal deposits. Further, the gut lumen may function as a reservoir of the virus.

3.4. Transmission

Transmission of SINV-2 to uninfected *S. invicta* colonies was successfully accomplished by feeding. SINV-2 was detectable in treated colonies within 7 days of exposure to a partially purified homogenate of SINV-2-infected fire ants (Fig. 2). The infection was detected for at least 28 days after treatment, indicating sustained infection among recipient colonies. No significant differences were noted in the percent infection among recipient colonies for weeks 1 through 4 ($p = 0.28$). Plus and minus SINV-2 genome strands were also detected in the recipient colonies. The number of plus and minus strands increased throughout the duration of the test (Fig. 2). Neither strand was detected among the control colonies treated with homogenate prepared from SINV-2-negative ants. These results show that both plus and minus strand RNA copies were produced *de novo* in the uninfected colonies and that virus replication was occurring. Ratios of plus to minus strand copies were 10:1 to 100:1, consistent with values detected in SINV-2 infected larvae and workers collected directly from the field (Hashimoto and Valles, unpublished data) and other replicating positive-strand RNA viruses (Regoes et al., 2005; Castillo et al., 2007). These data also suggest that SINV-2 is transmitted to nestmates horizontally by food-borne dispersion, probably by trophalaxis. Further studies on the transmission route are underway to test our hypothetical mode of SINV-2 dissemination in fire ants.

The fire ant viruses SINV-1 and SINV-2 are recent discoveries that hold promise as alternatives to traditional insecticides for controlling red imported fire ants in the United States. These viruses seem to fit the paradigm of positive-strand RNA viruses in honey bees—they are present in the population (often together within the same colony) as persistent, asymptomatic infections. However, under certain conditions, the virus begins to replicate resulting in discernable symptoms and often death. Therefore, with further study, SINV-1 and -2 may be able to be artificially manipulated to induce harmful colony behaviors or death in *S. invicta*.

Acknowledgments

We thank Drs. J. Becnel and D. Oi (USDA-ARS) for critical reviews of the manuscript. The use of trade, firm, or corporation names in this publication is for the information and convenience of the reader. Such use does not constitute an official endorsement or approval by the United States Department of Agriculture or the Agricultural Research Service of any product or service to the exclusion of others that may be suitable.

References

- Ackey, D.H., Beck, S.D., 1972. Nutrition of the pea aphid, *Acyrtosiphon pisum*: requirement for trace metals, sulphur, and cholesterol. *J. Insect Physiol.* 18, 1901–1914.
- Bailey, L., Ball, B.V., Perry, J.N., 1981. The prevalence of viruses of honey bees in Britain. *Ann. Gen. Virol.* 25, 175–186.
- Bailey, L., Ball, B.V., Perry, J.N., 1983. Association of viruses with two protozoal pathogens of the honey bee. *Ann. Appl. Biol.* 103, 13–20.
- Castillo, I., Rodriguez-Inigo, E., Bartolome, J., de Lucas, S., Ortiz-Movilla, N., Lopez-Alcorocho, J.M., Pardo, M., Carreno, V., 2007. Hepatitis C virus replicates in peripheral blood mononuclear cells of patients with occult hepatitis C virus infection. *Gut* 54, 682–685.
- Chen, Y.P., Pettis, J.S., Collins, A., Feldlaufer, M.F., 2006. Prevalence and transmission of honeybee viruses. *Appl. Environ. Microbiol.* 72, 606–611.
- Chen, Y.P., Siede, R., 2007. Honey bee viruses. *Adv. Virus Res.* 70, 33–80.
- Creighton, W.S., 1950. The ants of North America. *Bull. Mus. Comp. Zool. Harvard Univ.* 104, 1–585.
- Gildow, F.E., D'Arcy, C.J., 1990. Cytopathology and experimental host range of *Rhopalosiphum padi* virus, a small isometric RNA virus infecting cereal grain aphids. *J. Invertebr. Pathol.* 55, 245–257.
- Hashimoto, Y., Valles, S.M., 2007. *Solenopsis invicta* virus-1 tissue tropism and intra-colony infection rate in the imported fire ant: a quantitative PCR-based study. *J. Invertebr. Pathol.* 96, 156–161.
- Hashimoto, Y., Valles, S.M., 2008. Detection and quantitation of genomic and intermediary replicating viral RNA in the fire ants infected with *Solenopsis invicta* virus-2 by real-time PCR. *J. Invertebr. Pathol.* 98, 243–245.
- Hashimoto, Y., Valles, S.M., Strong, C.A., 2007. Detection of *Solenopsis invicta* virus in fire ants by real-time PCR. *J. Virol. Methods* 140, 132–139.
- Klein, D., Janda, P., Steinborn, R., Muller, M., Salmons, B., Gunzburg, W.H., 1999. Proviral load determination of different feline immunodeficiency virus isolates using real-time polymerase chain reaction: influence of mismatches on quantification. *Electrophoresis* 20, 291–299.
- Nakashima, N., Sasaki, J., Tsuda, K., Yasunaga, C., Noda, H., 1998. Properties of a new picorna-like virus of the brown-winged green bug, *plautia stali*. *J. Invertebr. Pathol.* 71, 151–158.
- Plus, N., Croizier, G., Jousset, F.X., David, J., 1975. Picornaviruses of laboratory and wild *Drosophila melanogaster*: geographical distribution and serotypic composition. *Ann. Microbiol.* 126A, 107–121.
- Porter, S.D., 1992. Frequency and distribution of polygynous fire ants (Hymenoptera: Formicidae) in Florida. *Fla. Entomol.* 75, 248–258.
- Porter, S.D., Fowler, H.G., Mackay, W.P., 1992. Fire ant mound densities in the United States and Brazil (Hymenoptera: Formicidae). *J. Econ. Entomol.* 85, 1154–1161.

- Porter, S.D., Williams, D.F., Patterson, R.S., Fowler, H.G., 1997. Intercontinental differences in the abundance of *Solenopsis* fire ants (Hymenoptera: Formicidae): escape from natural enemies? *Environ. Entomol.* 26, 373–384.
- Regoes, R.R., Crotty, S., Antia, R., Tanaka, M.M., 2005. Optimal replication of poliovirus within cells. *Am. Nat.* 165, 364–373.
- SAS Institute, 1988. SAS procedures guide for personal computers. SAS Institute, Cary, NC.
- Tentcheva, D., Gauthier, L., Zappulla, N., Dainat, B., Cousserans, F., Colin, M.E., Bergoin, M., 2004. Prevalence and seasonal variations of six bee viruses in *Apis mellifera* L. and *Varroa destructor* mite populations in France. *Appl. Environ. Microbiol.* 70, 7185–7191.
- Valles, S.M., Porter, S.D., 2003. Identification of polygynous and monogynous fire ant colonies (*Solenopsis invicta*) by multiplex PCR of *Gp-9* alleles. *Insect. Soc.* 50, 199–200.
- Valles, S.M., Strong, C.A., Dang, P.M., Hunter, W.B., Pereira, R.M., Oi, D.H., Shapiro, A.M., Williams, D.F., 2004. A picorna-like virus from the red imported fire ant, *Solenopsis invicta*: initial discovery, genome sequence, and characterization. *Virology* 328, 151–157.
- Valles, S.M., Strong, C.A., 2005. *Solenopsis invicta* virus-1A (SINV-1A): distinct species or genotype of SINV-1? *J. Invertebr. Pathol.* 88, 232–237.
- Valles, S.M., Strong, C.A., Hashimoto, Y., 2007a. A new positive-strand RNA virus with unique genome characteristics from the red imported fire ant, *Solenopsis invicta*. *Virology* 365, 457–463.
- Valles, S.M., Strong, C.A., Oi, D.H., Porter, S.D., Pereira, R.M., Vander Meer, R.K., Hashimoto, Y., Hooper-Bui, L.M., Sánchez-Arroyo, H., Davis, T., Karpakakunjarum, V., Davis, T., Vail, K.M., Fudd Graham, L.C., Briano, J.A., Calcaterra, L.A., Gilbert, L.E., Ward, R., Ward, K., Oliver, J.B., Taniguchi, G., Thompson, D.C., 2007b. Phenology, distribution, and host specificity of *Solenopsis invicta* virus-1. *J. Invertebr. Pathol.* 96, 18–27.
- van den Heuvel, J.F., Hummelen, H., Verbeek, M., Dulleman, A.M., van der Wilk, F., 1997. Characteristics of *acyrthosiphon pisum* virus, a newly identified virus infecting the pea aphid. *J. Invertebr. Pathol.* 70, 169–176.
- Williams, D.F., Collins, H.L., Oi, D.H., 2001. The red imported fire ant (Hymenoptera: Formicidae): an historical perspective of treatment programs and the development of chemical baits for control. *Am. Entomol.* 47, 146–159.
- Williams, D.F., Oi, D.H., Porter, S.D., Pereira, R.M., Briano, J.A., 2003. Biological control of imported fire ants (Hymenoptera: Formicidae). *Am. Entomol.* 49, 150–163.