

USDA, FDA, and ODS-NIH Database for the Iodine Content of Common Foods Release 3

Prepared by:
**Janet M. Roseland¹, Judith H. Spungen², Kristine Y. Patterson¹,
Abby G. Ershow³, Jaime J. Gahche³, Katherine Heydorn¹, Karen Regan³,
Pamela R. Pehrsson¹**

¹ Methods and Application of Food Composition Laboratory
Agricultural Research Service
U.S. Department of Agriculture

² Center for Food Safety and Applied Nutrition
U.S. Food and Drug Administration (FDA)

³ Office of Dietary Supplements
National Institutes of Health

March 2023

U.S. Department of Agriculture
Agricultural Research Service
Beltsville Human Nutrition Research Center
Methods and Application of Food Composition Laboratory (MAFCL)
10300 Baltimore Avenue
Building 005, Room 107, BARC – West
Beltsville, Maryland 20705 Tel. 301-504-0630
MAFCL web site: <http://www.ars.usda.gov/nutrientdata>
FoodData Central database web site: <https://fdc.nal.usda.gov/>

Supported by: National Institutes of Health-Office of Dietary Supplements (NIH-ODS), U.S. Department of Agriculture (USDA), U.S. Food and Drug Administration-Center for Food Safety and Nutrition (FDA-CFSAN)

The authors gratefully acknowledge Terry Cuncell, Dana Hoffman-Pennesi, Todor Todorov, and Mark Wirtz, of FDA's Center for Food Safety and Applied Nutrition, and the staff of FDA's Kansas City Laboratory for administration of the FDA Total Diet Study (TDS) program, for TDS data management, and for careful review of this document. The authors also acknowledge Katherine Phillips, Virginia Tech, for preparation of the USDA samples.

Introduction

Iodine is an essential mineral for human health, functioning as a component of thyroid hormones with important roles in growth and maturation, neurologic development, reproduction, and energy metabolism (Lee et al., 2016a; Rohner et al., 2014). Severe iodine deficiency during pregnancy and early childhood has serious effects on the developing fetus and infant and can cause permanent damage (Swanson et al., 2012). Goiter, enlargement of the thyroid gland, and other disturbances of thyroid function can be seen at all ages in individuals whose diets lack iodine at needed levels.

Adequate dietary iodine intake is crucial, with 150 µg/day as the daily recommended dietary allowance (RDA) for ages 14 years and older, 220 µg/day for pregnant women, and 290 µg/day for lactating women (Institute of Medicine, 2001; Rohner et al., 2014). On the other end of the spectrum, the tolerable upper intake level for iodine is 1100 µg/day for adults (Institute of Medicine, 2001). Excessive intake also can result in adverse health effects such as thyroid dysfunction (e.g., goiter, hyperthyroidism), autoimmune thyroid disease, and cancer (Luo et al., 2014; Ershow et al., 2016).

Iodine intake in the U.S. population overall is generally considered sufficient, but subgroups identified as at-risk for iodine insufficiency include women of reproductive age, young adults, and non-Hispanic blacks, with a trend of increasing deficiency over a 12-year period of the National Health and Nutrition Examination Survey (NHANES) from 2001-2012 (Lee et al., 2016a). Iodine insufficiency in the U.S. is a public health concern especially in pregnant women (Pearce, 2015). NHANES 2007-2010 results suggested that 55% of pregnant women and 37% of non-pregnant women of childbearing age had inadequate iodine intakes as measured by urinary iodine (Caldwell et al., 2013).

The USDA-ARS MAFCL FNR group is evaluating several chemicals possessing the ability to inhibit iodine uptake in the thyroid gland, including perchlorate and thiocyanate. Scientific background regarding these two iodine inhibitors is available in Appendices A and B.

To assess iodine status, iodine intake must be estimated, which necessitates the availability of data on iodine content and variability for individual foods (Swanson et al., 2012). Workshops that were convened by the Office of Dietary Supplements, National Institutes of Health (NIH) with other scientists in 2011 and 2014, confirmed the need for food composition tables on iodine content (Swanson et al., 2012; Ershow et al., 2018). Thus, the impetus for this project has been the need for updated and expanded data on the iodine content of foods to address continued public health concerns. The database presented here has been developed, through collaboration between the U.S. Department of Agriculture (USDA) Methods and Application of Food Composition Laboratory (MAFCL) in Beltsville, MD, the U.S. Food and Drug Administration (FDA) Center for Food Safety and Applied Nutrition in College Park, MD and the Office of Dietary Supplements, National Institutes of Health, Bethesda, MD. This release includes about 425 foods.

Background

The FDA collects about 270 kinds of food and beverage samples regionally as part of its Total Diet Study (TDS) and analyzes these samples for nutrient elements, including iodine. These data are available on-line (FDA, 2020). The TDS program began in 1961, and analysis of TDS samples for iodine began in 1973. The TDS currently monitors levels of over 400 contaminants and nutrients in the average U.S. diet; the number varies slightly from year to year. Juan et al. (2016) used the TDS data and data on iodine concentrations in supplements, water, and salt, along with What We Eat In America (WWEIA), the dietary

intake interview portion of NHANES, to estimate intakes of iodine from foods, water, and supplements for the U.S. population. Abt et al. (2018) used updated TDS iodine concentration data to estimate intakes of iodine from food sources.

More recently, the USDA has also been analyzing iodine in foods and dietary supplements obtained nationwide (Pehrsson et al., 2016; Ershow et al., 2016; USDA, 2017). These have included notable dietary food sources of iodine e.g., seaweed, fish and other seafood, dairy foods, iodized salt, eggs, (Rohner et al., 2014; Lee et al., 2016b; Pehrsson et al., 2016) and dietary supplements of vitamins and minerals for adults, children, and pregnant women (USDA, 2017).

A wide range of iodine concentrations in individual foods and variable use of iodized salt at the table and in food preparation make intake assessment especially challenging (Swanson et al., 2012). Varying levels in foods are due to factors such as the amount of iodine in soil where crops are grown, extent of iodine supplementation to animals, use of iodophors as sanitizing agents, and iodine-containing ingredients in processed foods (Ershow et al., 2018). Iodine is present naturally at relatively high levels in seaweed, many saltwater fishes, and other seafood due to the ability to concentrate iodine from their seawater environment (Rohner et al., 2014).

Analytical methodology and quality control

FDA's Kansas City Laboratory (Lenexa, KS), FDA's Maryland Laboratory (College Park, MD) and the USDA contract laboratory analyzed the foods for iodine using inductively coupled plasma mass spectrometry (ICP-MS). The USDA samples were solubilized using a strong base; a stabilizer was added followed by dilution and filtration prior to iodine analysis (Sullivan and Zywicki, 2012). The limit of quantitation (LOQ) for the analyses was 10 mcg/100g of iodine for most of the foods and 50 mcg/100g for the salt samples. FDA solubilized TDS samples using tetramethyl-ammonium hydroxide and a hot block extraction system at 85°C was used to extract the available iodine (Todorov and Gray 2017). FDA's limit of detection (LOD) for iodine ranges from 0.01 – 0.04 mcg/100g; the LOQ ranges from 0.1 – 0.5 mcg/100 g.

Quality control materials used for analysis of the USDA samples included certified standard reference materials (SRMs) from the National Institute of Standards and Technology (NIST, Gaithersburg MD 20899): SRM 1548a (a material representing a typical diet), SRM 1549 (non-fat milk powder), SRM 1849 (nutritional formula), and SRM 3530 (iodized table salt). Secondary reference materials from Virginia Tech (Phillips et al., 2006) that have been cross-validated against the SRMs were also used. FDA used SRM 1549a (whole milk powder) as a reference material for quality control of their iodine analyses. All data from the reference materials as well as the samples were examined either by the quality control committee at USDA or by FDA, depending on the source of the data before acceptance. As a crucial aspect of the data review process, a conservative approach was used. Every value that seemed illogical was investigated and was rejected or retested at the lab, if possible. Data were accepted only by consensus among the scientific panel after thorough consideration.

Source and handling of samples: USDA

Many of the samples analyzed by USDA came from the National Food and Nutrient Analysis Program (NFNAP) which has been in progress since 1999 (Haytowitz and Pehrsson, 2018). The foods collected under this study and other related studies were processed by Virginia Tech, Blacksburg, VA. Foods that

required preparation were made using directions on the packages. For most of the foods, other than milk, eggs, salt, almond beverage, and soy beverage, described below, samples from more than one location were combined.

Where appropriate, the samples were processed with liquid nitrogen, frozen and sent out for analysis with additional aliquots of the food samples stored long-term at -60°C. It was determined that iodine in the samples remains stable under such conditions, by having five samples with different matrices sent for iodine analysis and five years later analyzing aliquots of the same samples. The results did not show any loss of iodine over time and were within the limits of analytical uncertainty.

While this program has been running for over 20 years, many of the food samples are more recent. For example, the whole raw fresh eggs were collected and analyzed in 2019, and the other types of eggs, dried and frozen, primarily for commercial use, were collected in 2017. Non-flavored fluid milks as well as salt samples (including iodized table salt, iodized sea salt, and non-iodized sea salt) were also collected and analyzed in 2018-19.

In addition to examining the uptake of sodium by pasta boiled in salted water, Virginia Tech scientists had the pasta analyzed for iodine uptake since iodized salt was used during cooking (Bianchi et al., 2019). Included in the data are replicate samples using the recommended amount of salt in the cooking water based on the label instructions.

Some samples of white bread, whole wheat bread, and hamburger/hot dog rolls that were obtained in 2019 had labels listing potassium iodate or calcium iodate as a dough conditioner. These baked products were analyzed for iodine to provide information on the amount derived from the dough conditioner.

In an earlier study to determine the fluoride content of tap water in the U.S. (Pehrsson et al., 2006), USDA collected tap water samples from 144 locations around the country. A random subset of 40 of these archived water samples was analyzed for iodine by the FDA laboratory.

Source and handling of samples: FDA

TDS data included in this iodine content database were limited to results for samples collected beginning in 2016 because of a change in analytical methodology at that time. Prior to 2016, TDS samples were analyzed for iodine concentrations using ultraviolet-visible (UV-VIS) spectrophotometry through the catalysis of the Cesium +4/Arsenic +3 reaction (adapted from Fischer et al., 1986). The ICP-MS method in use for iodine analysis since 2016 has limits of detection (LODs) and LOQs that are lower (i.e., better) than the older method, and there is less interference from other signals; these factors affect estimates of element concentrations and exposures.

After receipt of samples, FDA prepares foods as for consumption; for example, apples are washed, bananas are peeled, and oatmeal is cooked. Deionized water is used in washing, cooking, and beverage preparation. Non-iodized salt was used in cooking food mixtures (e.g., cornbread, scrambled eggs, tuna casserole), but no salt was used in cooking single items (e.g., cooked cereals, vegetables).

Prior to the 2018 fiscal year (October 2017 – September 2018), TDS foods were obtained in four regional market basket (MB) collections per year. For each regional collection, products purchased in each of the three cities within the region were composited to form single analytical samples. All TDS foods were collected in each MB.

The TDS food list and sampling plan were modified at the beginning of the 2018 fiscal year as part of FDA's modernization of the program. Some TDS foods were dropped, and others were added. TDS foods were categorized as "regional" foods (possibly varying in nutrient or contaminant concentrations by region or season) or "national" foods (less likely to vary in nutrient and contaminant concentrations by region or season). Under the new sampling plan, regional TDS foods are collected in each of six U.S. regions, in each of two time periods (November – April and May – October); results are presented as collection #1 (October) through collection #12 (September). For each regional collection, products purchased in each of the three cities within a collection region are composited to form single analytical samples. National foods are collected once per year, in Lenexa, KS; the results are presented as collection #13.

In anticipation of the upcoming major change to the TDS sampling plan, FDA conducted a pilot study of the new sampling procedures in the last market basket of 2017 (July). Available data include results for 86 regional foods.

Recent soy beverage and almond beverage sampling was done by USDA with analyses by FDA as described above. Convenience samples of 5 different brands of shelf stable soy beverage and 6 different brands of shelf stable almond beverage were obtained at large grocery stores in Columbia, MD, and College Park, MD, in August 2021. These samples were analyzed along with reference materials at FDA's laboratory in College Park, MD. These results, combined with data from FDA's TDS, were used to obtain mean, standard deviation (SD), and range for each of the two beverage types. See the Data Discussion section below concerning these listings in the datasets.

Database formats, procedures, and notes

The tables provide a description for each food. The Nutrient Data Bank (NDB) numbers familiar to those using USDA's National Nutrient Database for Standard Reference (SR) file and the TDS food numbers from FDA are each shown for reference purposes. Database identification numbers are also provided in column "DB_ID". Descriptions of foods provided in the table may not exactly match SR or TDS descriptions. Foods listed without an SR or TDS number are foods that have not previously been reported in USDA or TDS composition data tables.

SR did not include data on the iodine content of foods. However, as previously noted, USDA arranged for stored NFNAP samples to be analyzed for iodine, focusing primarily on foods likely to have appreciable amounts of iodine. The results of those analyses are included in FoodData Central (FDC). FDA posts TDS data on iodine concentrations on its TDS website along with concentrations of other nutrient elements and contaminants (FDA, 2019).

When both FDA and USDA data were available for the same food types, the data were combined, as indicated in Data Source(s) column in the tables. Where only USDA or FDA is indicated, the data are solely from that source. The year in which the samples were chemically analyzed is shown, which is typically within a year of when the respective food samples were procured.

Footnotes are given where further descriptions or explanations of specific food descriptions are needed. The datasets also provide means, standard deviations, value ranges, and sample sizes.

Data are depicted in two different formats: 1) per 100 grams of food and 2) per serving. Per 100g of food is the unit used both in USDA's previous SR file (Haytowitz and Pehrsson, 2018) and in the current FoodData Central (USDA ARS, 2019). For amounts per serving, the reference which was used for determining each serving size is shown in the "Serving Size Reference" column. Serving sizes were determined primarily by using the FDA's Reference Amounts Customarily Consumed (RACC) (FDA, 2018). For items not specified in RACC guidance, other references were consulted including USDA FoodData Central's SR Legacy, FNDDS and Branded Foods databases (USDA/ARS, 2019), American Egg Board (American Egg Board, 2021), the U.S. Dietary Guidelines for Americans (USDA/USHHS, 2020) and serving sizes from product labels (e.g., infant formulas). "Similar USDA item" is referenced for a few foods where the weight of an identical food was not available in SR Legacy, so the weight of a comparable food in SR Legacy was used. For ease of use, some item weights per serving are rounded within a few grams.

Data discussion

Care should be taken in using data means, especially where the number of analytical samples (n) is very low. As seen in foods having very large numbers (n) of samples analyzed, the variability in the iodine content can be high. However, even where the number of food samples is small, these data provide an estimate of the iodine content and indicate where additional data would be useful, or conversely, where it would not be productive to allocate research resources.

Also affecting the reporting of data on variability is the combining of food samples before analysis. For some foods, both FDA and USDA have homogenized two or more food samples together and then analyzed the composite sample for iodine. The resulting mean values are the best estimates of content but lack information on variability. For the FDA national samples (n=1) and for the milk, egg and salt data reported by USDA, individual samples were analyzed and not combined with any other samples.

FDA's results for foods below the LOD are shown as zero (0), with values between the LOD and LOQ reported as the detected iodine concentrations. The USDA reports data below their LOQ as simply "<". When the USDA's LOQ is the only information available for a food, "<" along with the LOQ is given, but if FDA had data on the same food type, an estimate of the USDA value was made based on the FDA data and included in the calculations. This impacted only 14 foods and 5 of those had USDA values both above and below the LOQ. Using only the higher data would have biased the values, so the values below LOQ were estimated using business rules established for the USDA previous database for foods, SR. For foods where all values were below LOQ and the FDA data was also low, the business rule applied gave estimates with low but not zero values.

It is clear from the data that when iodate dough conditioner is used in bakery production of breads, iodine content is far higher than when that conditioner is not used. Therefore, we separated out the data for breads known or suspected to contain iodate conditioners from those that did not and gave them separate entries in the database. These are white bread, whole wheat bread, white bread buns, and one type of fast-food sandwich bun. All of these have descriptions indicating iodate dough conditioner has been used or is likely to have been used and have an NDB and/or TDS number with an asterisk. While high, none of the iodine values for these breads exceed the regulatory maximum based on the amount of iodate allowed by the FDA (FDA, 2019).

Our analyses of the convenience sampling of shelf-stable soy beverages revealed two samples were considerably higher in iodine content than the other four samples. The ingredient labels of the higher-iodine

samples indicated the presence of seaweed or its derivative, thus influencing iodine levels. Therefore, the seaweed-containing soy beverages have been reported separately in the datasets.

We have excluded a number of foods with high iodine levels that were likely due to presence of erythrosine (also known as FD&C Red No. 3). Erythrosine is a cherry-colored dye used in foods such as decorating gels, glaze cherries and other fruits, some candies, salmon spreads, bakery and snack foods, jellies, ice creams, and popsicles (Gupta et al., 2006; Wenlock and Buss, 1982).¹ Most forms of iodine in food are generally considered very bioavailable (easily absorbed), as high as 99% (Gonzali et al., 2017). The iodine content of erythrosine, however, has been found to have low bioavailability and thus unlikely to be a significant contributor to iodine status (Jahreis et al., 2001).

These data contain only analytical data for iodine content of the foods. In some cases, the food described may not be in the form as consumed, e.g. raw rather than cooked. A direct estimate from raw to prepared is difficult since iodine could, for example, be lost during heating or by moving into cooking water. Estimated prepared values would need to be calculated by the researcher. A retention factor is a way to quantitate the amount of a nutrient remaining in the food after preparation. The principle of using a nutrient retention factor is based on investigations by Reinivuo et al., 2009; Schakel et al., 1997; and Murphy et al., 1975. The appropriate retention factor reflects the effects of food preparation on the food's nutrient content (USDA, 2007). For foods of significant iodine content with only raw data available, additional studies are planned to obtain cooked data for future releases of this database as well as to determine retention factors.

Conclusions

The iodine dataset containing a total of 467 entries is a compilation of both USDA and FDA data having a wide variety of foods. As more data become available, subsequent releases with additional foods are anticipated. These data provide guidance for selecting additional foods for iodine analysis. These data also show where expanding the sample size with more analyses for foods with substantial iodine content would be beneficial.

References

Abt, E., Spungen, J., Pouillot, R., Gamalo-Siebers, M., Wirtz M. (2018). Update on dietary intake of perchlorate and iodine from U.S. Food and Drug Administration's total diet study: 2008–2012. *Journal of Exposure Science & Environmental Epidemiology*, 28:21-30. DOI:10.1038/jes.2016.78

American Egg Board. (2021). Shell Eggs Equivalency. <https://incredibleegg.org/professionals/foodservice/egg-and-egg-products/conversion-to-egg-products> (Accessed on 8 July 2021).

Bianchi, L.M., Phillips, K.M., McGinty, R.C., Ahuja, J.K., Pehrsson, P.R. (2019). Cooking parameters affect the sodium content of prepared pasta. *Food Chemistry*, 271:479-487. DOI:10.1016/j.foodchem.2018.07.198

Caldwell, K.L., Pan, Y., Mortensen, M.E., Makhmudov, A., Merrill, L., Moye, J. (2013). Iodine status in pregnant women in the National Children's Study and in U.S. women (15-44 years), National Health and Nutrition Examination Survey 2005-2010. *Thyroid*, 23(8):927-937. DOI:10.1087/thy.2013.0012

¹ Another widely used red food dye, FD&C Red No. 40, contains no iodine.

- Ershow, A.G., Goodman, G., Coates, P.M., Swanson, C.A. (2016). Research needs for assessing iodine intake, iodine status, and the effects of maternal iodine supplementation. *American Journal of Clinical Nutrition*, 104(Suppl):941S–949S. DOI:10.3945/ajcn.116.134858
- Ershow, A.G., Skeaff, S.A.; Merkel, J.M.; Pehrsson, P.R. (2018). Development of databases on iodine in foods and dietary supplements. *Nutrients*, 10, 100:1-20. DOI:10.3390/nu10010100
- Fischer, P.W., L'Abbe, M.R., Giroux, A. (1986). Colorimetric determination of total iodine in foods by iodide-catalyzed reduction of Ce⁴⁺. *Journal of AOAC International*, 69:687–689.
- Gonzali, S., Kiferle, C., Perata, P. (2017). Iodine biofortification of crops: agronomic biofortification, metabolic engineering and iodine bioavailability. *Current Opinion in Biotechnology*, 44:16-26. DOI:10.1016/j.copbio.2016.10.004
- Gupta, V. K., Mittal, A., Kurup, L., Mittal J. (2006). Adsorption of a hazardous dye, erythrosine, over hen feathers. *Journal of Colloid Interface Science*, 304:52-57. DOI:10.1016/j.jcis.2006.08.032
- Haytowitz, D.B., Pehrsson, P.R. (2018). USDA's National Food and Nutrient Analysis Program (NFNAP) produces high-quality data for USDA food composition databases: Two decades of collaboration, *Food Chemistry*, 238(1): 134-138. DOI:10.1016/j.foodchem.2016.11.082
- Institute of Medicine. (2001). Panel on Micronutrients. Dietary Reference Intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. National Academies Press: Washington, DC, USA; <https://www.ncbi.nlm.nih.gov/books/NBK222323>.
- Jahreis, G., Hausmann, W., Kiessling, G., Franke, K., Leiterer, M. (2001). Bioavailability of iodine from normal diets rich in dairy products – results of balance studies in women. *Experimental and Clinical Endocrinology and Diabetes*, 109:163-167. DOI:10.1055/s-2001-14840
- Juan, WY., Trumbo, P.R., Spungen, J.H., Dwyer, J.T.; Carriquiry, A.L., Zimmerman, T.P., Swanson, C.A., Murphy, S.P. (2016). Comparison of 2 methods for estimating the prevalences of inadequate and excessive iodine intakes. *American Journal of Clinical Nutrition*, 104(Suppl):888S–897S. DOI:10.3945/ajcn.115.110346
- Lee, K.W., Shin, D., Cho, M.S., Song, W.O. (2016a). Food group intakes as determinants of iodine status among US adult population. *Nutrients*, 8:325; DOI:10.3390/nu8060325.
- Lee, K.W., Cho, M.S., Shin, D., Song, W.O. (2016b). Changes in iodine status among US adults, 2001-2012. *International Journal of Food Science and Nutrition*, 67(2):184-194; DOI:10.3109/09637486.2016.1144717.
- Luo, Y., Kawashima, A., Ishido, Y., Yoshihara, A., Oda, K., Hiroi, N., Ito, T., Ishii, N., Suzuki, K. (2014). Iodine excess as an environmental risk factor for autoimmune thyroid disease. *International Journal of Molecular Science*, 15:12895-12912; DOI:10.3390/ijms150712895
- Murphy, E.W., Criner, P.E., Gray, B.C. (1975). Comparison of methods for determining retentions of nutrients in cooked foods. *Journal of Agricultural and Food Chemistry*, 23:1153-1157; DOI:10.1021/jf60202a021
- Pearce, E.N. (2015). Is iodine deficiency reemerging in the United States? *AACE Clinical Case Reports*, 1(1): e81-82. DOI:10.4158/EP14472.CO.
- Pehrsson, P.R., Patterson, K.Y., Spungen, J.H., Wirtz, M.S., Andrews, K.W., Dwyer, J.T., Swanson, C.A. (2016). Iodine in food- and dietary supplement–composition databases. *American Journal of Clinical Nutrition*, 104(Suppl):868S–876S. DOI:10.3945/ajcn.115.110064

- Pehrsson, P.R., Perry, C.R., Cutrufelli, R.C., Patterson, K.Y., Wilger, J., Haytowitz, D.B., Holden, J.M., Day, C.D., Himes, J.H., Harnack, L., Levy, S., Wefel, J., Heilman, J., Phillips, K.M., Rasor, A.S. (2006). Sampling and initial findings for a study of fluoride in drinking water in the United States. *Journal of Food Composition and Analysis*, 19(Suppl):S45-S51. DOI:10.1016/j.jfca.2005.11.004
- Phillips, K.M., Patterson, K.Y., Rasor, A.S., Exler, J., Haytowitz, D.B., Holden, J.M., Pehrsson, P.R. (2006). Quality control materials in the USDA National Food and Nutrient Analysis Program (NFNAP). *Analytical and Bioanalytical Chemistry*, 384(6):1341-1355. DOI:10.1007/s00216-005-0294-0
- Reinivuo, H., Bell, S., Ovaskainen, M-L. (2009). Harmonisation of recipe calculation procedures in European food composition databases. *Journal of Food Composition and Analysis*, 22:410-413. DOI:10.1016/j.jfca.2009.04.003
- Rohner, F., Zimmermann, M., Jooste, P., Pandav, C., Caldwell, K., Raghavan, R., Raiten, D.J. (2014). Biomarkers of nutrition for development—iodine review. *Journal of Nutrition*, 144:1322S–1342S. DOI:10.3945/jn.113.181974
- Schakel, S.F., Buzzard, I.M., Gebhardt, S.E. (1997). Procedures for estimating nutrient values for food composition databases. *Journal of Food Composition and Analysis*, 10:102-114.
- Sullivan, D., Zywicki, R. (2012). Determination of total iodine in foods and dietary supplements using inductively coupled plasma-mass spectrometry. *Journal of AOAC International*, 95(1):195-202. DOI:10.5740/jaoacint.11-350
- Swanson, C.A., Zimmermann, M.B., Skeaff, S., Pearce, E.N., Dwyer, J.T., Trumbo, P.R., Zehaluk, C., Andrews, K.W., Carriquiry, A., Caldwell, K.L., et al. (2012). Summary of an NIH workshop to identify research needs to improve the monitoring of iodine status in the United States and to inform the DRI. *Journal of Nutrition*, 142(Suppl): 1175S–1185S. DOI:10.3945/jn.111.156448
- Todorov, T.I., Gray, P.J. (2017). Inductively coupled plasma-mass spectrometric determination of iodine in food using tetramethyl ammonium hydroxide extraction. Elemental Analysis Manual Method 4.13. U.S. Food and Drug Administration. <https://www.fda.gov/media/103112/download> (accessed on 9 December 2019).
- U.S. Department of Agriculture (USDA), Agricultural Research Service (ARS). (2019). FoodData Central. <https://fdc.nal.usda.gov/> (accessed on 11 June 2019).
- U.S. Department of Agriculture (USDA), Agricultural Research Service (ARS). (2007). USDA Table of Nutrient Retention Factors, Release 6. Version current 2007. <http://www.ars.usda.gov/Main/docs.htm?docid=9448> (accessed on 28 January 2020).
- U.S. Department of Agriculture (USDA), Agricultural Research Service (ARS) and U.S. Department of Health and Human Services, National Institutes of Health, Office of Dietary Supplements. (2017) Dietary Supplement Ingredient Database (DSID) release 4.0. <https://dsid.usda.nih.gov> (accessed on 17 May 2020).
- U.S. Department of Agriculture (USDA) and U.S. Department of Health and Human Services (USHHS). (2020). Dietary Guidelines for Americans, 2020-2025. 9th Edition. <https://dietaryguidelines.gov/> (assessed on 10 August 2021)
- U.S. Food and Drug Administration (FDA). 2020. Total Diet Study. Available online: <https://www.fda.gov/food/science-research-food/total-diet-study> (accessed on 13 May 2020).

U.S. Food and Drug Administration (FDA). 2019. Code of Federal Regulations, Title 21 CFR136.110. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcr/CFRSearch.cfm?fr=136.110> (accessed on 13 May 2020).

U.S. Food and Drug Administration (FDA). 2018. Guidance for Industry: Reference Amounts Customarily Consumed (List of Products for Each Product Category). <https://www.fda.gov/media/102587> (accessed on 1 September 2021).

Wenlock, R.W., Buss, D.H. (1982). Trace nutrients 4*. Iodine in British food. *British Journal of Nutrition*, 47:381-390. DOI:10.1079/BJN19820049

Appendix A: Perchlorate

Perchlorate (ClO_4^-) is a stable anion which forms naturally in the atmosphere and in arid regions like the Southwestern US (Dasgupta et al., 2005). Perchlorate is also synthetically manufactured for use in solid rocket propellant, fireworks, and safety flares (Dasgupta et al., 2006). The U.S. Environmental Protection Agency (EPA) does not regulate perchlorate in drinking water, but based on current epidemiological evidence, maintains a daily oral exposure reference dose of 0.007 mg/kg-day (Greer et al., 2002; EPA, 2005; EPA, 2022a).

The thyroid gland is responsible for regulating metabolism and is critical during pregnancy for the growth and development of newborns. Excessive perchlorate exposure causes disruption to the sodium/iodide symporter (NIS), the plasma membrane used in the production of thyroid hormones (Wolff, 1998; Pleus and Corey, 2018). Perchlorate has the highest potency of all known iodine uptake inhibitors and competes against iodide to block the NIS directly, resulting in the depletion of thyroid hormones (Tonacchera et al., 2004). Because of its effectiveness, perchlorate has been used historically as a drug to treat hyperthyroidism (Wolff, 1998). Perchlorate's thyroid-suppressing effects are temporarily and subside once exposure is removed.

Due to monitoring efforts by the U.S. EPA, perchlorate drinking water levels are of minimal public health concern (EPA, 2022b). Therefore, dietary perchlorate is proposed as the main exposure source in the United States. The FDA maintains the largest dataset of perchlorate in foods and beverages. Perchlorate has been detected in many food categories including dairy, fruits, vegetables, meats, and legumes and grains (FDA, 2017; FDA, 2018) although amounts are very small relative to their effect on iodine uptake.

Nearly all exposure to dietary perchlorate is below the current reference dose and thus at safe levels. Furthermore, scientific evidence suggests no causal relationship exists between changes in thyroid hormone levels of the US population and environmental perchlorate exposure levels (Pleus and Corey, 2018; Charnley, 2008). Therefore, despite perchlorate's potency and thyroid disruption potential, it is likely a minor contributor to iodine inhibition in the thyroid gland.

The MAFCL FNR group's research will continue to evaluate the assessment of iodine uptake inhibitors for health outcomes.

References

- Charnley, G. (2008). Perchlorate: overview of risks and regulation. *Food and Chemical Toxicology*, 46(7), 2307–2315. DOI:10.1016/j.fct.2008.03.006
- Dasgupta, P. K., Martinelango, P. K., Jackson, W. A., Anderson, T. A., Tian, K., Tock, R. W., & Rajagopalan, S. (2005). The Origin of Naturally Occurring Perchlorate: The Role of Atmospheric Processes. *Environmental Science & Technology*, 39(6), 1569–1575. DOI:10.1021/es048612x
- Dasgupta, P. K., Dyke, J. V., Kirk, A. B., & Jackson, W. A. (2006). Perchlorate in the United States. Analysis of Relative Source Contributions to the Food Chain. *Environmental Science & Technology*, 40(21), 6608–6614. DOI:10.1021/es061321z
- Greer, M. A., Goodman, G., Pleus, R. C., & Greer, S. E. (2002). Health effects assessment for environmental perchlorate contamination: the dose response for inhibition of thyroidal radioiodine uptake in humans. *Environmental Health Perspectives*, 110(9), 927–937. DOI:10.1289/ehp.02110927
- Pleus, R. C., Corey, L. M. (2018). Environmental exposure to perchlorate: A review of toxicology and human health. *Toxicology and Applied Pharmacology*, 358, 102–109. DOI:10.1016/j.taap.2018.09.001

Tonacchera, M., Pinchera, A., Dimida, A., Ferrarini, E., Agretti, P., Vitti, P., Santini, F., Crump, K., & Gibbs, J. (2004). Relative potencies and additivity of perchlorate, thiocyanate, nitrate, and iodide on the inhibition of radioactive iodide uptake by the human sodium iodide symporter. *Thyroid*, 14(12), 1012–1019. DOI:10.1089/thy.2004.14.1012

U.S. Environmental Protection Agency (EPA). (2005, February 18). Perchlorate (ClO₄) and Perchlorate Salts. Integrated Risk Information System (IRIS). https://cfpub.epa.gov/ncea/iris2/chemicalLanding.cfm?substance_nmbr=1007 (assessed on 23 June 2022).

U.S. Environmental Protection Agency (EPA). (2022a, April 1). Perchlorate in Drinking Water. United States Environmental Protection Agency. <https://www.epa.gov/sdwa/perchlorate-drinking-water> (assessed on 12 August 2022).

U.S. Environmental Protection Agency (EPA). (2022b, March 31). EPA Announces Plan to Protect the Public from Perchlorate in Drinking Water. EPA. <https://www.epa.gov/newsreleases/epa-announces-plan-protect-public-perchlorate-drinking-water> (assessed on 22 June 2022).

U.S. Food and Drug Administration (FDA). (2017, December 27). Survey Data on Perchlorate in Food - 2005-2006 and 2008-2012 Total Diet Study Results. Center for Food Safety and Applied Nutrition. <https://www.fda.gov/food/chemical-contaminants-food/survey-data-perchlorate-food-2005-2006-and-2008-2012-total-diet-study-results> (assessed on 22 June 2022).

U.S. Food and Drug Administration (FDA). (2018, January 25). Exploratory Survey Data on Perchlorate in Food 2004-2005. Center for Food Safety and Applied Nutrition. <https://www.fda.gov/food/chemical-contaminants-food/exploratory-survey-data-perchlorate-food-2004-2005> (assessed on 22 June 2022).

Wolff, J. (1998). Perchlorate and the thyroid gland. *Pharmacological Reviews*, 50(1), 89–105.

Appendix B: Thiocyanate

Thiocyanate (SCN^-) is a moderately potent iodine uptake inhibitor and biotransformation metabolite (Tonacchera et al., 2004). Exposure sources conducive to thiocyanate generation are cyanide produced during cigarette smoking and the ingestion of two plant products: cyanogenic glycosides and glucosinolates (Laurberg et al., 2009). Thiocyanate is not subject to regulation in the United States because it is a naturally occurring metabolite.

Detoxification from exposure sources into thiocyanate occurs via three metabolic routes:

1. Cyanide from cigarette smoke is primarily hydrolyzed in the cyanide metabolic pathway, facilitated by the enzyme rhodanese, to produce thiocyanate (Logue et al., 2010).
2. Cyanogenic glycosides are a naturally occurring class of plant toxins. Disruption to plant tissue creates hydrogen cyanide (cyanogenesis) which is principally detoxified to thiocyanate (Gleadow and Møller, 2014).
3. Glucosinolates, secondary plant metabolites found in the Brassicaceae family, are complex compounds hypothesized to offer many protective health benefits. Plants in the Brassicaceae family also contain a defensive enzyme called myrosinase which breaks down glucosinolates. When Brassicaceae plant tissue is disrupted, myrosinase hydrolyzes glucosinolates to produce several compounds, including thiocyanate (Barba et al., 2016). Myrosinase is denatured when plants are cooked; therefore, glucosinolates may also be broken down naturally by gut microbiota (Barba et al., 2016; Agerbirk et al., 2008).

The thyroid gland is responsible for regulating metabolism and is critical during pregnancy for the growth and development of newborns. Thiocyanate disrupts at least two thyroid processes (Willemijn and Lumen, 2017). High levels of thiocyanate compete against iodide to temporarily block the sodium/iodide symporter (NIS) channel, the plasma membrane used in the production of thyroid hormones. Excess thiocyanate also disrupts biochemical processes in the thyroid gland involving iodine (organification). Thiocyanate's thyroid-suppressing effects are temporarily and subside once exposure is removed.

Due to their ubiquity in food, research suggests dietary exposure to thiocyanates and nitrates far outweighs dietary exposure to perchlorate (De Groef et al., 2006; Eisenbrand and Gelbke, 2016). However, quantitative data is scarce because thiocyanate is only formed after disrupting plant cells through chopping, cooking, and other processes. Thiocyanate has been detected in dairy products, infant formula, and vegetables (Laurberg et al., 2002; Niemann and Anderson, 2008; Sanchez et al., 2007). Epidemiological evidence suggests no association between thyroid function and urinary thiocyanate (Mortensen et al., 2016; Horton et al., 2015; Leung et al., 2011; Pearce et al., 2010; Steinmaus et al., 2013; Lewandowski, Peterson, and Charnley, 2015). Furthermore, negative effects of dietary thiocyanate may be mitigated by modest iodine fortification (Agerbirk et al., 2008). Therefore, dietary consumption levels of thiocyanates do not appear to pose an adverse effect to human health and are likely of low concern when evaluating the risk of iodine uptake inhibition.

The MAFCL FNR group's research will continue to evaluate the assessment of iodine uptake inhibitors for health outcomes.

References

Agerbirk, N., De Vos, M., Kim, J. H., & Jander, G. (2008). Indole glucosinolate breakdown and its biological effects. *Phytochemistry Reviews*, 8(1), 101–120. DOI:10.1007/s11101-008-9098-0

- Barba, F. J., Nikmaram, N., Roohinejad, S., Khelfa, A., Zhu, Z., & Koubaa, M. (2016). Bioavailability of Glucosinolates and Their Breakdown Products: Impact of Processing. *Frontiers in Nutrition*, 3. DOI:10.3389/fnut.2016.00024
- De Groef, B., Decallonne, B. R., Van der Geyten, S., Darras, V. M., & Bouillon, R. (2006). Perchlorate versus other environmental sodium/iodide symporter inhibitors: potential thyroid-related health effects. *European Journal of Endocrinology*, 155(1), 17–25. DOI:10.1530/eje.1.02190
- Eisenbrand, G., & Gelbke, H.-P. (2016). Assessing the potential impact on the thyroid axis of environmentally relevant food constituents/contaminants in humans. *Archives of Toxicology*, 90(8), 1841–1857. DOI:10.1007/s00204-016-1735-6
- Gleadow, R. M., & Møller, B. L. (2014). Cyanogenic Glycosides: Synthesis, Physiology, and Phenotypic Plasticity. *Annual Review of Plant Biology*, 65(1), 155–185. DOI:10.1146/annurev-arplant-050213-040027
- Horton, M. K., Blount, B. C., Valentin-Blasini, L., Wapner, R., Whyatt, R., Gennings, C., & Factor-Litvak, P. (2015). Co-occurring exposure to perchlorate, nitrate and thiocyanate alters thyroid function in healthy pregnant women. *Environmental Research*, 143, 1–9. DOI:10.1016/j.envres.2015.09.013
- Laurberg, P., Andersen, S., Knudsen, N., Ovesen, L., Nøhr, S. B., & Bülow Pedersen, I. (2002). Thiocyanate in food and iodine in milk: From domestic animal feeding to improved understanding of cretinism. *Thyroid*, 12(10), 897–902. DOI:10.1089/105072502761016520
- Laurberg, P., Pedersen, I. B., Carlé, A., Andersen, S., Knudsen, N., & Karmisholt, J. (2009). The Relationship between Thiocyanate and Iodine. *Comprehensive Handbook of Iodine: Nutritional, Biochemical, Pathological and Therapeutic Aspects*, 275–281. DOI:10.1016/b978-0-12-374135-6.00028-5
- Leung, A. M., LaMar, A., He, X., Braverman, L. E., & Pearce, E. N. (2011). Iodine status and thyroid function of Boston-area vegetarians and vegans. *The Journal of Clinical Endocrinology & Metabolism*, 96(8). DOI:10.1210/jc.2011-0256
- Lewandowski, T. A., Peterson, M. K., & Charnley, G. (2015). Iodine supplementation and drinking-water perchlorate mitigation. *Food and Chemical Toxicology*, 80, 261–270. DOI:10.1016/j.fct.2015.03.014
- Logue, B. A., Hinkens, D. M., Baskin, S. I., & Rockwood, G. A. (2010). The Analysis of Cyanide and its Breakdown Products in Biological Samples. *Critical Reviews in Analytical Chemistry*, 40(2), 122–147. DOI:10.1080/10408340903535315
- Mortensen, M. E., Birch, R., Wong, L.-Y., Valentin-Blasini, L., Boyle, E. B., Caldwell, K. L., Merrill, L. S., Moye, J., & Blount, B. C. (2016). Thyroid antagonists and thyroid indicators in U.S. pregnant women in the Vanguard Study of the National Children's Study. *Environmental Research*, 149, 179–188. DOI:10.1016/j.envres.2016.05.017
- Niemann, R. A., & Anderson, D. L. (2008). Determination of iodide and thiocyanate in powdered milk and infant formula by on-line enrichment ion chromatography with photodiode array detection. *Journal of Chromatography A*, 1200(2), 193–197. DOI:10.1016/j.chroma.2008.05.064
- Pearce, E. N., Lazarus, J. H., Smyth, P. P., He, X., Dall'Amico, D., Parkes, A. B., Burns, R., Smith, D. F., Maina, A., Bestwick, J. P., Jooman, M., Leung, A. M., & Braverman, L. E. (2010). Perchlorate and

Thiocyanate Exposure and Thyroid Function in First-Trimester Pregnant Women. *The Journal of Clinical Endocrinology & Metabolism*, 95(7), 3207–3215. DOI:10.1210/jc.2010-0014

Sanchez, C. A., Blount, B. C., Valentin-Blasini, L., & Krieger, R. I. (2007). Perchlorate, Thiocyanate, and Nitrate in Edible Cole Crops (*Brassica* sp.) Produced in the Lower Colorado River Region. *Bulletin of Environmental Contamination and Toxicology*, 79(6), 655–659. DOI:10.1007/s00128-007-9292-6

Steinmaus, C., Miller, M. D., Cushing, L., Blount, B. C., & Smith, A. H. (2013). Combined effects of perchlorate, thiocyanate, and iodine on thyroid function in the National Health and Nutrition Examination Survey 2007–08. *Environmental Research*, 123, 17–24. DOI:10.1016/j.envres.2013.01.005

Tonacchera, M., Pinchera, A., Dimida, A., Ferrarini, E., Agretti, P., Vitti, P., Santini, F., Crump, K., & Gibbs, J. (2004). Relative Potencies and Additivity of Perchlorate, Thiocyanate, Nitrate, and Iodide on the Inhibition of Radioactive Iodide Uptake by the Human Sodium Iodide Symporter. *Thyroid*, 14(12), 1012–1019. DOI:10.1089/thy.2004.14.1012

Willemin, M.-E., & Lumen, A. (2017). Thiocyanate: a review and evaluation of the kinetics and the modes of action for thyroid hormone perturbations. *Critical Reviews in Toxicology*, 47(7), 543–569. DOI:10.1080/10408444.2017.1281590

USDA, FDA and ODS-NIH Database for the Iodine Content of Common Foods mcg per 100 grams
Release 3, February 2023

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s) Analyzed
Dairy and Eggs										
91	01001	164	Butter, salted	11	4.6	0.6	3.7	5.9	FDA	2016-2020
359	01192		Cheese product, made with 2% reduced fat milk	1	54				USDA	2013
24	01253	10	Cheese, American, processed	29	58	18.9	35	112	FDA/USDA	2011,2016-2020
25	01009	12	Cheese, cheddar (sharp/mild)	38	46	13	27	106	FDA/USDA	2013,2016-2020
450	01019		Cheese, feta, whole milk, crumbled	8	48	21	30	76	USDA	2022
190	01025	391	Cheese, Monterey Jack	28	41	12	20	77	FDA	2017-2020
191	01029	392	Cheese, mozzarella	30	51	10	27	67	FDA/USDA	2013,2017-2020
349	01032		Cheese, parmesan, grated	9	82	38	45	175	USDA	2013,2022
348	43379		Cheese, processed, Swiss, low fat	1	60				USDA	2012
350	01036		Cheese, ricotta, whole milk	1	66				USDA	2017
127	01040	236	Cheese, Swiss	35	137	113	33	440	FDA	2016-2020
161	01015	332	Cottage cheese, creamed, reduced fat	11	37	7.2	25	47	FDA/USDA	2013,2016-2020
489	01012		Cottage cheese, full fat, small or large curd	10	46	13	20	63	USDA	2022
128	01017	237	Cream cheese	18	36	10	21	55	FDA/USDA	2016-2020,2022
94	01068	168	Cream substitute, non-dairy, fluid	10	0.6	0.2	0.4	1.0	FDA	2016-2020
351	01053		Cream, heavy whipping, fluid	9	21	7.0	5.3	30	USDA	2013,2022
93	01049	167	Cream, half & half	36	29	6.1	18	46	FDA/USDA	2013,2016-2020
362	01258		Egg, white, dried, stabilized, glucose reduced	15	34	11	22	59	USDA	2018
358	01172		Egg, white, raw, frozen, pasteurized	6	<10				USDA	2018

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
355	01133		Egg, whole, dried	17	274	52	202	407	USDA	2018
353	01123		Egg, whole, raw, fresh	24	49	21	27	115	USDA	2019
357	01171		Egg, whole, raw, frozen, pasteurized	14	62	13.9	45	81	USDA	2018
356	01137		Egg, yolk, dried	17	349	66	261	437	USDA	2018
354	01126		Egg, yolk, raw, frozen, pasteurized	15	177	52	76	267	USDA	2018
34	01129	37	Eggs, hard-boiled	35	61	29.6	14	120	FDA	2016-2020
336	01132	35	Eggs, scrambled with oil	7	43	13	20	57	FDA	2016-2017
192	19270	393	Ice cream, chocolate	28	31	4.8	24	42	FDA	2017-2020
341	01243	177	Ice cream, lowfat, vanilla	7	45	9.8	27	56	FDA	2016-2017
143	19095	286	Ice cream, regular (not lowfat), vanilla	34	38	5.8	26	53	FDA	2016-2020
446	01088		Milk, buttermilk, lowfat	8	34	5.3	26	42	USDA	2022
22	01103	3	Milk, chocolate, lowfat, fluid	37	30	6.8	17	44	FDA/USDA	2014,2016-2020
21	01079	2	Milk, reduced fat (2%), fluid	59	36	13	20	103	FDA/USDA	2016-2020
352	01082		Milk, lowfat (1%), fluid	24	36	15	13	85	USDA	2018
23	01085	4	Milk, skim, fluid	59	34	9.2	15	65	FDA/USDA	2016-2020
20	01077	1	Milk, whole, fluid	59	34	11	17	84	FDA/USDA	2016-2020
154	01056	300	Sour cream	18	31	5.8	22	48	FDA/USDA	2016-2020,2022
290		333	Sour cream dip, any flavor	7	25	6.1	16	31	FDA	2016-2017
295	01298	409	Yogurt, frozen, vanilla	27	46	9.4	26	76	FDA	2017-2020
447	01293		Yogurt, Greek, whole milk, plain	8	42	19	17	79	USDA	2022
361	01256		Yogurt, Greek, plain, nonfat	6	51	3.5	45	56	USDA	2013
363	01284		Yogurt, Greek, strawberry, lowfat	1	42				USDA	2014
364	01285		Yogurt, Greek, strawberry, nonfat	2	41	2.8	39	43	USDA	2007
365	01286		Yogurt, Greek, vanilla, nonfat	2	50	1.6	49	52	USDA	2014
126	01122	235	Yogurt, fruit-flavored, lowfat	11	31	6.5	23	43	FDA/USDA	2013,2016-2020
222	01220	521	Yogurt, vanilla, lowfat	3	34	3.0	31	37	FDA	2018-2020

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
448	01116		Yogurt, whole milk, plain	8	32	12	22	57	USDA	2022
Spices and Miscellaneous										
487		902	Baking powder	1	0.0				FDA	2020
488		910	Cocoa powder	1	0.0				FDA	2020
19	02047	909	Salt, table, iodized	26	5213	1329	3410	8900	FDA/USDA	2017-2018,2020
372			Sea salt, iodized	9	4663	1143	2250	6080	USDA	2018
373		904	Sea salt, non-iodized	28	48	9.4	0	50	FDA/USDA	2018,2020
Soups, Sauces, Gravies and Condiments										
273		532	Broth, chicken, ready-to-use from carton	3	0.7	0.2	0.5	0.9	FDA	2018-2020
180	06116	374	Brown gravy, canned or bottled	10	1.5	0.3	1.0	2.0	FDA	2016-2020
412	27052		Dip, salsa con queso (salsa with cheese)	6	19	14	10	47	USDA	2012-2013
98	11935	173	Ketchup, tomato, bottled	10	1.0	0.4	0.6	1.9	FDA	2016-2020
152	02046	298	Mustard, yellow, plain	10	0.4	0.4	0.0	1.0	FDA	2016-2020
177	06164	359	Salsa, tomato, bottled	10	1.6	0.8	0.0	2.9	FDA	2016-2020
203	06150	405	Sauce, barbecue, bottled	27	2.7	1.3	0.9	7.0	FDA	2018-2020
366	06179		Sauce, fish, ready-to-serve	1	21				USDA	2008
219	16123/1 6124	518	Sauce, soy	3	0.5	0.1	0.4	0.6	FDA	2018-2020
12	27050	373	Sauce, sweet & sour, bottled	7	0.7	0.4	0.4	1.5	FDA	2016-2017
221	06931	520	Sauce, tomato pasta, bottled	3	1.6	0.4	1.3	2.1	FDA	2018-2020
5	06404	283	Soup, bean with bacon/pork, canned, condensed, prepared with water	7	0.3	0.1	0.1	0.4	FDA	2016-2017
274		533	Soup, broccoli cheese, canned, ready to eat (RTE)	3	4.6	0.2	4.5	4.8	FDA	2018-2020
246		155	Soup, chicken noodle, canned, ready to eat (RTE)	10	0.9	0.2	0.7	1.5	FDA	2016-2020
230	06030	534	Soup, clam chowder, New England, canned, ready to eat (RTE)	3	11	7.9	6.3	20	FDA	2018-2020

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
319		285	Soup, clam chowder, New England, canned, condensed, prepared with whole milk	7	20	5.1	13	27	FDA	2016-2017
275		535	Soup, cream of mushroom, canned, ready to eat (RTE)	3	2.2	0.1	2.1	2.3	FDA	2018-2020
276		536	Soup, cream of potato, canned, ready to eat (RTE)	3	3.3	0.4	3.0	3.7	FDA	2018-2020
262		367	Soup, ramen noodles, prepared with water	10	0.3	0.4	0.0	0.9	FDA	2016-2020
247		156	Soup, tomato, canned, ready to eat (RTE)	10	0.7	0.2	0.3	1.1	FDA	2016-2020
231	06742	537	Soup, vegetable beef, canned, ready to eat (RTE)	3	1.8	1.1	0.8	2.9	FDA	2018-2020
312	06471	157	Soup, vegetable beef, canned, condensed, prepared with water	7	1.1	0.4	0.8	2.0	FDA	2016-2017
277		538	Soup, vegetable, canned, ready to eat (RTE)	3	0.9	0.0	0.9	0.9	FDA	2018-2020
Baby Foods										
302	03165	730	Baby food, apples with berries	8	0.1	0.1	0.0	0.3	FDA	2016-2017,2020
240	03163	731	Baby food, apples with fruit other than berries	15	0.2	0.2	0.0	0.5	FDA	2016-2020
122	03117	225	Baby food, applesauce	15	0.0	0.1	0.0	0.2	FDA	2016-2020
473		755	Baby food, apple and sweet potato with cinnamon, pouch	5	0.1	0.2	0.0	0.3	FDA	2019
470		751	Baby food, apple, spinach and avocado, bowl/pouch	5	2.5	0.2	2.3	2.7	FDA	2019
474		756	Baby food, apple, sweet potato and pineapple, pouch	5	0.5	0.2	0.4	0.7	FDA	2019
298	03143	717	Baby food, apricots with mixed fruit	7	0.3	0.1	0.2	0.4	FDA	2016-2017
157	43546	313	Baby food, bananas	15	0.2	0.4	0.0	1.7	FDA	2016-2020
463		744	Baby food, banana and blueberry, pouch	5	0.6	0.4	0.0	0.9	FDA	2019

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
462		743	Baby food, banana and strawberry, glass jar	5	0.2	0.2	0.0	0.4	FDA	2019
464		745	Baby food, banana, apple and pear, plastic jar	5	0.0	0.0	0.0	0.0	FDA	2019
461		742	Baby food, banana, blackberry and blueberry, plastic jar	5	0.2	0.4	0.0	0.8	FDA	2019
123	03131	226	Baby food, peaches	29	0.1	0.2	0.0	0.9	FDA	2016-2020
124	03133	227	Baby food, pears	15	0.0	0.1	0.0	0.1	FDA	2016-2020
296	03159	713	Baby food, pears and pineapple	7	0.2	0.1	0.0	0.3	FDA	2016-2017
465		746	Baby food, organic pears and spinach, pouch	5	0.7	0.2	0.5	1.0	FDA	2019
468		749	Baby food, pear, blueberry, apple and avocado, pouch	5	0.4	0.1	0.3	0.6	FDA	2019
478		760	Baby food, pear, mango, avocado, pouch	5	0.1	0.3	0.0	0.6	FDA	2019
297	03170	714	Baby food, plums/prunes with apples or pears	7	0.6	0.7	0.0	2.2	FDA	2016-2017
241	03139	736	Baby food, prunes	8	1.0	0.6	0.5	2.4	FDA	2018-2020
477		759	Baby food, mango, carrot and turmeric, bowl	2	0.7	0.0	0.7	0.7	FDA	2019
459		740	Baby food, mango, glass jar	5	0.4	0.0	0.3	0.4	FDA	2019
460		741	Baby food, mango, pouch	5	0.6	0.1	0.6	0.7	FDA	2019
479		761	Baby food, mango, yellow zucchini, corn and turmeric, pouch	2	1.5	0.2	1.3	1.6	FDA	2019
125	03166	230	Baby food, juice, apple	15	1.0	1.2	0.0	3.8	FDA	2016-2020
237	44074	712	Baby food, juice, grape	15	0.8	0.5	0.3	1.8	FDA	2016-2020
236	43408	711	Baby food, juice, pear	15	0.3	0.3	0.0	1.1	FDA	2016-2020
480		762	Baby food, juice, mixed fruit	3	0.0	0.0	0.0	0.0	FDA	2019
117	03100	218	Baby food, carrots	21	0.5	0.4	0.0	1.4	FDA	2016-2020
118	03092	219	Baby food, green beans	15	0.3	0.4	0.0	1.2	FDA	2016-2020
121	03121	223	Baby food, peas	15	0.1	0.1	0.0	0.2	FDA	2016-2020

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
466		747	Baby food, peas and spinach, glass jar	1	0.0				FDA	2019
469		750	Baby food, peas, green beans and avocado, pouch	5	0.4	0.1	0.3	0.5	FDA	2019
476		758	Baby food, pumpkin, banana, papaya and cardamom, bowl	3	0.2	0.2	0.0	0.3	FDA	2019
159	03105	320	Baby food, squash	15	0.4	0.4	0.0	1.2	FDA	2016-2020
120	03109	221	Baby food, sweet potatoes	15	0.4	0.2	0.0	0.8	FDA	2016-2020
475		757	Baby food, sweet potato, apple and corn, pouch	5	0.8	0.2	0.5	0.9	FDA	2019
467		748	Baby food, sweet potato, apple and spinach, pouch	5	1.1	0.3	0.8	1.3	FDA	2019
119	03279	220	Baby food, vegetables, mixed	15	0.4	0.2	0.0	0.9	FDA	2016-2020
482	03189	764	Baby food, cereal, oatmeal, dry	7	0.4	1.1	0.0	2.9	FDA	2019-2020
483		765	Baby food, cereal, rice, dry	9	0.0	0.0	0.0	0.0	FDA	2019-2020
484		766	Baby food, cereal, mixed, dry	7	1.0	1.7	0.0	3.9	FDA	2019-2020
278		701	Baby food, cereal, mixed, prepared with water	8	0.0	0.0	0.0	0.1	FDA	2016-2018
300		725	Baby food, cereal, oatmeal with fruit, prepared with water	7	0.0	0.1	0.0	0.1	FDA	2016-2017
258		323	Baby food, cereal, oatmeal, prepared with water	8	0.1	0.1	0.0	0.4	FDA	2016-2018
259		324	Baby food, cereal, rice, prepared with water	20	0.0	0.1	0.0	0.3	FDA	2016-2019
471		753	Baby food, organic yogurt, apple, pumpkin, cinnamon and quinoa, pouch	4	11	2.1	8.8	13	FDA	2019
238	43523	721	Baby food, fruit yogurt dessert	15	23	5.5	14	32	FDA	2016-2020
481		763	Baby food, yogurt, peach pear	3	33	3.8	30	37	FDA	2019
257		317	Baby food, biscuits, teething	15	20	8.0	3.0	28	FDA	2016-2020
299	03214	723	Baby food, cookies, arrowroot	8	8.9	2.2	6.1	13	FDA/USDA	2015-2017
280		733	Baby food, finger foods, puffed snack	8	0.3	0.9	0.0	2.6	FDA	2018-2020

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
432	42284		Baby food, finger food, baked product, cereal, fortified	1	<10				USDA	2014
114	03069	214	Baby food, chicken noodle dinner	15	0.5	0.7	0.0	2.6	FDA	2016-2020
301	43008	726	Baby food, chicken with rice	7	0.7	0.5	0.0	1.3	FDA	2016-2017
279		732	Baby food, macaroni and cheese with vegetables	14	7.0	1.3	4.2	8.8	FDA	2016-2020
115	03045	215	Baby food, pasta, tomato and beef	15	2.0	1.4	0.0	4.6	FDA	2016-2020
242	03046	737	Baby food, ravioli, cheese-filled, with tomato sauce	8	4.6	0.6	3.7	5.8	FDA	2018-2020
111	03002	205	Baby food, beef and broth/gravy	15	1.3	0.8	0.0	3.0	FDA	2016-2020
343	03013	207	Baby food, chicken and broth/gravy	7	28	8.2	11	35	FDA	2016-2017
160	03016	328	Baby food, turkey and broth/gravy	15	1.1	0.5	0.0	2.5	FDA	2016-2020
116	03083	216	Baby food, turkey and rice	15	0.5	0.4	0.0	1.8	FDA	2016-2020
472		754	Baby food, turkey, quinoa, apple and sweet potato, pouch	5	0.7	0.1	0.6	0.7	FDA	2019
112	03055	211	Baby food, vegetables and beef	10	0.5	0.4	0.0	1.1	FDA	2016-2020
113	03073	212	Baby food, vegetables and chicken	10	0.5	0.3	0.0	0.9	FDA	2016-2020
239	03085	728	Baby food, vegetables and turkey	14	2.1	2.4	0.0	6.7	FDA	2016-2020
485		767	Baby food, infant formula, milk-based, powder	2	115	21	100	130	FDA	2019-2020
486		768	Baby food, infant formula, soy-based, powder	2	98	0.7	97	98	FDA	2019-2020
249		202/734	Baby food, Infant formula, milk-based, iron fortified, ready-to-feed	8	13	1.7	12	16	FDA	2016-2018
321		309/735	Baby food, Infant formula, soy-based, ready-to-feed	8	11	4.4	0.6	15	FDA	2016-2018
283		738	Baby food, water, baby, bottled	8	0.0	0.0	0.0	0.0	FDA	2018-2020
Fats and Oils										
90	04691	162	Margarine, salted	10	1.2	1.5	0.0	3.9	FDA	2016-2020
92	04018	166	Mayonnaise, regular, bottled	10	5.3	1.6	2.6	7.6	FDA	2016-2020

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
182	04053	378	Oil, olive	10	0.2	0.2	0.0	0.5	FDA	2016-2020
183	04044	379	Oil, vegetable	10	0.0	0.0	0.0	0.0	FDA	2016-2020
433	43016		Salad dressing, coleslaw, bottled	1	<10				USDA	2017
329	43215	376	Salad dressing, creamy/buttermilk type, low-calorie, bottled	7	13	2.0	9.2	15	FDA	2016-2017
328		375	Salad dressing, creamy/buttermilk type, regular, bottled	7	8.4	1.6	7.1	12	FDA	2016-2017
181	04114	377	Salad dressing, Italian, regular, bottled	10	0.7	0.4	0.4	1.7	FDA	2016-2020
217	04640	516	Salad dressing, ranch, low-calorie, bottled	3	7.8	0.4	7.5	8.2	FDA	2018-2020
218	04639	517	Salad dressing, ranch, regular, bottled	6	10	4.9	5.9	19	FDA/USDA	2013,2018-2020
Fruit										
54	09003	78	Apple, red, with peel, raw	35	0.1	0.1	0.0	0.6	FDA	2016-2020
59	09020	84	Applesauce, bottled	10	0.1	0.1	0.0	0.4	FDA	2016-2020
324	09026	348	Apricots, canned in heavy/light syrup	7	6.2	5.0	1.3	15	FDA	2016-2017
67	09037	97	Avocado, raw	35	0.2	0.7	0.0	3.4	FDA	2016-2020
56	09040	80	Banana, raw	35	0.2	0.6	0.0	3.5	FDA	2016-2020
196	09050	398	Blueberries, raw	28	0.3	0.4	0.0	1.2	FDA	2017-2020
64	09181	89	Cantaloupe, raw/frozen	35	0.3	0.3	0.0	1.3	FDA	2016-2020
62	09099	87	Fruit cocktail, canned in light syrup	10	3.9	3.1	0.0	10	FDA	2016-2020
65	09111	92	Grapefruit, raw	35	0.1	0.2	0.0	0.7	FDA	2016-2020
63	09132	88	Grapes, seedless, red/green, raw	35	0.4	0.3	0.0	1.1	FDA	2016-2020
453	09176	412	Mango, raw/frozen	6	0.1	0.2	0.0	0.4	FDA	2020
153	09193	299	Olives, black, pitted	10	2.1	1.6	0.0	5.2	FDA	2016-2020
55	09200	79	Orange, raw	35	0.2	0.2	0.0	1.0	FDA	2016-2020
345	09239	254	Peach, canned in light/medium syrup	7	6.2	3.6	1.3	10	FDA	2016-2017
58	09236	83	Peach, raw/frozen	35	0.2	0.4	0.0	2.6	FDA	2016-2020
346	09256	255	Pear, canned in light syrup	7	0.1	0.0	0.1	0.2	FDA	2016-2017

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
60	09252	85	Pear, with peel, raw	35	0.1	0.2	0.0	0.7	FDA	2016-2020
309	09354	93	Pineapple, canned in juice	7	1.8	2.3	0.6	7.1	FDA	2016-2017
194	09266	396	Pineapple, raw/frozen	28	0.3	0.4	0.0	2.3	FDA	2017-2020
66	09298	95	Raisins	10	1.7	1.3	0.0	3.5	FDA	2016-2020
61	09316	86	Strawberries, raw/frozen	35	0.3	0.3	0.0	1.6	FDA	2016-2020
57	09326	81	Watermelon, raw/frozen	35	0.1	0.2	0.0	0.8	FDA	2016-2020
Vegetables										
74	11012	115	Asparagus, fresh/frozen, boiled	35	0.9	0.7	0.0	3.9	FDA	2016-2020
340	11084	131	Beets, canned	7	0.4	0.4	0.1	1.2	FDA	2016-2017
72	11742	113	Broccoli, fresh/frozen, boiled	35	0.5	0.6	0.0	2.7	FDA	2016-2020
140	11099	263	Brussels sprouts, fresh/frozen, boiled	35	0.0	0.1	0.0	0.5	FDA	2016-2020
331	11110	110	Cabbage, fresh, boiled	8	0.1	0.1	0.0	0.3	FDA	2016-2017
204	11109	406	Cabbage, raw	27	0.0	0.0	0.0	0.0	FDA	2018-2020
175	11960	356	Carrot, baby, raw	35	0.6	1.1	0.0	3.6	FDA	2016-2020
313	11125	259	Carrot, fresh, peeled, boiled	7	0.5	0.6	0.0	1.6	FDA	2016-2017
75	11136	116	Cauliflower, fresh/frozen, boiled	35	0.6	0.7	0.0	3.0	FDA	2016-2020
73	11143	114	Celery, raw	35	1.7	2.1	0.0	8.9	FDA	2016-2020
70	11162	108	Collards, fresh/frozen, boiled	35	0.8	1.8	0.0	7.4	FDA	2016-2020
45	11172	55	Corn, canned, drained solids	10	0.3	0.3	0.0	0.9	FDA	2016-2020
44	11179	54	Corn, frozen, boiled	35	0.2	1.2	0.0	7.0	FDA	2016-2020
79	11206	123	Cucumber, peeled, raw	35	0.5	0.4	0.0	2.1	FDA	2016-2020
315	11210	265	Eggplant, fresh, without peel, boiled	14	0.2	0.3	0.0	1.0	FDA	2016-2018
250		410	Eggplant, with peel, baked	21	0.6	0.5	0.0	1.4	FDA	2018-2020
454	11215	413	Garlic, raw	6	1.0	1.5	0.0	3.1	FDA	2020
78	11729	122	Green beans, canned, drained solids	10	0.3	0.5	0.0	1.2	FDA	2016-2020
77	11061	121	Green beans, fresh/frozen, boiled	35	0.6	0.7	0.0	2.6	FDA	2016-2020
452	11236	411	Kale, fresh, pan-cooked	6	1.8	2.1	0.0	4.4	FDA	2020
71	11252	109	Lettuce, iceberg, raw	35	0.2	0.4	0.0	1.8	FDA	2016-2020

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
176	11253	357	Lettuce, leaf, raw	35	1.1	1.4	0.0	8.0	FDA	2016-2020
36	11032	42	Lima beans, immature, frozen, boiled	9	0.2	0.4	0.0	0.9	FDA	2016-2019
142	11584	268	Mixed vegetables, frozen, boiled	10	0.2	0.3	0.0	0.9	FDA	2016-2020
141	11260	264	Mushrooms, raw	35	0.3	0.2	0.0	1.1	FDA	2016-2020
451		130	Mushrooms, canned, drained solids	1	0.0				FDA	2020
434			Nori, seaweed, dried	3	2317	618	1640	2850	USDA	2018
317	11279	267	Okra, fresh/frozen, boiled	7	1.0	0.6	0.3	1.8	FDA	2016-2017
82	11282	128	Onion, mature, raw	35	0.3	0.4	0.0	1.6	FDA	2016-2020
37	11313	46	Peas, green, fresh/frozen, boiled	10	0.1	0.2	0.0	0.5	FDA	2016-2020
80	11333	125	Pepper, bell, green, raw	35	0.2	0.5	0.0	2.7	FDA	2016-2020
89	11937	161	Pickles, dill, cucumber	10	0.7	0.5	0.3	1.7	FDA	2016-2020
83	11367	136	Potato, peeled, boiled ¹	34	0.0	0.1	0.0	0.6	FDA	2016-2020
84	11674	137	Potato, with peel, baked	35	0.5	1.4	0.0	7.0	FDA	2016-2020
293	11371	354	Potatoes, mashed, prepared from fresh	7	6.2	3.6	3.8	14	FDA	2016-2017
220	11549	119/51 9	Sauce, tomato, canned/bottled	10	0.9	0.5	0.0	1.5	FDA	2016-2020
1	11458	107	Spinach, fresh/frozen, boiled	8	3.9	2.9	1.0	8.1	FDA	2016-2017
206	11457	408	Spinach, raw	27	6.7	7.3	1.6	40	FDA	2018-2020
332	11468	124	Squash, summer, fresh/frozen, boiled	8	0.3	0.1	0.1	0.5	FDA	2016-2017
205	11478	407	Squash, summer, zucchini, fresh/frozen, boiled	27	0.6	1.0	0.0	4.6	FDA	2018-2020
81	11644	126	Squash, winter (Hubbard or acorn), fresh/frozen, boiled	35	0.8	1.1	0.0	5.3	FDA	2016-2020
195	11508	397	Sweet potato, baked, peel removed	28	1.5	1.5	0.0	6.2	FDA	2017-2020
325	11514	358	Sweet potato, canned	7	0.7	0.8	0.0	2.3	FDA	2016-2017
76	11529	117	Tomato, raw	35	0.2	0.3	0.0	1.2	FDA	2016-2020
316	11567	266	Turnip, fresh/frozen, boiled	7	0.3	0.2	0.0	0.5	FDA	2016-2017
Legumes										
270		527	Beans, black, canned, drained	3	0.0	0.0	0.0	0.0	FDA	2018-2020

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
456		545	Beans, garbanzo (chickpeas), canned, drained solids	1	0.0				FDA	2020
271		528	Beans, kidney, canned, drained	3	0.0	0.0	0.0	0.0	FDA	2018-2020
228	16146	529	Beans, pinto, canned, drained	3	0.9	1.6	0.0	2.8	FDA	2018-2020
304	16043	38	Beans, pinto, from dry, boiled	7	0.0	0.1	0.0	0.2	FDA	2016-2017
167	16103	341	Beans, refried, canned	10	0.8	0.8	0.0	2.5	FDA	2016-2020
272		530	Beans, white, canned, drained	3	0.9	1.5	0.0	2.6	FDA	2018-2020
323	16050	342	Beans, white, from dry, boiled	7	0.1	0.2	0.0	0.4	FDA	2016-2017
35	16009	39	Beans, with pork, canned	10	0.2	0.2	0.0	0.5	FDA	2016-2020
457		546	Lentils, dry, cooked	1	0.0				FDA	2020
38	16167	47	Peanut butter, smooth/creamy	10	0.4	0.4	0.0	1.2	FDA	2016-2020
39	16090	48	Peanuts, dry roasted, salted	10	0.5	0.5	0.0	1.1	FDA	2016-2020
458	16126	547	Tofu, firm, plain, drained solids	1	0.0				FDA	2020
18	43137	544	Veggie burger, soy based, baked	3	9.5	1.6	7.9	11	FDA	2018-2020
Nuts and Seeds										
227	12061	526	Almonds, shelled	3	0.0	0.0	0.0	0.0	FDA	2018-2020
455	12586	414	Nuts, cashew, oil roasted, with salt added	1	0.0				FDA	2020
168	12537	343	Seeds, sunflower, shelled, salted, roasted	10	0.1	0.3	0.0	0.9	FDA	2016-2020
201	12155	403	Walnuts, shelled	23	0.0	0.0	0.0	0.0	FDA	2017-2020
Beverages including Juices										
110	14003	198	Alcohol, beer	10	0.9	0.5	0.2	2.1	FDA	2016-2020
207	14051	500	Alcohol, distilled, vodka	3	0.0	0.0	0.0	0.0	FDA	2018-2020
248		200	Alcohol, distilled, whiskey/scotch	3	0.0	0.0	0.0	0.0	FDA	2018-2020
342	14084	199/501/502	Alcohol, wine, red/white	13	1.6	0.7	0.7	3.3	FDA	2016-2020
210	14091	503	Beverage, almond	9	0.4	0.3	0.1	0.9	FDA/USDA	2020-2022
211	14090	504	Beverage, coconut water	3	0.6	0.2	0.4	0.8	FDA	2018-2020
212	14060	505	Beverage, energy	3	0.1	0.2	0.0	0.4	FDA	2018-2020

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
213	16222	506	Beverage, soy, shelf stable	6	1.3	1.2	0.3	3	FDA/USDA	2019-2022
445			Beverage, soy, shelf stable (with seaweed or seaweed derivative)	2	4.9	2.9	2.9	6.9	FDA/USDA	2018,2022
264		507	Beverage, sports	3	0.0	0.0	0.0	0.0	FDA	2018-2020
374	14164		Beverage, chocolate malt powder, prepared with 1% milk, fortified	1	28				USDA	2014
108	14416	194	Carbonated beverage, cola, low-calorie	10	0.3	0.2	0.0	0.6	FDA	2016-2020
106	14148	191	Carbonated beverage, cola, regular	10	0.2	0.2	0.0	0.5	FDA	2016-2020
320	14144	306/508	Carbonated beverage, fruit-flavored	10	0.1	0.2	0.0	0.5	FDA	2016-2020
155	14209	305	Coffee, brewed from ground	10	0.1	0.1	0.0	0.3	FDA	2016-2020
13	14201	381	Coffee, decaffeinated, from ground	7	0.2	0.1	0.1	0.4	FDA	2016-2017
173	14242	351	Cranberry juice cocktail, canned/bottled	10	0.2	0.3	0.0	0.8	FDA	2016-2020
156	14651	307	Fruit drink (5% - 25% juice), canned/bottled	10	0.7	0.3	0.3	1.2	FDA	2016-2020
107	14541	193	Fruit drink from powder, reconstituted	10	0.1	0.2	0.0	0.5	FDA	2016-2020
68	09400	99	Juice, apple, bottled	10	0.4	0.6	0.0	1.9	FDA	2016-2020
172	09444	350	Juice, fruit blend (100% juice), canned/bottled	10	0.8	0.3	0.4	1.4	FDA	2016-2020
138	09130	257	Juice, grape, bottled	10	0.3	0.2	0.1	0.8	FDA	2016-2020
69	09123	100	Juice, grapefruit, bottle/carton	10	5.1	14.4	0.0	46	FDA	2016-2020
225	09152	524	Juice, lemon, bottled	3	1.5	1.6	0.0	3.2	FDA	2018-2020
174	09207	352	Juice, orange, bottle/carton	10	8.5	7.9	1.6	24	FDA	2016-2020
338	09209	98	Juice, orange, from frozen concentrate, reconstituted	7	0.4	0.2	0.2	0.7	FDA	2016-2017
137	09409	256	Juice, pineapple, canned	10	0.2	0.2	0.0	0.6	FDA	2016-2020
339	09294	103	Juice, prune, bottled	7	1.3	1.3	0.4	3.2	FDA	2016-2017
314	11886	261	Juice, tomato, bottled	7	1.1	0.3	0.7	1.4	FDA	2016-2017
226	11578	525	Juice, tomato-vegetable, bottled	3	0.8	0.5	0.4	1.3	FDA	2018-2020
310	14293	105	Lemonade, from frozen concentrate, reconstituted	7	0.1	0.1	0.0	0.2	FDA	2016-2017

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
322		331, 509	Meal replacement, liquid, ready to drink (RTD), assorted flavors	10	39	41	18	143	FDA	2016-2020
215	14066	510	Powder, protein	3	98	46	67	150	FDA	2018-2020
109	14355	197	Tea, brewed from tea bag	10	0.1	0.1	0.0	0.4	FDA	2016-2020
14	14352	382	Tea, decaffeinated, brewed from tea bag	7	0.1	0.1	0.0	0.2	FDA	2016-2017
184	14555	380	Water, mineral/spring, bottled	35	0.2	0.3	0.0	1.3	FDA	2016-2020
439	14411		Water, tap, drinking ²	40	1.1	2.0	0.0	11	USDA	2003
Seafood										
389	15141		Crustaceans, crab, blue, canned	3	38	12	27	52	USDA	2008
390	15148		Crustaceans, lobster, northern, cooked, moist heat	3	185	44	139	226	USDA	2008
391	15149		Crustaceans, shrimp, mixed species, raw	9	16	9.7	10	39	USDA	2008,2016
132	15151	244	Crustaceans, shrimp, precooked, shell removed, no tail	39	15	10	3.4	55	FDA/USDA	2008,2016-2020
33	15027	34	Fish sticks or patty, frozen, oven-cooked	14	67	28	25	134	FDA/USDA	2012-2013, 2016-2020
165	15235	339	Fish, catfish, pan-cooked with oil	35	3.2	2.5	0.0	8.1	FDA	2016-2020
186	15192	387	Fish, cod, baked	28	172	63	88	330	FDA	2017-2020
375	15019		Fish, cod, Pacific, raw	7	131	67	51	241	USDA	2007-2008
376	15028		Fish, flatfish (flounder and sole species), raw	4	14	3.9	11	20	USDA	2008
377	15033		Fish, haddock, raw	3	227	88	138	314	USDA	2008
378	15036		Fish, halibut, Atlantic and Pacific, raw	5	10	0.1	10.0	10	USDA	2003,2008
379	15057		Fish, ocean perch, Atlantic, raw	4	11	0.7	10.0	12	USDA	2008
396	15266		Fish, pollock, Alaska, raw	3	44	8.7	37	54	USDA	2008
380	15070		Fish, rockfish, Pacific, mixed species, raw	3	14	4.8	11	20	USDA	2008
421	35171		Fish, salmon, chum, dried (Alaska Native)	1	38				USDA	2004
395	15260		Fish, salmon, pink, canned, drained solids	2	15	7.4	10	21	USDA	2008,2010
381	15083		Fish, salmon, pink, raw	5	26	15.7	12	50	USDA	2007-2008

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
384	15087		Fish, salmon, sockeye, canned, drained solids	1	22				USDA	2010
382	15085		Fish, salmon, sockeye, raw	2	14	3.8	12	17	USDA	2008
158	15237	318	Fish, salmon, steaks/fillets, baked	35	13	7.1	0.0	29	FDA	2016-2020
422	35184		Fish, smelt, dried (Alaska Native)	1	216				USDA	2004
371			Fish, swai, cooked	2	<10				USDA	2017
385	15110		Fish, swordfish, raw	4	19	3.8	16	25	USDA	2008
187	15262	388	Fish, tilapia, baked	28	6.9	15	0.0	81	FDA	2017-2020
386	15115		Fish, trout, rainbow, wild, raw	1	<10				USDA	2008
166	15121	340	Fish, tuna, canned in water, drained	15	9.4	1.6	6.6	12	FDA/USDA	2008,2012,2016-2020
388	15118		Fish, tuna, fresh, bluefin, cooked, dry heat	4	23	4.4	18	28	USDA	2008
387	15117		Fish, tuna, fresh, bluefin, raw	4	18	5.9	12	26	USDA	2008
420	35091		Fish, whitefish, broad, including head, eyes, cheeks and soft bones (Alaska Native), raw	1	34				USDA	2005
392	15160		Mollusks, clam, mixed species, canned, drained solids	4	66	9.1	58	78	USDA	2007
393	15169		Mollusks, oyster, eastern, wild, cooked, moist heat	3	109	26	81	132	USDA	2008
394	15172		Mollusks, scallop, mixed species, raw	4	<10				USDA	2008
Grains³										
16	20081	900	Flour, white, all-purpose	3	0.8	1.4	0.0	2.5	FDA	2018-2020
53	20110	69	Noodles, egg, enriched, boiled	10	3.7	1.2	1.7	5.4	FDA	2016-2020
347	20321		Pasta, enriched, boiled in water with iodized salt using amount per pasta label	4	22	6.6	13	28	USDA	2017
232	20134	539	Pasta, rice noodles, cooked	3	0.9	1.5	0.0	2.6	FDA	2018-2020
171	20121	347	Pasta, spaghetti, enriched, boiled	10	0.3	0.9	0.0	2.8	FDA	2016-2020
233	20125	540	Pasta, whole wheat, cooked	3	0.0	0.0	0.0	0.0	FDA	2018-2020
223	20137	522	Quinoa, cooked	3	0.0	0.0	0.0	0.0	FDA	2018-2020

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
200	20041	402	Rice, brown, cooked	28	0.0	0.0	0.0	0.0	FDA	2017-2020
40	20051	50	Rice, white, enriched, cooked	34	0.2	1.1	0.0	6.3	FDA	2016-2020
Cereals										
254		74	Cereal, bran with raisins	10	0.7	0.6	0.0	1.7	FDA	2016-2020
253		71	Cereal, corn flakes	10	0.2	0.4	0.0	1.2	FDA	2016-2020
43	08091	53	Cereal, corn/hominy grits, enriched, cooked	10	0.0	0.1	0.0	0.2	FDA	2016-2020
42	08169	52	Cereal, cream of wheat (farina), enriched, cooked	10	0.0	0.1	0.0	0.2	FDA	2016-2020
255		75	Cereal, crisped rice	10	0.3	0.3	0.0	0.8	FDA	2016-2020
306		72	Cereal, fruit-flavored, sweetened ⁴	6	0.5	0.3	0.2	0.9	FDA	2016-2017
266		512	Cereal, granola	3	11	2.4	8.3	13	FDA	2018-2020
308		76	Cereal, granola with raisins	7	14	4.0	9.4	20	FDA	2016-2017
256		77	Cereal, oat ring	10	0.7	0.5	0.0	1.2	FDA	2016-2020
267		513	Cereal, oat ring, honey	3	0.0	0.0	0.0	0.0	FDA	2018-2020
41	08121	51	Cereal, oatmeal, plain, cooked	10	0.2	0.4	0.0	1.1	FDA	2016-2020
307		73	Cereal, shredded wheat	7	0.1	0.2	0.0	0.6	FDA	2016-2017
268		514	Cereal, shredded wheat, frosted	3	0.0	0.0	0.0	0.0	FDA	2018-2020
269		515	Cereal, whole wheat, cooked	3	0.0	0.0	0.0	0.0	FDA	2018-2020
Baked Products										
133	18001	249	Bagel, plain, toasted	35	0.6	1.7	0.0	9.3	FDA	2016-2020
305	18015	61	Biscuits, refrigerated-type, baked	8	3.5	3.0	0.7	10	FDA/USDA	2013,2016-2017
344	28397	248	Bread, multigrain ⁵	4	1.8	2.0	0.0	4.4	FDA	2016-2017
337	18060	64	Bread, rye	7	0.6	0.3	0.0	0.9	FDA	2016-2017
199	18350	401	Bread, white roll/bun (hamburger/hotdog) ⁶	16	2.4	3.3	0.0	10	FDA/USDA	2017-2020
442	18350*	401*	Bread, white roll/bun (hamburger/hotdog), with iodate dough conditioner	16	1060	342	480	1730	FDA/USDA	2017-2020

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
46	18069	58	Bread, white, enriched, pre-sliced ⁷	15	1.8	2.7	0.0	9.5	FDA/USDA	2011,2013,2016-2020
440	18069*	58*	Bread, white, enriched, pre-sliced, with iodate dough conditioner	19	592	198	350	1178	FDA/USDA	2013,2016-2020
48	18075	62	Bread, whole-wheat, commercially prepared ⁸	21	1.9	2.5	0.0	9.5	FDA/USDA	2013,2016-2020
441	18075*	62*	Bread, whole-wheat, commercially prepared, with iodate dough conditioner	8	546	214	340	1010	FDA/USDA	2016,2018-2019
15	18079	903	Breadcrumbs	3	112	68	45	180	FDA	2018-2020
170	18362	345	Breakfast tart/toaster pastry ⁹	9	2.3	2.2	0.0	5.4	FDA	2016-2020
146	18151	291	Brownie, commercially prepared	10	9.7	12	0.9	42	FDA	2016-2020
99	18096	178	Cake, chocolate with chocolate icing, commercially prepared ¹⁰	34	8.1	3	3.1	17	FDA	2016-2020
263		369	Cake, white with white icing, commercially prepared ¹¹	33	13	15	2.7	72	FDA	2016-2020
399	18140		Cake, yellow with chocolate frosting, commercially prepared	3	<10				USDA	2011
185	18964	386	Cinnamon roll, from package, iced	30	3.8	3.3	0.0	12	FDA	2017-2020
100	18160	183	Cookies, from package, chocolate chip	11	1.9	2.9	0.0	10	FDA/USDA	2012,2016-2020
101	18166	184	Cookies, from package, sandwich with crème filling	10	0.3	0.4	0.0	1.1	FDA	2016-2020
147	18204	292	Cookies, from package, sugar ¹²	8	4.7	2.7	0.0	8.9	FDA	2016-2020
47	18024	60	Cornbread, homemade	10	27	12.2	12	49	FDA	2016-2020
398	18023		Cornbread, from dry mix, prepared with 2% milk, margarine, and eggs	1	17				USDA	2012
136	18229	252	Crackers, butter-type	10	0.3	0.4	0.0	1.1	FDA	2016-2020
229	18214	531	Crackers, cheese, regular	4	11	5.5	4.7	18	FDA/USDA	2009,2018-2020
135	18173	251	Crackers, graham	10	0.9	0.8	0.0	2.6	FDA	2016-2020
51	18228	66	Crackers, saltine	10	0.3	0.4	0.0	0.8	FDA	2016-2020

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
145	18248	290	Doughnut, cake-type, plain, commercially prepared	10	15	15	5.1	41	FDA	2016-2020
134	18437	250	English muffin, plain, toasted	35	1.1	2.8	0.0	15	FDA	2016-2020
50	18274	65	Muffin, blueberry, commercially prepared	35	12	4.8	5.3	35	FDA	2016-2020
169	18936	344	Pancakes, frozen, heated	12	7.1	2.2	3.2	10	FDA/USDA	2013,2016-2020
234	18335	541	Pie crust, commercially prepared	3	1.2	2.0	0.0	3.5	FDA	2018-2020
102	18443	185	Pie, apple, fresh/frozen, commercially prepared	10	0.4	0.5	0.0	1.2	FDA	2016-2020
103	18326	186	Pie, pumpkin, fresh/frozen, commercially prepared	10	22	3.4	17.5	30	FDA	2016-2020
3	18244	182	Sweet roll/Danish pastry	4	6.0	1.4	4.4	7.7	FDA	2016
198	18363	400	Tortilla, corn	28	0.3	0.9	0.0	3.6	FDA	2017-2020
49	18970	63	Tortilla, flour	34	0.4	1.4	0.0	7.5	FDA	2016-2020
Sweets and Snacks										
216	19919	511	Candy, fruit snacks	3	1.6	2.8	0.0	4.9	FDA	2018-2020
178	19155	371	Candy bar (chocolate, nougat, and nuts)	10	17	2.9	14	22	FDA	2016-2020
104	19120	187	Candy bar, milk chocolate, plain	10	46	5.9	35	54	FDA	2016-2020
148	19107	293	Candy, hard, assorted flavors	10	7.8	12.6	0.0	35	FDA	2016-2020
85	19411	138	Chips, potato	10	2.3	3.6	0.0	9.1	FDA	2016-2020
402	19412		Chips, potato, made from dried potatoes, cheese-flavor	1	<10				USDA	2012
52	25028	67	Chips, corn/tortilla	10	1.8	1.6	0.1	4.5	FDA	2016-2020
105	19173	190	Gelatin dessert, assorted flavors, prepared	10	0.5	0.4	0.1	1.4	FDA	2016-2020
224	19015	523	Granola bar	3	1.9	1.6	0.0	3.0	FDA	2018-2020
327		370	Granola bar, with raisins	7	2.0	1.6	0.6	4.6	FDA	2016-2017
97	19296	172	Honey	10	0.7	0.4	0.3	1.7	FDA	2016-2020
151	19300	296	Jelly, assorted flavors	10	1.0	0.6	0.2	2.1	FDA	2016-2020

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
144	19283	288	Popsicle, fruit-flavored ¹³	3	1.1	1.2	0.3	2.5	FDA	2016-2020
179	25026	372	Popcorn, microwave, butter-flavored	10	0.4	0.8	0.0	2.5	FDA	2016-2020
149	19047	294	Pretzels, hard, salted	10	0.5	0.4	0.0	1.2	FDA	2016-2020
235	19183	542	Pudding, ready-to-eat, chocolate	3	14	1.2	13	15	FDA	2018-2020
326		368	Pudding, ready-to-eat, assorted flavors other than chocolate	7	4.7	1.9	3.0	7.6	FDA	2016-2017
401	19193		Pudding, rice, ready-to-eat	1	28				USDA	2012
289	19097	287	Sherbet, fruit-flavored ¹⁴	3	14	0.7	14	15	FDA	2016
330		385	Sorbet, fruit-flavored	6	0.8	0.6	0.0	1.8	FDA	2017-2020
95	19335	169	Sugar, white, granulated	10	0.0	0.1	0.0	0.3	FDA	2016-2020
150	19348	295	Syrup, chocolate	10	0.7	0.6	0.3	2.0	FDA	2016-2020
96	19129	170	Syrup, pancake	10	0.2	0.1	0.0	0.4	FDA	2016-2020
Mixed Dishes										
284		269	Beef stroganoff with noodles, homemade	7	6.2	1.0	5.3	8.1	FDA	2016-2017
88	22906	152	Chicken potpie, frozen, heated	10	3.6	2.1	1.9	9.0	FDA	2016-2020
86	22904	145	Chili con carne with beans, canned	10	1.7	1.0	0.0	2.8	FDA	2016-2020
294	21127	355	Coleslaw, mayonnaise-type, from grocery/deli	7	1.8	0.8	1.0	2.8	FDA	2016-2017
411	22973		Corn dogs, frozen, prepared	1	24				USDA	2011
6	22977	361	Lasagna with meat, frozen, heated	7	8.4	1.1	7.0	9.8	FDA	2016-2017
418	32019		Lasagna, cheese, frozen, unprepared	5	12	2.2	10	15	USDA	2012-2013
87	22960	146	Macaroni and cheese, prepared from box mix	17	19	4.5	11	29	FDA/USDA	2012,2016-2020
291		346	Macaroni salad, from grocery/deli	7	3.2	1.2	1.9	5.3	FDA	2016-2017
245		148	Meatloaf, beef, homemade	7	16	11	8.1	34	FDA	2016-2017
416	32008		Pasta mix, classic cheeseburger macaroni, unprepared	1	<10				USDA	2013
417	32010		Pasta mix, Italian lasagna, unprepared	2	13	3.6	10	15	USDA	2013

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
409	21505		Pizza, cheese topping, thin crust, frozen, baked	7	23	5.0	16	29	USDA	2017
410	22971		Potato salad with egg	5	<10				USDA	2010,2017
292	21140	353	Potato salad, mayonnaise-type, from grocery/deli	7	2.6	1.7	0.0	5.0	FDA	2016-2017
244		142	Spaghetti with meat sauce, homemade	7	1.8	0.6	1.0	2.6	FDA	2016-2017
438			Sushi, California roll	3	31	9.5	20	37	USDA	2009
415	32007		Taquitos, frozen, beef and cheese, oven-heated	1	<10				USDA	2012
414	32006		Taquitos, frozen, chicken and cheese, oven-heated	1	<10				USDA	2012
285		272	Tuna noodle casserole, homemade	7	16	9.3	11	36	FDA	2016-2017
419	32037		Turnover, pepperoni pizza	1	21				USDA	2017
Fast Foods										
197	21142	399	Biscuits, fast food	32	9.1	2.4	6.0	17	FDA/USDA	2017-2020
10	21064	365	Burrito with beef, beans and cheese, from Mexican carry-out	7	5.4	2.2	2.7	9.8	FDA	2016-2017
162	21469	336	Chicken breast, fried, with skin, fast food ¹⁵	33	3.8	2.1	0.0	10	FDA	2016-2020
164	21470	338	Chicken leg, fried, with skin, fast food	35	2.3	1.7	0.0	7.7	FDA	2016-2020
131	21229	241	Chicken nuggets, fast food	35	0.9	1.6	0.0	7.3	FDA	2016-2020
303	01110	7	Milk shake, chocolate, fast food	7	35	9.4	23	45	FDA	2016-2017
193	01111	394	Milk shake, vanilla, fast food	28	36	5.8	22	47	FDA	2017-2020
318	21302	281	Pizza, cheese and pepperoni, regular crust, from pizza carry-out	11	12	4.2	7.1	20	FDA/USDA	2010,2016-2017
202	21299	404	Pizza, cheese, fast food	30	12	2.9	5.7	17	FDA/USDA	2010,2017-2020
435			Pizza, crust only, fast food	9	<10				USDA	2017
408	21484		Pizza, with sausage topping, regular crust, fast food	1	13				USDA	2010
139	21138	258	Potatoes, French fries, fast food	35	1.1	2.1	0.0	10	FDA	2016-2020

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
2	21112	147	Sandwich, hamburger, single, large patty, fast food	7	3.3	1.4	2.0	5.9	FDA	2016-2017
406	21092		Sandwich, cheeseburger, double, regular patty, plain, fast food	8	12	1.9	10	15	USDA	2017
443	21092*		Sandwich, cheeseburger, double, regular patty, fast food (bun w iodate dough conditioner)	4	279	96	164	392	USDA	2017
286	21096	275	Sandwich, cheeseburger, single, large patty, fast food	9	11	1.9	8.6	15	FDA/USDA	2013,2016-2017
11	21490	366	Sandwich, chicken filet (broiled) sandwich, fast food	7	1.8	1.6	0.4	4.5	FDA	2016-2017
288	21021	278	Sandwich, egg, cheese, and ham on English muffin, fast food	8	24	5.9	16	32	FDA/USDA	2012,2016-2017
287	21105	276	Sandwich, fish, fast food	10	36	12	23	60	FDA/USDA	2013,2016-2017
405	21059		Shrimp, breaded and fried, fast food	2	<10				USDA	2010
4	21486	279	Taco/tostada with beef and cheese, from Mexican carry-out	7	7.0	2.7	4.4	12	FDA	2016-2017
Restaurant Foods (excluding Fast Food Restaurants)										
7	36603	362	Beef with vegetables in sauce, restaurant, Chinese	7	4.3	5.2	0.6	16	FDA	2016-2017
428	36050		Cheese enchilada, restaurant, Mexican	1	16				USDA	2013
429	36052		Cheese quesadilla, restaurant, Mexican	1	23				USDA	2013
430	36055		Cheese ravioli with marinara sauce, restaurant, Italian	1	22				USDA	2013
8	36626	363	Chicken with vegetables in sauce, restaurant, Chinese	7	3.8	4.6	0.8	13	FDA	2016-2017
444	36033		Fish fillet, parmesan crusted tilapia, restaurant, family style	1	13				USDA	2013
436			Fish, salmon, grilled, restaurant	1	<10				USDA	2017
423	36012		Fried mozzarella sticks, restaurant, family style	3	38	4.1	34	42	USDA	2013

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
9	36602	364	Fried rice, meatless, restaurant, Chinese	7	5.5	5.6	1.2	17	FDA	2016-2017
427	36041		Lasagna with meat, restaurant, Italian	1	28				USDA	2013
425	36017		Macaroni & cheese, from kids' menu, restaurant, family style	10	22	7.4	13	32	USDA	2013,2017
431	36620		Shrimp and vegetables, restaurant, Chinese	1	<10				USDA	2009
Sausage and Luncheon Meats										
32	07959	29	Bologna (beef/pork)	10	25	9.7	13	44	FDA	2016-2020
31	07949	28	Frankfurter, beef/pork, boiled ¹⁶	28	8.8	12	0.0	60	FDA	2016-2020
369	07945		Frankfurter, beef, heated	2	<10				USDA	2010
261	07081	335/543	Luncheon meat (chicken/turkey)	10	3.5	1.7	0.0	5.4	FDA	2016-2020
129	07028	239	Luncheon meat, ham	10	1.2	0.9	0.0	3.0	FDA	2016-2020
188	07072	389	Salami, dry/hard	31	10	9.4	2.5	48	FDA/USDA	2011,2017-2020
243		30	Salami, luncheon-meat type (not hard)	7	12	8.0	3.5	23	FDA	2016-2017
367	07019		Sausage, pork, chorizo, raw	1	30				USDA	2017
28	07953	19	Sausage, pork, oven-cooked	35	4.2	2.9	0.0	11	FDA	2016-2020
Meats										
130	05064	240	Chicken breast, oven-roasted (skin removed)	35	1.2	1.9	0.0	8.1	FDA	2016-2020
163	05098	337	Chicken thigh, oven-roasted (skin removed)	35	0.9	1.8	0.0	9.5	FDA	2016-2020
30	05192	26	Turkey breast, oven-roasted	35	4.8	13.2	0.0	61	FDA	2016-2020
189	05306	390	Turkey, ground, pan-cooked	28	3.9	4.6	0.0	17	FDA	2017-2020
26	10136	17	Ham, cured (not canned), baked	35	0.7	1.4	0.0	5.9	FDA	2016-2020
29	10860	20	Pork bacon, oven-cooked	35	1.1	1.7	0.0	7.7	FDA	2016-2020
27	10179	18	Pork chop, pan-cooked with oil	35	0.5	1.3	0.0	6.8	FDA	2016-2020
333		21	Pork roast, loin, oven-roasted	7	0.4	0.3	0.0	0.9	FDA	2016-2017
335	17204	27	Beef/calf, liver, pan-cooked with oil	7	16	5.0	7.7	22	FDA	2016-2017

DB_ID	NDB No.	TDS No.	Description	n	Iodine mcg/100g	SD	Min	Max	Source(s)	Year(s)
251		13	Beef, ground, pan-cooked	35	7.5	3.0	3.3	19	FDA	2016-2020
334		14	Beef roast, chuck, oven-roasted	8	3.8	2.2	1.8	8.2	FDA	2016-2017
260		334	Beef steak, loin/sirloin, broiled	34	4.7	1.7	0.0	8.4	FDA	2016-2020
252		22	Lamb chop, pan-cooked with oil	35	2.5	2.7	0.0	12	FDA	2016-2020

*TDS and/or NDB identifiers with asterisks indicate products that differ from the generic product in that they contain iodate dough conditioners

¹Omitted sample with value of 64 mcg I/100g as an outlier

²USDA water samples analyzed by FDA

³Grain products prepared without salt unless otherwise noted

⁴Omitted sample with 386 mcg I/100g – value probably from FD&C Red No. 3 food coloring

⁵Omitted 3 samples with values from 72 – 224 mcg I/100g – probably mixture of breads with and without iodate dough conditioners

⁶Omitted 6 samples with values from 130 – 416 mcg I/100g – probably mixture of breads with and without iodate dough conditioners

⁷Omitted 11 samples with values from 38 – 310 mcg I/100g – probably mixture of breads with and without iodate dough conditioners

⁸Omitted 10 samples with values from 110 – 290 mcg I/100g – probably mixture of breads with and without iodate dough conditioners

⁹Omitted sample with 92.4 mcg I/100g – value probably from FD&C Red No. 3 food coloring

¹⁰Omitted sample with 120 mcg I/100g – value probably from FD&C Red No. 3 food coloring

¹¹Omitted 3 samples with 310 and 335 mcg I/100g – values probably from FD&C Red No. 3 food coloring

¹²Omitted 2 samples with 42 and 973 mcg I/100g – values probably from FD&C Red No. 3 food coloring

¹³Omitted 9 samples from 150 to 621 mcg I/100g – values probably from FD&C Red No. 3 food coloring

¹⁴Omitted sample with 545 mcg I/100g – value probably from FD&C Red No. 3 food coloring

¹⁵Omitted samples with values of 38 and 42 mcg I/100g – values may reflect the use of disinfectant during poultry cleaning, but we were unable to confirm this use

¹⁶Omitted sample with value of 1000 mcg I/100g as an outlier, although value was confirmed by the lab