

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/376681996>

A SURVEY OF PHYTOPARASITIC NEMATODES ON SOME PLANT CROPS IN NORTHERN EGYPT

Article in *Nematropica* · December 2023

CITATIONS

3

READS

69

4 authors, including:



Mihail Kantor

Pennsylvania State University

69 PUBLICATIONS 329 CITATIONS

[SEE PROFILE](#)



Alemayehu Habteweld

United States Department of Agriculture

28 PUBLICATIONS 146 CITATIONS

[SEE PROFILE](#)



Zafar Handoo

United States Department of Agriculture

249 PUBLICATIONS 3,105 CITATIONS

[SEE PROFILE](#)

RESEARCH NOTE

A SURVEY OF PHYTOPARASITIC NEMATODES ON SOME PLANT CROPS IN NORTHERN EGYPT

I. K. A. Ibrahim¹, M. Kantor², A. Habteweld³, and Z. A. Handoo³

¹Department of Plant Pathology, Faculty of Agriculture, Alexandria University, Alexandria, Egypt; ²Plant Pathology & Environmental Microbiology Department, The Pennsylvania State University, University Park, PA 16802, U.S.A; ³Mycology and Nematology Genetic Diversity & Biology Laboratory, USDA, ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705, U.S.A. *Corresponding author: zafar.handoo@usda.gov

ABSTRACT

Ibrahim, I. K. A, M. Kantor, A. Habteweld, and Z. A. Handoo. 2023. A survey of phytoparasitic nematodes on some plant crops in northern Egypt. *Nematropica* 53:104-109.

A nematode survey was conducted in Alexandria and El-Behera Governorates in northern Egypt during the 2019-2022 cropping seasons to study the occurrence, population density, host association and distribution of phytoparasitic nematodes associated with seven crops. A total of 604 soil and root samples containing mixed population of 21 genera and 25 species of phytoparasitic nematodes were analyzed. Root-knot nematodes (*Meloidogyne incognita* and *M. javanica*) with 36-62% frequency of occurrence, respectively, were the most frequently encountered nematodes. The citrus nematode (*Tylenchulus semipenetrans*) had 94% frequency of occurrence on citrus trees, while the golden potato cyst nematode (*Globodera rostochiensis*) had a high (54%) frequency of occurrence on potato in El-Behera governorate. The genera *Criconema* spp., *Hemicriconemoides* spp., *Longidorus* spp., *Merlinius* spp., *Mesocriconema* spp., *Paratylenchus* spp., *Psilenchus* spp., and *Rotylenchus* spp. were less common, with a low (8-12%) frequency of occurrence.

Key words: Northern Egypt, occurrence, phytoparasitic nematodes, survey, citrus

RESUMEN

Ibrahim, I. K. A, M. Kantor, A. Habteweld, and Z. A. Handoo. 2023. Un estudio sobre nematodos fitoparásitos en algunos cultivos en el norte de Egipto. *Nematropica* 53:104-109.

Se realizó un estudio de nematodos en Alexandria y El-Behera, en el norte de Egipto durante las estaciones de producción entre el 2019 a 2022 para estudiar la ocurrencia, la densidad poblacional, la asociación de hospederos y la distribución de nematodos fitoparásitos asociados con siete cultivos. Se analizaron un total de 604 muestras de suelo y raíces que contenían poblaciones mixtas de 21 géneros y 25 especies de nematodos fitoparásitos. Los nematodos agalladores (*Meloidogyne incognita* y *M. javanica*), con una frecuencia de ocurrencia del 36 y 62%, respectivamente, fueron los nematodos más frecuentemente encontrados. El nemátoro del cítrico (*Tylenchulus semipenetrans*) tuvo una frecuencia de ocurrencia del 94% en los árboles de cítricos que se muestrearon, mientras que el nematodo dorado de la papa (*Globodera rostochiensis*) tuvo una alta frecuencia de ocurrencia (54%) en papa en El-Behera. Los géneros *Criconema* spp., *Hemicriconemoides* spp., *Longidorus* spp., *Merlinius* spp., *Mesocriconema* spp., *Paratylenchus* spp., *Psilenchus* spp., y *Rotylenchus* spp. fueron los menos comunes, con niveles bajos (frecuencia de ocurrencia

entre 8-12%).

Palabras clave: Cultivos, norte de Egipto, ocurrencia, nematodos fitoparásitos, studio

Information concerning the occurrence and distribution of phytoparasitic nematodes in Egypt is very important since many nematodes, such as citrus (*Tylenchulus semipenetrans*), cyst (*Globodera* spp., *Heterodera* spp.), dagger (*Xiphinema* spp.), lance (*Hoplolaimus* spp.), lesion (*Pratylenchus* spp.), root-knot (*Meloidogyne* spp.), reniform (*Rotylenchulus reniformis*), ring (*Mesocriconema* spp.), spiral (*Helicotylenchus* spp.) and stunt (*Tylenchorhynchus* spp.) may occur at high densities and cause economic damage to many crop plants (Oteifa, 1962; Tarjan, 1964; Oteifa and Tarjan, 1965; Ibrahim, 1990; Ismail and Eissa, 1993; Lamberti et al., 1996; Oteifa et al., 1997; Ibrahim et al., 2000, 2010; Ibrahim and Handoo, 2007; Adam et al., 2013; Handoo et al., 2015). The present nematode survey was carried out in Alexandria and El-Behera governorates in north Egypt, which has a Mediterranean climate with rainy cool winters and dry hot summers that favors and encourages the cultivation of several economically important agricultural crops. The leading agricultural crops in this region are wheat, corn, rice, clover, potato, tomato, sugar beet, guava, banana, orange, grape, and date palm. Certain important crops from this region, like potatoes and oranges, are underperforming, with low yields and poor development due to high infestations with phytoparasitic nematodes.

Recent studies in Egypt have shown the presence of about 60 genera and 170 species of phytoparasitic nematodes associated with many cultivated plants, grasses, and weeds (Ibrahim et al., 2010, 2023; Adam et al., 2013; Ibrahim and Handoo, 2016). The objective of this study was to identify phytoparasitic nematodes associated with certain crops in northern Egypt and provide additional information on the occurrence and distribution of genera and species of phytoparasitic nematodes that have the potential to significantly impact agricultural production in Egypt.

From 2019-2022, 604 samples were collected from a variety of cultivated plants from different localities in Alexandria and El-Behera governorates in northern Egypt (Fig. 1). From Alexandria governorate, 86 soil samples were collected from the ornamental shrub lantana

(*Lantana camara* L.) and 84 samples from tomato (*Lycopersicon esculentum* Mill.). From El-Behera governorate, 92, 84, 56, 86, and 116 soil samples were collected from citrus (*Citrus aurantium* L.), guava (*Psidium guajava* L.), potato (*Solanum tuberosum* L.), spinach (*Spinacia oleracea* L.), and sugar beet (*Beta vulgaris* L.), respectively. Soil and root samples were collected from the rhizosphere region of the surveyed host plants at a depth of 15-40 cm below the soil surface. Soil samples were collected using a standard (2.5 cm diam., 30 cm depth) soil probe. The soil types in Alexandria are sandy clay and in El-Behera are silt, clay, sand, or calcareous soil. Root samples were washed free of soil and examined for cyst and root-knot nematode infections. Root-knot nematodes were isolated from galled roots and identified by examination of perineal patterns of adult females as well as the characters of the second-stage juveniles (Taylor and Sasser, 1987; Hunt and Handoo, 2009). Cysts and females of cyst nematodes were extracted from sugar beet soil and root samples and identified by morphological characteristics (Mulvey and Golden, 1983; Golden, 1986). Nematodes from a composite sample of 250 cm³ soil were extracted using Cobb's wet-sieving and centrifugal sugar flotation techniques (Ayoub, 1980; Ibrahim et al., 2000). Nematodes were fixed in 3% formaldehyde solution then identified to genus and counted using a binocular stereomicroscope (Goodey, 1963; Mai and Lyon, 1975). Some nematode specimens were processed in 5% formaldehyde solution followed by anhydrous glycerin (Seinhorst, 1959) and examined under a compound microscope for species identification by morphological analysis using taxonomic keys (Sher, 1966; Golden, 1986; Handoo and Golden, 1989; Raski, 1991; Handoo, 2000; Handoo et al., 2007; Geraert, 2008; Hunt and Handoo, 2009). Frequency of occurrence (FO = [number of positive samples/numbers of total samples] x 100) and population densities (PD) were determined for the identified nematodes in composite samples and recorded (Table 1). Some nematode samples were fixed in 2% formaldehyde solution and sent to the Mycology and Nematology Genetic Diversity & Biology Laboratory, USDA (MNGDBL) in Beltsville, Maryland, USA for



Figure 1. Map of Egypt governorates. Soil samples were collected from Alexandria and El-Behra governorates.

species identification using compound microscopes.

The FO and PD of phytoparasitic nematode genera and species found in the rhizosphere soil samples collected from the surveyed crop plants in Alexandria and El-Behra governorates in northern Egypt are shown in Table 1. A total of 21 genera and 25 species of phytoparasitic nematodes were detected in the soil samples of the surveyed plants. Root-knot nematodes (*Meloidogyne incognita* and *M. javanica*) were the most frequent nematodes encountered with 32-62% FO and PD of 284-366 nematodes/250 cm³ soil. They were found in all soil samples of the surveyed plants except those collected from citrus trees. *Helicotylenchus* spp., *Pratylenchus* spp., *Tylenchorhynchus* spp. and *Tylenchus* spp. were common and observed in soil samples from all surveyed plants with 10 - 42% FO and 72 - 260 nematodes/250 cm³ soil. *Criconema* spp., *Hemicriconemooides* spp., *Longidorus* spp., *Merlinius* spp., *Paratylenchus* spp. and

Rotylenchus spp. were less common with 8-12% FO.

In soil samples from the Alexandria governorate, 12 nematode genera were identified associated with lantana and tomato (Table 1). *Meloidogyne incognita* and *M. javanica* were the most frequently found nematodes in these crops with 36-62% FO and PD of 284-366 nematodes/250 cm³ soil. *Aphelenchoides* spp., *Merlinius* spp. and *Rotylenchulus reniformis* were observed on lantana while *Criconema* spp., *Hoplolaimus* spp. and *Trichodorus* spp. occurred on tomato with 12-18% FO and PD 54-184 nematodes/250 cm³ soil. A previous nematode survey in Alexandria governorate showed the occurrence of 21 genera and 27 species of phytoparasitic nematodes associated with certain ornamental plants, weeds, and grasses (Ibrahim *et al.*, 2000).

Nematodes in 21 genera were extracted from

Table 1. Genera and species of phytoparasitic nematodes and their frequency of occurrence% (FO) and average nematode population density (PD) on the associated host plants in Alexandria and El-Behra governorates.

Nematode species	Alexandria				El-Behra			
	Lantana (86) ^x FO ^y	Tomato (84) FO PD	Citrus (92) FO PD	Guava (84) FO PD	Potato (56) FO PD	Spinach (86) FO PD	Sugar Beet (116) FO PD	
<i>Aphelenchoides</i> spp.	14	154	-	8	11	20	136	16
<i>Cricotylenchus</i> spp.	-	-	12	140	12	132	-	-
<i>Ditylenchus</i> spp.	-	-	-	-	10	184	24	10
<i>Globodera rostochiensis</i>	-	-	-	40	192	20	160	54
<i>Helicotylenchus</i> spp.	42	160	22	120	-	-	28	84
<i>Hemicricotylenchoides</i> spp	-	-	-	-	-	-	-	382
<i>Heterodera schachtii</i>	-	-	-	-	-	-	-	-
<i>Heterodera trifolii</i>	-	-	-	-	-	-	-	-
<i>Hoplolaimus</i> spp.	-	-	18	160	16	158	20	182
<i>Longidorus</i> spp.	-	-	-	10	84	-	-	-
<i>Meloidogyne incognita</i>	54	320	62	366	-	52	282	24
<i>Meloidogyne javanica</i>	36	248	62	320	-	40	274	-
<i>Merlinius</i> spp.	12	132	-	-	-	-	-	-
<i>Mesocriconema</i> spp.	-	-	-	-	-	-	-	-
<i>Paratylenchus</i> spp.	-	-	-	-	-	-	-	-
<i>Pratylenchus</i> spp.	36	120	32	136	20	142	24	164
<i>Psilenchus</i> spp.	-	-	-	-	-	-	-	-
<i>Rotylenchulus reniformis</i>	16	184	-	-	-	-	-	-
<i>Rotylenchus</i> spp.	-	-	-	-	10	74	-	-
<i>Trichodorus</i> spp.	-	-	18	96	-	16	140	-
<i>Tylenchorhynchus</i> spp.	16	84	22	112	18	124	16	74
<i>Tylenchulus semipenetrans</i>	-	-	-	-	94	5,640	-	-
<i>Tylenchus</i> spp.	14	96	10	114	16	74	10	72
<i>Xiphinema</i> spp.	12	74	12	66	14	112	8	74
<i>Xiphinema rivesi</i>	-	-	-	-	36	342	-	-

^xTotal numbers of soil samples.

^yFO = Number of positive samples/total number of collected soil samples 100.

^zPD = Numbers of nematodes per 250 cm³ soil.

from soil samples collected from the El-Behera governorate (Table 1). *Helicotylenchus* spp., *Hoplolaimus* spp., *Pratylenchus* spp., *Tylenchorhynchus* spp. and *Tylenchus* spp. were found on all surveyed host plants. *Meloidogyne incognita* and *M. javanica* were found in all surveyed plants except citrus with 40-64% FO and PO of 282-364 nematodes/250 cm³ soil.

In the El-Behera governorate, *Heterodera schachtii*, was found on spinach and sugar beet with 20-48% FO and PD of 264-274 nematodes/250 cm³ soil while *H. trifolii* was observed on spinach with 14% FO and an average PD of 182 nematodes/250 cm³ soil. *Globodera rostochiensis*, was identified on potato with 54% FO and an average PD of 382 nematodes/250 cm³ of soil. This cyst nematode was widespread in El-Nitron Valley and El-Noaria region, southwest of El Behera governorate. *Tylenchulus semipenetrans* was very common in all citrus trees surveyed with high (94%) FO and an average PD of 5,646 nematodes/250 cm³ soil. Recently the pathogenicity and control of this nematode on citrus and other fruit trees were studied by Ibrahim *et al.* (2022). *Xiphinema* spp. was identified on citrus, guava, and spinach with 8-14% FO and PD of 66-112 nematodes/250 cm³ soil while *X. rivesi* was found on citrus trees in El-Noaria region with 36% FO and an average PD of 342 nematodes/250 cm³ soil. In a previous nematode survey in El-Behera governorates (Ibrahim *et al.*, 2000), 14 nematode genera were recorded on ornamental palm trees, grasses, and weeds. The present results are supportive of previous studies (Handoo *et al.*, 2015; Ibrahim *et al.*, 2017, 2023) regarding the occurrence of *X. rivesi* on citrus and *G. rostochiensis* on potato in El-Behera governorate, Egypt.

The results of this survey provide insight concerning important phytoparasitic nematode genera and species associated with certain important crops grown in northern Egypt. Surveyed plant crops were selected either because their economic importance or because they showed stunted growth symptoms with leaf yellowing on citrus, potato and sugar beet that were infested with *T. semipenetrans*, *G. rostochiensis* and *H. schachtii*. In conclusion, this survey establishes a foundation for assessing the extent and intensity of these phytoparasitic nematode-related challenges, thus aiding in the formulation of effective mitigation strategies for sustainable plant health.

ACKNOWLEDGMENTS

Alemayehu Habteweld was supported, in part, by an appointment to the Research Participation Program at the Mycology and Nematology Genetic Diversity and Biology Laboratory USDA, ARS, Northeast Area, Beltsville, MD, administered by the Oak Ridge Institute for Science and Education through an interagency agreement between the U.S. Department of Energy and USDA-ARS.

LITERATURE CITED

- Adam, M., H. Heuer, E. M. Ramadan, M. A. Hussein, and J. Hallmann. 2013. Occurrence of plant-parasitic nematodes in organic farming in Egypt. International Journal of Nematology 23:82-90.
- Ayoub, S. M. 1980. Plant nematology, an agriculture training aid. Sacramento, California: Nema Aid Publications.
- Geraert, E. 2008. The Tylenchidae of the world. Identification of the family Tylenchidae (Nematoda). Gent, Belgium: Academia Press.
- Goodey, J. B. 1963. Soil and freshwater nematodes. New York: John Wiley and Sons, Inc.
- Golden, A. M. 1986. Morphology and identification of cyst nematodes. Pp. 23-45 in F. Lamberti and C. E. Taylor, (eds.) Cyst Nematodes. New York, NY: Plenum Press.
- Handoo, Z. A. 2000. A key and diagnostic compendium to the species of the genus *Tylenchorhynchus* Cobb, 1913 (Nematoda: Belonolaimidae). Journal of Nematology 32:20-34.
- Handoo, Z. A., and A. M. Golden. 1989. A key and diagnostic compendium to the species of the genus *Pratylenchus* Filipjev, 1936 (lesion nematodes). Journal of Nematology 21:202-218.
- Handoo, Z. A., A. Khan, and S. Islam. 2007. A key and diagnostic compendium to the species of the genus *Merlinius* Siddiqi, 1970 (Nematode: Tylenchida) with description of *Merlinius khuzdarensis* n. sp. associated with date palm. Nematology 9:251-260.
- Handoo, Z. A., and I. K. A. Ibrahim. 2002. Description and SEM observation of a new species of cyst nematode *Heterodera goldeni* (Nematoda: Heterodidae) attacking *Panicum coloratum* in Egypt. Journal of Nematology

- 34:312-318.
- Handoo, Z. A., I. K. A. Ibrahim, D. J. Chitwood, and A. A. Mokbel. 2015. First record of *Xiphinema revesi* Dalmasso, 1969 on citrus in northern Egypt. *Pakistan Journal of Nematology* 33:161-165.
- Hunt, D. J., and Z. A. Handoo 2009. Taxonomy, identification, and principal species. Pp. 55-97 in Perry, R. N., Moens, M. and Starr, J. (eds.). Root-knot Nematodes. Wallingford, UK: CABI Publishing
- Ibrahim, I. K. A. 1990. The status of phytoparasitic nematodes and the associated host plants in Egypt. *International Nematology Network Newsletter* 7:33-38.
- Ibrahim, I. K. A., and Z. A. Handoo. 2007. A survey of cyst nematodes (*Heterodera spp.*) in Northern Egypt. *Pakistan Journal of Nematology* 25:335-337.
- Ibrahim, I. K. A., and Z. A. Handoo. 2016. Occurrence of phytoparasitic nematodes on some crop plants in northern Egypt. *Pakistan Journal of Nematology* 34:163-169.
- Ibrahim, I. K. A., Z. A. Handoo, and A. A. El-Sherbini. 2000. A survey of phytoparasitic nematodes on cultivated and non-cultivated plants in northern Egypt. Supplement of *Journal Nematology* 32:478-485.
- Ibrahim, I. K. A., A. A. Mokbel, and Z. A. Handoo. 2010. Current status of phytoparasitic nematodes and their host plants in Egypt. *Nematropica* 40:239-262.
- Ibrahim, I. K. A., Z. A. Handoo, and A. B. A. Basyony. 2017. The cyst nematodes *Heterodera* and *Globodera* species in Egypt. *Pakistan Journal of Nematology* 35:151-154.
- Ibrahim, I. K. A., A. H. A. Abu-Habib, M. Kantor, and Z.A. Handoo. 2022. Pathogenicity and control of the citrus nematode *Tylenchulus semipenetrans* on citrus, grape, olive, loquat, and persimmon species and cultivars. *Nematropica* 52:79- 84.
- Ibrahim, I. K. A., M. Kantor, and Z. A. Handoo. 2023. Current status of genera and species of phytoparasitic nematodes in Egypt. *Nematropica* 53:16-29.
- Ismail, A. E., and M. F. M. Eissa. 1993. Plant-parasitic nematodes associated with ornamental palms in three botanic gardens in Egypt. *Pakistan Journal of Nematology* 11:53-59.
- Lamberti, F., A. Agostinelli, and V. Radicci. 1996. Longidorid nematodes from northern Egypt. *Nematologia Mediterranea* 24:307-339.
- Mai, W. F., and H. H. Lyon. 1975. Pictorial key to genera of plant-parasitic nematodes. Fourth Edition. Ithaca, NY: Comstock Publishing Associates, Cornell University Press.
- Mulvey, R. H., and A. M. Golden. 1983. An illustrated key to the cyst-forming genera and species of Heteroderidae in the western hemisphere with species morphometrics and distribution. *Journal of Nematology* 15:1-59.
- Oteifa, B. A. 1962. Species of root-lesion nematodes commonly associated with economic crops in the Delta of U.A.R. *Plant Disease Reporter* 46:572-575.
- Oteifa, B. A., M. M. Shams El-Dean, and M. H. El-Hamawi. 1997. A preliminary compiled study on the biodiversity of free-living, plant and insect parasitic nematodes in Egypt. *Egyptian Journal of Agronematology* 1:1-36.
- Oteifa, B. A., and A. C. Tarjan. 1965. Potentially important plant parasitic nematodes present in established orchards of newly-reclaimed sandy areas of the U.A.R. *Plant Disease Reporter* 49:596-597.
- Raski, D. J. 1991. Tylenchulidae in agricultural soils. Pp. 761-794 in Nickle, W. R. (ed.) *Manual of Agricultural Nematology*. New York, NY: Marcel Dekker, Inc.
- Seinhorst, J. W. 1959. A rapid method for the transfer of nematodes from fixative to anhydrous glycerine. *Nematologica* 4:67-69.
- Sher, S. A. 1966. Revision of the Hoplolaiminae (Nematoda). *Helicotylenchus* Steiner, 1945 *Nematologica* 12:1-56.
- Tarjan, A.C. 1964. Plant parasitic nematodes in the United Arab Republic. FAO Plant Protection Bulletin 12:49-56.
- Taylor, A. L., and J. N. Sasser. 1987. Biology, identification, and control of root-knot nematodes (*Meloidogyne spp.*). Raleigh, NC: North Carolina State University Graphics.

Received:

Accepted for publication:

20/VI/2023

28/VIII/2023

Recibido:

Aceptado para publicación: