

# Climate change reduces the net sink of CH<sub>4</sub> and N<sub>2</sub>O in a semiarid grassland

FEIKE A. DIJKSTRA\*, JACK A. MORGAN†, RONALD F. FOLLETT‡ and DANIEL R. LECAIN†

\*Department of Environmental Sciences and Centre for Carbon, Water and Food, Faculty of Agriculture and Environment, The University of Sydney, Camden, NSW 2570, Australia, †Rangeland Resources Research Unit, USDA-ARS, Fort Collins, CO 80526, USA, ‡Soil, Plant, and Nutrient Research Unit, USDA-ARS, Fort Collins, CO 80526, USA

## Abstract

Atmospheric concentrations of methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) have increased over the last 150 years because of human activity. Soils are important sources and sinks of both potent greenhouse gases where their production and consumption are largely regulated by biological processes. Climate change could alter these processes thereby affecting both rate and direction of their exchange with the atmosphere. We examined how a rise in atmospheric CO<sub>2</sub> and temperature affected CH<sub>4</sub> and N<sub>2</sub>O fluxes in a well-drained upland soil (volumetric water content ranging between 6% and 23%) in a semiarid grassland during five growing seasons. We hypothesized that responses of CH<sub>4</sub> and N<sub>2</sub>O fluxes to elevated CO<sub>2</sub> and warming would be driven primarily by treatment effects on soil moisture. Previously we showed that elevated CO<sub>2</sub> increased and warming decreased soil moisture in this grassland. We therefore expected that elevated CO<sub>2</sub> and warming would have opposing effects on CH<sub>4</sub> and N<sub>2</sub>O fluxes. Methane was taken up throughout the growing season in all 5 years. A bell-shaped relationship was observed with soil moisture with highest CH<sub>4</sub> uptake at intermediate soil moisture. Both N<sub>2</sub>O emission and uptake occurred at our site with some years showing cumulative N<sub>2</sub>O emission and other years showing cumulative N<sub>2</sub>O uptake. Nitrous oxide exchange switched from net uptake to net emission with increasing soil moisture. In contrast to our hypothesis, both elevated CO<sub>2</sub> and warming reduced the sink of CH<sub>4</sub> and N<sub>2</sub>O expressed in CO<sub>2</sub> equivalents (across 5 years by 7% and 11% for elevated CO<sub>2</sub> and warming respectively) suggesting that soil moisture changes were not solely responsible for this reduction. We conclude that in a future climate this semiarid grassland may become a smaller sink for atmospheric CH<sub>4</sub> and N<sub>2</sub>O expressed in CO<sub>2</sub>-equivalents.

**Keywords:** climate change, global warming potential, multifactor experiment, PHACE, positive feedback, water availability

Received 6 February 2013; revised version received 6 February 2013 and accepted 17 February 2013

## Introduction

Methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) are powerful greenhouse gases that are 25 and 298 times more potent than carbon dioxide (CO<sub>2</sub>) over a 100 year lifespan (Forster *et al.*, 2007). Terrestrial ecosystems are large sources and sinks of CH<sub>4</sub> and N<sub>2</sub>O. Climate change may alter the source and sink strength of CH<sub>4</sub> and N<sub>2</sub>O in terrestrial ecosystems with potentially important feedbacks to the global climate system (King, 1997; Xu-Ri *et al.*, 2012). Based on a meta-analysis including field, greenhouse and growth chamber studies, it was estimated that a rise in atmospheric CO<sub>2</sub> concentration projected for the mid- to end of this century would increase CH<sub>4</sub> and N<sub>2</sub>O emission from terrestrial ecosystems equivalent to 1.1 Pg CO<sub>2</sub> per yr globally, thereby offsetting 16.6% or more of the concurrent increase in terrestrial C storage (van Groenigen *et al.*, 2011). Similarly, using a process-based model, it was estimated

that combined CH<sub>4</sub> and N<sub>2</sub>O emission from terrestrial ecosystems in North America alone would increase by 0.7–1.3 Pg CO<sub>2</sub> eq. per yr by the end of this century in response to future climate change scenarios that were derived from three different global climate models (elevated CO<sub>2</sub> effects not included, Tian *et al.*, 2012).

Results from manipulative field experiments indicate that elevated CO<sub>2</sub> and warming effects on CH<sub>4</sub> and N<sub>2</sub>O exchange with the atmosphere vary widely among different systems (Dijkstra *et al.*, 2012). Elevated CO<sub>2</sub> reduced CH<sub>4</sub> uptake in a temperate forest (Phillips *et al.*, 2001a; Dubbs & Whalen, 2010) and in a temperate grassland (Ineson *et al.*, 1998), but had no effect in other temperate and semiarid grasslands (Mosier *et al.*, 2002; Blankinship *et al.*, 2010). Reduced CH<sub>4</sub> uptake was associated with increased soil moisture impeding the supply of atmospheric CH<sub>4</sub> for oxidation by methanotrophs in the soil, and increasing CH<sub>4</sub> production by methanogens (Ineson *et al.*, 1998; Phillips *et al.*, 2001a). Elevated CO<sub>2</sub> increased N<sub>2</sub>O emission in grasslands and agroecosystems that were fertilized with N (Ineson *et al.*, 1998; Baggs *et al.*, 2003; Kammann *et al.*, 2008;

Correspondence: Feike A. Dijkstra, tel. + 61 2 9351 1817, fax + 61 2 8627 1099, e-mail: feike.dijkstra@sydney.edu.au

Lam *et al.*, 2011), but generally had no effect in non-fertilized systems (Phillips *et al.*, 2001b; Billings *et al.*, 2002; Mosier *et al.*, 2002; Larsen *et al.*, 2011; Niboyet *et al.*, 2011). Greater labile C inputs under elevated CO<sub>2</sub> could stimulate N<sub>2</sub>O production by denitrifiers, at least when inorganic N is also abundant (Dijkstra & Morgan, 2012; Dijkstra *et al.*, 2012). However, in systems where N availability is low, elevated CO<sub>2</sub> could reduce N availability thereby constraining N<sub>2</sub>O production (Hungate *et al.*, 1997; Mosier *et al.*, 2002). Warming increased CH<sub>4</sub> uptake in a temperate forest and several subarctic systems (Peterjohn *et al.*, 1994; Sjögersten & Wookey, 2002), most likely because the soil drying effect of warming enhanced diffusivity thereby stimulating the oxidation of atmospheric CH<sub>4</sub> in the soil. Warming increased N<sub>2</sub>O emission in urban lawns and a heath land (Bijoor *et al.*, 2008; Larsen *et al.*, 2011), decreased N<sub>2</sub>O emission in a wheat field and an alpine meadow (Hantschel *et al.*, 1995; Hu *et al.*, 2010), and had no effect on N<sub>2</sub>O exchange in an annual grassland and temperate forests (Peterjohn *et al.*, 1994; McHale *et al.*, 1998; Niboyet *et al.*, 2011). Inconsistent warming effects could arise because warming affects multiple processes that may cause opposing effects on N<sub>2</sub>O exchange. For instance, an increase in soil temperature could directly enhance the activity of nitrifiers and denitrifiers thereby stimulating N<sub>2</sub>O emission, but a decrease in soil moisture could reduce microbial activity (McHale *et al.*, 1998; Bijoor *et al.*, 2008).

In semiarid grasslands, which comprise roughly 11% of the global land surface (Bailey, 1979), one of the most important drivers for biological activity is soil moisture, including the production and uptake of CH<sub>4</sub> and N<sub>2</sub>O (Mosier *et al.*, 2008). Because soil moisture is low in most of the year, production and uptake of CH<sub>4</sub> and N<sub>2</sub>O in semiarid grasslands can be very different from more mesic environments. For instance, in mesic environments net CH<sub>4</sub> uptake is often negatively related to soil moisture because soil moisture reduces CH<sub>4</sub> diffusivity into the soil for oxidation by methanotrophs, or increases CH<sub>4</sub> emission by methanogens (Del Grosso *et al.*, 2000; Phillips *et al.*, 2001a). In semiarid environments however, soils are often so dry that soil moisture effects on methanotroph activity can become more important than effects on CH<sub>4</sub> diffusivity or methanogen activity (von Fischer *et al.*, 2009). Therefore, under dry conditions net CH<sub>4</sub> uptake can increase with increased soil moisture (Mosier *et al.*, 2008; Dijkstra *et al.*, 2011). In mesic and wet environments N<sub>2</sub>O emissions can be high because intermediate to high levels of soil moisture are conducive to nitrification and denitrification, both of which contribute to N<sub>2</sub>O emissions (Klemetsson *et al.*, 1988; Davidson, 1992). Net N<sub>2</sub>O emission in mesic environments is sometimes

restrained, however, because of N<sub>2</sub>O consumption by denitrifiers (Holtgrieve *et al.*, 2006; Chapuis-lardy *et al.*, 2007). Consumption of N<sub>2</sub>O has also been observed under dry conditions resulting in a net N<sub>2</sub>O sink (Donoso *et al.*, 1993; Goldberg & Gebauer, 2009; Grover *et al.*, 2012; Stewart *et al.*, 2012). While the underlying mechanisms are still unclear, a possible explanation is that drier soil conditions enhance N<sub>2</sub>O diffusion from the atmosphere into the soil where under limited inorganic N supply from the soil, atmospheric N<sub>2</sub>O is reduced to N<sub>2</sub> by denitrifiers (Chapuis-lardy *et al.*, 2007).

Because biological processes strongly depend on soil moisture in semiarid grasslands we postulate that climate change effects on CH<sub>4</sub> and N<sub>2</sub>O in these systems are largely mediated by changes in soil moisture. Previously we have shown that elevated CO<sub>2</sub> increased soil moisture in a semiarid grassland in Wyoming, USA, because of reduced stomatal conductance, while warming decrease soil moisture because of desiccation effects (Morgan *et al.*, 2011). During the first two growing seasons of the experiment CH<sub>4</sub> uptake was strongly controlled by soil moisture, and elevated CO<sub>2</sub> and warming effects could largely be explained by treatment effects on soil moisture (Dijkstra *et al.*, 2011). The drier soil conditions with warming reduced CH<sub>4</sub> uptake over the whole growing season. However, the increase in soil moisture under elevated CO<sub>2</sub> enhanced CH<sub>4</sub> uptake during dry periods of the growing season, but reduced CH<sub>4</sub> uptake during wet periods. When elevated CO<sub>2</sub> effects on CH<sub>4</sub> uptake were considered for the whole growing season, elevated CO<sub>2</sub> had no net effect. Interactive CO<sub>2</sub> × warming effects were also not observed during the first 2 years. Here, we present single and combined effects of elevated CO<sub>2</sub> and warming on CH<sub>4</sub> and N<sub>2</sub>O fluxes during five growing seasons. None of the five growing seasons were exceptionally dry or wet (Table 1). We hypothesized that an increase in soil moisture under elevated CO<sub>2</sub> would have no effect on CH<sub>4</sub> uptake, but that a decrease in soil moisture with warming would decrease CH<sub>4</sub> uptake, similar to what we found during the first 2 years. We further hypothesized that elevated CO<sub>2</sub> would have no effect on N<sub>2</sub>O emission because of a tighter N cycle under elevated CO<sub>2</sub> (Dijkstra *et al.*, 2010). On the other hand, we

**Table 1** Total precipitation and mean temperature during the growing season for each year and averaged across all 5 years

	2007	2008	2009	2010	2011	Average
Precip. (mm)	288	349	353	242	363	319
Mean temp (°C)	17.4	15.9	12.5	13.3	13.9	14.6

hypothesized that drier soil conditions with warming would reduce N<sub>2</sub>O emission, and with limited N available, potentially turning this system into an N<sub>2</sub>O sink. Because the decrease in net N<sub>2</sub>O emission would offset the decrease in net CH<sub>4</sub> uptake with warming, combined CH<sub>4</sub> and N<sub>2</sub>O fluxes in this grassland expressed in CO<sub>2</sub> equivalents would be insensitive to climate change.

## Materials and methods

### Site description and experimental design

We conducted our study in the Prairie Heating And CO<sub>2</sub> Enrichment (PHACE) experiment in a semiarid grassland in Wyoming, USA (41° 11' N, 104° 54' W). Vegetation at the site is a northern mixed prairie dominated by the C<sub>4</sub> grass *Bouteloua gracilis* (H.B.K) Lag and the C<sub>3</sub> grasses *Pascopyrum smithii* (Rydb.) A. Love and *Hesperostipa comata* Trin and Rupr. Other species include the sedge *Carex eleocharis* L. Bailey, the forb *Sphaeralcea coccinea* (Nutt.) Rydb., and the subshrub *Artemisia frigida* Willd. The soil is a fine-loamy, mixed, mesic Aridic Argiustoll with a pH of 7.0 and a total organic C and N content of 1.7% and 0.16%, respectively, in the top 15 cm. Mean air temperature in January is -2.5 °C and 17.5 °C in July and the mean annual precipitation is 384 mm (132-year mean). Methane and N<sub>2</sub>O fluxes were measured during the growing season (April–October) from 2007 to 2011 (see below). Total precipitation during the growing season varied between 242 mm in 2010 and 363 mm in 2011, while mean air temperature varied between 12.5 °C in 2009 and 17.4 °C in 2007 (Table 1).

In 2005 twenty 3.4 m diameter plots were established in a 2.5 ha fenced-off area that had been moderately grazed until 2005. A plastic flange was dug into the ground 60-cm deep around each plot preventing lateral water flow. We used Free Air CO<sub>2</sub> Enrichment technology (Miglietta *et al.*, 2001) to increase the atmospheric CO<sub>2</sub> concentration to 600 ppmv (±40 ppmv). The CO<sub>2</sub> was injected into the plot from a plastic pipe, perforated with 300-µm laser-drilled holes, surrounding the plot just above the plant canopy. The CO<sub>2</sub> treatment started in April 2006 and only occurred when plants were photosynthetically active (during light hours and during the growing season from April through October). We used ceramic infrared heaters (1000 W; Mor Electric Heating Assoc., Inc., Comstock Park, MI, USA) controlled by a proportional-integral-derivative feed-back loop (Kimball *et al.*, 2008) to increase the canopy temperature by 1.5 °C above ambient during the day and by 3 °C during the night. Heaters were attached to a metal triangular frame 1.5 m above the ground surface (six heaters per plot). The warming treatment started in April 2007 and ran continuously throughout the year. The CO<sub>2</sub> and warming treatments were implemented in a full factorial design with five replicates of each of the treatment combinations (ct: ambient CO<sub>2</sub> and ambient temperature, cT: ambient CO<sub>2</sub> and elevated temperature, Ct: elevated CO<sub>2</sub> and ambient temperature, and CT: elevated CO<sub>2</sub> and elevated temperature). Detailed information about the site and CO<sub>2</sub> and

warming treatments can be found elsewhere (Dijkstra *et al.*, 2010; Morgan *et al.*, 2011). In June 2008 we established 0.4 m<sup>-2</sup> subplots without plants in each of the 20 plots. These subplots were separated from the main plot by a metal sheet buried 30 cm into the ground. The vegetation was killed by lightly spraying with the broad spectrum systemic herbicide glyphosate. The N that we added with the glyphosate (<0.3 g N m<sup>-2</sup> compared to approximately 0.5 g N m<sup>-2</sup> as NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> in the top 15 cm of the soil, Dijkstra *et al.*, 2010) may have had some temporary impacts on soil N availability but most likely no or minor impacts in the long-term. New seedlings after spraying were regularly removed by hand.

### CH<sub>4</sub> and N<sub>2</sub>O flux measurements

We measured CH<sub>4</sub> and N<sub>2</sub>O fluxes approximately every other week during five consecutive growing seasons (from April through October), starting in 2007 until 2011 (between 12 and 16 measurements per season) using static chambers (Hutchinson & Mosier, 1981). We inserted chamber anchors (diameter 20 cm, height 10 cm) 8 cm into the soil, two in each plot (one in an area with vegetation intact (with plants), and one in the area where we killed the vegetation (without plants)). Anchors were installed at least 1 month before the first measurements were taken. Measurements were taken during midmorning. After placement of the chambers on the anchors, gas samples were taken from the headspace after 0, 15, 30 and 45 min. Gas samples were analysed for CH<sub>4</sub> and N<sub>2</sub>O concentration on a gas chromatograph (GC) equipped with a flame ionization detector for CH<sub>4</sub> and an electron capture detector for N<sub>2</sub>O (Varian 3800; Varian, Inc., Palo Alto, CA, USA) usually within 2 days after sampling. The precision of the GC was about 1 ppb for both gases. The CH<sub>4</sub> and N<sub>2</sub>O fluxes were calculated as the slope of linear regressions from the measured gas concentrations with time. The *r*<sup>2</sup> value of the regressions was sometimes below 0.4, particularly when fluxes were low. Because we did not want to create a bias against low fluxes, we did not discard fluxes with low *r*<sup>2</sup> values, unless CO<sub>2</sub> concentrations measured at the same time as CH<sub>4</sub> and N<sub>2</sub>O did not show a clear increasing trend with time. We calculated cumulative fluxes of CH<sub>4</sub> (in mg C m<sup>-2</sup>) and N<sub>2</sub>O (in mg N m<sup>-2</sup>) produced/consumed over the growing season by multiplying the average flux measured at two consecutive dates by the time interval, and by summing up the cumulative fluxes calculated for each time interval of the growing season. Because we did not start with the CH<sub>4</sub> and N<sub>2</sub>O flux measurements in the subplots without plants until June in 2008, we only calculated cumulative fluxes of CH<sub>4</sub> and N<sub>2</sub>O in the subplots without plants for 2009, 2010 and 2011. We calculated the cumulative Global Warming Potential from CH<sub>4</sub> and N<sub>2</sub>O (in g CO<sub>2</sub> eq. per m<sup>2</sup>) by adding the cumulative GWP (Global Warming Potential) from CH<sub>4</sub> (cumulative flux of CH<sub>4</sub> in g CH<sub>4</sub> per m<sup>2</sup> multiplied by 25) and the cumulative GWP from N<sub>2</sub>O (cumulative flux of N<sub>2</sub>O in g N<sub>2</sub>O per m<sup>2</sup> multiplied by 298).

### Soil moisture, temperature and plot greenness

Soil moisture at 10-cm soil depth was measured in each plot (with plants only) with EnviroSMART probes (Sentek Sensor

Technologies, Stepney, Australia). Water filled pore space (WFPS) in the top 15 cm of the soil was calculated based on soil moisture measured at 10-cm soil depth and bulk densities measured at 0–5 and 5–15 cm soil depth in 2005. Soil temperature at 10-cm soil depth was measured in each plot (with plants only) with thermocouples. Soil moisture and temperature data were logged every hour throughout the year (CR10X data loggers; Campbell Scientific, Logan, UT, USA). We calculated seasonal average WFPS and soil temperature for each year by averaging the WFPS and soil temperature values recorded at the time of flux measurements.

We measured plot greenness as a degree of photosynthetically active plant biomass inside the static chambers at the time of CH<sub>4</sub> and N<sub>2</sub>O flux measurements. In each plot (with plants only) digital photographs were taken of the surface area inside the anchors with a camera attached to a tripod from 50 cm above the ground. Photographs were taken directly after each time we sampled for CH<sub>4</sub> and N<sub>2</sub>O. Photographs were then analysed for the total green area as a percentage of the total surface area inside the anchor (where the area that was not green was either bare soil or senesced plant material) using the software program SamplePoint (Booth *et al.*, 2006). Plot greenness varied between 0 (in early April and late October) and 62% (July–August) during the season (Fig. S2). Plot greenness measured in mid-July of 2007–2011 was positively related to aboveground green biomass harvested in mid-July of each year ( $n = 100$ ,  $P < 0.0001$ ,  $r = 0.65$ ), suggesting that plot greenness provided a reasonable measure of photosynthetically active plant biomass.

### Statistical analyses

We used a repeated measures analysis of variance (repeated measures ANOVA) to test for main effects of CO<sub>2</sub> (ambient vs. elevated), temperature (no warming vs. warming, both between-subjects factors), year (2007–2011, within-subjects factor), and their interactive effects on cumulative CH<sub>4</sub>, N<sub>2</sub>O and GWP from CH<sub>4</sub> and N<sub>2</sub>O. We ran the repeated measures analysis separately for measurements with and without plants. We used *post-hoc* tests (Tukey's HSD) to test for differences in cumulative CH<sub>4</sub>, N<sub>2</sub>O and GWP from CH<sub>4</sub> and N<sub>2</sub>O among the different CO<sub>2</sub> and warming treatment combinations. Repeated measures ANOVA was further used for WFPS, soil temperature and plot greenness measured in the subplots with plants only. We used linear and nonlinear regressions to relate CH<sub>4</sub> and N<sub>2</sub>O fluxes to WFPS, soil temperature and plot greenness using data that were aggregated by date (i.e., averaged across the CO<sub>2</sub> and warming treatments) and by date and treatment (i.e., average of the five replicates of each treatment on each date). We further tested if relationships differed among the CO<sub>2</sub> and warming treatments using analysis of covariance (ANCOVA) with the CO<sub>2</sub> and warming treatment as main effects and either WFPS, soil temperature and plot greenness as the covariate. Because CH<sub>4</sub> fluxes showed bell-shaped relationships with WFPS and soil temperature, we included a quadratic term of the covariate in the ANCOVAs (Dijkstra *et al.*, 2011). Significant interactions between main effects and the covariate indicate that relationships between


CH<sub>4</sub> or N<sub>2</sub>O fluxes with WFPS, soil temperature, or plot greenness were altered by the CO<sub>2</sub> and/or warming treatment. When necessary, data were log-transformed to reduce heteroscedasticity and improve assumptions of normality. All statistical analyses were performed with JMP (version 4.0.4; SAS Institute, Cary, NC, USA).

### Results


Soils were a net sink for CH<sub>4</sub> throughout all five growing seasons (as indicated by negative fluxes). Variable CH<sub>4</sub> fluxes were observed within each season although CH<sub>4</sub> uptake tended to be highest mid to late summer and smallest at the start and end of the growing season (Fig. S1). In the presence of plants, cumulative CH<sub>4</sub> uptake during the growing season was smallest in 2007 (90 mg C m<sup>-2</sup> averaged across the CO<sub>2</sub> and warming treatments) and greatest in 2011 (130 mg C m<sup>-2</sup>, Fig. 1). Elevated CO<sub>2</sub> increased cumulative CH<sub>4</sub> uptake in 2007 by 15%, but decreased it in all other years up to 12% in 2011 causing a significant CO<sub>2</sub> × year interactive effect ( $P = 0.02$ ). Warming significantly reduced cumulative CH<sub>4</sub> uptake across all years ( $P < 0.0001$ ) with an average decrease of 15%. Warming also significantly reduced cumulative CH<sub>4</sub> uptake without plants ( $P = 0.01$ ), while elevated CO<sub>2</sub> had no effect on cumulative CH<sub>4</sub> uptake without plants. Cumulative CH<sub>4</sub> uptake was higher without than with plants.

Nitrous oxide fluxes in the presence of plants were small throughout the growing season and were often negative, particularly during mid to late summer (Fig. S1), suggesting soil uptake of N<sub>2</sub>O at those times. Cumulative fluxes of N<sub>2</sub>O were also sometimes negative (N<sub>2</sub>O uptake), particularly during the last 2 years (Fig. 2a). Across years there was no significant elevated CO<sub>2</sub> effect on cumulative N<sub>2</sub>O, but during the last 2 years N<sub>2</sub>O uptake decreased under elevated CO<sub>2</sub> causing a significant CO<sub>2</sub> × year interactive effect ( $P = 0.002$ ). There was no warming effect on cumulative fluxes of N<sub>2</sub>O. Cumulative fluxes of N<sub>2</sub>O were much larger without than with plants, particularly in 2009 and 2011 (Fig. 2b). Elevated CO<sub>2</sub> significantly reduced the cumulative flux of N<sub>2</sub>O production without plants ( $P = 0.006$ ).

Seasonal variability in WFPS, soil temperature and plot greenness was large (Fig. S2); where WFPS ranged between 11% and 41%, soil temperature between 2 and 29 °C, and plot greenness between 0% and 62% during the growing season. Growing season averages of WFPS were significantly higher under elevated CO<sub>2</sub> (absolute increase of 3.8% across all years,  $P < 0.0001$ ) and significantly lower with warming (absolute decrease of 2.1% across all years,  $P < 0.0001$ , Table 2). Soil temperature was significantly higher with warming (increase of



**Fig. 1** Cumulative  $\text{CH}_4$  in  $\text{mg C m}^{-2}$  (left  $y$ -axis) and in  $\text{g CO}_2 \text{ eq. per m}^2$  (right  $y$ -axis) in plots with (a) and without plants (b) in response to elevated  $\text{CO}_2$  and warming. Negative values indicate sinks. Treatments: ct: ambient  $\text{CO}_2$  and ambient temperature; cT: ambient  $\text{CO}_2$  and elevated temperature; Ct: elevated  $\text{CO}_2$  and ambient temperature; CT: elevated  $\text{CO}_2$  and elevated temperature. W: warming treatment. Error bars indicate  $\pm 1$  SE. ANOVA  $P$ -values are reported when  $P < 0.05$  (in bold) or  $P < 0.1$  (in italics). Different letters above bars indicate significant differences among the elevated  $\text{CO}_2$  and warming treatments for each year separately ( $P < 0.05$ , Tukey's HSD test).



**Fig. 2** Cumulative  $\text{N}_2\text{O}$  in  $\text{mg N m}^{-2}$  (left  $y$ -axis) and in  $\text{g CO}_2 \text{ eq. per m}^2$  (right  $y$ -axis) in plots with (a) and without plants (b) in response to elevated  $\text{CO}_2$  and warming. Negative values indicate sinks. Treatments: ct: ambient  $\text{CO}_2$  and ambient temperature; cT: ambient  $\text{CO}_2$  and elevated temperature; Ct: elevated  $\text{CO}_2$  and ambient temperature; CT: elevated  $\text{CO}_2$  and elevated temperature. W: warming treatment. Error bars indicate  $\pm 1$  SE. ANOVA  $P$ -values are reported when  $P < 0.05$  (in bold) or  $P < 0.1$  (in italics). Different letters above bars indicate significant differences among the elevated  $\text{CO}_2$  and warming treatments for each year separately ( $P < 0.05$ , Tukey's HSD test).

1.9 °C,  $P < 0.0001$ ), but was not affected by elevated  $\text{CO}_2$ . Elevated  $\text{CO}_2$  and warming also had no significant effect on plot greenness.

The temporal variation in  $\text{CH}_4$  and  $\text{N}_2\text{O}$  fluxes in subplots with plants could to a large degree be explained by the temporal variation in WFPS (Fig. 3a and b). We observed a significant bell-shaped relationship for  $\text{CH}_4$  with the largest  $\text{CH}_4$  uptake at 24% WFPS and smaller uptake at higher and lower WFPS. We further observed significant interactions between WFPS and the  $\text{CO}_2$  and warming treatments (Table S1), indicating that the relationships differed among the  $\text{CO}_2$  and warming treatments. Individual regressions for each treatment showed that elevated  $\text{CO}_2$  increased the

optimum WFPS (on average by 3.6%), while warming showed reduced  $\text{CH}_4$  uptake at optimum WFPS (on average by  $3.1 \mu\text{g C m}^{-2} \text{ hr}^{-1}$ , Fig. S3, Table S1). Similar results were obtained during the first 2 years of the experiment (Dijkstra *et al.*, 2011). On the other hand the temporal  $\text{N}_2\text{O}$  fluxes showed a significant positive linear relationship with WFPS (Fig. 3b), while the slope of this relationship was lower under elevated  $\text{CO}_2$  (Fig. S3, Table S1). Despite the large range in soil temperatures at the time of our measurements, soil temperature had no effect on  $\text{CH}_4$  when data were aggregated across the  $\text{CO}_2$  and warming treatment (Fig. 3c). However,  $\text{CH}_4$  fluxes showed bell-shaped relationships with soil temperature, similar to the relationships with

**Table 2** Water filled pore space (WFPS), soil temperature and plot greenness averaged by the elevated CO<sub>2</sub> and warming treatments measured during trace gas flux measurements

Year	WFPS (%)			Soil temp (°C)			Plot greenness (%)					
	ct	cT	Ct	CT	ct	cT	Ct	CT	ct	cT	Ct	CT
2007	21.8 ± 1.5	20.1 ± 0.7	26.1 ± 1.9	24.6 ± 2.0	16.3 ± 0.6	18.5 ± 0.6	16.3 ± 0.5	18.2 ± 0.6	20.2 ± 2.0	21.7 ± 1.7	21.7 ± 0.8	21.9 ± 1.1
2008	24.5 ± 1.2	21.6 ± 0.5	27.6 ± 1.2	25.4 ± 1.3	15.5 ± 1.0	17.6 ± 0.5	15.7 ± 0.6	16.9 ± 0.6	23.4 ± 1.6	23.7 ± 1.1	21.3 ± 0.8	22.8 ± 0.7
2009	24.3 ± 1.2	22.5 ± 0.8	29.1 ± 1.4	26.6 ± 1.2	14.5 ± 0.9	17.3 ± 1.0	15.0 ± 1.0	16.0 ± 0.7	21.8 ± 1.4	22.5 ± 1.6	18.6 ± 0.6	20.3 ± 1.2
2010	21.3 ± 0.9	20.2 ± 0.4	24.7 ± 0.9	23.2 ± 0.9	14.3 ± 0.5	16.6 ± 0.5	14.6 ± 0.5	16.0 ± 0.5	15.7 ± 0.8	18.3 ± 1.5	16.7 ± 1.7	16.0 ± 1.1
2011	24.3 ± 1.1	22.2 ± 0.3	28.3 ± 0.7	25.1 ± 1.0	13.4 ± 0.4	15.5 ± 0.5	13.3 ± 0.5	15.2 ± 0.1	23.4 ± 0.8	27.1 ± 1.4	25.7 ± 2.9	22.0 ± 0.6
ANOVA <i>P</i> -values												
CO <sub>2</sub>	<0.0001				0.24				0.16			
W	<0.0001				<0.0001				0.32			
CO <sub>2</sub> × W	0.74				0.06				0.21			
Year	0.35				<0.0001				0.92			
CO <sub>2</sub> × year	0.37				0.87				0.51			
W × year	0.75				0.99				0.77			

Ct, ambient CO<sub>2</sub> and ambient temperature; cT, ambient CO<sub>2</sub> and elevated temperature; Ct, elevated CO<sub>2</sub> and ambient temperature; CT, elevated CO<sub>2</sub> and elevated temperature; W, warming treatment.

*P*-values are in bold when *P* < 0.05 and in italics when *P* < 0.1.

WFPS, when regressions were performed for each CO<sub>2</sub> and warming treatment (Fig. S3, Table S1). Interestingly, N<sub>2</sub>O fluxes were significantly negatively related to soil temperature (Fig. 3d), while this relationship was not affected by the CO<sub>2</sub> and warming treatments (Fig. S3, Table S1). However, it should be noted that WFPS and soil temperature were also significantly negatively correlated (*P* < 0.0001, *r* = 0.51). Plot greenness had no effect on N<sub>2</sub>O fluxes, but was significantly correlated with CH<sub>4</sub> fluxes (Fig. 3e and f) without differences among the CO<sub>2</sub> and warming treatments (Fig. S3, Table S1). Plot greenness was not correlated with WFPS (*P* > 0.1) but was positively related to soil temperature (*P* < 0.0001, *r* = 0.57).


This semiarid grassland was at all times a net sink regarding the combined cumulative GWP from CH<sub>4</sub> and N<sub>2</sub>O (Fig. 4). Both elevated CO<sub>2</sub> and warming significantly reduced this sink (*P* = 0.04 for elevated CO<sub>2</sub> and *P* = 0.002 for warming). The reduced net sink of CH<sub>4</sub> and N<sub>2</sub>O under elevated CO<sub>2</sub> increased with time (significant CO<sub>2</sub> × year interactive effect, *P* = 0.0003). Across years, elevated CO<sub>2</sub> and warming reduced the net sink for CH<sub>4</sub> and N<sub>2</sub>O by 7.2% and 11%, respectively, with the largest reduction of 17% when elevated CO<sub>2</sub> and warming were combined.

**Discussion**

This grassland was a sink for CH<sub>4</sub> at all times during the growing season, while it was sometimes a source and sometimes a sink for N<sub>2</sub>O (i.e., net N<sub>2</sub>O uptake). When CH<sub>4</sub> and N<sub>2</sub>O were combined in terms of CO<sub>2</sub> equivalents, this grassland was always a sink. Contrary to our hypothesis, both elevated CO<sub>2</sub> and warming reduced the net sink of CH<sub>4</sub> and N<sub>2</sub>O in this grassland, and the largest net sink reductions occurred when elevated CO<sub>2</sub> and warming were combined (Fig. 4). Combined effects of elevated CO<sub>2</sub> and warming resulted in a significant decrease in the net sink of CH<sub>4</sub> and N<sub>2</sub>O by 28% and 24% in 2010 and 2011 respectively. Our results suggest that this semiarid grassland causes a positive feedback to global warming by reducing the net sink of CH<sub>4</sub> and N<sub>2</sub>O.

*CH<sub>4</sub>*

Both elevated CO<sub>2</sub> and warming reduced CH<sub>4</sub> uptake in this semiarid grassland in most of the years; CH<sub>4</sub> uptake in plots exposed to elevated CO<sub>2</sub> and warming together was significantly lower than in the control plots in three of the 5 years (2008, 2010, and 2011, Fig. 1). The reduced CH<sub>4</sub> uptake under elevated CO<sub>2</sub> is in contrast to observations in a similar semiarid grassland where CH<sub>4</sub> uptake was not affected by elevated




**Fig. 3** Fluxes of CH<sub>4</sub> (a, c, e) and N<sub>2</sub>O (b, d, f) as a function of water filled pore space (WFPS, a, b), soil temperature (c, d) and plot greenness (e, f). Each data point is the average CH<sub>4</sub> or N<sub>2</sub>O flux, and average WFPS, soil temperature, or plot greenness measured across all treatments at a specific date between 2007 and 2011. Regression lines are only shown when significant ( $P < 0.05$ ).

CO<sub>2</sub> (Mosier *et al.*, 2002), but supports results from studies in mesic environments (Ineson *et al.*, 1998; Phillips *et al.*, 2001a; Dubbs & Whalen, 2010). On the other hand, the reduced CH<sub>4</sub> uptake with warming in this semiarid grassland differed from studies in wetter environments where no or increased CH<sub>4</sub> uptake was found in response to warming (Peterjohn *et al.*, 1994; Sjögersten & Wookey, 2002; Blankinship *et al.*, 2010).

Treatment effects on CH<sub>4</sub> uptake may have been mediated by their effect on soil moisture. The CH<sub>4</sub> flux

showed a bell-shaped relationship with WFPS with an optimum CH<sub>4</sub> uptake rate at 24% (Fig. 3a). Soil moisture consistently increased under elevated CO<sub>2</sub> and consistently decreased with warming (Table 2), which could have caused opposing effects on CH<sub>4</sub> uptake depending on what side of the curve the change in soil moisture occurred (Dijkstra *et al.*, 2011). A soil moisture increase under elevated CO<sub>2</sub> could increase CH<sub>4</sub> uptake (through stimulating methanotroph activity) when soils are relatively dry, or decrease CH<sub>4</sub> uptake (through



**Fig. 4** Cumulative Global Warming Potential (GWP) in g CO<sub>2</sub> eq. per m<sup>2</sup> in plots with plants in response to elevated CO<sub>2</sub> and warming. Treatments: ct: ambient CO<sub>2</sub> and ambient temperature; cT: ambient CO<sub>2</sub> and elevated temperature; Ct: elevated CO<sub>2</sub> and ambient temperature; CT: elevated CO<sub>2</sub> and elevated temperature. W: warming treatment. Error bars indicate  $\pm 1$  SE. ANOVA *P*-values are reported when *P* < 0.05 (in bold) or *P* < 0.1 (in italics). Different letters above bars indicate significant differences among the elevated CO<sub>2</sub> and warming treatments for each year separately (*P* < 0.05, Tukey's HSD test).

decreasing CH<sub>4</sub> diffusivity into the soil) when soils are relatively wet, while a soil moisture decrease with warming should have the opposite effect. In contrast, we observed reductions in CH<sub>4</sub> uptake with elevated CO<sub>2</sub> and warming that are hard to explain by the treatment effects on soil moisture alone. For instance, when elevated CO<sub>2</sub> and warming were combined, average WFPS was only slightly higher than in the control plots (Table 2) because of opposing effects of the two treatments on soil moisture (Morgan *et al.*, 2011). Yet, plots exposed to elevated CO<sub>2</sub> and warming (CT plots) often showed the largest reduction in CH<sub>4</sub> uptake. The relationship between CH<sub>4</sub> uptake and WFPS differed somewhat among the treatments (Fig. S3, Table S1), further suggesting that other drivers than soil moisture alone must have contributed to the reduction in CH<sub>4</sub> uptake under elevated CO<sub>2</sub> and warming.

Another driver that could have influenced treatment effects on CH<sub>4</sub> uptake is soil temperature. An increase in soil temperature could enhance methanotroph activity, but could also enhance methanogen activity that tends to be more responsive to temperature (Topp & Pattey, 1997). Both processes usually occur at the same time (Yavitt *et al.*, 1995; von Fischer & Hedin, 2007), and stimulation of both processes by soil temperature may have cancelled out each other's effect. We observed no relationship between CH<sub>4</sub> uptake and soil temperature when the data were aggregated across all treatments (Fig. 3c), but found similar bell-shaped relationships as observed between CH<sub>4</sub> uptake and WFPS (Fig. S3). Because of the strong correlation between

WFPS and soil temperature, the relationships between CH<sub>4</sub> uptake and soil temperature for each treatment may have been driven partially by soil moisture. Nevertheless, as with WFPS, the reduced CH<sub>4</sub> uptake with elevated CO<sub>2</sub> and warming combined are difficult to explain by treatment effects on soil temperature alone (note that elevated CO<sub>2</sub> had no effect on soil temperature, Table 2).

Treatment effects on N cycling may also have affected CH<sub>4</sub> uptake. Ammonium (NH<sub>4</sub><sup>+</sup>) can suppress CH<sub>4</sub> oxidation in the soil because some of the CH<sub>4</sub> oxidizing microbes switch to oxidizing NH<sub>4</sub><sup>+</sup> with increased availability of NH<sub>4</sub><sup>+</sup> (Stuedler *et al.*, 1989; Hanson & Hanson, 1996). Elevated CO<sub>2</sub> significantly increased NH<sub>4</sub><sup>+</sup> concentrations in our experiment (Carrillo *et al.*, 2012), which could have contributed to reduced CH<sub>4</sub> uptake that we observed in some years under elevated CO<sub>2</sub>. It has also been suggested that soil nitrate (NO<sub>3</sub><sup>-</sup>) could stimulate CH<sub>4</sub> uptake at low CH<sub>4</sub> concentrations due to shifts in the CH<sub>4</sub> oxidizing bacteria community (Jang *et al.*, 2011). We observed a significant reduction in soil NO<sub>3</sub><sup>-</sup> under elevated CO<sub>2</sub> (Carrillo *et al.*, 2012), suggesting that this too may have contributed to the reduced CH<sub>4</sub> uptake under elevated CO<sub>2</sub>, particularly with a progressively tighter N cycle under elevated CO<sub>2</sub> (Dijkstra *et al.*, 2010). Soil NO<sub>3</sub><sup>-</sup> was much higher in the subplots without plants (Carrillo *et al.*, 2012), suggesting that soil NO<sub>3</sub><sup>-</sup> may also have played a role in the higher CH<sub>4</sub> uptake without plants (Fig. 1).

## N<sub>2</sub>O

The N<sub>2</sub>O fluxes in this grassland were small (cumulative N<sub>2</sub>O never exceeded 2 mg N m<sup>-2</sup>), while both production and uptake of N<sub>2</sub>O occurred (Fig. 2a). Because of the small fluxes, elevated CO<sub>2</sub> and warming had no effect on N<sub>2</sub>O across all years. However, elevated CO<sub>2</sub> significantly reduced the N<sub>2</sub>O sink in the last 2 years. This is in contrast to studies conducted in a similar semiarid grassland where no elevated CO<sub>2</sub> effects on N<sub>2</sub>O fluxes were observed (Mosier *et al.*, 2002). In temperate grasslands that received N fertilization elevated CO<sub>2</sub> increased N<sub>2</sub>O emission (Ineson *et al.*, 1998; Baggs *et al.*, 2003; Kammann *et al.*, 2008). In these grasslands N fertilization may have alleviated N constraints, while at the same time increased supply of labile C under elevated CO<sub>2</sub> may have reduced C constraints on denitrification thereby causing the increase in N<sub>2</sub>O emission (Dijkstra & Morgan, 2012; Dijkstra *et al.*, 2012). On the other hand, in systems without N fertilization and where N availability is limiting denitrification, an increase in labile C under elevated CO<sub>2</sub> may not increase N<sub>2</sub>O emission. At our site, N<sub>2</sub>O emission



appeared to be limited by N availability rather than by C availability. The absence of plant N uptake in subplots without plants significantly increased N availability (Carrillo *et al.*, 2012), which may have caused the significant increase in N<sub>2</sub>O emission in these plots (Fig. 3). We found no relationship between plot greenness and N<sub>2</sub>O fluxes (Fig. 3f). Assuming that plot greenness correlates with photosynthetically active plant biomass, and therefore with plant input of labile C (Leake *et al.*, 2006), the lack of a relationship between plot greenness and N<sub>2</sub>O exchange suggests that seasonal variation in plant input of labile C had no effect on N<sub>2</sub>O fluxes (although plot greenness may have related poorly to labile C input into the soil at the time of measurement). Furthermore, we often observed uptake of N<sub>2</sub>O in the plots with plants, particularly during the last 2 years, suggesting low levels of available N in these plots. The process of N<sub>2</sub>O uptake under dry conditions is still little understood (Chapuis-lardy *et al.*, 2007), but dry soil conditions may have enhanced diffusion of atmospheric N<sub>2</sub>O into the soil where in the absence of NO<sub>3</sub><sup>-</sup>, N<sub>2</sub>O was used as the electron acceptor for denitrification (Stewart *et al.*, 2012). The positive relationship between WFPS and N<sub>2</sub>O flux (Fig. 3b) suggests that elevated CO<sub>2</sub>-induced increases in soil moisture may have reduced the diffusion of N<sub>2</sub>O into the soil and therefore reduced N<sub>2</sub>O uptake in the last 2 years.

Nitrous oxide fluxes were negatively related to soil temperature (Fig. 3d). Often N<sub>2</sub>O emissions increase with increased temperature (Dobbie & Smith, 2001; Mosier *et al.*, 2008). Most likely, N<sub>2</sub>O fluxes in our study were driven more by soil moisture than by temperature, as WFPS was significantly negatively correlated with soil temperature ( $P < 0.0001$ ,  $r = 0.51$ ). Warming had also no effect on the N<sub>2</sub>O flux. Possibly, the direct stimulatory effect of increased soil temperature with warming on the N<sub>2</sub>O flux was offset by the indirect inhibitory effect of reduced soil moisture with warming (McHale *et al.*, 1998; Bijoor *et al.*, 2008).

### Cumulative GWP

Both elevated CO<sub>2</sub> and warming significantly increased the cumulative GWP from CH<sub>4</sub> and N<sub>2</sub>O combined. The largest increase in the cumulative GWP occurred when elevated CO<sub>2</sub> and warming were combined, resulting in a significant increase in 2010 and 2011 of 1.6 and 1.4 g CO<sub>2</sub> eq. per m<sup>2</sup>, respectively, compared with the control plots (Fig. 4). Methane was the largest contributor to the increase in cumulative GWP in response to combined effects of elevated CO<sub>2</sub> and warming; 61% and 74% of the total increase in cumulative GWP in 2010 and 2011, respectively, was caused by reductions in CH<sub>4</sub> uptake. We hypothesized that

elevated CO<sub>2</sub> and warming would have opposing effects on the cumulative GWP from CH<sub>4</sub> and N<sub>2</sub>O because of their opposing effects on soil moisture (Morgan *et al.*, 2011). Because CH<sub>4</sub> was the dominant contributor to the cumulative GWP, and because average soil moisture contents were on average near optimum CH<sub>4</sub> uptake rates in control plots, both elevated CO<sub>2</sub>-induced increases and warming-induced decreases in soil moisture increased the cumulative GWP. Other drivers, such as soil N availability may also have played a role in causing synergistic rather than antagonistic effects of elevated CO<sub>2</sub> and warming on the net sink of CH<sub>4</sub> and N<sub>2</sub>O in this system.

Our results show that, when expressed in CO<sub>2</sub>-equivalents, both elevated CO<sub>2</sub> and warming reduced the net sink of CH<sub>4</sub> and N<sub>2</sub>O in this semiarid grassland. Elevated CO<sub>2</sub> and warming effects on this sink may be different in exceptionally dry or wet seasons (i.e., seasons with less than 242 mm observed in 2010 or more than 363 mm observed in 2011), or in other semiarid grasslands with different rainfall regimes. Nevertheless, when we extrapolate our results of the last two seasons (which were the driest and wettest season during the 5-year period that we measured) to the global land surface of semiarid grasslands (11% of the global land surface or 16 383 400 km<sup>2</sup>, Bailey, 1979), then between 22 and 26 Tg CO<sub>2</sub> eq. per yr less will be taken up as CH<sub>4</sub> and N<sub>2</sub>O in response to elevated CO<sub>2</sub> and temperature at levels that are predicted for the mid- to end of this century. Combined effects of elevated CO<sub>2</sub> and warming also caused some of the largest losses in soil C at our site (E. Pendall, J. L. Heisler-White, D. G. Williams, F. A. Dijkstra, Y. Carrillo, J. A. Morgan, D. R. LeCain, in review). These results together with our results suggest that semiarid grasslands cause an important positive feedback to climate change.

### Acknowledgements

We thank Erik Hardy, Valerie O'Neill, David Smith, Mary Smith, Katie Tylka, Megan Nix, and Ally Eden for technical assistance. This project was supported by the USDA-ARS Climate Change, Soils & Emissions and GRACEnet Programs, a USDA-CSREES Soil Processes Program (2008-35107-18655), the US Department of Energy's Office of Science (BER), and by NSF (DEB# 1021559). FAD acknowledges support from the Australian Research Council (FT100100779).

### References

- Baggs EM, Richter M, Hartwig UA, Cadisch G (2003) Nitrous oxide emissions from grass swards during the eighth year of elevated atmospheric pCO<sub>2</sub> (Swiss FACE). *Global Change Biology*, **9**, 1214–1222.
- Bailey HP (1979) Semiarid climates: their definition and distribution. In: *Agriculture in Semiarid Environments* (eds Hall AE, Cannell GH, Lawton HW), pp. 73–97. Springer-Verlag, New York.

- Bijoor NS, Czimeczik CI, Pataki DE, Billings SA (2008) Effects of temperature and fertilization on nitrogen cycling and community composition of an urban lawn. *Global Change Biology*, **14**, 2119–2131.
- Billings SA, Schaeffer SM, Evans RD (2002) Trace N gas losses and N mineralization in Mojave desert soils exposed to elevated CO<sub>2</sub>. *Soil Biology and Biochemistry*, **34**, 1777–1784.
- Blankinship JC, Brown JR, Dijkstra P, Hungate BA (2010) Effects of interactive global changes on methane uptake in an annual grassland. *Journal of Geophysical Research*, **115**, G02008.
- Booth DT, Cox SE, Berryman RD (2006) Point sampling digital imagery with 'Sample-point'. *Environmental Monitoring and Assessment*, **123**, 97–108.
- Carrillo Y, Dijkstra FA, Pendall E, Morgan JA, Blumenthal DM (2012) Controls over soil nitrogen pools in a semiarid grassland under elevated CO<sub>2</sub> and warming. *Ecosystems*, **15**, 761–774.
- Chapuis-lardy L, Wrage N, Metay A, Chotte JL, Bernoux M (2007) Soils, a sink for N<sub>2</sub>O? A review. *Global Change Biology*, **13**, 1–17.
- Davidson EA (1992) Sources of nitric oxide and nitrous oxide following wetting of dry soil. *Soil Science Society of America Journal*, **56**, 95–102.
- Del Grosso SJ, Parton WJ, Mosier AR *et al.* (2000) General CH<sub>4</sub> oxidation model and comparisons of CH<sub>4</sub> oxidation in natural and managed systems. *Global Biogeochemical Cycles*, **14**, 999–1019.
- Dijkstra FA, Morgan JA (2012) Elevated CO<sub>2</sub> and warming effects on soil carbon sequestration and greenhouse gas exchange in agroecosystems: a review. In *Managing Agricultural Greenhouse Gases* (eds Liebig MA, Follett RF, Franzluebbers AJ), pp. 467–486. Elsevier, Amsterdam.
- Dijkstra FA, Blumenthal D, Morgan JA, Pendall E, Carrillo Y, Follett RF (2010) Contrasting effects of elevated CO<sub>2</sub> and warming on nitrogen cycling in a semiarid grassland. *New Phytologist*, **187**, 426–437.
- Dijkstra FA, Morgan JA, Von Fischer JC, Follett RF (2011) Elevated CO<sub>2</sub> and warming effects on CH<sub>4</sub> uptake in a semiarid grassland below optimum soil moisture. *Journal of Geophysical Research - Biogeosciences*, **116**, G01007, doi: 10.1029/2010JG001288
- Dijkstra FA, Prior SA, Runion GB, Torbert HA, Tian H, Lu C, Venterea RT (2012) Effects of elevated CO<sub>2</sub> and warming on methane and nitrous oxide fluxes: evidence from field experiments. *Frontiers in Ecology and the Environment*, **10**, 520–527.
- Dobbie KE, Smith KA (2001) The effects of temperature, water-filled pore space and land use on N<sub>2</sub>O emissions from an imperfectly drained gleysol. *European Journal of Soil Science*, **52**, 667–673.
- Donoso L, Santana R, Sanhueza E (1993) Seasonal variation of N<sub>2</sub>O fluxes at a tropical savannah site: soil consumption of N<sub>2</sub>O during the dry season. *Geophysical Research Letters*, **20**, 1379–1382.
- Dubbs LL, Whalen SC (2010) Reduced net atmospheric CH<sub>4</sub> consumption is a sustained response to elevated CO<sub>2</sub> in a temperate forest. *Biology and Fertility of Soils*, **46**, 597–606.
- von Fischer JC, Hedin LO (2007) Controls on soil methane fluxes: tests of biophysical mechanisms using stable isotope tracers. *Global Biogeochemical Cycles*, **21**, doi: 10.1029/2006GB002687
- von Fischer JC, Butters G, Duchateau PC, Thelwell RJ, Siller R (2009) In situ measures of methanotroph activity in upland soils: a reaction-diffusion model and field observation of water stress. *Journal of Geophysical Research G: Biogeosciences*, **114**, G01015. doi:10.1029/2008JG000731.
- Forster P, Ramaswamy V, Artaxo P *et al.* (2007) Changes in atmospheric constituents and in radiative forcing. In *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL) pp. 130–234. Cambridge University Press, Cambridge.
- Goldberg SD, Gebauer G (2009) Drought turns a Central European Norway spruce forest soil from an N<sub>2</sub>O source to a transient N<sub>2</sub>O sink. *Global Change Biology*, **15**, 850–860.
- van Groenigen KJ, Osenberg CW, Hungate BA (2011) Increased soil emissions of potent greenhouse gases under increased atmospheric CO<sub>2</sub>. *Nature*, **475**, 214–216.
- Grover SPP, Livesley SJ, Hutley LB *et al.* (2012) Land use change and the impact on greenhouse gas exchange in north Australian savanna soils. *Biogeosciences*, **9**, 423–437.
- Hanson RS, Hanson TE (1996) Methanotrophic bacteria. *Microbiological Reviews*, **60**, 439–471.
- Hantschel RE, Kamp T, Beese F (1995) Increasing the soil temperature to study global warming effects on the soil nitrogen cycle in agroecosystems. *Journal of Biogeography*, **22**, 375–380.
- Holtgrieve GW, Jewett PK, Matson PA (2006) Variations in soil N cycling and trace gas emissions in wet tropical forests. *Oecologia*, **146**, 584–594.
- Hu Y, Chang X, Lin X *et al.* (2010) Effects of warming and grazing on N<sub>2</sub>O fluxes in an alpine meadow ecosystem on the Tibetan plateau. *Soil Biology and Biochemistry*, **42**, 944–952.
- Hungate BA, Lund CP, Pearson HL, Chapin FS III (1997) Elevated CO<sub>2</sub> and nutrient addition alter soil N cycling and N trace gas fluxes with early season wet-up in a California annual grassland. *Biogeochemistry*, **37**, 89–109.
- Hutchinson GL, Mosier AR (1981) Improved soil cover method for field measurement of nitrous oxide fluxes. *Soil Science Society of America Journal*, **45**, 311–316.
- Ineson P, Coward PA, Hartwig UA (1998) Soil gas fluxes of N<sub>2</sub>O, CH<sub>4</sub> and CO<sub>2</sub> beneath *Lolium perenne* under elevated CO<sub>2</sub>: the Swiss free air carbon dioxide enrichment experiment. *Plant and Soil*, **198**, 89–95.
- Jang I, Lee S, Zoh KD, Kang H (2011) Methane concentrations and methanotrophic community structure influence the response of soil methane oxidation to nitrogen content in a temperate forest. *Soil Biology and Biochemistry*, **43**, 620–627.
- Kammann C, Müller C, Grünhage L, Jäger H-J (2008) Elevated CO<sub>2</sub> stimulates N<sub>2</sub>O emissions in permanent grassland. *Soil Biology and Biochemistry*, **40**, 2194–2205.
- Kimball BA, Conley MM, Wang S, Lin X, Luo C, Morgan J, Smith D (2008) Infrared heater arrays for warming ecosystem field plots. *Global Change Biology*, **14**, 309–320.
- King GM (1997) Responses of atmospheric methane consumption by soils to global climate change. *Global Change Biology*, **3**, 351–362.
- Klemmedtsen L, Svensson BH, Rosswall T (1988) Relationships between soil moisture content and nitrous oxide production during nitrification and denitrification. *Biology and Fertility of Soils*, **6**, 106–111.
- Lam SK, Lin E, Norton R, Chen D (2011) The effect of increased atmospheric carbon dioxide concentration on emissions of nitrous oxide, carbon dioxide and methane from a wheat field in a semi-arid environment in northern China. *Soil Biology and Biochemistry*, **43**, 458–461.
- Larsen KS, Andresen LC, Beier C *et al.* (2011) Reduced N cycling in response to elevated CO<sub>2</sub>, warming, and drought in a Danish heathland: synthesizing results of the CLIMAITE project after two years of treatments. *Global Change Biology*, **17**, 1884–1899.
- Leake JR, Ostle NJ, Rangel-Castro JI, Johnson D (2006) Carbon fluxes from plants through soil organisms determined by field <sup>13</sup>C<sub>2</sub> pulse-labelling in an upland grassland. *Applied Soil Ecology*, **33**, 152–175.
- McHale PJ, Mitchell MJ, Bowles FP, Bowles PP (1998) Soil warming in a northern hardwood forest: trace gas fluxes and leaf litter decomposition. *Canadian Journal of Forest Research*, **28**, 1365–1372.
- Miglietta F, Hoosbeek MR, Foot J *et al.* (2001) Spatial and temporal performance of the miniFACE (Free Air CO<sub>2</sub> Enrichment) system on bog ecosystems in northern and central Europe. *Environmental Monitoring and Assessment*, **66**, 107–127.
- Morgan JA, Lecain DR, Pendall E *et al.* (2011) C<sub>4</sub> grasses prosper as carbon dioxide eliminates desiccation in warmed semi-arid grassland. *Nature*, **476**, 202–205.
- Mosier AR, Morgan JA, King JY, LeCain D, Milchunas DG (2002) Soil-atmosphere exchange of CH<sub>4</sub>, CO<sub>2</sub>, NO<sub>x</sub>, and N<sub>2</sub>O in the Colorado shortgrass steppe under elevated CO<sub>2</sub>. *Plant and Soil*, **240**, 201–211.
- Mosier AR, Parton WJ, Martin RE *et al.* (2008) Soil-atmosphere exchange of trace gases in the Colorado shortgrass steppe. In: *Ecology of the Shortgrass Steppe. A Long-Term Perspective* (eds Lauenroth WK, Burke IC), pp. 342–372. Oxford University Press, Oxford.
- Niboyet A, Brown JR, Dijkstra P *et al.* (2011) Global change could amplify fire effects on soil greenhouse gas emissions. *PLoS ONE*, **6**, e20105.
- Peterjohn WT, Melillo JM, Steudler PA, Newkirk KM, Bowles FP, Aber JD (1994) Responses of trace gas fluxes and N availability to experimentally elevated soil temperatures. *Ecological Applications*, **4**, 617–625.
- Phillips RL, Whalen SC, Schlesinger WH (2001a) Influence of atmospheric CO<sub>2</sub> enrichment on methane consumption in a temperate forest soil. *Global Change Biology*, **7**, 557–563.
- Phillips RL, Whalen SC, Schlesinger WH (2001b) Influence of atmospheric CO<sub>2</sub> enrichment on nitrous oxide flux in a temperate forest ecosystem. *Global Biogeochemical Cycles*, **15**, 741–752.
- Sjögersten S, Wookey PA (2002) Spatio-temporal variability and environmental controls of methane fluxes at the forest-tundra ecotone in the Fennoscandian mountains. *Global Change Biology*, **8**, 885–894.
- Steudler PA, Bowden RD, Melillo JM, Aber JD (1989) Influence of nitrogen fertilization on methane uptake in temperate forest soils. *Nature*, **341**, 314–316.
- Stewart KJ, Brummell ME, Farrell RE, Siciliano SD (2012) N<sub>2</sub>O flux from plant-soil systems in polar deserts switch between sources and sinks under different light conditions. *Soil Biology and Biochemistry*, **48**, 69–77.

Tian H, Lu C, Chen G *et al.* (2012) Contemporary and projected biogenic fluxes of methane and nitrous oxide in terrestrial ecosystems of North America. *Frontiers in Ecology and the Environment*, **10**, 528–536.

Topp E, Patey E (1997) Soils as sources and sinks for atmospheric methane. *Canadian Journal of Soil Science*, **77**, 167–178.

Xu-Ri, Prentice IC, Spahni R, Niu HS (2012) Modelling terrestrial nitrous oxide emissions and implications for climate feedback. *New Phytologist*, **196**, 472–488.

Yavitt JB, Fahey TJ, Simmons JA (1995) Methane and carbon dioxide dynamics in a northern hardwood ecosystem. *Soil Science Society of America Journal*, **59**, 796–804.

## Supporting Information

Additional Supporting Information may be found in the online version of this article:

**Figure S1.** Seasonal fluxes of CH<sub>4</sub> (a) and N<sub>2</sub>O (b) during the growing season in 2007–2011 in response to elevated CO<sub>2</sub> and warming. Negative values indicate fluxes from the atmosphere to the soil. Treatments: ct: ambient CO<sub>2</sub> and ambient temperature; cT: ambient CO<sub>2</sub> and elevated temperature; Ct: elevated CO<sub>2</sub> and ambient temperature; CT: elevated CO<sub>2</sub> and elevated temperature. Error bars indicate Tukey's HSD.

**Figure S2.** Seasonal water filled pore space (WFPS) (a), soil temperature (b), and plot greenness (c) during the growing season in 2007–2011 in response to elevated CO<sub>2</sub> and warming. Treatments: ct: ambient CO<sub>2</sub> and ambient temperature; cT: ambient CO<sub>2</sub> and elevated temperature; Ct: elevated CO<sub>2</sub> and ambient temperature; CT: elevated CO<sub>2</sub> and elevated temperature. Error bars indicate Tukey's HSD.

**Figure S3.** Fluxes of CH<sub>4</sub> (a, c, e) and N<sub>2</sub>O (b, d, f) as a function of water filled pore space (WFPS, a, b), soil temperature (c, d) and plot greenness (e, f). Each data point is the average CH<sub>4</sub> or N<sub>2</sub>O flux, and average WFPS, soil temperature, or plot greenness measured across all five replicates for each treatment at a specific date between 2007 and 2011. Treatments: ct: ambient CO<sub>2</sub> and ambient temperature; cT: ambient CO<sub>2</sub> and elevated temperature; Ct: elevated CO<sub>2</sub> and ambient temperature; CT: elevated CO<sub>2</sub> and elevated temperature. Regression lines are only shown when significant ( $P < 0.05$ , Table S1).

**Table S1.** Results from ANCOVAs with water filled pore space (WFPS), soil temperature (Temp), and plot greenness (Greenness) as covariates, and as independent variables in regression analyses to predict CH<sub>4</sub> and N<sub>2</sub>O fluxes for each treatment (ct: ambient CO<sub>2</sub> and ambient temperature; cT: ambient CO<sub>2</sub> and elevated temperature; Ct: elevated CO<sub>2</sub> and ambient temperature; CT: elevated CO<sub>2</sub> and elevated temperature).