

CORN ROOT INFLUENCE ON AUTOMATED MEASUREMENT OF SOIL CARBON DIOXIDE CONCENTRATIONS

D. Chen¹, J. A. E. Molina¹, C. E. Clapp², R. T. Venterea², and A. J. Palazzo³

Carbon dioxide (CO₂) production is a more desirable indicator of soil carbon (C) dynamics than CO₂ flux at the soil-air interface, which is significantly influenced by the gas-transport condition of the soil. Production of CO₂ can be computed from CO₂ concentrations if high-temporal measurements are made. Our objective was to design, implement, and test an automated CO₂ measurement system that requires low maintenance but provides high-temporal resolution of CO₂ concentrations in soil. The CO₂ sensors were located at different soil depths from 10 to 60 cm, with and without roots, to measure the effect of corn (*Zea mays*, L.) root activities on CO₂ concentrations over time. A direct comparison indicates that soil CO₂ measured with the automated measurement system represents CO₂ of soil surrounding the sensor. Computed CO₂ production was highest in the soils above 20 cm. A peak of soil CO₂ concentration occurred after each of the major rain events. The amplitude of the peaks decreased with depth. Differences between the CO₂ concentrations in the root and root-excluded soils were small between rainfall events and large at and after rain. Soil CO₂ concentration showed diurnal variations at the 10-, 20-, and 40-cm depths, whereas it was hardly detectable at the 60-cm depth. The automated CO₂ measurement system is a useful tool for gaining knowledge of CO₂ production in soils over time and across depth and of contributions from roots and bulk soil to total CO₂ production in soils. (Soil Science 2005;170:779-787)

Key words: CO₂ sensor, soil CO₂ respiration, corn root rhizosphere, CO₂ production.

SOIL respiration, which is estimated to be one order of magnitude higher than fossil fuel emission, is a significant component of the global C balance (Raich and Schlesinger, 1992). Several researchers have recognized that CO₂ production is a more desirable indicator of soil C dynamics than CO₂ flux at the soil-air interface, which is significantly influenced by soil gas-

transport conditions. Methods have been developed to compute CO₂ production using concentrations at various depths of soil (Davidson and Trumbore, 1995; De Jong and Schappert, 1972; Risk et al., 2002). One problem causing concern, however, is the introduction of computed negative CO₂ production, probably caused by the low-temporal resolution of the data (De Jong and Schappert, 1972; Risk et al., 2002).

Most researchers have determined CO₂ concentration in soil air by extracting soil gas samples and analyzing them by gas chromatography (GC) or other instruments (Burton and Beauchamp, 1994; Buyanovsky and Wagner, 1983; Davidson and Trumbore, 1995; De Jong and Schappert, 1972; Risk et al., 2002). Soil gas sampling is a difficult and cumbersome procedure that limits the frequency of CO₂ measurements. Gas sampling may also cause artificial

¹Department of Soil, Water and Climate, Borlaug Hall, 1991 Upper Buford Circle, University of Minnesota, St. Paul, MN 55108. Dr. Chen is corresponding author. Current address: California Department of Water Resources, P.O. Box 942836, Sacramento, CA 94236-0001. E-mail: dchen@water.ca.gov

²USDA-ARS and Department of Soil, Water, and Climate, University of Minnesota, St. Paul, MN 55108.

³USACOE, Cold Regions Research and Engineering Laboratory, Hanover, NH 03755.

Received Jan. 25, 2005; accepted May 17, 2005.

DOI: 10.1097/01.ss.0000190512.41298.fc

airflow within the soil, depending on how much and how often sampling is carried out. Soil CO₂ concentrations can change rapidly with events such as rainfall or tillage, and are subject to both diurnal and seasonal variations. Data collected weekly or even biweekly may not provide an accurate and dynamic measure of soil CO₂ concentration. Recently, however, newer *in situ* measurements of soil CO₂ have been reported. Hirano et al., (2003), Tang et al., (2003), and Jassal et al. (2004) used buried infrared gas analyzers and solid-state CO₂ sensors to measure CO₂ concentration every 30 minutes, 30 seconds, and hourly, respectively, and provided detailed variations of soil CO₂ concentration at multiple depths.

Carbon dioxide production should be highest in the upper soil layers where organic matter accumulates and most of the metabolic activity is located. Yet, studies have shown that CO₂ concentration generally increases with depth (e.g., Davidson and Trumbore, 1995; De Jong and Schappert, 1972; Reardon et al., 1979). However, not all studies agreed on the origin of the CO₂ in the deeper soil. Some studies suggest that the origin is surface soils (Reardon et al., 1979; Risk et al., 2002) and others imply deeper soils (Davidson and Trumbore, 1995; De Jong and Schappert, 1972; Goulden et al., 1998). Seasonal variations of CO₂ concentration can penetrate into the deep soils and may also involve exchanging CO₂ between the vadose zone and groundwater (Reardon et al., 1979). It is also known that soil CO₂ concentrations undergo large changes after rains (Buyanovsky and Wagner, 1983).

Carbon dioxide is generated due to both live root activities and microbial activities. As roots penetrate the soil, they produce large amounts of organic C which could be another source of CO₂ production due to root activities in the rhizosphere, in addition to live root respiration (Bolinder et al., 1999). The contribution of root respiration to total soil respiration, estimated with various methods, varies from below 10% to more than 90% (Hanson et al., 2000). CO₂ production increased faster with increase of soil temperature in rooted soil than in root-excluded soil (Boone et al., 1998).

Our objective was to design, implement, and test an automated CO₂ measurement system that requires low maintenance but provides high-temporal resolution of CO₂ concentrations in soil. Several goals were kept in mind in designing the system. One was to minimize soil

disturbance during installation of the system in the field. Another was to allow for an easy and rapid replacement of the sensors in case of malfunction. The sensors were installed at different depths, in soils with and without roots, to document the effect of corn (*Zea mays* L.) root activity on CO₂ concentrations.

MATERIALS AND METHODS

Experimental Site

All experiments were carried out in two 6 × 9 m corn plots, plots III-7 and IV-5, of a long-term continuous corn field study started in 1980 at the University of Minnesota Research and Outreach Center, Rosemount, Minnesota (44 degrees 45' N, 93 degrees 04' W) (Clapp et al., 2000; Clay et al., 1990; Clay et al., 1985; Layese et al., 2002). The soil is a Waukegan silt loam (fine-silty over sandy or sandy-skeletal, mixed, superactive, mesic Typic Hapludoll) with an underlying layer of gravel at about 90 cm below the soil surface. The climate of the region is subhumid. The weather station located at the Research Center provides air temperature and precipitation data. Thermistors (107 Temperature Probe, Campbell Scientific, Inc., Logan, UT) buried at 10, 20, 40, and 60 cm depth provide soil temperature every 30 minutes. Corn was planted on day of the year (DOY) 136. Ammonium sulfate was broadcast at a rate of 150 kg N ha⁻¹ one week after planting. Row-spacing was 75 cm and planting density was 80,000 ha⁻¹. At the end of the growing season, corn grain and stover were harvested and removed from the plots; the soil was then rototilled to a depth of about 15 cm.

CO₂ Probes

Carbon dioxide concentrations were measured with CO₂ transmitters (model GMT220, Vaisala, Inc., Woburn, MA), consisting of a sensor, a box containing voltage output terminals, and a cable connecting the sensor to the box. The voltage output terminals were connected to a datalogger (model CR10X, Campbell Scientific, Inc., Logan, UT), which was programmed to read and record the voltage every 30 minutes. The sensor is a cylinder 1.85 cm in diameter and 10.9 cm in length (Fig. 1). Each sensor was placed at the bottom of a PVC pipe (3.7 cm i.d.), which was pushed into the void space left by removal of a soil core 8 months before CO₂ concentration measurements were started on DOY 183. The pipe




Fig. 1. Diagram of the CO₂ sensor unit in the PVC pipe.

was sealed at both ends with rubber stoppers. The pipe was further divided into upper and lower chambers by a donut-shaped rubber septum and its supporting plastic disk. Silicone sealant was added on top of the supporting disk to ensure no leaks between the upper and lower chamber. Only the tip of the sensor, where CO₂ was actually measured, was enclosed in the lower chamber. The soil gas diffused into the lower chamber through holes drilled in the wall of the pipe. The holes were covered with waterproof breathable fabric to prevent liquid water from getting into the PVC pipe. The empty space of the lower chamber was kept as small as possible to minimize differences between the soil and chamber CO₂ concentrations. We confirmed no leakage between the upper and lower chambers by introducing a few pieces of dry ice into the upper chamber and found no increase in CO₂ in the lower chamber. When replacing a sensor is required, only the sensor, septum and supporting disk are pulled out, and the PVC pipe stays in the soil. It takes <20 minutes to replace a problem sensor without disturbing the soil and creating a gap in data collection.

The transmitters were installed both in root-active (RA) and between root-excluded (RE) corn rows at 10-, 20-, 40-, and 60-cm depths, for a total of 8 transmitters in each plot. The between-row sensors in their PVC pipes were

set inside larger PVC pipes (20.2 cm i.d.). The bottom edge of the large PVC pipe was sharpened and pushed into the ground with a tractor bucket loader. The upper rim of the larger PVC pipe was flush with the soil surface. The bottom of the larger PVC pipes were set 20 cm deeper than the bottom of the small PVC pipes. Care was taken to prevent weed growth within the large pipe.

To confirm that the air inside the sensor was representative of the surrounding soil, paired gas samples were taken from the lower chamber and from soil at the same depth as the sensor 5 to 15 cm from the pipe. Soil gas samples were taken through 0.16 cm i.d. Teflon tubing inserted in the soil. The same type of tubing had been inserted into the 3.7 cm i.d. PVC pipe when the CO₂ sensor was installed. The gas samples were taken weekly over a period of 5 weeks. These samples were analyzed for CO₂ by GC against standard concentrations. The GC is equipped with a thermal conductivity detector. How closely the CO₂ concentration in the bulk soil is represented by the CO₂ concentration measured inside the PVC pipe was assessed by the root mean square error (RMSE):

$$\text{RMSE} = \sqrt{\frac{1}{K} \sum_{i=1}^K (S_i - P_i)^2} \quad (1)$$

where S_i and P_i are CO₂ concentrations in the soil and inside the PVC pipe, respectively, and K is the total pairs of measurements.

CO₂ Production

Production of CO₂ was computed in the RA soil of plot III-7 from DOY 222 to 228, during water redistribution after a rainfall on DOY 218. On DOYs 223 and 226, the decrease in water content during water distribution was determined gravimetrically in soil samples taken at 10 cm intervals from the soil surface to the 60-cm depth. Water contents between DOYs 222 and 228 were estimated by linear extrapolation.

Production of CO₂ in soil can be calculated by rearranging the one-dimensional CO₂ transport equation of Šimůnek and Suarez (1993):

$$S = \frac{\partial(\varepsilon C_G + \theta C_L)}{\partial t} - \frac{\partial}{\partial z} \left(D_G(\theta) \frac{\partial C_G}{\partial z} + D_L(\theta) \frac{\partial C_L}{\partial z} \right) \quad (2)$$

where S is the CO₂ source/sink term (g cm⁻³ h⁻¹), t is time (h), z is soil depth (cm), ε and θ are soil air and water contents (cm³ cm⁻³),

respectively, C_G and C_L are CO_2 concentrations in soil air and water (g cm^{-3}), respectively, and D_G and D_L are diffusivity of CO_2 in soil air and water ($\text{cm}^2 \text{h}^{-1}$), respectively. Convective fluxes are ignored in the above equation. C_L can be related to C_G through Henry's law, $C_L = K_H \cdot C_G$, where K_H is Henry's law constant in a CO_2 - H_2O system (dimensionless) (Plummer and Busenberg, 1982). Substituting C_G for C_L in Eq.(2), and defining retardation factor $R = \varepsilon + \theta \cdot K_H$ and effective diffusivity $D = D_G + D_L \cdot K_H$, Eq.(2) becomes:

$$S = R \frac{\partial C_G}{\partial t} - \frac{\partial}{\partial z} (D \frac{\partial C_G}{\partial z}) \quad (3)$$

In the calculation, time is divided into steps at times t^j ($j = 1, 2, 3, \dots$) when C_G is measured and the soil depth is divided into layers at depths z_i ($i = 1, 2, 3, \dots$) where C_G is measured. C_G measured at depth z_i and time t^j is noted as C_G^{ij} . Other parameters, S , R , and D , are similarly noted. Carbon dioxide production in soil is then approximated:

$$S_i^j = \frac{R_i^{j+1} + R_i^j C_i^{j+1} - C_i^j}{2} \frac{2}{t^{j+1} - t^j} - \frac{2}{z_{i+1} - z_{i-1}} \left(D_{i+1/2}^j \frac{C_{i+1}^j - C_i^j}{z_{i+1} - z_i} - D_{i-1/2}^j \frac{C_i^j - C_{i-1}^j}{z_i - z_{i-1}} \right) \quad (4)$$

where C_G is written as C for simplification.

Diffusivities of CO_2 in soil air and water are calculated using the Millington and Quirk (1961) equation:

$$\frac{D}{D_0} = \frac{\lambda^{10/3}}{\Phi^2} \quad (5)$$

where Φ is soil porosity. When Eq.(5) is used to calculate CO_2 diffusivity in soil air, $D = D_G$, $\lambda = \varepsilon$, D_0 is CO_2 diffusivity in free air (no soil), and its values at different temperatures can be calculated with the equation of Fuller et al. (1966). For calculating diffusivity in soil water, $D = D_L$, $\lambda = \theta$, D_0 is CO_2 diffusivity in free water (no soil), and its values at different temperatures can be calculated with the equation of Bird et al. (1960) (p. 515).

To minimize noise of measured CO_2 concentrations being amplified in the calculation of CO_2 production, the original measured data were smoothed with a Savitzky-Golay digital filter, which smoothes noisy data but preserves peak height and widths (Press et al., 1992; Savitzky and Golay, 1964):

$$X_k = \left(\sum_{n=-m}^{n=m} B_n \cdot x_{k+n} \right) / N \quad (6)$$

where x_{k+n} ($n = -m$ to m) are members of a subset of original series of values x_o ($o = 1, 2, 3, \dots, M$, and M is total number of members in the original series). For our data, x_o is time series of measured soil CO_2 concentrations. X_k is the smoothed value. The coefficients of B_n ($n = -m$ to m) are convoluting integers, and the denominator N is a normalizing factor. B_n and N vary, depending on degrees of polynomial and total points ($2m + 1$) used in deriving Eq.(6). Savitzky and Golay (1964) had listed B_n and N for up to quintic polynomials and $m = 2$ to 12 points of original values. Applying Eq.(6) repeatedly for $k = m + 1$ to $M - m$, a new, or smoothed, series of values is constructed. We used the B_n of a cubic polynomial and $m = 8$ to smooth the measured CO_2 concentrations.

RESULTS

A comparison of CO_2 concentrations inside the lower chamber and in the surrounding soil showed no discernible bias at lower concentrations (Fig. 2). The root mean square error (RMSE) of CO_2 concentration between soil gas samples taken from the bulk soil and inside the PVC pipe is about 830 ppm. At higher CO_2 concentrations, gas samples taken from the soil tended to be lower than the concentration of samples taken inside the lower chamber, possibly due to the reduced gas diffusivity in deep and wet soil where the concentrations tended to be higher. Overall, the soil CO_2 concentrations in the lower chamber were very close to the concentrations in the adjacent soil.

The time series of CO_2 concentration from DOY 183 to 258 in RA soils and RE soils of plots III-7 and IV-5 are shown in Fig. 3. Soil CO_2 concentrations showed an increase with depth, surges after rains, and diurnal variations. A peak of soil CO_2 concentration occurred at each of the major rain events. The CO_2 surge began immediately after the rain at the 10- and 20-cm depths but was delayed for several hours at the 40- and 60-cm depths. The amplitude of the peaks decreased with depth. The overall soil CO_2 concentrations gradually declined over the entire study period. The downward trend may be caused by the gradual decline in soil water content during a dry summer. Differences between the CO_2 concentrations in RA and




Fig. 2. Comparison of CO₂ concentrations by gas chromatography inside lower chamber and its surrounding soil.

RE soils were small between rain events and greater during and after rain. After a rain, CO₂ concentrations of RA soils at the 10 and 20 cm depths increased faster and to a higher peak concentration than those of the RE soils.

Soil CO₂ concentrations showed diurnal variations at the 10-, 20-, and 40-cm depths, whereas similar variations were hardly detectable at the 60-cm depth except for the early days of RE soil of plot IV-5 (Fig. 3 and 4). Soil CO₂ concentrations and temperature were in phase at the 10- and 20-cm depths and out of phase at 40-cm. Soil temperatures peaked a little earlier at the 10-cm depth than at the 20-cm depth, as did CO₂. At the 40-cm depth, CO₂ peaked several hours earlier than did temperature. The timings of maximum and minimum CO₂ concentration varied by about 2 hours on different days.

The computed CO₂ production in the RA soil of plot III-7 from DOY 222 to 228 during water redistribution was maximum at the 10-cm depth, and minimal at 40-cm, although higher CO₂ concentrations were found in the deeper soil (Fig. 5). Average CO₂ production rates were 2.2, 0.56, and 0.22 g m⁻³ hr⁻¹ at depths of 10, 20, and 40 cm, respectively. Production of CO₂ also showed diurnal variations.

DISCUSSION

The peaks of CO₂ concentration were likely due to the sealing of surface soil by rain. The

peaks were higher in the RA soil than in the RE soil. Rates of CO₂ respiration are usually higher in soils with plant roots (Boone et al., 1998). Corn roots feed the microbial population in the rhizosphere with exuded nutrients and sloughed-off material, which are rapidly decomposed with attendant production of CO₂ (Molina et al., 2005; Molina et al., 2001; Paterson, 2003). In addition, live roots respire and contribute to total CO₂ production. Thus, rates of CO₂ respiration in the RA soil should have been higher than in the RE soil. This increased activity coupled with a reduced CO₂ diffusion in the upper soil layers saturated with water caused the observed higher peaks of CO₂ concentrations in the RA soil. Subsequently, as the water was redistributed in the soil profile, gas diffusion increased and differences in the CO₂ concentrations between the RA and RE soils were less pronounced.

Jassal et al. (2004) had suggested that the large increase in CO₂ concentration at the 20-cm depth after rainfall in a Douglas fir stand was the result of a sudden flush in soil microbial activity with a small increase in soil water content in a dry soil. We suggest that this mechanism did not apply to our corn field because our data indicated that soil CO₂ concentration increased drastically whether soil was dry or wet before rain. Lee et al. (2004) reported that a sudden increase in moisture content in the litter layer in a mixed species forest

caused an instant increase in CO_2 release. This is unlikely the mechanism that caused the large increase of soil CO_2 concentration in our experiment because corn stover of previous years had been removed from the plots. The CO_2 production rate varies with change of soil moisture content, especially in the dry state (Lee et al., 2004). The CO_2 peaks observed in our plots may be the result of production rate increase and sealing of surface soil by rain, but the main cause is likely to be the sealing of surface soil by rain.

Computed CO_2 production in the RA soil was highest in the upper soil layers and was similar in magnitude to that observed in virgin prairie soil at 10 cm (De Jong and Schappert, 1972). Measured soil CO_2 concentration decreased over time due to the increase in soil gas diffusivity in the period in which soil moisture was continuously decreasing after rain-

fall; however, the computed CO_2 production held at a relatively steady rate except the diurnal variation (Fig. 5). Soil CO_2 concentration does not necessarily reflect the soil CO_2 production.

Diurnal variations of soil CO_2 concentrations have been reported by Makarov (1958) where maximum CO_2 output occurred at mid-day and 3 pm. Edwards and Riggs (2003) used an automated monitoring chamber system that showed detailed diurnal variations in CO_2 efflux at the soil surface. De Jong and Schappert (1972) reported a diurnal variation in CO_2 concentration at a 15-cm depth, but negligible change at a 30-cm depth. They suggested that the effect of diurnal variations on soil respiration can be accounted for by always sampling at the same time of day. Various correlations have also been established between the diurnal variations of CO_2 concentration and temperature. Tang et al. (2003) observed that soil CO_2 concentrations at




Fig. 3. Soil CO_2 concentrations at various depths, in root-active (RA) and root-excluded (RE) soils of plots III-7 and IV-5. A, Precipitation; B, CO_2 concentration in RA soils; C, CO_2 concentration in RE soils. Numbers in the graphs are soil depths.




Fig. 4. Diurnal variations of the soil CO₂ concentration (A) and temperature (B). The numbers in the graphs are soil depths. In (A): the dots are measured values and lines are smoothed values using the Savitzky-Golay filter.

depths of 8 and 16 cm in oak-grass savanna peaked at a time around 14:30 to 16:30 when soil temperatures were highest. In a cool-temperate deciduous broadleaf forest, CO₂ concentrations reached a minimum in early afternoon (Hirano et al., 2003). In a boreal forest, soil CO₂ concentrations were out of phase with temperature (Hirsch et al., 2003). Jassal et al. (2004) reported that in a 54-year-old Douglas fir stand diurnal variation of CO₂ efflux followed variation of soil temperature at the 2-cm depth. The different timings suggest different driving forces behind the CO₂ diurnal variations. When CO₂ is in phase with temperature, the driving force is temperature (Tang et al., 2003), provided that variation in gas transport conditions during the day is small. When CO₂ is not in phase with temperature, different forces are involved. Hirano et al. (2003) suggest that CO₂ variation in the trunk space between a forest floor and canopy is the driving force. Hirsch et al. (2003) suggest that wind speed changes between morning and evening could be the driving force.

In our experiment, diurnal variations at 10 and 20 cm are in phase with diurnal variation of soil temperature; variation at 40 cm is out of phase with soil temperature. It can be seen from Fig. 4 that peaks and valleys of CO₂ concentration at 40 cm follow those at 20 cm, therefore, it was likely that CO₂ diurnal variation at the 40-cm depth is driven by CO₂ variation in soils above it.

In conclusion, CO₂ concentrations measured by the automated CO₂ measurement system represent those of adjacent soils. The high-temporal resolution of CO₂ concentration at multiple depths permits calculation of CO₂ production in soil. CO₂ production provides the information whereby CO₂ is generated in the soil and excludes the influence of gas-transport conditions. It will be advantageous, in predicting effects of temperature increase on soil respiration, to correlate soil temperature with CO₂ production instead of CO₂ flux at the soil-air interface which is also influenced by gas-transport conditions of the soil. CO₂ flux at the soil-air interface provides information of the




Fig. 5. Comparison of the CO₂ concentration (A) and production (B). The numbers in the graphs are soil depths. In (A): Dots are measured values and lines are smoothed values using the Savitzky-Golay filter.

interaction between soil and atmosphere, but less detailed information of CO₂ generation below ground. Combined measurements of both, flux and production, would provide a more complete understanding of CO₂ generation and transport in soil. Most of the CO₂ in our field was produced in the surface soil and the CO₂ production decreased with depth. Corn roots contributed significantly to CO₂ production in soil that is characterized by a faster increase and a higher peak of soil CO₂ concentration during and after rain events in RA soil than in RE soil. Soil CO₂ concentration changed dramatically during and after rain events. Carbon dioxide diurnal variation of surface soil in the field was caused by diurnal variation of soil temperature, whereas CO₂ variation at 40 cm was probably driven by CO₂ variation above it. For future work, non-destructive methods of determining soil water content, such as time domain reflectometry, could be used to provide long-term and frequent measurements of soil water content for both RA and RE soils. Comparison of CO₂

productions of RA and RE soils over the long-term can certainly be achieved.

ACKNOWLEDGMENTS

This research was partially supported by a grant from BARD, the USA-Israel Binational Agricultural Research Foundation. We are grateful for the assistance of Ryan Vial and Daniel Lee in the field and laboratory phases of this study.

REFERENCES

- Bird, R. B., W. E. Stewaet, and E. N. Lightfoot. 1960. Transport phenomena. John Wiley & Sons, New York.
- Bolinder, M. A., D. A. Angers, M. Giroux, and M. R. Laverdière. 1999. Estimating C inputs retained as soil organic matter from corn (*Zea mays* L.). *Plant Soil*. 215:85-91.
- Boone, R. D., K. J. Nadelhoffer, J. D. Canary, and J. P. Kaye. 1998. Roots exert a strong influence on the temperature sensitivity of soil respiration. *Nature*. 396:570-572.
- Burton, D. L., and E. G. Beauchamp. 1994. Profile nitrous oxide and carbon dioxide concentrations

- in a soil subject to freezing. *Soil Sci. Soc. Am. J.* 58:115–122.
- Buyanovsky, G. A., and G. H. Wagner. 1983. Annual cycles of carbon dioxide level in soil air. *Soil Sci. Soc. Am. J.* 47:1139–1145.
- Clapp, C. E., R. R. Allmaras, M. F. Layese, D. R. Linden, and R. H. Dowdy. 2000. Soil organic carbon and ¹³C abundance as related to tillage, crop residue, and nitrogen fertilization under continuous corn management in Minnesota. *Soil Tillage Res.* 55:127–142.
- Clay, D. E., C. E. Clapp, J. A. E. Molina, and R. H. Dowdy. 1990. Influence of nitrogen fertilization, tillage, and residue management on a soil nitrogen mineralization index. *Commun. Soil Sci. Plant Anal.* 21:323–335.
- Clay, D. E., C. E. Clapp, J. A. E. Molina, and D. R. Linden. 1985. Nitrogen-tillage-residue management I. Simulating soil and plant behavior by the model NCSWAP. *Plant Soil.* 84:67–77.
- Davidson, E. A., and S. E. Trumbore. 1995. Gas diffusivity and production of CO₂ in deep soils of the eastern Amazon. *Tellus.* 47B:550–565.
- De Jong, E., and H. J. V. Schappert. 1972. Calculation of soil respiration and activity from CO₂ profiles in the soil. *Soil Sci.* 113:328–333.
- Edwards, N. T., and J. S. Riggs. 2003. Automated monitoring of soil respiration: A moving chamber design. *Soil Sci. Soc. Am. J.* 67:1266–1271.
- Fuller, E. N., P. D. Schettler, and J. C. Giddings. 1966. A new method for prediction of binary gas-phase diffusion coefficients. *Indust. Eng. Chem.* 58:19–27.
- Goulden, M. L., S. C. Wofsy, J. W. Harden, S. E. Trumbore, P. M. Crill, S. T. Gower, T. Fries, B. C. Daube, S.-M. Fan, D. J. Sutton, A. Bazzaz, and J. W. Munger. 1998. Sensitivity of boreal forest carbon balance to soil thaw. *Science* 279:214–217.
- Hanson, P. J., N. T. Edwards, C. T. Garten, and J. A. Andrews. 2000. Separating root and soil microbial contributions to soil respiration: A review of methods and observation. *Biogeochemistry.* 48:115–146.
- Hirano, T., H. Kim, and Y. Tanaka. 2003. Long-term half-hourly measurement of soil CO₂ concentration and soil respiration in a temperate deciduous forest. *J. Geophys. Res.* 108:7–1 to 7–13.
- Hirsch, A. I., S. E. Trumbore, and M. L. Goulden. 2003. Direct measurement of the deep soil respiration accompanying seasonal thawing of a boreal forest soil. *J. Geophys. Res.* 108:2–1 to 2–10.
- Jassal, R. S., T. A. Black, G. B. Drewitt, M. D. Novak, D. Gaumont-Guay, and Z. Nescic. 2004. A model of the production and transport of CO₂ in soil: predicting soil CO₂ concentrations and CO₂ efflux from a forest floor. *Agric. For. Meteorol.* 124:219–236.
- Layese, M. F., C. E. Clapp, R. R. Allmaras, D. R. Linden, S. M. Copeland, J. A. E. Molina, and R. H. Dowdy. 2002. Current and relic carbon using natural abundance carbon-13. *Soil Sci.* 167:315–326.
- Lee, X., H. J. Wu, J. Sigler, C. Oishi, and T. Siccama. 2004. Rapid and transient response of soil respiration to rain. *Global Change Biol.* 10:1017–1026.
- Makarov, B. N. 1958. Diurnal variation in soil respiration and in carbon-dioxide content of the layer of air next to the soil. *Soils Fert.* 21:168.
- Millington, R. J., and J. P. Quirk. 1961. Permeability of porous solids. *Trans. Faraday Soc.* 57:1200–1207.
- Molina, J. A. E., C. E. Clapp, R. R. Allmaras, and M. F. Layese. 2005. Simulation of nitrogen rhizodeposition and assimilation back into (*Zea mays* L.) roots. *Soil Biol. Biochem.* 37:93–100.
- Molina, J. A. E., C. E. Clapp, D. R. Linden, R. R. Allmaras, M. F. Layese, R. H. Dowdy, and H. H. Cheng. 2001. Modeling the incorporation of corn (*Zea mays* L.) carbon from roots and rhizodeposition into soil organic matter. *Soil Biol. Biochem.* 33:83–92.
- Paterson, E. 2003. Importance of rhizodeposition in the coupling of plant and microbial productivity. *Eur. J. Soil Sci.* 54:741–750.
- Plummer, L. N., and E. Busenberg. 1982. The solubilities of calcite, aragonite and vaterite in CO₂-H₂O solutions between 0 and 90 °C, and an evaluation of the aqueous model for the system CaCO₃-CO₂-H₂O. *Geochim. Cosmochim. Acta.* 46:1011–1040.
- Press, W. H., S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery. 1992. *Numerical recipes in FORTRAN: The art of scientific computing*. Cambridge University Press, Cambridge, England.
- Raich, J. W., and W. H. Schlesinger. 1992. The global carbon dioxide flux in soil respiration and its relationship to vegetation and climate. *Tellus.* 44B:81–99.
- Reardon, E. J., G. B. Allison, and P. Fritz. 1979. Seasonal chemical and isotopic variations of soil CO₂ at Trout Creek, Ontario. *J. Hydrol.* 43:355–371.
- Risk, D., L. Kellman, and H. Beltrami. 2002. Soil CO₂ production and surface flux at four climate observatories in eastern Canada. *Global Biogeochem. Cycles.* 16:69–1 to 69–12.
- Savitzky, A., and M. J. E. Golay. 1964. Smoothing and differentiation of data by simplified least squares procedures. *Anal. Chem.* 36:1627–1639.
- Šimůnek, J., and D. L. Suarez. Modeling of carbon dioxide transport and production in soil, I: Model development. *Water Resour. Res.* 29:487–497, 1993.
- Tang, J., D. D. Baldocchi, Y. Qi, and L. Xu. 2003. Assessing soil CO₂ efflux using continuous measurements of CO₂ profiles in soils with small solid-state sensors. *Agric. For. Meteorol.* 118:207–220.