

Will extreme climatic events facilitate biological invasions?

Jeffrey M Diez^{1*}, Carla M D'Antonio², Jeffrey S Dukes³, Edwin D Grosholz⁴, Julian D Olden⁵, Cascade JB Sorte⁶, Dana M Blumenthal⁷, Bethany A Bradley⁸, Regan Early⁹, Inés Ibáñez¹⁰, Sierra J Jones¹¹, Joshua J Lawler¹², and Luke P Miller¹³

Extreme climatic events (ECEs) – such as unusual heat waves, hurricanes, floods, and droughts – can dramatically affect ecological and evolutionary processes, and these events are projected to become more frequent and more intense with ongoing climate change. However, the implications of ECEs for biological invasions remain poorly understood. Using concepts and empirical evidence from invasion ecology, we identify mechanisms by which ECEs may influence the invasion process, from initial introduction through establishment and spread. We summarize how ECEs can enhance invasions by promoting the transport of propagules into new regions, by decreasing the resistance of native communities to establishment, and also sometimes by putting existing non-native species at a competitive disadvantage. Finally, we outline priority research areas and management approaches for anticipating future risks of unwanted invasions following ECEs. Given predicted increases in both ECE occurrence and rates of species introductions around the globe during the coming decades, there is an urgent need to understand how these two processes interact to affect ecosystem composition and functioning.

Front Ecol Environ 2012; 10(5): 249–257, doi:10.1890/110137 (published online 21 May 2012)

Ecosystems are experiencing not only gradual shifts in mean climate conditions but also dramatic changes in climate variability and prevalence of extreme climatic events (ECEs). ECEs such as droughts, floods, severe storms, and heat waves are changing in frequency, magnitude, timing, and duration, depending on the region and the specific climate event (WebPanel 1; Easterling *et al.* 2000; Karl *et al.* 2008). These extreme events can strongly influence terrestrial and aquatic ecosystems (Parmesan *et al.* 2000; Thibault and Brown 2008) and may combine synergistically with other agents of environmental change, such as species invasions, to dramatically change ecosystems.

Previous studies have shown that changes in broad climatic conditions may influence the probability of species

invasions, while highlighting that the effects of changing climate are likely to be diverse and context-dependent (Rahel and Olden 2008; Walther *et al.* 2009; Bradley *et al.* 2010). However, the potential for ECEs in particular to promote species invasions has not been systematically assessed. Here, we use theory from community ecology and invasion biology to identify mechanisms and pathways by which ECEs may affect the establishment and spread of introduced species in recipient ecosystems. We then review current evidence for ECE impacts on invasions and assess the potential for changes in the frequency and magnitude of ECEs to affect species introductions in the future. By focusing on mechanisms of invasion following an ECE, we assess whether generalizations about invasion risk are possible across species, ecosystems, and ECE type. Finally, we highlight critical areas of research necessary for a better understanding of these processes, and suggest strategies for anticipating and subsequently managing invasions associated with ECEs.

In a nutshell:

- Extreme climatic events, such as intense heat waves, hurricanes, floods, and droughts, may facilitate biological invasions, leading to new science, management, and policy challenges
- These events often influence invasions through increased movements of non-native species and decreased biotic resistance of native communities to invader establishment, but specific outcomes depend on the ecosystem and type of event
- Efforts to minimize impacts of invasive species in a changing climate must include preparation for extreme events

■ What is an extreme climatic event?

Climatic events can be defined as “extreme” according to two broad perspectives. First, extreme events are episodes that fall within the statistical tails of a climate parameter's historical range; a commonly used threshold is the most extreme 1% of annual values, equivalent to an annual event happening once per 100 years (eg a “100-year flood”). The frequency of such events will shift as the mean or variance of a climate variable changes (Meehl *et al.* 2000). Second, an extreme event can be defined by organism-based criteria, such as the conditions

¹Institute of Integrative Biology, ETH, Zurich, Switzerland * (jeffrey.diez@env.ethz.ch); ²Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara, CA; ³Department of Forestry and Natural Resources and Department of Biological Sciences, Purdue University, West Lafayette, IN; (continued on p257)

substantially exceeding the acclimation capacity of an organism (Gutschick and BassiriRad 2003). According to this definition, extreme events are specific to particular organisms or ecosystems because of differing physiological tolerances and evolutionary histories. This definition can also account for the relative “abruptness” of events (the magnitude of change per unit time), which will depend on the length of species life cycles and the successional stages of ecosystems (Jentsch *et al.* 2007). Here, we follow Smith (2011), who integrated these perspectives by defining an ECE as “an episode or occurrence in which a statistically rare or unusual climatic period alters ecosystem structure and/or function well outside the bounds of what is considered typical or normal variability”. This characterization allows us to draw from a diverse body of literature, while recognizing that the definition of an ECE has both statistical and ecological dimensions.

■ Mechanisms by which ECEs may influence biological invasions

The invasion process can be considered a progression of events in which individuals of a species are introduced to a novel location, establish a self-sustaining population, spread across the landscape, and reach levels of local and





regional abundance that can eventually impact resident species and ecosystems (Theoharides and Dukes 2007). Therefore, for an ECE to have an effect on invasion, it must influence at least one of these stages (Table 1). Most research to date has focused on the establishment and population growth stages, but there is also evidence that ECEs may affect the transport of propagules into new regions and/or subsequent spread within a region. The manner in which ECEs influence the ecological impact of an invasion event is the least understood stage.

Previous empirical and theoretical work suggests that ECEs may act via different mechanisms across these stages. Identifying the mechanisms at the transport stage is relatively straightforward: ECEs such as storms (discussed below) can increase the chances of movement of non-native species propagules over long distances or across physical barriers. The effects of ECEs on establishment or population growth stages are more complex, but we suggest these will occur via two general pathways (Figure 1), both of which are based on the creation of an “invasion window”. First, extreme events may cause abrupt and widespread mortality of resident species, thus acting as disturbances or “punctuated killing events” (Sousa 1984). Disturbances increase the availability of resources (eg nutrients, water, prey, space) to other

species and may provide “resource opportunities” for introduced species (Shea and Chesson 2002). The duration of the invasion window is a function of the resilience of the native community as well as the magnitude, duration, frequency, and timing of the event(s) (Figure 1). Second, extreme events may stress (but not uniformly kill) resident individuals, decreasing their capacity to utilize resources and thereby limiting their growth and/or reproduction. Until the stressor ceases, conditions may be unsuitable for both native and non-native species (noted by the “abiotic resistance threshold” line in Figure 1). If residents do not immediately recover when the stressor ceases (eg when a severe drought ends), then resources become available that invaders could potentially access (Figure 1).

Mounting evidence suggests that non-native plant species may be favored following ECEs because they tend to have broader environmental tolerances than co-occurring native species (Dukes and Mooney 1999) as well as traits that favor rapid resource acquisition, growth, and colonization of disturbed areas (Pyšek and

Table 1. Extreme climatic events may influence different stages of invasion

Stage of invasion	Mechanisms of ECE effects	Management actions
	<ul style="list-style-type: none"> Storm-driven large-scale movements, as from: <ul style="list-style-type: none"> o strong winds o storm surges 	<ul style="list-style-type: none"> Identify vulnerable habitats
	<ul style="list-style-type: none"> Reduced biotic resistance via: <ul style="list-style-type: none"> o disturbances o stressors 	<ul style="list-style-type: none"> Habitat restoration to increase resilience following ECEs and resistance to invaders
	<ul style="list-style-type: none"> Overcoming dispersal barriers Reduced biotic resistance (as for establishment) 	<ul style="list-style-type: none"> Early detection and eradication of non-native individuals Bolster barriers between invaded and uninvaded areas
	<ul style="list-style-type: none"> Changing per capita effects of invaders (eg drought leading to more concentrated density of non-native fish predators, or greater fuel build up due to drought-caused death of native woody plants leading to higher fire intensity) 	<ul style="list-style-type: none"> Increased awareness/effort post-ECE Unique opportunities for eradication provided by some ECEs

Notes: Transport from an initial source population (orange circle) may initiate invasions in novel environments (large blue circle). Once introduced to a new area, additional ECEs may enhance progression through the stages of invasion, from establishment through population spread and ultimately impacts on native species (small blue circles). Targeted management actions can help limit the probabilities of advancement through each stage of invasion.

Richardson 2007). Although no systematic reviews have assessed whether non-native species are favored by environmental extremes, disturbance theory and empirical data suggest that this is likely. Ultimately, the effects of changing ECEs on communities will depend on the combination of species-specific tolerances of non-native and native species (Figure 2).

Underlying these proposed mechanisms for ECE effects on invasion is the assumption that biological interactions (with pathogens, predators, herbivores, and competitors) or abiotic conditions can provide resistance to invasions. However, the importance of biotic resistance in shaping species invasions continues to be debated in the ecological literature and can be complicated to tease apart from other factors, such as resource heterogeneity (Fridley *et al.* 2007; Melbourne *et al.* 2007). Moreover, evaluating biotic resistance clearly requires unambiguous specification of spatial scales. Native and non-native species richness are often positively correlated at large scales but negatively correlated at small scales, particularly those scales used in experimental studies (Fridley *et al.* 2007). ECEs may affect the invasion of communities both by reducing biotic resistance at local scales and also through effects on abiotic and biotic heterogeneity at landscape scales (Melbourne *et al.* 2007). Thus, even if communities do not exhibit strong biotic resistance, ECEs may influence invasions by changing resource heterogeneity and by resetting the pool of competing species through disturbance.

Below, we provide an overview of how ECEs can affect the abundance and impact of non-native species – through dispersal opportunities, disturbances, resource pulses, and physiological stress. As we highlight in the following sections, these are not mutually exclusive categories because ECEs often change conditions in a manner that spans multiple determinants of species invasions.

Dispersal opportunities

The unprecedented movements of species around the globe are largely driven by human activity, but emerging evidence suggests that ECEs may further promote the transport and introduction of non-native species (Table 1). For example, flooding events can facilitate dispersal of invasive species during unusually high periods of precipitation or rapid melting of snow and glaciers. Flooding events have been linked to numerous

aquatic invasions, including the first known introduction of black carp (*Mylopharyngodon piceus*) into the Missouri River when floodwaters allowed fish to escape from hatchery ponds and subsequently spread downstream (Nico *et al.* 2005). Floods have also been implicated in the spread of other cultured fish species, such as tilapia (a group of species in the Cichlidae family) in Southeast




Figure 1. Conceptual diagram of how extreme disturbance or stress events affect environmental conditions and resistance of an ecosystem to invasion. (a) Disturbance events immediately shift environmental conditions outside those where some or all of the species in the community can survive. As the resident community declines, biotic resistance due to competition and predation is relaxed. When biotic resistance declines past a threshold for a given invasive species, it creates an “invasion window” (shaded areas). The likelihood of increased invasive species dominance depends on the degree of reduction of biotic resistance and the rate at which the community recovers, restoring biotic resistance. Thus, the different sized invasion windows resulting from “fast” and “slow” recoveries are shown here. The frequency of disturbance (return time) and timing of the invasion window within a season (not pictured) can also be important for determining whether and which invasive species benefit. (b) Stress events can also lead to windows of opportunity for invasive species if there is a delay between when abiotic conditions become suitable for an invasive species and when native communities recover. Although the invasion window typically begins immediately following a disturbance event, it can occur at the end of a period of extreme stress. Stressful abiotic conditions (such as a prolonged drought) may increase resistance to invasion (light blue curve) but can also decrease the native community’s ability to resist invasion (black curve). Thus, an invasion window may begin when the abiotic stress has improved to within suitable conditions for the invasive species (eg after rain ends a drought) but before the native community recovers.




Figure 2. Differences among species tolerance curves will determine community responses to disturbance or stress events. (a) Climate variables such as temperature and precipitation are expected to shift in both their mean and variance, and may create different frequencies of what are considered ECEs for a given community. Here, the threshold of what is extreme for this community is shown with a dashed vertical line and the differences in current and future frequencies of these events are shown with blue and red shading, respectively. (b) Communities contain both native and non-native species exhibiting a diversity of tolerances to particular climate factors. In this figure, curves represent how hypothetical native (blue) and non-native (orange) species respond to variation in a particular climate variable. When climate events fall outside of the tolerance curve of a species, it may be stressed or suffer high mortality. Ultimately, the ecological consequences of the shifting climate in (a) will depend on the specific combination of resident and introduced species response curves in (b).

Asia (Canonico *et al.* 2005). In addition, severe flooding in 1993 resulted in extensive dispersal of zebra mussels (*Dreissena polymorpha*) to many new river drainages in the Mississippi River watershed (Tucker 1996). Non-native species have also increased after flooding in systems with upstream dams when fish are transported downstream over reservoir spillways (Schultz *et al.* 2003).

The transport of invasive species may also be promoted by the strong winds, large waves, and storm surges associated with high-magnitude storms. At large scales, the increasing frequency of extreme, large-scale storms that transport airborne dust particles between continents presents opportunities for the dispersal of non-native viruses, bacteria, and fungi (Kellogg and Griffin 2006); the deposition of this dust may also stress native communities. At more regional scales, storm events can facilitate the release of captive non-native species. Terrestrial organisms believed to have benefited from storm-assisted movements include the cactus moth (*Cactoblastis cactorum*; from the Caribbean to Mexico), the red palm mite (*Raoiella indica*; within the Caribbean), and the agricul-

tural weed *Parthenium hysterophorus* across Swaziland (Burgiel and Muir 2010).

Disturbances and resource pulses

ECEs often create resource pulses (eg of light or space) that non-native species are able to utilize. These disturbances and resource pulses can facilitate both the establishment and spread phases of invasion (Figure 1). Storm events are among the most important of these disturbances, sometimes allowing non-native species to rapidly establish and spread (WebTable 1). For example, following Hurricane Andrew in Florida in 1992, non-native vines benefited from wind-driven tree canopy loss and spread widely across the state, despite the presence of native vines (Horvitz *et al.* 1998). Similar responses were observed following Hurricane Katrina in Louisiana in 2005 (Brown *et al.* 2011). In marine systems, storm surges and wave action can also leave communities susceptible to invasion (Denny *et al.* 2009). Pre- and post-hurricane surveys in Dominica recorded the burial of native seagrass beds by unusually large sediment loads as a result of storm surges associated with Hurricane Omar in 2008, which led to the establishment of invasive seagrasses (*Halophila* spp; Steiner *et al.* 2010).

Extreme floods may also act as disturbance agents with variable effects on invasions in aquatic systems. The physical changes to aquatic systems caused by floods can create opportunities for colonization, establishment, and spread by non-native species (Table 1), while suppression of flood regimes can also facilitate invasion. For example, Kercher and Zedler (2004) reported that the invasive reed canarygrass (*Phalaris arundinacea*) and broadleaf cattail (*Typha latifolia*) outgrew other native (and non-native) perennial species in experimentally manipulated cyclic and constant flooding and drought events. By contrast, in arid streams of the American Southwest, flooding has been shown to benefit native fishes. Floods allow many native species that evolved in areas where precipitation is often torrential – and where the onset of flooding can occur in seconds to minutes – to displace non-native fishes that evolved in mesic systems with more constant hydrology, where floods build over periods of hours to days (Meffe 1984).

Heat waves can cause persistent stress (discussed below) and also abrupt mortality of resident species. Both stress and mortality can facilitate the establishment and spread stages of invasion, when non-native species are able to either tolerate higher temperatures or more rapidly take up the resources made available by the deaths or reduced performance of native residents. In a marine coastal ecosystem in New Zealand, a heat wave caused the weekly high temperature to exceed the maximum from previous years by 7°C; this led to mortality that was an order of magnitude higher for a native mussel than for a non-native mussel species (Petes *et al.* 2007). In France's Rhône River, increases in non-native and eury-tolerant (tolerant of a broad range of conditions)

macroinvertebrates were temporally correlated with the 2003 European heat wave, as well as with severe flood events that may have contributed to increased resources (Daufresne *et al.* 2007).

Experimental manipulations corroborate the trends observed during and following heat waves. Sorte *et al.* (2010) simulated a heat wave in laboratory mesocosms of a marine epibenthic fouling community from California and found that the only species able to tolerate this disturbance were non-native in origin. Furthermore, the increased dominance of these non-natives persisted through 3 months of community development in the field, as a result of the non-natives' ability to rapidly capitalize on the increased resources (ie open space) both by growth and colonization of new recruits. Similarly, Song *et al.* (2010) showed that a heat wave led to a decrease in biomass of a native herbaceous *Wedelia* species (Asteraceae) but not its non-native congener, whereas White *et al.* (2001) found decreased native biomass and increased invasibility of a grassland system following extreme heat events. If indicative of responses to ECEs, these examples suggest that extreme disturbances have the potential to favor invaders because of differences in tolerances and non-native species abilities to preempt available resources (eg by growth and colonization) more quickly than native species.

Stressors

A third important mechanism by which extreme events may affect establishment of invasive species is by creating stressful conditions that reduce the biotic resistance of a community over longer time periods (Table 1). These stressful conditions may sometimes be punctuated by a large die-off at the end of the period, resulting in conditions similar to disturbances. The distinction is that stress events may not always lead to death, and the intermediate period of reduced competition and predation may lower the biotic resistance of communities. As with disturbances, these periods of stress may facilitate establishment or spread of invasive species (Figure 1).

Extreme heat waves and droughts are among the clearest examples of stressful conditions imposed by ECEs. The severe droughts predicted to increase in frequency with climate change can exceed the tolerances of resident species, leading to reduced vigor and widespread mortality events (Allen *et al.* 2010). Extreme droughts may involve either exceptional duration of drought conditions or shifted timing of drought relative to critical life-history stages. There are currently few data on which plant species are favored by drought-driven tree die-offs within forests, but non-native species are poised to increase in some drought-stricken ecosystems. For example, invasive grasses (eg *Bromus* spp) that are widespread in western North America may be suppressed temporarily by drought but can recover rapidly, and could then invade areas of pinyon pine (*Pinus* spp) or juniper (*Juniperus* spp) die-off

(Kane *et al.* 2011). Drought has also contributed to the increasing prevalence of the invasive tamarisk (*Tamarix ramosissima*) in riparian areas across western North America (Stromberg 1998), where declining water tables select against native species. Both climatically induced drought and water extraction and water-table reduction by humans decrease water access for riparian plants and increase accumulation of salt in surface soils; these conditions act in concert to promote the more drought- and salt-tolerant *Tamarix* (Vandersande *et al.* 2001).

Although invasive species may initially be better able to colonize areas where drought has reduced biotic resistance, native species in drought-prone ecosystems can be better adapted to prolonged drought conditions than non-native species. For example, experimental drought reduced water-use efficiency much more in invasive dandelions (*Taraxacum officinale*) than in a native congener (*Taraxacum ceratophorum*) (Brock and Galen 2005). In Hawaii, the native grass *Heteropogon contortus* is more tolerant of drought than the dominant invader, *Pennisetum setaceum*, with which it typically competes (Goergen and Daehler 2002). In Venezuela, native grasses are more tolerant of high vapor-pressure deficits and water-stressed conditions than two African grasses that invade native savannas (Baruch and Jackson 2005). The many invasive plant species with high growth rates, leaf areas, and water use (Cavaleri and Sack 2010) may be at a disadvantage if drought conditions recur or persist after an ECE.

Extreme drought events can also lead to unusually low stream flows, which have been shown to promote resistance to aquatic invasions. The introduced brown trout (*Salmo trutta*) in New Zealand is more susceptible than native galaxiid fish to stresses associated with low flows; such events may therefore prevent trout from eliminating galaxiid fish in low gradient streams (Leprieur *et al.* 2006). In this case, water diversions caused the low-flow events, but we would expect similar outcomes under climate-induced drought conditions. Likewise, during a recent extreme drought in Victoria, Australia, many streams were reduced to small, isolated pools with depleted oxygen levels and high water temperatures; invasive common carp (*Cyprinus carpio*) were unable to survive these conditions and local populations were extirpated, whereas the native galaxiids survived (Lake 2003). Thus, ECEs that increase environmental stress levels may at times favor either non-native or native species in a manner that is taxon- and system-specific.

■ Synergies among stressors

The consequences of ECEs for biological invasions will sometimes depend on other aspects of global change driven by anthropogenic stressors. For example, gradual climate change may force many organisms closer to their physiological tolerance limits, thereby reducing their competitive ability and resilience to extreme events. In the western US, the compounding effects of recurrent



Figure 3. (a) Mortality events due to hurricane damage, such as that in Mississippi due to Hurricane Katrina, can open up resources that will support the establishment of non-native species. (b) Prolonged drought in a high-elevation woodland on Mauna Kea, Hawaii, where invasive grasses are recovering more quickly than native species. (c) Palm oasis in the Nevada desert that burned as a result of the synergism of *Tamarix* invasion, increasing drought, and severe fire weather. (d) An old bridge normally hidden beneath the surface waters of Llyn-on Reservoir, Wales, is clearly visible during the extreme drought of 2010. Droughts can create isolated pool habitats, such as those in the foreground, that intensify interactions between introduced and native species. Whether these changes favor native or introduced species may depend on the system and the specific tolerances and behavior of resident species.

droughts and gradual climate change may induce higher tree mortality during an ECE (Adams *et al.* 2009). The resulting increase in resource availability (eg water, light, space) after a drought could provide a window of opportunity for introduced species better adapted to changing conditions. As discussed above, *T ramosissima* invasions of riparian zones in southwestern North America are facilitated by drought; however, native *Populus* spp can outcompete *T ramosissima* after flood events when soil moisture is high (Sher *et al.* 2000). Thus, the net outcomes of ECEs for these riparian ecosystems are likely to depend on interacting factors, such as alteration of groundwater and flow regimes by dams, in combination with ECEs and even fire occurrence.

The effects of ECEs are also likely to interact with ongoing changes in propagule pressure of non-native species. The expanding number and volume of commercial trade routes will continue to spread invasive species

across the globe. This means that invasive species are more likely to be present in regional species pools, poised to take advantage of windows of opportunity created by ECEs (Olden *et al.* 2011). Additional synergistic interactions between ECEs and invasions could take place through positive feedbacks between non-native species and the physical environment, disturbance regimes, biogeochemical cycling, and biotic composition of invaded ecosystems (Crooks 2002). For example, in Hawaii, extreme drought causes mortality among some of the dominant native woody species (Figure 3; Lohse *et al.* 1995), thereby facilitating the dominance of non-native grasses. This increases the likelihood of fire and could potentially shift the ecosystem toward a grass-dominated state (CMD'A pers observation). Even previously fire-prone ecosystems may experience damaging fire frequencies and/or intensities through the interaction of ECEs and established invaders. The cogongrass (*Imperata cylin-*

drica) invasion in Florida pine ecosystems has increased fire intensity, causing greater native species mortality and facilitating further invasion (Lippincott 2000). Although such interactions have rarely been studied, these examples highlight how positive feedbacks could allow invasive species to “transform” some ecosystems to new, possibly persistent states.

Similar effects may ripple across trophic levels. For example, changes in vegetation structure caused by hurricanes have been shown to precipitate shifts in bird communities resulting from new nesting opportunities in invasive vines (Brown *et al.* 2011). In the Bahamas, Schoener *et al.* (2001) found that predation by introduced lizards resulted in local extinction of the smaller native lizard species only after storm events reduced the natives’ population sizes. The effects of ECEs on invasions at one trophic level may therefore depend on changes due to ECEs at other trophic levels.

■ Future research and management efforts

Research to date has shown that ECEs can affect the invasion process but that this relationship is complex and context-dependent. Although many studies suggest that ECEs can enhance dispersal and reduce biotic resistance to non-native species establishment (WebTable 1), few have examined how ECEs affect the magnitude of invader impacts on native systems. Also, ECEs do not uniformly favor non-native species, and many non-natives that benefit may have no substantial ecological effect. In some systems, events like extreme floods or persistent droughts may negatively affect established invaders, thereby providing opportunities for restoration of native species. Although this growing body of research allows us to identify some mechanisms by which ECEs influence invasions, we still lack sufficient information to make definitive predictions about how increasing ECEs will interact with the invasion process and with species impacts in most ecosystems.

Several areas of research could help improve both our basic understanding of invasive species responses to ECEs and our ability to anticipate and mitigate their effects in the future. First, it is important to determine whether there are specific characteristics of introduced species and recipient native communities that are likely to lead to invasion after ECEs. Are there traits and demographic characteristics of introduced species that would allow them to tolerate extremes better than native species? What makes some native communities more vulnerable to invasion post-ECE than others? Are communities that have evolved with higher natural climatic variability more resilient to ECEs? These are basic ecological questions with clear management implications. However, despite these remaining uncertainties about ECE effects on invasions, our review points to concrete actions that managers can take to prepare for and to mitigate the impacts of extreme events (WebPanel 2). Resource man-

agers can begin to target their efforts toward species that appear more likely to invade after an ECE, such as introduced vines following hurricanes (Horvitz *et al.* 1998). Similarly, efforts can be made to increase resilience of communities that may be susceptible to ECEs and subsequent invasions. Riparian zones of riverine ecosystems that are prone to flooding are a clear example of where targeted restoration efforts may help increase community resilience to invasion after flooding events.

A second research area is to identify biological thresholds of responses to extreme events. It is currently unclear where the critical tolerance thresholds are, which, once crossed, will cause dramatic change in a community’s trajectory. Part of this challenge is to understand how synergies among global change drivers affect system thresholds, especially because extreme conditions can arise as a result of the co-occurrence of unusual, though not extreme, values of two or more climate variables (Denny *et al.* 2009). A more mechanistic understanding of the conditions that lead to changes in a system is necessary if we are to anticipate the results of ECEs in different ecosystems.

Progress on these questions will require a range of approaches, including manipulative experiments, observational studies, and modeling. Controlled experiments are most useful for isolating mechanisms and are becoming more powerful as approaches are developed for manipulating extreme values for two climate variables concurrently (Smith 2011). Observational studies are critical for understanding dynamics over larger spatial and temporal scales and for systems not amenable to controlled manipulation. Successful observational studies of the impacts of ECEs will rely on long-term monitoring of areas prone to ECEs (eg hurricane corridors, drought-prone areas) and rapid mobilization of research efforts to take advantage of chance events. Managers have a critical role to play in this research by identifying susceptible and control areas for study. Coupling observational studies directly to management needs and actions will allow comparisons among events of varying magnitudes, durations, and frequencies at broad spatial and temporal scales. Finally, new modeling approaches are needed to integrate experimental and observational data with ecological theory in order to build risk assessments of future invasions in response to ECEs. Currently, most theoretical investigations of how biological systems shift among alternative states (eg between uninvaded and invaded states) have focused on gradual changes in drivers such as climate variables (Scheffer and Carpenter 2003) with the goal of identifying early-warning signals that critical thresholds are being approached (Scheffer *et al.* 2009). Only recently have modeling efforts focused on shifts among alternate states in response to stochastic climatic events (Schooler *et al.* 2011) and the modeling of transient dynamics over ecologically relevant timescales (Hastings 2010). Such modeling can also be extremely useful for managers by evaluating alternative management scenarios in the face of both gradual climate change and ECEs.

In summary, the intersection between ECEs and biological invasions represents an important focus for understanding and predicting future changes in a wide range of ecosystems. We suggest that future studies of climate change and species invasions should not only consider future trends in mean values of climatic factors but instead should elucidate the mechanisms by which ECEs influence invasions across a broad range of taxa, trophic levels, ecosystems, and biogeographic regions. Management of natural areas will increasingly require planning for and mitigation of the effects of ECEs in an ever more invaded world.

■ Acknowledgements

This review was conducted as part of the Climate Change & Invasive Species Working Group supported by the National Center for Ecological Analysis and Synthesis, a Center funded by the National Science Foundation (grant #EF-0553768), the University of California, Santa Barbara, and the State of California.


■ References

- Adams HD, Guardiola-Claramonte M, Barron-Gafford GA, *et al.* 2009. Temperature sensitivity of drought-induced tree mortality portends increased regional die-off under global-change-type drought. *P Natl Acad Sci USA* **106**: 7063–66.
- Allen CD, Macalady AK, Chenchouni H, *et al.* 2010. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecol Manag* **259**: 660–84.
- Baruch Z and Jackson RB. 2005. Responses of tropical native and invader C-4 grasses to water stress, clipping and increased atmospheric CO₂ concentration. *Oecologia* **145**: 522–32.
- Bradley BA, Blumenthal DM, Wilcove DS, and Ziska LH. 2010. Predicting plant invasions in an era of global change. *Trends Ecol Evol* **25**: 310–18.
- Brock MT and Galen C. 2005. Drought tolerance in the alpine dandelion, *Taraxacum ceratophorum* (Asteraceae), its exotic congener *T. officinale*, and interspecific hybrids under natural and experimental conditions. *Am J Bot* **92**: 1311–21.
- Brown DR, Sherry TW, and Harris J. 2011. Hurricane Katrina impacts the breeding bird community in a bottomland hardwood forest of the Pearl River basin, Louisiana. *Forest Ecol Manag* **261**: 111–19.
- Burgiel SW and Muir AA. 2010. Invasive species, climate change and ecosystem-based adaptation: addressing multiple drivers of global change. Washington, DC, and Nairobi, Kenya: Global Invasive Species Programme.
- Canonico GC, Arthington A, McCrary JK, and Thieme ML. 2005. The effects of introduced tilapias on native biodiversity. *Aquat Conserv* **15**: 463–83.
- Cavaleri MA and Sack L. 2010. Comparative water use of native and invasive plants at multiple scales: a global meta-analysis. *Ecology* **91**: 2705–15.
- Crooks JA. 2002. Characterizing ecosystem-level consequences of biological invasions: the role of ecosystem engineers. *Oikos* **97**: 153–66.
- Daufresne M, Bady P, and Fruguet JF. 2007. Impacts of global changes and extreme hydroclimatic events on macroinvertebrate community structures in the French Rhône River. *Oecologia* **151**: 544–59.
- Denny MW, Hunt LJH, Miller LP, and Harley CDG. 2009. On the prediction of extreme ecological events. *Ecol Monogr* **79**: 397–421.
- Dukes JS and Mooney HA. 1999. Does global change increase the success of biological invaders? *Trends Ecol Evol* **14**: 135–39.
- Easterling D, Meehl G, Parmesan C, *et al.* 2000. Climate extremes: observations, modeling, and impacts. *Science* **289**: 2068–74.
- Fridley JD, Stachowicz JJ, Naeem S, *et al.* 2007. The invasion paradox: reconciling pattern and process in species invasions. *Ecology* **88**: 3–17.
- Goergen E and Daehler CC. 2002. Factors affecting seedling recruitment in an invasive grass (*Pennisetum setaceum*) and a native grass (*Heteropogon contortus*) in the Hawaiian Islands. *Plant Ecol* **161**: 147–56.
- Gutschick VP and BassiriRad H. 2003. Extreme events as shaping physiology, ecology, and evolution of plants: toward a unified definition and evaluation of their consequences. *New Phytol* **160**: 21–42.
- Hastings A. 2010. Timescales, dynamics, and ecological understanding. *Ecology* **91**: 3471–80.
- Horvitz CC, Pascarella JB, McMann S, *et al.* 1998. Functional roles of invasive non-indigenous plants in hurricane-affected subtropical hardwood forests. *Ecol Appl* **8**: 947–74.
- Jentsch A, Kreyling J, and Beierkuhnlein C. 2007. A new generation of climate-change experiments: events, not trends. *Front Ecol Environ* **5**: 365–74.
- Kane JM, Meinhardt KA, Chang T, *et al.* 2011. Drought-induced mortality of a foundation species (*Juniperus monosperma*) promotes positive afterlife effects in understory vegetation. *Plant Ecol* **212**: 733–41.
- Karl TR, Meehl GA, Miller CD, *et al.* (Eds). 2008. Weather and climate extremes in a changing climate. Washington, DC: NOAA National Climatic Data Center.
- Kellogg CA and Griffin DW. 2006. Aerobiology and the global transport of desert dust. *Trends Ecol Evol* **21**: 638–44.
- Kercher SM and Zedler JB. 2004. Flood tolerance in wetland angiosperms: a comparison of invasive and noninvasive species. *Aquat Bot* **80**: 89–102.
- Lake PS. 2003. Ecological effects of perturbation by drought in flowing waters. *Freshwater Biol* **48**: 1161–72.
- Leprieux F, Hickey MA, Arbuckle CJ, *et al.* 2006. Hydrological disturbance benefits a native fish at the expense of an exotic fish. *J Appl Ecol* **43**: 930–39.
- Lippincott CL. 2000. Effects of *Imperata cylindrica* (L) Beauv (cogongrass) invasion on fire regime in Florida sandhill (USA). *Nat Area J* **20**: 140–49.
- Lohse KA, Nullet D, and Vitousek PM. 1995. Effects of extreme drought on vegetation of a single lava flow on Mauna Loa, Hawai'i. *Pac Sci* **49**: 212–20.
- Meehl G, Zwiers F, Evans J, *et al.* 2000. Trends in extreme weather and climate events: issues related to modeling extremes in projections of future climate change. *Bull Amer Meteor Soc* **81**: 427–36.
- Meffe GK. 1984. Effects of abiotic disturbance on coexistence of predator-prey fish species. *Ecology* **65**: 1525–34.
- Melbourne BA, Cornell HV, Davies KF, *et al.* 2007. Invasion in a heterogeneous world: resistance, coexistence or hostile takeover? *Ecol Lett* **10**: 77–94.
- Nico LG, Williams JD, and Jelks HL. 2005. Black carp: biological synopsis and risk assessment of an introduced fish. Bethesda, MD: American Fisheries Society. Special Publication 32.
- Olden JD, Lockwood JL, and Parr CL. 2011. Species invasions and the biotic homogenization of faunas and floras. In: Ladle RJ and Whittaker RJ (Eds). Conservation biogeography. West Sussex, UK: Wiley-Blackwell.
- Parmesan C, Root T, and Willig M. 2000. Impacts of extreme weather and climate on terrestrial biota. *Bull Amer Meteor Soc* **81**: 443–50.

- Petes LE, Menge BA, and Murphy GD. 2007. Environmental stress decreases survival, growth, and reproduction in New Zealand mussels. *J Exp Mar Biol Ecol* **351**: 83–91.
- Pyšek P and Richardson DM. 2007. Traits associated with invasiveness in alien plants: where do we stand? In: Nentwig W (Ed). *Biological invasions*. Berlin, Germany, and Heidelberg, Germany: Springer-Verlag.
- Rahel FJ and Olden JD. 2008. Assessing the effects of climate change on aquatic invasive species. *Conserv Biol* **22**: 521–33.
- Scheffer M, Bascompte J, Brock WA, *et al.* 2009. Early-warning signals for critical transitions. *Nature* **461**: 53–59.
- Scheffer M and Carpenter SR. 2003. Catastrophic regime shifts in ecosystems: linking theory to observation. *Trends Ecol Evol* **18**: 648–56.
- Schoener TW, Spiller DA, and Losos JB. 2001. Predators increase the risk of catastrophic extinction of prey populations. *Nature* **412**: 183–86.
- Schooler SS, Salau B, Julien MH, and Ives AR. 2011. Alternative stable states explain unpredictable biological control of *Salvinia molesta* in Kakadu. *Nature* **470**: 86–89.
- Schultz AA, Maughan OE, Bonar SA, and Matter WJ. 2003. Effects of flooding on abundance of native and nonnative fishes downstream from a small impoundment. *N Am J Fish Manage* **23**: 503–11.
- Shea K and Chesson P. 2002. Community ecology theory as a framework for biological invasions. *Trends Ecol Evol* **17**: 170–76.
- Sher AA, Marshall DL, and Gilbert SA. 2000. Competition between native *Populus deltoides* and invasive *Tamarix ramosissima* and the implications for reestablishing flooding disturbance. *Conserv Biol* **14**: 1744–54.
- Smith MD. 2011. An ecological perspective on extreme climatic events: a synthetic definition and framework to guide future research. *J Ecol* **99**: 656–63.
- Song LY, Chow WS, Sun LL, *et al.* 2010. Acclimation of photosystem II to high temperature in two *Wedelia* species from different geographical origins: implications for biological invasions upon global warming. *J Exp Bot* **61**: 4087–96.
- Sorte CJB, Fuller A, and Bracken MES. 2010. Impacts of a simulated heat wave on composition of a marine community. *Oikos* **119**: 1909–18.
- Sousa WP. 1984. The role of disturbance in natural communities. *Annu Rev Ecol Syst* **15**: 353–91.
- Steiner SCC, Macfarlane KJ, Price LM, and Willette DA. 2010. The distribution of seagrasses in Dominica, Lesser Antilles. *Rev Biol Trop* **58**: 89–98.
- Stromberg JC. 1998. Dynamics of Fremont cottonwood (*Populus fremontii*) and saltcedar (*Tamarix chinensis*) populations along the San Pedro River, Arizona. *J Arid Environ* **40**: 133–55.
- Theoharides KA and Dukes JS. 2007. Plant invasion across space and time: factors affecting nonindigenous species success during four stages of invasion. *New Phytol* **176**: 256–73.
- Thibault KM and Brown JH. 2008. Impact of an extreme climatic event on community assembly. *P Natl Acad Sci USA* **105**: 3410–15.
- Tucker JK. 1996. Post-flood strandings of unionid mussels. *J Freshwater Ecol* **11**: 433–38.
- Vandersande MW, Glenn EP, and Walworth JL. 2001. Tolerance of five riparian plants from the lower Colorado River to salinity drought and inundation. *J Arid Environ* **49**: 147–59.
- Walther GR, Roques A, Hulme PE, *et al.* 2009. Alien species in a warmer world: risks and opportunities. *Trends Ecol Evol* **24**: 686–93.
- White TA, Campbell BD, Kemp PD, and Hunt CL. 2001. Impacts of extreme climatic events on competition during grassland invasions. *Glob Change Biol* **7**: 1–13.

⁴Department of Environmental Science and Policy, University of California, Davis, CA; ⁵School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA; ⁶Department of Environmental, Earth and Ocean Sciences, University of Massachusetts, Boston, MA; ⁷Rangeland Resources Research Unit, USDA Agricultural Research Service, Fort Collins, CO; ⁸Department of Environmental Conservation, University of Massachusetts, Amherst, MA; ⁹Cátedra Rui Nabeiro, Universidade de Évora, Évora, Portugal; ¹⁰School of Natural Resources, University of Michigan, Ann Arbor, MI; ¹¹Department of Biological Sciences, University of South Carolina, Columbia, SC; ¹²School of Forest Resources, University of Washington, Seattle, WA; ¹³Marine Science Center, Northeastern University, Nahant, MA

ESA's *Issues in Ecology* series



Each publication in this series:

- Reports a consensus of a panel of scientific experts
- Uses clear, commonly understood language
- Focuses on issues related to the environment
- Is intended for scientists, educators, students, and decision-makers
- Is reviewed by external experts for technical content

Issue #14: “Setting Limits: Using Air Pollution Thresholds to Protect and Restore U.S. Ecosystems”

For more information, visit:

www.esa.org/issues

WebPanel 1. Linking climate change to extreme events

The frequency, duration, timing, and magnitude of ECEs have changed markedly over recent years, a large part of which can be linked to climate change. In this panel, we provide a brief review of the expected changes in temperature, precipitation, and storm-related ECEs (for thorough reviews, see Tebaldi et al. 2006; Meehl et al. 2007; Karl et al. 2008).

Temperature

Changing climate is expected to alter extreme temperature events, such as heat waves and unseasonal frosts, which have important ecological implications.

Heat waves

In the continental US, weather stations monitored since the 1900s reveal more frequent high-temperature extremes (temperatures above the 95th percentile) in the late 20th century, a trend that is predicted to continue under current global-warming scenarios.

Unseasonal frosts

Warm early season temperatures due to climate change are leading to an earlier onset of spring. The frequency of frost events is expected to decline; however, late (post-spring-warming) frosts are likely to increase with increasing temperature variation.

Precipitation

Climate models generally predict precipitation to become more “peaked” with climate change. This may be associated with altered frequency and magnitude of droughts in many regions, as well as severe precipitation events and associated flooding.

Drought


Although climate projections for precipitation have higher uncertainty than temperature projections, models consistently forecast dry areas (eg semi-arid and arid subtropics) to become drier. For example, model ensembles show a consistent drying trend in the southwestern US, particularly during summer months. Within the US, these extreme droughts are likely to be similar to the drought that caused the Dust Bowl in the 1930s. An analysis of the Mediterranean region projected a consistent decrease in precipitation during spring and summer months (Gao and Giorgi 2008). In both of these cases, decreased precipitation is projected to result in more extreme droughts and, consequently, an expansion of arid lands.

Flooding

Extreme flood events are of a magnitude, in space and duration, that exceeds the tolerance limits of most ecosystems affected. Some areas of the US are experiencing an increase in precipitation, particularly since the 1970s, including an increase in severe precipitation events. Milly et al. (2002) used a model ensemble to estimate future 100-year flood events and projected an increase in frequency in several regions of the world, including high northern latitudes (boreal forest), tropics in South America and Africa, and large portions of China, as well as the upper midwestern US.

Storms

The rise in ocean temperatures and the water vapor content of the atmosphere is affecting the formation and magnitude of tropical storms. Maximum wind speeds of cyclones have substantially increased over the past 30 years, and hurricanes are predicted to continue increasing in intensity as climate changes. As sea level is also rising, the impact of these storms will increase, intensifying the risk of flooding and storm damage.



WebFigure 1. Examples of predicted global changes in dry days and heat waves from the late 20th century to the late 21st century under IPCC scenario A1B (from Tebaldi et al. 2006). The “dry days” index is defined as the maximum number of consecutive dry days in a year, and “heat waves” are the maximum periods of at least five consecutive days with maximum temperature at least 5 °C higher than the 1961–1990 mean for the same calendar day. Values are changes in multi-model averages in standardized units. Stippled regions show where five of the nine models used in the analysis predict statistically significant changes. Tebaldi et al. (2006) found general trends toward increasing extremes of ten indicators of temperature and extreme precipitation events, but with substantial regional variability and differences among models. The coarse resolution of available global models (varying in these nine from 1–5 °C resolution) also suggests that there is more local heterogeneity than depicted by these maps. With kind permission from Springer Science and Business Media.

WebPanel 2. Planning and managing for extreme events and invasions**Anticipate events and responses**

Planning for ECEs will require fundamental knowledge of how and where ECEs are likely to facilitate future species invasions. Managers will need to (a) understand the historical and likely future nature and patterns of ECEs, (b) identify which areas and types of habitats will be particularly susceptible to ECEs (eg low-lying or exposed sites), (c) determine which taxa will be most likely to respond to ECEs (eg introduced vines following hurricanes; Horvitz *et al.* 1998), and (d) identify control areas to monitor following ECEs.

Set goals and targets

A basic understanding of changes in ECEs and likely responses by potentially ecosystem-transforming invasive species will allow managers to delineate areas for monitoring and/or control and eradication of invaders following ECEs, and to prioritize or even triage species and sites.

Develop agreements among management authorities

Coordinating efforts across agencies, individuals, and landscapes will be important for addressing climate-change effects in general and for addressing ECEs in particular. If cooperative agreements that clearly delineate responsibilities are put in place in advance, management responses will be faster and more effective.

Develop capacity for restoration

Responding to ECEs will require resources that include stores of native species in the form of seeds, seedlings, eggs, or individuals. If facilities are developed and capacities to produce and store these resources are expanded, managers will have a much greater chance of preserving ecosystem structure and function. Implementing these measures will depend on allocation of sufficient funding in advance of the ECEs.

Preventing/reducing impacts**Reduce exposure to extreme events**

Several climate-change adaptation strategies have been proposed for reducing the exposure of systems to extreme events. For example, coastal restoration can reduce the exposure of coastal and near-coastal systems to storm surges, and riparian restoration can reduce the exposure of streams to extreme heat events.

Improve the ability of a system to respond to extreme events

Other adaptation strategies can increase the resilience of a system, allowing it to better respond to ECEs. For example, floodplain restoration may increase biotic resistance to invasion following the arrival of non-native propagules under extremely high flows.

Minimize propagule pressure and maximize biotic resistance

By reducing sources of non-native propagules, restoring native populations to increase biotic resistance, and managing pathways of invasion, it will be possible to reduce the ability of some introduced species to take advantage of ECEs.

Responding to events**Develop early detection/rapid response following ECEs**

Successful responses to ECEs will require quick action facilitated by early detection and organized efforts.

Engage volunteer networks for post-ECE monitoring and management activities

Volunteer networks and community groups can provide the large teams needed to both monitor for invasions following ECEs and respond to them once detected.

WebTable 1. Examples from the literature documenting effects of extreme events on species invasions

Type of event	Mechanism	System	Location	Organism	Citation	Biological response
Storms						
Waves	Disturbance	Rocky intertidal	South Africa	Marine invertebrates	Erlandsson <i>et al.</i> (2006)	Following wave disturbance, an invasive mussel recolonized faster and attained three times the cover than the native species
Hurricane – storm surge	Disturbance	Seagrass bed	Dominica	Seagrass	Steiner <i>et al.</i> (2010)	An invasive seagrass grew at a wider range of depths than native seagrass, allowing it to survive strong storm surges caused by Hurricane Omar
	Disturbance	Low-elevation islands	The Bahamas	Lizards	Schoener <i>et al.</i> (2001)	Storm exacerbated impacts of experimentally introduced lizards, driving native lizards to extinction
Hurricane – winds	Disturbance, resource pulse	Forest	Florida	Plants	Horvitz <i>et al.</i> (1998)	Hurricane Andrew accelerated invasion by exotics that inhibited regeneration of native species
	Disturbance, resource pulse	Forest	Louisiana	Plants	Brown <i>et al.</i> (2011)	Damage from Hurricane Katrina was followed by a 15-fold increase in invasive blackberries
	Disturbance, resource pulse	Forest	Puerto Rico	Plants	Flynn <i>et al.</i> (2010)	Exotic species were more susceptible to hurricane damage than co-occurring natives
	Disturbance, resource pulse	Forest	Mississippi	Plants	Chabreck and Palmisan (1973)	Hurricane Camille facilitated the increase of invasive understory shrubs
	Disturbance, resource pulse	Forest	Jamaica	Plants	Bellingham <i>et al.</i> (2005)	Hurricane Gilbert facilitated the rapid increase of an invasive tree
Tornado – winds	Disturbance, resource pulse	Forest	Texas	Plants	Glitzenstein and Harcombe (1988)	Tornado increased abundance of exotic Chinese tallow
Cyclone – winds	Disturbance, resource pulse	Forest	Madagascar	Plants	Brown and Gurevitch (2004)	Cyclone facilitated exotic trees
Temperature						
Heat wave	Stress	Rocky intertidal	New Zealand	Marine invertebrates	Petes <i>et al.</i> (2007)	Resistance (survival) was higher for an invasive than a native mussel
	Stress	Marine subtidal	California	Marine invertebrates	Sorte <i>et al.</i> (2010)	Invasive species were more resistant and resilient than natives to a simulated heat wave
	Stress	Open terrestrial habitats	China	Plants	Song <i>et al.</i> (2010)	Invasive species were more resistant (sustained higher biomass) than natives to a simulated heat wave

continued

WebTable 1. – Continued

Type of event	Mechanism	System	Location	Organism	Citation	Biological response
	Stress	Grassland	New Zealand	Plants	White <i>et al.</i> (2001)	Experimental heating suppressed competition by resident C ₃ perennials and promoted invasion by C ₄ annuals
Heat wave and floods	Disturbance, stress	River	France	Freshwater invertebrates	Daufresne <i>et al.</i> (2007)	Multiple events converged, leading to a shift in community composition toward increased invasives
Precipitation: drought and floods						
Drought	Stress	Stream	Arkansas/Missouri	Freshwater invertebrates	Larson <i>et al.</i> (2009)	Invasive crayfish were more tolerant of desiccation, leading to an increase in numbers with decreased stream discharge, where natives decrease
	Stress	Stream	Colorado	Fishes	Labbe and Fausch (2000)	Drought-induced habitat loss in stream ecosystems can confine native fishes to remnant pools and intensify rates of predation by invasive fishes
	Stress	Stream	Kansas	Plants	Nippert <i>et al.</i> (2010)	Invasive <i>Tamarix</i> rapidly shifted water sources in response to declines in the water table, thereby minimizing leaf-level water stress during drought periods
	Stress	Lake	Central Europe	Clam	Werner and Rothhaupt (2008)	Drought conditions decreased lake volume and winter temperatures, causing a massive die-off of the invasive <i>Corbicula fluminea</i>
Drought and flood	Dispersal	Stream	Brazil	Copepod ectoparasite	Medeiros and Maltchik (1999)	Floods disseminated <i>Lernaea cyprinacea</i> in a fish community and drought intensified the ectoparasite infestation
Flood	Disturbance, resource pulse	Estuary	California	Marine invertebrates	Nichols <i>et al.</i> (1990)	Seasonal flooding scoured soft-bottom habitat, followed by the introduction and subsequent dominance of the Asian clam <i>Potamocorbula amurensis</i>
Flood followed by drought	Disturbance, stress	Restored vernal pools	California	Plants	Collinge <i>et al.</i> (2011)	Extreme flooding reduced plant abundance in one year; extreme drought allowed faster rebound of non-natives in following year

■ WebReferences

- Bellingham PJ, Tanner EVJ, and Healey JR. 2005. Hurricane disturbance accelerates invasion by the alien tree *Pittosporum undulatum* in Jamaican montane rain forests. *J Veg Sci* **16**: 675–84.
- Brown DR, Sherry TW, and Harris J. 2011. Hurricane Katrina impacts the breeding bird community in a bottomland hardwood forest of the Pearl River basin, Louisiana. *Forest Ecol Manag* **261**: 111–19.
- Brown KA and Gurevitch J. 2004. Long-term impacts of logging on forest diversity in Madagascar. *P Natl Acad Sci USA* **101**: 6045–49.
- Chabreck RH and Palmisan AW. 1973. Effects of Hurricane Camille on the marshes of the Mississippi River delta. *Ecology* **54**: 1118–23.
- Collinge SK, Ray C, and Gerhardt F. 2011. Long-term dynamics of biotic and abiotic resistance to exotic species invasion in restored vernal pool plant communities. *Ecol Appl* **21**: 2105–18.
- Daufresne M, Bady P, and Fruget JF. 2007. Impacts of global changes and extreme hydroclimatic events on macroinvertebrate community structures in the French Rhône River. *Oecologia* **151**: 544–59.
- Erlandsson J, Pal P, and McQuaid CD. 2006. Re-colonisation rate differs between co-existing indigenous and invasive intertidal mussels following major disturbance. *Mar Ecol-Prog Ser* **320**: 169–76.
- Flynn DFB, Uriarte M, Crk T, *et al.* 2010. Hurricane disturbance alters secondary forest recovery in Puerto Rico. *Biotropica* **42**: 149–57.
- Gao XJ and Giorgi F. 2008. Increased aridity in the Mediterranean region under greenhouse gas forcing estimated from high resolution simulations with a regional climate model. *Global Planet Change* **62**: 195–209.
- Glitzenstein JS and Harcombe PA. 1988. Effects of the December 1983 tornado on forest vegetation of the big thicket, south-east Texas, USA. *Forest Ecol Manag* **25**: 269–90.
- Horvitz CC, Pascarella JB, McMann S, *et al.* 1998. Functional roles of invasive non-indigenous plants in hurricane-affected subtropical hardwood forests. *Ecol Appl* **8**: 947–74.
- Karl TR, Meehl GA, Miller CD, *et al.* (Eds). 2008. Weather and climate extremes in a changing climate. Washington, DC: US Climate Change Science Program and the Subcommittee on Global Change Research, US Department of Commerce, NOAA National Climatic Data Center.
- Labbe TR and Fausch KD. 2000. Dynamics of intermittent stream habitat regulate persistence of a threatened fish at multiple scales. *Ecol Appl* **10**: 1774–91.
- Larson ER, Magoulick D, Turner C, and Laycock KH. 2009. Disturbance and species displacement: different tolerances to stream drying and desiccation in a native and an invasive crayfish. *Freshwater Biol* **54**: 1899–908.
- Medeiros ESF and Maltchik L. 1999. The effects of hydrological disturbance on the intensity of infestation of *Lernaea cyprinacea* in an intermittent stream fish community. *J Arid Environ* **43**: 351–56.
- Meehl GA, Stocker TF, Collins WD, *et al.* 2007. Global climate projections. In: Solomon S, Qin D, Manning M, *et al.* (Eds). Climate change 2007: the physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge, UK, and New York, NY: Cambridge University Press.
- Milly PCD, Wetherald RT, Dunne KA, and Delworth TL. 2002. Increasing risk of great floods in a changing climate. *Nature* **415**: 514–17.
- Nichols FH, Thompson JK, and Schemel LE. 1990. Remarkable invasion of San Francisco Bay (California, USA) by the Asian clam (*Potamocorbula amurensis*). II. Displacement of a former community. *Mar Ecol-Prog Ser* **66**: 95–101.
- Nippert JB, Butler JJ, Kluitenberg GJ, *et al.* 2010. Patterns of *Tamarix* water use during a record drought. *Oecologia* **162**: 283–92.
- Petes LE, Menge BA, and Murphy GD. 2007. Environmental stress decreases survival, growth, and reproduction in New Zealand mussels. *J Exp Mar Biol Ecol* **351**: 83–91.
- Schoener TW, Spiller DA, and Losos JB. 2001. Predators increase the risk of catastrophic extinction of prey populations. *Nature* **412**: 183–86.
- Song LY, Chow WS, Sun LL, *et al.* 2010. Acclimation of photosystem II to high temperature in two *Wedelia* species from different geographical origins: implications for biological invasions upon global warming. *J Exp Bot* **61**: 4087–96.
- Sorte CJB, Fuller A, and Bracken MES. 2010. Impacts of a simulated heat wave on composition of a marine community. *Oikos* **119**: 1909–18.
- Steiner SCC, Macfarlane KJ, Price LM, and Willette DA. 2010. The distribution of seagrasses in Dominica, Lesser Antilles. *Revista De Biologia Tropical* **58**: 89–98.
- Tebaldi C, Hayhoe K, Arblaster JM, and Meehl GA. 2006. Going to the extremes: an intercomparison of model-simulated historical and future changes in extreme events. *Climatic Change* **75**: 185–211.
- Werner S and Rothhaupt KO. 2008. Mass mortality of the invasive bivalve *Corbicula fluminea* induced by a severe low-water event and associated low water temperatures. *Hydrobiologia* **613**: 143–50.
- White TA, Campbell BD, Kemp PD, and Hunt CL. 2001. Impacts of extreme climatic events on competition during grassland invasions. *Glob Change Biol* **7**: 1–13.