

Involvement of facultative apomixis in inheritance of *EPSPS* gene amplification in glyphosate-resistant *Amaranthus palmeri*

Daniela N. Ribeiro · Zhiqiang Pan · Stephen O. Duke ·
Vijay K. Nandula · Brian S. Baldwin · David R. Shaw ·
Franck E. Dayan

Received: 20 August 2013 / Accepted: 28 September 2013 / Published online: 20 October 2013
© Springer-Verlag Berlin Heidelberg 2013

Abstract The inheritance of glyphosate resistance in two *Amaranthus palmeri* populations (R1 and R2) was examined in reciprocal crosses (RC) and second reciprocal crosses (2RC) between glyphosate-resistant (R) and -susceptible (S) parents of this dioecious species. R populations and Female-R × Male-S crosses contain higher 5-enolpyruvylshikimate-3-phosphate synthase (*EPSPS*) gene copy numbers than the S population. *EPSPS* expression, *EPSPS* enzyme activity, *EPSPS* protein quantity, and level of resistance to glyphosate correlated positively with genomic *EPSPS* relative copy number. Transfer of resistance was more influenced by the female than the male parent in spite of the fact that the multiple copies of *EPSPS* are amplified in the nuclear genome. This led us to hypothesize that this perplexing pattern of inheritance may result from apomictic seed production in *A. palmeri*. We confirmed that reproductively isolated R and S female plants produced seeds, indicating that *A. palmeri* can produce seeds both sexually and apomictically (facultative apomixis). This apomictic trait accounts for the low copy number inheritance in the Female-S × Male-R offsprings.

Apomixis may also enhance the stability of the glyphosate resistance trait in the R populations in the absence of reproductive partners.

Keywords *Amaranthus palmeri* · Apomixis · Asexual reproduction · 5-Enolpyruvylshikimate-3-phosphate synthase (*EPSPS*) · Facultative apomixis · Glyphosate · Herbicide resistance · Palmer amaranth

Abbreviations

ALS	Acetolactate synthase
CNT	Counts per mm ²
EPSPS	5-Enolpyruvylshikimate-3-phosphate synthase
GR	Glyphosate-resistant
GS	Glyphosate-susceptible

Introduction

The non-selective herbicide glyphosate (*N*-(phosphonomethyl) glycine) has become the leading herbicide worldwide, largely because of its use with transgenic, glyphosate-resistant (GR) crops (Duke and Powles 2008). Glyphosate inhibits 5-enolpyruvylshikimate-3-phosphate synthase (*EPSPS*), an enzyme of the shikimate pathway, thereby, preventing the biosynthesis of the aromatic amino acids phenylalanine, tyrosine and tryptophan (Siehl et al. 1997; Steinrücken and Amrhein 1980). GR crop technology was utilized on ca. 128 million ha in 2012 worldwide (James 2012), boosting farm income globally by \$32.2 billion (Brookes and Barfoot 2013) and significantly reducing environmental impact of weed management (Bonny 2011; Gardner and Nelson 2008; Kleter and Kuiper 2003).

Although strong arguments were made against the likelihood of weeds evolving resistance to glyphosate (Bradshaw

Electronic supplementary material The online version of this article (doi:10.1007/s00425-013-1972-3) contains supplementary material, which is available to authorized users.

D. N. Ribeiro · B. S. Baldwin · D. R. Shaw
Department of Plant and Soil Sciences, Mississippi State
University, Starkville, MS 39762, USA

Z. Pan · S. O. Duke · F. E. Dayan (✉)
Natural Products Utilization Research Unit, United States
Department of Agriculture, Oxford, MS 38677, USA
e-mail: fdayan@olemiss.edu; franck.dayan@ars.usda.gov

V. K. Nandula
Crop Production Systems Research Unit, United States
Department of Agriculture, Stoneville, MS 38776, USA

et al. 1997), the first case of a GR weed was reported in *Lolium rigidum* (Gaud) about 20 years following the introduction of the herbicide glyphosate (Heap 2013; Powles et al. 1998; Pratley et al. 1999). Resistance to glyphosate has now been confirmed in 23 other weed species (Heap 2013). Evolved glyphosate resistance has recently emerged in *Amaranthus palmeri* (S. Wats.) populations (Culpepper et al. 2006).

Amaranthus palmeri is a tall, upright, and dioecious summer annual species that is native to the Mexican states of Sonora and Baja California, as well as parts of southern Arizona and California, USA (Ehleringer 1983). *A. palmeri* was first described from specimens collected from San Diego County, CA, and along the banks of the Rio Grande River (Watson 1877). The species has since spread into the southeastern United States, where it has become a common and competitive weed in row crop production. Interference from *A. palmeri* affects the growth and yield of several crops that are now mostly GR crops in the USA: maize (*Zea mays* L.) (Massinga et al. 2001), soybean [*Glycine max* (L.) Merr.] (Bensch et al. 2003), and cotton (*Gossypium hirsutum* L.) (Morgan et al. 2001). Currently, GR *A. palmeri* infests more than two million ha in 17 US states (Heap 2013). GR *A. palmeri* has become the single greatest threat to the economic sustainability of cotton production in infested fields, largely due to the lack of adequate control provided by available post emergence herbicides (Mayo et al. 1995).

The steady increase in the number of weed species with evolved resistance to glyphosate has been exacerbated with the introduction and widespread adoption of GR crops (Duke and Powles 2009; James 2012). In evolved resistance, single base pair mutations in *EPSPS* have not provided a high level of resistance (Baerson et al. 2002; Ng et al. 2003; Perez-Jones and Mallory-Smith 2010). Alteration of more than one codon, as with site-directed mutagenesis to produce the GA21 version of maize *EPSPS*, has resulted in commercial GR maize with a much higher level of resistance (Green 2009). GA21-type mutations resulting from the double Thr to Ile101 and Pro to Ser 106 (TIPS) mutation was recently reported in a Malaysian goosegrass biotype (Jalaludin et al. 2013). The most recently evolved GR weed populations have other mechanisms of resistance based on amplification of the *EPSPS* gene(s) (Gaines et al. 2010) or sequestration of glyphosate in the vacuole (Ge et al. 2010, 2011, 2012).

Gene amplification of target site genes is a known resistance mechanism for insecticides (Bass and Field 2011) and fungicides (Selmecki et al. 2008). In the laboratory, step-wise increases in glyphosate concentrations of plant cell cultures has been selected for gene amplification of *EPSPS* (Pline-Srnic 2006). Amplification of the *EPSPS* gene is a mechanism for field-evolved glyphosate

resistance in *A. palmeri* (Gaines et al. 2010), *Lolium perenne* L. ssp. *multiflorum* (Lam.) Husnot (Salas et al. 2012) and *Kochia scoparia* (L.) Schrad. (Wiersma 2012). Glyphosate is the only herbicide for which there is field-evolved amplification of a target gene known to confer resistance.

In the case of *A. palmeri*, glyphosate resistance correlates with increased copy number of *EPSPS* genes, transcripts, protein content, and enzymatic activity (Gaines et al. 2011). These correlations indicate that between 30 and 50 *EPSPS* copies endow complete resistance to the normal field rate of glyphosate (0.5–1 kg ha⁻¹). These extra copies are found throughout the genome, on every chromosome (Gaines et al. 2010). The mechanism leading to the multiplication of gene copies has recently been linked to sequences with homology to miniature inverted-repeat transposable elements (MITEs) flanking the *EPSPS* gene copies in R individuals (Gaines et al. 2013). Other genetic elements such as a putative activator (Ac) transposase and a repetitive sequence region were also reported, though the precise mechanism of gene amplification is yet to be fully determined. Studies by Gaines and co-workers were all conducted on an *A. palmeri* population that evolved in the southeastern coastal plain of the USA. GR populations of this species were later found in 17 other US states extending throughout the south and as far west as California and as far north as Michigan (Heap 2013). This GR trait can be transferred via pollen to at least three other *Amaranthus* species (Gaines et al. 2012).

Very little is understood about the inheritance of this novel mode of resistance to glyphosate. The objectives of this research were to investigate the molecular mechanism of resistance to glyphosate in *A. palmeri* populations from Mississippi and investigate the mode of inheritance of this novel mechanism of resistance in this GR weed.

Materials and methods

Plant material and general experimental conditions

Seeds of suspected GR (R) populations of *A. palmeri* were collected from 35 GPS site coordinates across the northwestern region of Mississippi (MS), USA, in 2007 (Nandula et al. 2012). Each location corresponded to a population, with each population defined as a group of seed collected from 10 to 20 randomly selected female plants. A preliminary screening with 0.84-kg glyphosate ha⁻¹ was conducted on these 35 populations. The C1B1 and T4B1 accessions were selected for subsequent research because they had the greatest number of surviving individuals. A known susceptible (S) population, collected in Washington County, MS, was included

for comparison in all experiments. Seeds were stored at 10 °C until further use.

Experiments were carried out as described by Nandula et al. (2012). Seeds were planted at 1-cm depth in 50-cm by 20-cm by 6-cm plastic trays with holes containing a commercial potting mix (Metro-Mix 360®, Sun Gro Horticulture, Bellevue, WA, USA). Two weeks after emergence, *A. palmeri* plants were transplanted into 6-cm by 6-cm by 6-cm pots containing potting mix. Plants were greenhouse grown under 30/20 °C day/night temperature with a 14-h photoperiod, sub-irrigated as needed and fertilized with a nutrient solution (All Purpose Miracle-Gro®, The Scotts Company, Marysville, OH, USA) containing 200 mg L⁻¹ each of N, P₂O₅, and K₂O 1 week after transplanting and then once per month. Plants were sprayed at the four- to six-leaf stage with glyphosate (Roundup WeatherMAX®, Monsanto Company, St. Louis, MO, USA) using an 8002E nozzle and overhead compressed-air sprayer calibrated to deliver 140 L ha⁻¹ at 280 kPa.

Production of genetic populations

The first controlled crosses (Female-R × Male-R, R/R) were produced by first treating the C1B1 or T4B1 accessions (Nandula et al. 2012) with 0.84-kg glyphosate ha⁻¹, as previously described. At least 100 surviving individuals of each gender were selected and grown together in isolation from other populations in different greenhouses to ensure genetic purity of each controlled cross. Pollen from the male plants was spread on the female plants every morning over a period of 2 months. *A. palmeri* inflorescence spikes were hand-harvested where the majority of the seed-possessed coats were black in color (seed maturity). Seeds were air-dried in a greenhouse (25/20 °C day/night, 12-h photoperiod under natural sunlight conditions) for 7 days, cleaned, and stored at 10 °C until further use. Resistance was confirmed by treating a subset of whole plant first R/R controlled cross progeny with 0.84-kg glyphosate ha⁻¹ before proceeding with the second controlled crosses (data not shown). Second controlled crosses (R/R) were developed using at least 100 first R/R controlled crosses surviving individuals that were treated with 0.84-kg glyphosate ha⁻¹, as described for the development of the first R/R controlled crosses. A subset of second R/R controlled cross plants was grown and their resistance to glyphosate was confirmed following the previously described procedure (data not shown). Seeds from the C1B1 or T4B1 accessions having been through the first and second controlled crosses were, respectively, defined as resistant parents R1 and R2 accessions to generate subsequent genetic populations.

Reciprocal crosses (Female-S × Male-R, S/R, and Female-R × Male-S, R/S) were made by growing S

females in isolation with each R male (R1 and R2), and vice versa in separated greenhouses with the aim of generating four reciprocal crossed (RC) populations, hereafter referred to as S/R1, S/R2, R1/S, and R2/S. Plants were pollinated as previously described and resistance confirmed by the application of glyphosate at 0.42 kg ae ha⁻¹ to a subset of the RC generations. The majority (≥85 %) of S/R1 and S/R2 cross plants treated with glyphosate at the described rate were susceptible (data not shown) indicating homozygous susceptible individuals.

Second reciprocal crosses (Female-S/R × Male-S/R, S/R//S/R, and Female-R/S × Male-R/S, R/S//R/S) were made by growing each RC female in isolation with each RC male in separated greenhouses with the aim of creating four second reciprocal crossed (2RC) populations, hereafter referred to as S/R1//S/R1, S/R2//S/R2, R1/S//R1/S, and R2/S//R2/S. Plants were pollinated as previously described and resistance confirmed by the application of glyphosate at the low rate of 0.42 kg ae ha⁻¹ to a subset of the 2RC generations. The majority (≥70 %) of plants of S/R1//S/R1 and S/R2//S/R2 populations were susceptible, some plants of all four 2RC populations had an intermediate level of injury, and the majority (≥80 %) of plants of R1/S//R1/S and R2/S//R2/S were resistant (data not shown).

These experiments were designed to determine the inheritance of glyphosate resistance at a population level, rather than at the level of individuals. Therefore, crosses were performed with groups of plants with predetermined sensitivity or resistance to glyphosate. For these experiments, we assumed that little or no reproductive fitness penalty resulted in generating these crosses.

DNA, RNA isolation and cDNA synthesis

Total DNA and RNA were extracted from frozen 2- to 4-leaf stage tissue of S, R1, R2, RC, and 2RC plants. Genomic DNA (gDNA) was extracted using DNeasy Plant Mini kit (Qiagen, Valencia, CA, USA). Total RNA was isolated with the RNeasy Plant Mini Kit (Qiagen) according to the manufacturer's instructions and treated with the RNase-Free DNase (Qiagen). The quality and quantity of prepared genomic DNA and total RNA were assessed according to the MIQE Guidelines (Bustin et al. 2009). The DNA and RNA were quantified using a NanoDrop (ND-1000) spectrophotometer (Thermo Scientific, Wilmington, DE 19810) and checked for quality and integrity by TAE agarose gel electrophoresis. For quantitative real-time PCR (qPCR), cDNA was synthesized from 2 µg of total RNA in a 20 µL reaction volume using a proprietary blend of oligo(dT) and random hexamer primers according to the manufacturer's instructions (iScript cDNA Synthesis Kit, Bio-Rad Laboratories, Hercules, CA, USA).

EPSPS cDNA sequencing

Total RNAs for cDNA cloning were isolated from S, R1 and R2 populations as described above. First strand cDNA synthesis was then performed using 1 μg total RNAs and M-MuLV reverse transcriptase in a final volume of 20 μL using oligo(dT)₂₃VN according to the manufacturer's instructions (New England BioLabs, Ipswich, MA, USA). A pair of primers (sense: 5'-TGGCTCAAGCTACTACCATC AAC-3'; antisense: 5'-ATATAGCTACTCAATGCTTGGCG AAC-3') was designed based on the *EPSPS* coding sequence from *A. palmeri* (GenBank accession number FJ861242) (Gaines et al. 2010). PCR reaction contained 1 μL cDNA; 0.1 mM each of forward and reverse primers; 0.2 mM each of dATP, dCTP, dGTP, and dTTP; 2 mM MgCl_2 ; and 1 U of proof reading PfuUltra high-fidelity DNA polymerase (Stratagene, La Jolla, CA, USA) with a 1 \times concentration of supplied buffer in a final volume of 50 μL . The cycling conditions were 2 min at 95 °C followed by 30 cycles of 20 s at 95 °C, 20 s at 55 °C, and 1 min at 72 °C, with a final extension of 3 min at 72 °C. PCR products were ligated into pCR Blunt TOPO vector using Zero Blunt TOPO PCR Cloning Kit (Invitrogen, Carlsbad, CA, USA), transformed into *Escherichia coli* TOP10 cells and grown on selection media. Single colonies of six transformants of R1 and R2, and 11 transformants from S were cultured overnight in liquid LB media for plasmid extraction. Plasmid DNAs were sequenced and assembled using Lasergene version 10.0 SeqMan (DNASTAR, Madison, WI, USA). Multiple DNA sequence alignments of *EPSPS*, including sequences from both R and S *A. palmeri* populations (Gaines et al. 2010), GenBank accession numbers FJ861242 and FJ861243, were constructed using ClustalW version 10.0 (DNASTAR). RNA extraction and amplification of the *EPSPS* gene was performed on pooled leaf samples from S, R1 and R2 populations.

Quantitative PCR

Quantitative real-time PCR (qPCR) was used to measure *EPSPS* genomic copy number relative to the gene for acetolactate synthase (*ALS*) and cDNA expression level of *EPSPS* relative to *ALS* in S, R1, R2, RC, and 2RC *A. palmeri* plants according to previously described procedures (Gaines et al. 2010). The *ALS* gene was used as a low-copy control gene with known monogenic inheritance in other *Amaranthus* species (Trucco et al. 2005). The first experiment measured *EPSPS* genomic copy number and expression level in the population (i.e., combined leaf material from at least ten individuals per population studied). The second experiment measured *EPSPS* genomic copy number by individuals of each population (i.e., 30 individuals per population were analyzed separately).

Protein extraction and EPSPS enzyme activity assay

Ten grams of fresh leaf tissue was frozen in a -80 °C refrigerator prior to being ground to fine powder in a chilled mortar. Total soluble protein (TSP) was extracted from 2- to 4-leaf stage tissue of S, R1, R2, RC, and 2RC plants. Protein extraction and EPSPS assay were conducted following published protocols (Sammons et al. 2007; Webb 1992) and described in detail in Dayan et al. (2013).

The specific activity of EPSPS from *A. palmeri* plants was determined using EnzCheck phosphate assay kit (Invitrogen) with a UV-3101 spectrophotometer (Shimadzu North America, Columbia, MD, USA) as described in detail elsewhere (Dayan et al. 2013). The effect of glyphosate was tested in triplicates at concentrations ranging from 0.3 to 1,000 μM in half-log intervals. Two experiments were conducted with three replications per population. The collected samples consisted of combined leaf material from at least ten individuals per population studied.

Western blot analysis for the detection of EPSPS protein

Total cellular protein was isolated from 2- to 4-leaf stage tissue of S, R1, R2, RC, and 2RC plants. A quantity of 0.25 g of fresh leaf tissue was ground to fine powder in a chilled mortar as described above. Powdered tissues were homogenized in 500 μL cold extraction buffer (100 mM MOPS, 5 mM EDTA, 10 % glycerin, and 50 mM KCl, pH 7.0) with freshly added protease inhibitor (1 tablet per 10 mL of buffer) (Roche Applied Science, Indianapolis, IN, USA). The samples were thawed at room temperature and vortexed for 10 s. The mixtures were kept cold and placed on a Geno/Grinder mechanical shaker (SPEX SamplePrep, Metuchen, NJ, USA) for 1 min at 1,750 strokes per min. This step was repeated three times. Subsequently, the extract was centrifuged for 5 min at 18,000g. The supernatant was transferred to a new vial and it was centrifuged for 5 min at 18,400g. This step was repeated one more time and extract was stored at -80 °C. Protein concentrations were determined using a Bradford assay (Bradford 1976).

Western blot analysis for the detection of EPSPS protein was performed diluting soluble protein in 950- μL Laemmli premixed protein sample buffer (Bio-Rad Laboratories) and 0.71 M β -mercaptoethanol to reach a final concentration of 15 μg μL^{-1} . The sample was transferred to a heating block at 90 °C for 4 min and allowed to cool to room temperature. The soluble protein was separated on a 12 % Tris-HCl precast polyacrylamide gel (Bio-Rad Laboratories). After separation, the proteins were blotted onto 0.45 μM Immobilon-FL PVDF membranes (EMD Chemicals, San Diego, CA, USA) and the membrane was equilibrated with transfer buffer [0.2 % methanol and Tris/glycine buffer (2.5 mM Tris, 19.2 mM glycine, pH 8.3)]. The membrane

was washed with Tris-buffered saline (TBS) (20 mM Tris, pH 7.5 and 500 mM NaCl) and incubated overnight at 4 °C with 3 % gelatin from cold water fish skin. Western blot was probed with an EPSPS specific antibody (2 mg mL⁻¹) developed against recombinant maize EPSPS (Monsanto Company) at dilution 1:2,000 and re-probed against Alexa Fluor 635 goat anti-rabbit (Invitrogen) at dilution 1:2,000. Within 20 min of the final wash with TBS buffer and Tween-20, the array was scanned and analyzed on a Pharos FX Plus Molecular Imager system equipped with an external 635-nm laser and quantified by densitometric analysis using Quantity One analysis software (Bio-Rad Laboratories) and represented as CNT (counts mm⁻²). The collected samples consisted of combined leaf material from at least ten individuals per population studied with three replications per population.

Glyphosate dose–response bioassay

Plants from parental lines (R1 and R2) and from the first and second reciprocal crosses were submitted to a dose–response experiment by applying glyphosate at 0, 0.052, 0.105, 0.21, 0.42, 0.84, 1.68, 3.36, and 6.72 kg ae ha⁻¹ at the 3- to 4-leaf growth stage. Two weeks after treatment (WAT) shoots were clipped at soil level and fresh weight was determined. Biomass data are reported as percentage of the non-treated control. The experiment was conducted two times and arranged in a completely randomized design with triplicates.

Shikimate bioassay

Shikimate content of tissues of *A. palmeri* populations was determined spectrophotometrically in microtiter plates according to Shaner et al. (2005). Glyphosate concentrations ranged from 10 to 333 µM. The absorbance at 380 nm was measured within 30 min in a PowerWave XS microplate reader (Biotek, Winooski, VT, USA). Shikimate in µg mL⁻¹ was determined based on a standard curve. Two experiments were conducted with four replications per population.

Testing for facultative apomixis

To test for apomixis in *A. palmeri* parent (S, R1 and R2) populations, pollination bags were placed over main inflorescences prior to emergence and were examined daily to determine their sex. The male plants were discarded and the female plants were isolated in different geographically located greenhouses to ensure repeatability of the apomixis test and exclude external sources of pollen contamination. The inflorescences were enclosed in pollination bags for the whole duration of the experiments. A total of 44 S

individuals were grown, Spring/Summer, 2011, in Oxford (MS), 36 R1 individuals in Starkville (MS), and 38 R2 individuals in Abbeville (MS). *A. palmeri* inflorescence spikes were hand-harvested where the majority of the seed-possessed coats were black in color (seed maturity). *A. palmeri* seeds were air-dried, cleaned, and stored at 10 °C as previously described. The sex segregation ratio of the offsprings was determined by germinating the apomictically produced seeds and growing the plants until their sex could be determined by observing their flowers. A similar experiment was carried out during the winter, when no *A. palmeri* plants remained alive outside the greenhouse, to insure that no pollen could accidentally fertilize our reproductively isolated female plants. The inflorescence were also enclosed in pollination bags to further prevent pollination.

Intraspecific genetic diversity and relationships

Genetic markers that might confirm apomixis in *A. palmeri* were selected based on information from available literature (Chan and Sun 1997; Chandi et al. 2013; Lanoue et al. 1996; Lee et al. 2009; Ma et al. 2008; Popa et al. 2010; Wassom and Tranel 2005; Wetzel et al. 1999; Xu et al. 2011). RAPD (random-amplified polymorphic DNA) and ISSR (inter-simple sequence repeat) methods were used because they can provide up to a population level of identification of intraspecific variation.

DNA extraction was carried out as previously described from three different plants of the R1 population ($n = 3$) and diluted to 2 ng µL⁻¹. RAPD analyses were performed using 12 decamer primers (Online Resource 1) randomly selected from a list of 100 primers (NAPS Unit, University of British Columbia, Biotechnology Laboratory, Vancouver, BC, Canada). ISSR analyses were performed using 15 primers (Online Resource 1) randomly selected from a list of 100 primers (NAPS Unit, University of British Columbia) and five provided by Natascha Techen (National Center for Natural Products, University of Mississippi, Oxford, USA). PCR reaction contained 10 ng DNA; 0.1 µM each of forward and reverse primers; 200 µM each of dATP, dCTP, dGTP, and dTTP (Promega, Madison, WI, USA); 1.5 mM MgCl₂; and 1 unit of Platinum Taq DNA polymerase (Invitrogen) with a 1× concentration of supplied buffer in a final volume of 13.6 µL. The cycling conditions included 3 min at 94 °C followed by 45 cycles of 30 s at 94 °C, 30 s at 50 °C, and 3 min at 72 °C, with a final extension of 7 min at 72 °C in a thermal cycler (PTC-225, MJ Research, Watertown, MA, USA). PCR products were analyzed by electrophoresis on a 2 % TAE agarose gel stained with ethidium bromide and visualized under UV light using Quantity One (Bio-Rad, version 4.3.0). The sizes of the PCR products were compared to the molecular size standard 1 kb plus DNA ladder (Invitrogen).

Statistical analysis

Data from *EPSPS* genomic copy number and gene expression, *EPSPS* protein quantity and shikimate levels were analyzed by ANOVA via the PROC GLM statement using SAS software (version 9.3, SAS Institute, Cary, NC, USA) to determine the main effects and interactions of the factors at $P < 0.05$. No significant experiment effect was observed in repeated experiments; therefore, data from those experiments were pooled.

Where ANOVA indicated significant differences between treatments, means separation were performed using Fisher's protected least significant difference (LSD) test at $P = 0.05$ using SAS software. The data from *EPSPS* relative copy number were regressed against *EPSPS* cDNA relative expression level and against *EPSPS* enzyme activity.

The whole plant glyphosate dose–responses were analyzed using a four parameters log-logistic model (Seefeldt et al. 1995), whereas the in vitro dose–responses were analyzed with a three parameters log-logistic model using R software (version 2.15.2, R Foundation for Statistical Computing, Vienna, Austria) using package *drc*, *drm* and *modelFit* function. The graphs were generated with Sigma Plot (version 11, Systat Software, San Jose, CA, USA). Subsequently, the Student *t* test at $P < 0.05$ was used to determine any differences in potency between populations and the S population at the respective GR_{50} or IC_{50} effect level. The null hypothesis, H_0 : GR_{50} or IC_{50} populations/ GR_{50} or IC_{50} S population = 1, was tested against the alternative hypothesis, H_a : GR_{50} or IC_{50} populations/ GR_{50} or IC_{50} S population \neq 1. This test was performed using the open-source R software using package *drc*, *drm* function, and the comparisons were given by means of the selectivity index (SI) function.

Results

EPSPS cDNA sequencing

There was little difference among *EPSPS* sequences from the R1 (six clones) and R2 (six clones) populations. Consequently, the consensus sequence represents residues common to all GR sequences (R_consensus) (Online Resource 2). Alignment of consensus sequences from R and S individuals had several polymorphisms in all S sequences in the alignment contig when compared with the reference S sequence (FJ861242, Gaines et al. 2010). This may be the result of combining seeds from several plants at the time of initial collection of this population.

The cDNA sequence analysis of the *EPSPS* gene in both GR (R1 and R2) populations revealed several nucleotide substitutions resulting in silent mutations when compared with the S population sequence. The S and R sequences had 99 and 100 % homology to the *A. palmeri* reference sequence FJ861242 (S) of Gaines et al. (2010), respectively (online resource 2). Only one of the three single nucleotide polymorphisms (SNP) between S and R clones (contig position 866) resulted in a non-synonymous mutation (arginine to lysine amino acid substitution) at position 215, based on the maize mature *EPSPS* numbering system (Fig. 1), in the GS *A. palmeri* (Lebrun et al. 1997).

EPSPS gene copy number correlates with *EPSPS* gene expression

Analysis of the genomic DNA extracted from pooled samples of several individuals for each population showed that on average, R1 and R2 populations contained 59 and 33 copies of the *EPSPS* gene, whereas the S population contained a single copy (Fig. 2b). Quantitative RT-PCR on cDNA revealed that the single copy of *EPSPS* gene was proportionally



Fig. 1 Partial and deduced amino acid sequence alignment of the 5-enolpyruvylshikimate-3-phosphate synthase (*EPSPS*) gene of glyphosate-susceptible (S) and glyphosate-resistant (R) *A. palmeri* populations. The boxed codon shows the substitutions of arginine (R) to lysine (K) at amino acid 215 and no substitution at proline (P) 106, when using the maize mature *EPSPS* numbering system. ^aFJ861243,

glyphosate-resistant *A. palmeri* from Georgia (Gaines et al. 2010) sequence reference, sequence information can be found at <http://www.ncbi.nlm.nih.gov/nuccore/> ^bFJ861242, glyphosate-susceptible *A. palmeri* from Georgia (Gaines et al. 2010) sequence reference, sequence information can be found at <http://www.ncbi.nlm.nih.gov/nuccore/>




Fig. 2 Glyphosate effect (a), genomic copy number of *EPSPS* (b), and relative transcription level of *EPSPS* (c) in glyphosate-susceptible (S), -resistant (R1 and R2), and first (R/S and S/R) and second (R1/S//R1/S and S/R1//S/R1) reciprocal crosses of *A. palmeri* populations. Vertical bars represent \pm standard error of the mean ($n = 20$). Letters represent statistical difference based on Fisher’s LSD at $P = 0.05$

expressed, with a 1:1 correlation to *EPSPS* gene copy:*EPSPS* transcript richness (Figs. 2, 3). The *EPSPS* copy number was positively correlated with the gene expression level ($r = 0.94$, $P < 0.0001$, excluding R2 data, Fig. 3), and a pattern was observed where the reciprocal crosses generated by Female-S \times Male-R (S/R) and by Female-S/R \times Male-S/R (S/R//S/R) had lower *EPSPS* expression level than the reciprocal crosses generated by Female-R \times Male-S (R/S) and by Female-R/S \times Male-R/S (R/S//R/S). While the R2 population is shown in the graph (Fig. 3), it was excluded from the data used for the regression because it did not have a strong correlation between its relatively high number of gene copies and unexpectedly low number of mRNA transcript.

When individuals from each population were analyzed separately, there was a strong ($P < 0.0001$) relationship between *EPSPS* copy number and the populations (Fig. 4),




Fig. 3 Positive correlation between increase in 5-enolpyruvylshikimate-3-phosphate synthase (*EPSPS*) relative to acetolactate synthase (*ALS*) genomic copy number with increase in *EPSPS*: *ALS* cDNA expression levels in glyphosate-susceptible (S), -resistant (R1 and R2), and first reciprocal crosses (R/S and S/R) of *A. palmeri* populations. Regression of the entire dataset indicated a good relationship between mRNA levels and copy numbers (solid line, $r = 0.87$, $P < 0.0001$). The R2 population did not fit the model as well as the other populations. Excluding this population from the dataset improved the strength of that relationship (dotted line, $r = 0.94$, $P < 0.0001$)

with S, S/R1 and S/R2 with the fewest copy numbers. In fact, all of the S and 73 % of S/R1 and 70 % of S/R2 individuals analyzed had a single copy of the *EPSPS* gene. R1 and R2 populations had the highest copy numbers, and R1/S and R2/S had intermediate numbers (Fig. 4).

EPSPS gene copy number correlates with *EPSPS* enzyme activity and protein quantity

EPSPS specific activity in each *A. palmeri* population was inhibited by glyphosate concentrations (Online Resource 3, Fig. 5). The amounts of glyphosate needed to reduce the *EPSPS* activity by 50 % (IC_{50}) ranged from 5.5 to 55 μ M glyphosate (Online Resource 3), but there was no clear relationship between the IC_{50} values at the enzymatic level and in vivo sensitivity to glyphosate (i.e., the sensitive biotype had the highest IC_{50} value). Consequently, the R/S ratios for all populations were smaller than 1, suggesting that *EPSPS* from the GR plants was more sensitive to glyphosate than in the susceptible population.

The specific activity of *EPSPS* varied greatly between populations in the absence of glyphosate. The S biotype had




Fig. 4 Genomic copy number of 5-enolpyruvylshikimate-3-phosphate synthase (*EPSPS*) relative to acetolactate synthase (*ALS*) genomic copy of 30 ($n = 30$) sampled individuals per *A. palmeri* population (dark circle), glyphosate-susceptible (*S*), -resistant (*R1* and *R2*), and first (*R/S* and *S/R*) reciprocal crosses. Means of *EPSPS* relative gene copy number followed by the same letter are not significantly different by LSD test at 0.05. The *EPSPS* copy number segregation pattern observed in Fig. 2b, when combined sample population, is observed when averaging sampled individuals (dark gray square)




Fig. 5 Dose–response of glyphosate against *EPSPS* (5-enolpyruvylshikimate-3-phosphate synthase) enzyme activity of glyphosate-susceptible (*S*), -resistant (*R1* and *R2*), and first reciprocal cross (*R/S* and *S/R*) of *A. palmeri* populations. Glyphosate inhibition assays were normalized for total soluble protein (TSP) quantity. *S* black cross, 1 relative *EPSPS* copy, IC_{50} (glyphosate concentration that reduced enzyme activity by 50%) = 55 μM ; *R1* dark gray square, 59 relative copies, $IC_{50} = 16 \mu\text{M}$; *R2* black triangle, 33 relative copies, $IC_{50} = 21 \mu\text{M}$; *R1/S* light gray square, 43 relative copies, $IC_{50} = 15 \mu\text{M}$; *S/R1* open square, 19 relative copies, $IC_{50} = 7 \mu\text{M}$; *R2/S* light gray triangle, 30 relative copies, $IC_{50} = 6 \mu\text{M}$; *S/R2* open triangle, 1 relative copy, $IC_{50} = 26 \mu\text{M}$. *Pi* inorganic phosphate. Vertical bars represent \pm standard error of the mean ($n = 6$)




Fig. 6 Positive correlation among *EPSPS* (5-enolpyruvylshikimate-3-phosphate synthase) genomic copy number (a), *EPSPS* quantity (b), and specific activity of *EPSPS* enzyme (absence of inhibitor glyphosate) in glyphosate-susceptible (*S*), -resistant (*R1* and *R2*), and first (*R/S* and *S/R*) and second (*R/S//R/S* and *S/R//S/R*) reciprocal crosses of *A. palmeri* populations. *Pi* inorganic phosphate; *TSP* total soluble protein. Vertical bars represent \pm standard error of the mean ($n = 6$). Means of *EPSPS* activity with the same letter are not significantly different by Fisher's LSD test at 0.05

the lowest activity ($0.02 \mu\text{mol} \mu\text{g}^{-1} \text{protein min}^{-1}$), whereas both *R1* and *R2* parent populations had high activity (0.19 to $0.12 \mu\text{mol} \mu\text{g}^{-1} \text{protein min}^{-1}$, respectively) (Fig. 6c). In the RC populations, the *R1/S* and *R2/S* had intermediate *EPSPS* activity and had consistently higher activity than the *S/R1* and *S/R2* crosses ($r = 0.87$, $P < 0.0001$, including and excluding *R2* data) and *EPSPS* expression level relative to *ALS* ($r = 0.97$, $P < 0.0001$, excluding *R2* data; $r = 0.84$, $P < 0.0001$, including *R2* data) (Fig. 7). In the second reciprocal crosses (2RC), *R1/S//R1/S* had a level of *EPSPS* activity similar to *R1*. The other 2RC populations had intermediate levels of *EPSPS* specific activity, which was positively correlated with *EPSPS* relative copy number. Consistent with these observations, *EPSPS* protein level, as measured with immunoblotting, correlated strongly with other *EPSPS*-related parameters such as *EPSPS* gene copy number and enzymatic activity (Fig. 6).

EPSPS gene copy number correlates with level of glyphosate resistance

The shoot biomass of all *A. palmeri* population decreased in a dose-dependent manner with increasing glyphosate dose (Online Resource 4). Consistent with our preliminary data, the *S* population was the most sensitive to glyphosate,




Fig. 7 Positive correlation between specific activity of EPSPS (5-enolpyruvylshikimate-3-phosphate synthase) enzyme and *EPSPS* relative to acetolactate synthase (*ALS*) genomic copy number (**a**) and *EPSPS*: *ALS* cDNA expression levels (**b**) in glyphosate-susceptible (S), -resistant (R1 and R2), and first reciprocal cross (R/S and S/R) of *A. palmeri* populations. **a** regression of the entire dataset indicated a strong relationship between enzyme activity and copy number (solid and dotted line, $r = 0.87$, $P < 0.0001$). **b** Regression of the entire dataset indicated a good relationship between mRNA levels and enzyme activity (line not shown $r = 0.84$, $P < 0.0001$). The R2 population did not fit the model as well as the other populations. Excluding this population from the dataset improved the strength of that relationship (solid line, $r = 0.97$, $P < 0.0001$)

with a GR_{50} of 94.4 g ae ha^{-1} (Table 1). The R1 and R2 populations were 17 and 14.5 times more resistant, respectively, to glyphosate than S.

The S biotype had the highest shikimate accumulation at all glyphosate treatment concentrations, differing from the grouped R1 and R2 populations, which had

no accumulation except for the R2 at 333- μM glyphosate (Fig. 8). Shikimate accumulation was highly dependent on glyphosate concentration and between populations ($P < 0.005$ for both).

In all cases, the progeny of the RC and 2RC crosses had less resistance to glyphosate (lower GR_{50}) than the resistant parent (Table 1). Interestingly, the GR_{50} for the RC were closer to their respective maternal parent than the midpoint (Table 1). The 2RC dose–response was intermediate between those of the R and S populations, containing both highly susceptible and highly resistant individuals and a range of intermediate phenotypes (Table 1). Consistent with the pattern observed with the number of gene copy numbers, mRNA levels and protein expression, the greenhouse data suggest that the mode of inheritance of GR trait was maternally affected.

Facultative apomixis and intraspecific genetic diversity

Involvement of apomixis in glyphosate resistance inheritance was tested by measuring seed production in reproductively isolated female plants. In that experiment, reproductively isolated female individuals (44 S, 36 R1 and 38 R2) were studied. In all cases, seeds were produced with the exception of one R1 plant. From 60 to 100 % (depending on the population) of individuals studied produced 1–1,000 seeds, and some S individuals produced as many as 6,000 seeds (Table 2). The female:male ratio of the apomictically produced offspring was close to 1:1.

Attempts to monitor apomixis using RAPD and ISSR genetic markers were unsuccessful (Online Resources 5 and 6). No significant RAPD and ISSR polymorphism was observed in a preliminary test experiment on a subset of three R1 individuals. Therefore, none of these markers would have been useful in differentiating apomictically produced seeds from seeds resulting from a fertilized egg through sexual reproduction (Chandrika et al. 2010; Lata et al. 2010, 2011).

Discussion

Analysis of the *EPSPS* sequences from the S and R populations of *A. palmeri* from Mississippi revealed several polymorphisms. Only one of these SNPs resulted in a non-synonymous mutation (arginine to lysine amino acid substitution) at position 215 (Fig. 1). However, this mutation was in the *EPSPS* sequence from the glyphosate sensitive wild-type and is therefore not linked to resistance. Importantly, all the sequences possessed the normal proline 106 residue.

These *A. palmeri* populations from Mississippi evolved resistance to glyphosate by gene amplification of *EPSPS* as

Table 1 Glyphosate dose–response parameters and variables in the log-logistic model estimates for parents, reciprocal crosses and second reciprocal crosses of *A. palmeri* at 14 days after treatment

Population code ^{a, **}	R^2	% Fresh weight reduction (SE) ^d			GR_{50}^c g ae ha ⁻¹ (SE) ^d	GR_{50} ratio ^c
		L^c	U^c	s^c		
R1	0.99	16.56* (2.59)	100.24* (1.18)	3.42* (0.51)	1,623.3* (68.5)	17.2
R1/S	0.99	8.05* (2.07)	97.92* (1.21)	3.70* (0.41)	1,138.1* (44.7)	12.1
S/R1	0.99	3.69 (2.29)	100.75* (1.90)	1.69* (0.16)	464.2* (29.1)	4.9
R1/S//R1/S	0.98	6.63* (2.74)	97.98* (1.41)	2.20* (0.24)	976.4* (56.7)	10.4
S/R1//S/R1	0.99	7.08* (2.01)	98.02* (1.48)	2.65* (0.32)	663.6* (29.9)	7.0
R2	0.98	16.85* (2.72)	100.67* (1.23)	2.85* (0.37)	1,368.8* (71.3)	14.5
R2/S	0.99	3.54 (2.12)	97.85* (1.48)	2.35* (0.23)	758.9* (36.5)	8.0
S/R2	0.99	1.78 (1.79)	99.89* (1.88)	2.02* (0.18)	363.0* (18.7)	3.9
R2/S//R2/S	0.99	4.81* (1.91)	98.65* (1.51)	2.49* (0.25)	569.7* (26.5)	6.0
S/R2//S/R2	0.99	2.08 (1.55)	97.65* (1.53)	3.35* (0.41)	433.1* (14.8)	4.6
S	0.99	1.07 (1.37)	98.86* (2.58)	1.75* (0.16)	94.36* (5.6)	–

Model proposed by Seefeldt et al. (1995): y [fresh weight (% of untreated control)] = $L + \{(U - L)/[1 + (D/GR_{50})^s]\}$

^a Resistant parents (R1 and R2), susceptible parent (S), reciprocal crosses (Female-S × Male-R, S/R, and Female-R × Male-S, R/S), second reciprocal crosses (Female-S/R × Male-S/R, S/R//S/R, and Female-R/S × Male-R/S, R/S//R/S)

^c The parameter estimates are L , lower limit of response; U , upper limit of response; s , slope of the curve around the point of inflexion (GR_{50}); GR_{50} , glyphosate dose required to cause a 50 % reduction in plant growth and GR_{50} dose was estimated using responses to nine glyphosate doses (D) (0, 52, 105, 210, 420, 840, 1,680, 3,360, and 6,720 g ha⁻¹); and GR_{50} ratio, GR_{50} populations/ GR_{50} susceptible population

^d SE represents the standard error of the mean where $n = 6$ (polled data from two experiments)

* Estimated parameters of the log-logistic model are different according to Student t test at $P < 0.05$; accept alternative hypothesis, H_a : parameter $\neq 0$

** Relative potencies between populations and susceptible population at GR_{50} response level are different according to Student t test at $P < 0.05$; accept alternative hypothesis, H_a : relative potency $\neq 1$. The two exceptions, where the relative potencies of S/R1 vs. S/R2//S/R2 and R2/S vs. S/R1//S/R1 at GR_{50} response level are not different according to Student t test at $P < 0.05$; accept null hypothesis, H_0 : relative potency = 1




Fig. 8 Effect of glyphosate concentration on shikimate levels in excised leaf discs of glyphosate-resistant (R1 and R2) and -susceptible (S) *A. palmeri* populations. Means of shikimate level with the same letter are not significantly different by Fisher's LSD test at 0.05. Vertical bars represent \pm standard error of the mean ($n = 8$)

Table 2 Seed production of reproductively isolated female plants due to the effect of agamospermy/apomixis of glyphosate-susceptible (S) and -resistant (R1 and R2) *A. palmeri* plants used as parents to generate first reciprocal crosses (R/S and S/R)

Number of seeds	Biotypes		
	Number of individuals		
	S	R1	R2
0	0	0	0
1–999	29	25	38
1,000–1,999	8	5	0
2,000–2,999	3	4	0
>3,000	4	2	0

first reported by Gaines et al. (2010) in Georgia and in other species by others (Salas et al. 2012; Wiersma 2012). The populations have as many as 59 copies of *EPSPS* (Fig. 2b), which is similar to that reported in other GR biotypes of *A. palmeri*. The biotypes from Georgia had individual plants with as many as 160 copies of the *EPSPS* gene (Gaines et al. 2010), and six individuals from the S × R cross had 1–39 copies (Gaines et al. 2011). A GR population from North Carolina possessed 22–63 copies of *EPSPS* gene (Chandi et al. 2012).

As with other cases where resistance is associated in amplification of multiple *EPSPS* gene copy numbers, the specific activity of *EPSPS* in our R1 and R2 populations was six- to tenfold higher than in S plants (Fig. 6c and Online Resource 3). Similar differences in specific enzyme activity were observed in GR and GS *A. palmeri* from Georgia (Gaines et al. 2010), *L. perenne* L. ssp. *multiflorum* from Arkansas (Salas et al. 2012) and *K. scoparia* from the US central Great Plains (Wiersma 2012). R1 and R2 plants had a nine- to sixfold increase, respectively, in *EPSPS* enzyme activity relative to that of the S plants. Other *A. palmeri* populations had a 16-fold increase in specific activity between GR and GS (Gaines et al. 2010) and there was an average sixfold higher basal enzyme activity in GR *L. perenne* compared to the GS biotype (Salas et al. 2012).

In our study, the 2RC genomic copy number varied greatly and was an intermediate value between those of the R and S populations (R1/S//R1/S: 53, S/R1//S/R1: 5, R2/S//R2/S: 15, and S/R2//S/R2: 6), containing both highly susceptible and highly resistant individuals and a range of intermediate genotypes (Fig. 2b). The molecular mechanism leading to the insertion of multiple *EPSPS* copies throughout the genome of *A. palmeri* (Gaines et al. 2010) is not fully understood, but it is linked to the presence of MITEs flanking the *EPSPS* gene copies in R individuals (Gaines et al. 2013). Furthermore, this process is very dynamic, where one individual *A. palmeri* (S/R × S/R) had a higher relative copy number than the sum of their relative copy number from both parents (Gaines et al. 2010).

Reciprocal crosses generated by Female-R × Male-S (R/S) and by Female-R/S × Male-R/S (R/S//RS) of the populations tended to have higher *EPSPS* copy number than the other crosses (Fig. 2b). This pattern of inheritance where the direction of the cross affected the level of resistance suggests a maternal influence.

While the transfer of glyphosate resistance via pollen movement is evident in *A. palmeri*, the rate of transfer is inversely proportional to the distance from a R male parent (Sosnoskie et al. 2012). In fact, that study in which only GS female plants surrounded a central group of GR male plants, no resistance was transferred to the seeds of female plants located 600 m or more from the resistant male pollen donor. Nevertheless, GS seeds were produced by plants at all distances from the GR male plants (40 % at 5 m to 80 % at 250 m), and there was little difference in the number of seeds produced at distances of 50 m and farther from the pollen source. The authors proposed that these findings may be due to fertilization by autopolllination or from pollen providers outside the boundary of the experiment. These authors discounted agamospermy (apomixis) because the proportion of male to female offsprings approximated 1:1. However, all but one of the reproductively isolated *A. palmeri* plants in our study underwent

apomixis and produced seeds (Table 2) that yielded a similar ratio of male to female offspring.

Little is known about the inheritance of this incompletely dominant, nuclear encoded GR trait, but it might follow a polygenic additive pattern in populations of *A. palmeri* from Georgia and North Carolina (Chandi et al. 2012; Gaines 2009). However, both of these studies observed an unpredictable behavior in the inheritance of glyphosate resistance in some populations, which may be the result of facultative apomixis, even though the overall contribution of this process may be small at the population level under most conditions. Indeed, a relatively small number of seeds are produced by apomixis (Table 2) when compared with ‘normal’ seed production per female plant in the field reaching 200,000–600,000 seeds (Keeley et al. 1987). Nonetheless, production of apomictic seeds would be sufficient to maintain the resistant trait in a population.

Our observations that *A. palmeri* can produce seeds both sexually and apomictically (facultative apomixis) are consistent with some of the data by other groups (Chandi et al. 2012; Gaines 2009; Gaines et al. 2010, 2011; Sosnoskie et al. 2012). The occurrence of agamospermy in *A. palmeri* is also evident in several studies demonstrating that crosses between *A. palmeri* and *A. rudis* can yield non-hybrid progeny (Franssen et al. 2001; Steinau et al. 2003; Trucco et al. 2007; Wetzel et al. 1999). The ability of plants to reproduce apomictically is particularly advantageous because it “guarantees a protracted existence to the individual genotypes, over long periods and over wide areas” (Gustafsson 1947).

The current understanding of sex determination in dioecious species suggests that the change from unisexuality to bisexuality has been a short step in evolution (Chattopadhyay and Sharma 1991). Apomictic seeds derived from unreduced cells (apomeiosis) should be 100 % female. However, reproductively isolated *A. palmeri* individuals produced 1:1 female:male ratio of offspring apomictically. The production of both male and female apomictic offspring would require meiotic events and subsequent fusion (to restore the somatic number). Such events are known among apomictic species as synergid or antipodal apogamy. If the Y male sex determinant in *A. palmeri* were a non-heteromorphic chromosome, then it would segregate and assort during meiosis I with the other chromosomes of the genome. If gender of this dioecious species were determined by homozygous presence or absence of genes, then synergids or antipodals with W (conferring femaleness) or Y (conferring maleness) would produce genotypes; WW (a female) or YY (a male) in a 1:1 phenotypic ratio. While Westergaard (1958) reports the normal (sexual) male of *Acnida* spp. (an amaranth) to be heterozygous (XY); he also reports 8 of 18 plant families investigated showing possibility of the homozygous genotype (however, viability

is unknown for this amaranth species). It is possible that the offsprings we are observing in this 1:1 ratio are homozygous (normal) females and homozygous males.

Alternatively, if gender were determined by an X:autosome balance as found in *Rumex* and *Humulus* (Parker and Clark 1991; Jacobsen 1957, respectively), a 1:1 ratio of male to female would be possible in apomictic offspring. In this system, the X:autosome ratio determines the sexual fate of floral primordia, and thus gender of the plant, rather than the presence or absence of the Y chromosome. In diploid *Rumex*, XX are female and XYY are male. In polyploidy species of the genus *Rumex*, an X:autosome ratio of 1.0 or higher is female; a ratio of 0.5 or lower is male (Parker and Clark 1991). In domestic hops (*Humulus* spp.) a XY system is known. XX is female while XY is male. Polyploid variants (multiple X) of hops show XXXX to be female and XXYY to be male. In the above synergid/antipodal apogamy system proposed, meiosis I could change the ratio by eliminating the X chromosome from (half) the megaspore mother cells being generated, thereby generating males (00:autosomes) and females (XX:autosomes) in a 1:1 ratio.

A third possibility is that the amplification of *EPSPS* copy number is altering gene sequences that determine gender. Since *A. palmeri* does not have a heteromorphic sex chromosome (Grant 1959a; Westergaard 1958), it is possible that glyphosate-induced *EPSPS* gene copy number amplification occurs concurrent with or adjacent to segments of the genome that cause female/maleness. Selection for high *EPSPS* copy number would alter the ratio of sex factor:autosome resulting in the production of male and female apomictic offspring, however, a fixed ratio of 1:1 is unlikely.

However, *A. palmeri/A. rudis* non-hybrid apomictic offsprings were all females (Trucco et al. 2007). *A. palmeri* does not have a heteromorphic sex chromosome (Grant 1959a; Westergaard 1958), and the factors that determine sex are unknown. The dioecious condition may have evolved in *Amaranthus* after the split of 32/34 chromosomes, and sex expression is likely under the control of one or more genes (Grant 1959b). It is possible that the sex determination factors are present in both female and male dioecious plants. Glyphosate could affect sex expression by effects on sex-modifying hormones. Environmental stress on female *A. palmeri* is known to modulate hormonal balance (abscisic acid levels) in seeds (Jha et al. 2010). Therefore, the strong selection pressure imposed by glyphosate combined with its ability to reduce of indole-3-acetic acid (IAA) levels (e.g., Lee 1984) may contribute to this phenomenon.

The level of R alleles in a population under the selection pressure of glyphosate is dependent on the mode of inheritance of the resistance trait. Facultative apomixis

would maintain the presence of individuals with high levels of *EPSPS* gene copy number in *A. palmeri* populations with GR individuals, even in the presence of overwhelming amounts of GS pollen or absence of pollen donors. A deeper understanding of the mechanism of gene amplification (Gaines et al. 2013) and the regulation of apomixis and sex determination factors in *A. palmeri* populations are essential for understanding the stability of multiple *EPSPS* gene copies in populations.

Acknowledgments This research was funded in part by Monsanto Company. We thank Dr. Natasha Techen for her advice with the genetic markers. Susan B. Watson, Marilyn Ruscoe and J'Lynn Howell provided excellent technical support. Dr. R. Douglas Sammons and Dr. Dafu Wang's help with the Western blot analysis was greatly appreciated.

References

- Baerson SR, Rodriguez DJ, Tran M, Feng Y, Best NA, Dill GM (2002) Glyphosate-resistant goosegrass: identification of a mutation in the target enzyme 5-enolpyruvylshikimate-3-phosphate synthase. *Plant Physiol* 129:1265–1275
- Bass C, Field LM (2011) Gene amplification and insecticide resistance. *Pest Manag Sci* 67:886–890
- Bensch CN, Horak MJ, Peterson D (2003) Interference of redroot pigweed (*Amaranthus retroflexus*), Palmer amaranth (*A. palmeri*), and common waterhemp (*A. rudis*) in soybean. *Weed Sci* 51:37–43
- Bonny S (2011) Herbicide-tolerant transgenic soybean over 15 years of cultivation: pesticide use, weed resistance, and some economic issues. The case of the USA. *Sustainability* 3:1302–1322
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. *Anal Biochem* 72:248–254
- Bradshaw LD, Padgett SR, Kimball SL, Wells BH (1997) Perspectives on glyphosate resistance. *Weed Technol* 11:189–198
- Brookes G, Barfoot P (2013) The global income and production effects of genetically modified (GM) crops 1996–2011. *GM Crops Food* 4:1–10
- Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT (2009) The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 55:611–622
- Chan KF, Sun M (1997) Genetic diversity and relationships detected by isozyme and RAPD analysis of crop and wild species of *Amaranthus*. *Theor Appl Genet* 95:865–873
- Chandi A, Milla-Lewis SR, Giacomini D, Westra P, Preston C, Jordan DL, York AC, Burton JD, Whitaker JR (2012) Inheritance of evolved glyphosate resistance in a North Carolina Palmer amaranth (*Amaranthus palmeri*) biotype. *Int J Agron*. doi:10.1155/2012/176108
- Chandi A, Jordan DL, Milla-Lewis SR, Jordan DL, York AC, Burton JD, Zuleta CM, Whitaker JR, Culpepper AS (2013) Use of AFLP markers to assess genetic diversity in Palmer amaranth (*Amaranthus palmeri*) populations from North Carolina and Georgia. *Weed Sci* 61:136–145
- Chandrika M, Rai VR, Thoyajaksha V (2010) ISSR marker based analysis of micropropagated plantlets of *Nothapodytes foetida*. *Biol Plant* 54:561–565
- Chattopadhyay D, Sharma AK (1991) Sex determination in dioecious species of plants. *Feddes Repertorium* 102:29–55

- Culpepper AS, Grey TL, Vencill WK, Kichler JM, Webster TM, Brown SM, York AC, Davis JW, Hanna WW (2006) Glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) confirmed in Georgia. *Weed Sci* 54:620–626
- Dayan FE, Owens DK, Corniani N, Silva FML, Watson SB, Howell J, Shaner DL (2013) Biochemical markers and enzyme assays for herbicide mode of action and resistance studies. *Weed Sci*. doi:10.1614/WS-D-11-00194.1
- Duke SO, Powles SB (2008) Glyphosate: a once in a century herbicide. *Pest Manag Sci* 64:319–325
- Duke SO, Powles SB (2009) Glyphosate-resistant crops and weeds: now and in the future. *AgBioForum* 12:346–357
- Ehleringer J (1983) Ecophysiology of *Amaranthus palmeri*, a sonoran desert summer annual. *Oecologia* 57:107–112
- Franssen AS, Skinner DZ, Al-Khatib K, Horak MJ, Kulakow PA (2001) Interspecific hybridization and gene flow of ALS resistance in *Amaranthus* species. *Weed Sci* 49:598–606
- Gaines TA (2009) Molecular genetics of glyphosate resistance in Palmer amaranth (*Amaranthus palmeri* L.). Ph.D. Dissertation, Colorado State University, Fort Collins
- Gaines TA, Zhang W, Wang D, Bukun B, Chisholm ST, Shaner DL, Nissen SJ, Patzoldt WL, Tranel PJ, Culpepper AS, Grey TL, Webster TM, Vencill WK, Sammons RD, Jiang J, Preston C, Leach JE, Westra P (2010) Gene amplification confers glyphosate resistance in *Amaranthus palmeri*. *Proc Natl Acad Sci USA* 107:1029–1034
- Gaines TA, Shaner DL, Ward SM, Leach JE, Preston C, Westra P (2011) Mechanism of resistance of evolved glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*). *J Agric Food Chem* 59:5886–5889
- Gaines TA, Ward SM, Bukun B, Preston C, Leach JE, Westra P (2012) Interspecific hybridization transfers a previously unknown glyphosate resistance mechanism in *Amaranthus* species. *Evol Appl* 5:29–38
- Gaines TA, Wright AA, Molin WT, Lorentz L, Riggins CW, Tranel PJ, Beffa R, Westra P, Powles SB (2013) Identification of genetic elements associated with EPSPS gene amplification. *PLoS One* 8:e65819
- Gardner JG, Nelson GC (2008) Herbicides, glyphosate-resistance and acute mammalian toxicity: simulating an environmental effect of glyphosate-resistant weeds in the USA. *Pest Manag Sci* 64:470–478
- Ge X, d'Avignon DA, Ackerman JJH, Sammons RD (2010) Rapid vacuolar sequestration: the horseweed glyphosate resistance mechanism. *Pest Manag Sci* 66:345–348
- Ge X, d'Avignon DA, Ackerman JJH, Duncan B, Spaur MB, Sammons RD (2011) Glyphosate-resistant horseweed made sensitive to glyphosate: low-temperature suppression of glyphosate vacuolar sequestration revealed by ^{31}P NMR. *Pest Manag Sci* 67:1215–1221
- Ge X, d'Avignon DA, Ackerman JJH, Collavo A, Sattin M, Ostrander EL, Hall EL, Sammons RD (2012) Vacuolar glyphosate-sequestration correlates with glyphosate resistance in ryegrass (*Lolium* spp.) from Australia, South America, and Europe: A ^{31}P NMR investigation. *J Agric Food Chem* 60:1243–1250
- Grant WF (1959a) Cytogenetic studies in *Amaranthus*. III. Chromosome numbers and phylogenetic aspects. *Can J Genet Cytol* 1:313–328
- Grant WF (1959b) Cytogenetic studies in *Amaranthus*. I. Cytological aspects of sex determination in dioecious species. *Can J Bot* 37:413–417
- Green JM (2009) Evolution of glyphosate-resistant crop technology. *Weed Sci* 57:108–117
- Gustafsson A (1947) Apomixis in higher plants. III. Biotype and species formation. *Lunds Univ Arsskr* 43:71–178
- Heap I (2013) International Survey of Herbicide Resistant Weeds. Online. Internet. www.weedscience.org. Accessed 7 August 2013
- Jacobsen P (1957) The sex chromosomes in *Humulus* L. *Hereditas* 43:357–370
- Jalaludin A, Han H, Powles S (2013) Evolution in action: a double amino acid substitution in the EPSPS gene endows high-level glyphosate resistance. In: Proceedings of the global herbicide resistance challenge conference, Perth, p 35
- James C (2012) 2012 ISAAA report on global status of Biotech/GM Crops. International Service for the Acquisition of Agri-Biotech Applications Brief 44-2012, Ithaca
- Jha P, Norsworthy JK, Riley MB, Bridges W (2010) Shade and plant location effects on germination and hormone content of Palmer amaranth (*Amaranthus palmeri*) seed. *Weed Sci* 58:16–21
- Keeley PE, Carter CH, Thullen RM (1987) Influence of planting date on growth of Palmer amaranth (*Amaranthus palmeri*). *Weed Sci* 35:199–204
- Kleter GA, Kuiper HA (2003) Environmental fate and impact considerations related to the use of transgenic crops. In: Voss G, Ramos G (eds) Chemistry of plant protection. Wiley-VCH, Weinheim, pp 304–321
- Lanoue KZ, Wolf PG, Browning S, Hood EE (1996) Phylogenetic analysis of restriction-site variation in wild and cultivated *Amaranthus* species (*Amaranthaceae*). *Theor Appl Genet* 93:722–732
- Lata H, Chandra S, Techen N, Khan IA, ElSohly MA (2010) Assessment of the genetic stability of micropropagated plants of *Cannabis sativa* by ISSR markers. *Planta Med* 76:97–100
- Lata H, Chandra S, Techen N, Khan IA, ElSohly MA (2011) Molecular analysis of genetic fidelity in *Cannabis sativa* L. plants grown from synthetic (encapsulated) seeds following in vitro storage. *Biotech Lett* 33:2503–2508
- Lebrun M, Sailland A, Freyssinet G (1997) Mutated 5-enol pyruvylshikimate-3-phosphate synthase, gene coding for said protein and transformed plants containing said gene. World Patent Application No. WO9704103
- Lee TT (1984) Release of lateral buds from apical dominance by glyphosate in soybean and pea seedlings. *J Plant Growth Regul* 3:227–235
- Lee RM, Thimmapuram J, Thinglum KA, George G, Hernandez AG, Wright CL, Klim RW, Mikel MA, Tranel PJ (2009) Sampling the waterhemp (*Amaranthus tuberculatus*) genome using pyrosequencing technology. *Weed Sci* 57:463–469
- Ma GG, Park YJ, Lee GA, Lee JR, Lee SY, Kwak JG, Kim TS, Cho EG (2008) SSR primer sets for identification of *Amaranthus* species. Rep Korean Kongkae Taeho Kongbo Patent Oct 18, 2008, KR 2008092187, A20081015
- Massinga RA, Currie RS, Horak MJ, Boyer J (2001) Interference of Palmer amaranth in corn. *Weed Sci* 49:202–208
- Mayo CM, Horak MJ, Peterson DE, Boyer JE (1995) Differential control of four *Amaranthus* species by six postemergence herbicides in soybean (*Glycine max*). *Weed Technol* 9:141–147
- Morgan GD, Baumann PA, Chandler JM (2001) Competitive impact of Palmer amaranth (*Amaranthus palmeri*) on cotton (*Gossypium hirsutum*) development and yield. *Weed Technol* 15:408–412
- Nandula VK, Reddy KN, Koger CH, Poston DH, Rimando AM, Duke SO, Bond JA, Ribeiro DN (2012) Multiple resistance to glyphosate and pyriithiobac in Palmer amaranth (*Amaranthus palmeri*) from Mississippi and response to flumiclorac. *Weed Sci* 60:179–188
- Ng CH, Wickneswary R, Salmijah S, Teng YT, Ismail BS (2003) Gene polymorphisms in glyphosate-resistant and -susceptible biotypes of *Eleusine indica* from Malaysia. *Weed Res* 43:108–115
- Parker JS, Clark MS (1991) Dosage sex-chromosome systems in plants. *Plant Sci* 80:79–92
- Perez-Jones A, Mallory-Smith C (2010) Biochemical mechanisms and molecular basis of evolved glyphosate resistance in weed species. In: Nandula VK (ed) Glyphosate resistance in crops and weeds. Wiley, Hoboken, pp 119–140

- Pline-Srnic W (2006) Physiological mechanisms of glyphosate resistance. *Weed Technol* 20:290–300
- Popa G, Cornea CP, Ciuca M, Babeanu N, Popa O, Marin D (2010) Studies on genetic diversity in *Amaranthus* species using the RAPD markers. *Analele Universității din Oradea-Fascicula Biologie* 17:280–285
- Powles SB, Lorraine-Colwill DF, Delow JJ, Preston C (1998) Evolved resistance to glyphosate in rigid ryegrass (*Lolium rigidum*) in Australia. *Weed Sci* 46:604–607
- Pratley J, Urwin N, Stanton R, Baines P, Broster J, Cullis K, Schafer D, Bohn J, Krueger R (1999) Resistance to glyphosate in *Lolium rigidum*. I. Bioevaluation. *Weed Sci* 47:405–411
- Salas RA, Dayan FE, Pan Z, Watson SB, Dickson JW, Scott RC, Burgos NR (2012) EPSPS gene amplification in glyphosate-resistant Italian ryegrass (*Lolium perenne* ssp. *multiflorum*) from Arkansas. *Pest Manag Sci* 68:1223–1230
- Sammons RD, Meyer J, Hall E, Ostrander E, Schrader S (2007) A simple continuous assay for EPSP synthase from plant tissue. <http://www.cottoninc.com/2007-Glyphosate-Resistant-Palmer-Amaranth/11a-Industry-Sammons-NCWSS07-poster.pdf>. Accessed 20 January 2012
- Seefeldt SS, Jensen JE, Fuerst EP (1995) Log-logistic analysis of herbicide dose-response relationships. *Weed Technol* 9:218–227
- Selmecki A, Gerami-Nejad M, Paulson C, Forche A, Berman J (2008) An isochromosome confers drug resistance in vivo by amplification of two genes, ERG11 and TAC1. *Mol Microbiol* 68:624–641
- Shaner DL, Nadler-Hassar T, Henry WB, Koger CH (2005) A rapid in vivo shikimate accumulation assay with excised leaf discs. *Weed Sci* 53:769–774
- Siehl DL, Subramaniam MV, Walters EW, Blanding JH, Niderman T, Weinmann C (1997) Evaluating anthranilate synthase as a herbicide target. *Weed Sci* 45:628–633
- Sosnoskie LM, Webster MTM, Kichler JM, MacRae AW, Grey TL, Culpepper AS (2012) Pollen-mediated dispersal of glyphosate-resistance in Palmer amaranth under field conditions. *Weed Sci* 60:366–373
- Steinau AN, Skinner DZ, Steinau M (2003) Mechanism of extreme genetic recombination in weedy *Amaranthus* hybrids. *Weed Sci* 51:696–701
- Steinrücken H, Amrhein N (1980) The herbicide glyphosate is a potent inhibitor of 5-enolpyruvylshikimic acid-3-phosphate synthase. *Biochem Biophys Res Com* 94:1207–1212
- Trucco F, Jeschke MR, Rayburn AL, Tranel PJ (2005) Promiscuity in weedy amaranths: high frequency of female tall waterhemp (*Amaranthus tuberculatus*) × smooth pigweed (*A. hybridus*) hybridization under field conditions. *Weed Sci* 53:46–54
- Trucco F, Zheng D, Woodyard AJ, Walter JR, Tatum TC, Rayburn AL, Tranel PJ (2007) Nonhybrid progeny from crosses of dioecious Amaranths: implications for gene-flow research. *Weed Sci* 55:119–122
- Wassom JJ, Tranel PJ (2005) Amplified fragment length polymorphism-based genetic relationships among weedy *Amaranthus* species. *J Hered* 96:410–416
- Watson S (1877) Descriptions of new species of plants, with revisions of certain genera. *Proc Am Acad Arts Sci* 12:246–278
- Webb MR (1992) A continuous spectrophotometric assay for inorganic phosphate and for measuring phosphate release kinetics in biological systems. *Proc Natl Acad Sci USA* 89:4884–4887
- Westergaard M (1958) The mechanism of sex determination in dioecious plants. *Adv Genet* 9:217–281
- Wetzel DK, Horak MJ, Skinner DZ (1999) Use of PCR-based molecular markers to identify weedy *Amaranthus* species. *Weed Sci* 47:518–523
- Wiersma A (2012) Regional whole plant and molecular response of *Kochia scoparia* to glyphosate. M.S. thesis, Colorado State University, p 55
- Xu ZL, Ali Z, Yi JX, He XL, Zhang DY, Yu GJ, Khan AA, Khan IA, Ma HX (2011) Expressed sequence tag-simple sequence repeat-based molecular variance in two *Salicornia* (*Amaranthaceae*) populations. *Genet Mol Res* 10:1262–1276