

Cucurbit Genetics Cooperative

Report No. 16

July 1993



Table of Contents (article titles linked to html files)

- **Introduction**

- Comments from the CGC Coordinating Committee
- Comments from the CGC Gene List Committee
- Comments from the CGC Gene Curators
- 16th Annual CGC Business Meeting
- 1992 Cucurbit Conference
- 1993 Watermelon Research Group Meeting
- US Cucurbit Crop Advisory Committee 1992-93 Update
- Upcoming Meetings of Interest to Cucurbit Researchers
- Cucurbitaceae '94

- **Cucumber (*Cucumis sativus*)**

1. **Correlations between years for foliar gummy stem blight disease ratings on field grown cucumbers**
Paul C. St. Amand and Todd C. Wehner (USA)
CGC 16:1-2 (1993)
2. **Turgid flowers are essential for good fruit and seed set in cucumber**
R. Szegedi, I. Csernia and P. Milotay (Hungary)
CGC 16:3-4 (1993)
3. **Vine rolling vs. conventional multiple harvest of cucumbers in North Carolina**
Todd C. Wehner and Conrad H. Miller (USA)
CGC 16:5-7 (1993)
4. **Observations in fruit netting in cucumber**
Michael S. Uchneat and Todd C. Wehner (USA)
CGC 16:8-9 (1993)
5. **Leaf structure and photosynthetic relations in *Cucumis sativus* var. *sativus* and *Cucumis sativus* var. *hardwickii***
J.E. Staub and B.E. Struckmeyer (USA)
CGC 16:10-13 (1993)
6. **Cucumber (*Cucumis sativus* L.) induced mutations: A *Phaseolus* leaf mutant**
M. Rucinska, E. Bergier, K. Niemirowicz-Szczytt and A. Korzeniewska (Poland)
CGC 16:14-17 (1993)
7. **Independence of fruit length and 10 other characters in cucumber**
Nick E. Fanourakis (Greece)
CGC 16:18-21 (1993)
8. **Diallel analysis of cucumber germination at optimum and suboptimal temperatures**
P. Milotay (Hungary)
CGC 16:22-26 (1993)
9. **Application of factor analysis to cucumber breeding**
Meng Zhang and Hongwen Cui (P.R. China)
CGC 16:27-29 (1993)
10. **Application of narrow sense canonical characters to cucumber breeding**
Hongwen Cui and Yongtao Qi (P.R. China)
CGC 16:30-34 (1993)

- **Melon (*Cucumis melo*)**

11. ***Cucumis* germplasm: 1992 collection expedition in India**
J.D. McCreight, J.E. Staub, N.M. Koppa and U.C. Srivastaba (USA and India)

- CGC 16:35-36 (1993)
12. **Resistance to melon dieback in Spanish landraces of melon**
F. Nuez and J. Esteva (Spain)
CGC 16:37-38 (1993)
 13. **Club mildew: working group on resistance of melon to powder mildew**
James D. McCreight and Michel Pitrat (USA and France)
CGC 16:39 (1993)
 14. **Downy mildew (*Pseudoperonospora cubensis* B & C) resistance in melon (*Cucumis melo* L.)**
R.G. Somkuwar and T.A. More (India)
CGC 16:40-41 (1993)
 15. **Gene for resistance to fusarium wilt race 1 in Oriental pickling melon**
Harry S. Paris, Ron Cohen, Yael Danin-Poleg and Shoshanna Schreiber (Israel)
CGC 16:42-43 (1993)
 16. **Breeding and development of cucumber green mottle mosaic virus (CGMMV) resistant lines in melon (*Cucumis melo* L.)**
T.A. More, A. Varma, V.S. Seshadri, R.G. Somkuwar and L. Rajamony (India)
CGC 16:44-46 (1993)
 17. **The role of viruses in sudden wilt of melons in New York**
Henry M. Munger (USA)
CGC 16:47-48 (1993)
 18. **Screening of melons for sweetpotato whitefly resistance: 1992**
James D. McCreight (USA)
CGC 16:49-52 (1993)
 19. **Genotypic control of regeneration potential in *Cucumis melo***
J. Jain and T.A. More (India)
CGC 16:53-54 (1993)
- **Squash and Pumpkin (*Cucurbita* spp.)**
 20. **Genetic parthenocarpy in *Cucurbita pepo* L.**
R.W. Robinson (USA)
CGC 16:55-57 (1993)
 21. **Productivity of naked seed squash, *Cucurbita pepo* L.**
Warid A. Warid, Jaime J. Martinez and Juan M. Loaiza (Mexico)
CGC 16:58-60 (1993)
 22. **The chromoplast ultrastructure of two isogenic lines of *Cucurbita pepo* fruits at different developmental stages**
H.T. Lim and C.D. Boyer (Rep. Korea and USA)
CGC 16:61-63 (1993)
 23. **A case of extremely low expressivity of gene *B2* in *C. moschata***
Oved Shifriss (USA)
CGC 16:64-67 (1993)
 24. **Inheritance of immature fruit color in *C. moschata***
Antonio I.I. Cardoso, Paulo T. Della Vecchia and Norberto Silva (Brazil)
CGC 16:68-69 (1993)
 25. **Breeding bush types of *C. moschata* with field resistance to PRSV-W**
Paulo T. Della Vecchia, Pedro Terenciano S. and Amadeu Terenciano (Brazil)
CGC 16:70-72 (1993)
 26. **Powdery and down mildew resistance in *Cucurbita moschata* accessions**
Linda Wessel-Beaver (USA)
CGC 16:73-74 (1993)
 27. **Leaf silvering of squash: a brief review**
Harry S. Paris (Israel)
CGC 16:75-76 (1993)
 - **Watermelon (*Citrullus lanatus*)**
 28. **Inheritance of resistance to races 0, 1 and 2 of *Fusarium oxysporum* f.sp. *niveum* in watermelon (*Citrullus* sp. PI 296341)**
Xingping Zhang and Bill Rhodes (USA)
CGC 16:77-78 (1993)

29. **Male sterile, *ms*, in watermelon not linked to delayed green, *dg* and *I-dg***
Xingping Zhang and Bill Rhodes (USA)
CGC 16:79 (1993)
30. **Fertility shift in an *in vitro* regenerated male-sterile line in watermelon**
Xingping Zhang and Bill Rhodes (USA)
CGC 16:80 (1993)
31. **Protoplast isolation and culture of watermelon cotyledons**
Caiping Ma, X.P. Zhang, B.B. Rhodes and J.W. Adelberg (USA)
CGC 16:81-83 (1993)
- **Other Genera and Species**
 32. **Crossability between *Momordica charantia* and *Momordica dioica***
M. Abdul Vahab and K.V. Peter (India)
CGC 16:84-86 (1993)
 33. **Preliminary screening of cucurbits species for *Bemisia tabaci* Genn. whitefly resistance**
V. Moreno, J.L. Gomez Aguilera, C. Guerau de Arellano and L.A. Roig (Spain)
CGC 16:87-89 (1993)
- **Gene Lists and Germplasm**
 1. **Cucumber, melon and squash germplasm from the Cornell collection**
Molly Kyle, George Moriarty and Henry M. Munger (USA)
CGC 16:90-91 (1993)
 2. **Gene list update for cucumber**
Todd C. Wehner (USA)
CGC 16:92-97 (1993)
 3. **Gene nomenclature for the Cucurbitaceae**
CGC Gene List Committee
CGC 16:98 (1993)
- **Appendices**
 - 1993 CGC membership directory
 - Geographical distribution of CGC members in the USA
 - Geographical distribution of CGC members outside the USA
 - Covenant and by-laws of the Cucurbit Genetics Cooperative
 - CGC Financial Statement (31 December 1992)

Cucurbit Genetics Cooperative Report 16: vi-ix (Introduction) 1993

Introduction

Comments from the CGC Coordinating Committee

The Call for Papers for the 1994 Report (CGC Report No. 17) will be mailed in August 1993. Papers should be submitted to the respective Coordinating Committee members by 31 December 1993. The report will be published by June 1994. As always, we are eager to hear from CGC members regarding our current activities and the future direction of CGC.

- Gary W. Elmstrom (melon)
- Dennis T. Ray (watermelon)
- Mark G. Hutton (other genera)
- Jack E. Staub (cucumber)
- J. Brent Loy (*Cucurbita* spp.)
- Timothy J. Ng, Chairman

Comments from CGC Gene List Committee

Lists of known genes for the Cucurbitaceae have been published previously in Hortscience and in reports of the Cucurbit Genetics Cooperative. CGC is currently publishing complete lists of known genes for cucumber (*Cucumis sativus*), melon (*Cucumis melo*), watermelon (*Citrullus lanatus*), and *Cucurbita* spp. on a rotating basis.

It is hoped that scientists will consult these lists as well as the rules of gene nomenclature for the Cucurbitaceae before choosing a gene name and symbol. Thus, inadvertent duplication of gene names and symbols will be prevented. The rules of gene nomenclature were adopted in order to provide guidelines for the naming and symbolizing of genes previously reported and those which will be reported in the future. Scientists are urged to contact members of the Gene List Committee regarding questions in interpreting the nomenclature rules and in naming and symbolizing new genes.

- Cucumber: Todd C. Wehner
- Melon: Michael Pitrat
- Watermelon: Warren R. Henderson
- *Cucurbita* spp.: Mark G. Hutton and Richard W. Robinson
- Other Genera: Richard W. Robinson

Comments from the CGC Gene Curators

CGC has appointed Curators for the four major cultivated groups: cucumber, melon, watermelon and *Cucurbita* spp. A back-up Curator for the "Other genera" category is needed; any one wishing to take on this responsibility should contact the Chair.

Curators are responsible for collecting, maintaining and distributing upon request stocks of the known marker genes. CGC members are requested to forward samples of currently held gene stocks to the respective Curator.

- Cucumber: Todd C. Wehner and Jack E. Staub
- Melon: J.D. McCreight and Michel Pitrat
- Other genera: Richard W. Robinson
- Watermelon: Gary W. Elmstrom, E. Glen Price and Billy B. Rhodes
- *Cucurbita* spp.: Mark G. Hutton and Richard W. Robinson

16th Annual CGC Business Meeting

The 16th Annual Business Meeting of the Cucurbit Genetics Cooperative was held on 21 December 1992 in conjunction with the 1992 Cucurbit Conference in Raleigh, North Carolina. Todd Wehner chaired the meeting in Tim Ng's absence, and 21 members and guests (representing six different countries) were in attendance.

An update of CGC Report and CGC membership statistics was presented. Jack Staub was nominated and elected as CGC Coordinating Committee member for cucumber (*Cucumis sativus*). Jack replaces Todd Wehner, who had completed his ten year term as a member of the Coordinating Committee.

Two amendments to the CC By-laws were proposed by the CGC Chair: (1) to change the By-laws to allow for more than five members on the CGC Gene List Committee, and (2) to limit guaranteed availability of back issues of the CGC Report to only the most recent five issues. The first proposed amendment reflects both the increasing complexity of maintaining gene lists and the willingness of CGC members to volunteer their services in this activity. The second proposed amendment addresses the physical difficulty of storing and maintaining increasing stocks of back issues, the costs of mechanical reproduction of exhausted back issues, and the plans for storing older reports as digitized images.

There was little discussion of the first issues. However, with the second proposed amendment there was concern expressed that CGC should keep all back issues available until a concerted effort was made to get libraries to pick up the entire set for future reference. As CGC By-laws require that any amendments will be discussed further at the 1993 CGC Business Meeting and mail ballots will be included in the August "Call for Papers for CGC Report No. 17" mailing to the membership.

1992 Cucurbit Conference

The 1992 Cucurbit Conference was held 20-23 September 1992 in Raleigh, North Carolina. Participating organizations included the National Cucumber Conference (NCC), the Pickling Cucumber Improvement Committee (PCIC), the Cucurbit Genetics Cooperative (CGC), the Watermelon Research Group (WRG), the Cucurbit Crop Advisory Committee (CCAC), the National Melon Research Group (NMRG), and the Squash and Pumpkin research Group (SPRG). Tom Monaco coordinated the Cucumber Field Day.

There was an informal decision to have a US Cucurbit Conference in even years alternating with EUCARPIA, beginning in 1994. Many of the commodity groups present at the conference agreed to combine into a single cucurbit organization for coordination of the meetings.

1993 Watermelon Research Group Meeting

Gary Elmstrom, University of Florida, Leesburg, FL USA

The 12th annual meeting of the Watermelon Research Group (WRG) was held in Tulsa, Oklahoma, on 31 January 1993. Attendance was up due to a number of plant pathologists who were attending the watermelon fruit blotch workshop. Glen Price (American Sunmelon) made local arrangements and provided watermelon for all. Gary Elmstrom announced that he would be stepping down as WRG Chair after 12 years of service in this capacity, and Ray Martyn (Texas A&M) will be assuming these duties.

US Cucurbit Crop Advisory Committee 1992-1993 Update

J.D. McCreight, USDA-ARS, Salinas, CA USA

On 23 September 1992, the US Cucurbit Crop Advisory Committee (CCAC) met in Raleigh, North Carolina in conjunction with the Cucurbit Conference organized by the Department of Horticultural Science, North Carolina State University.

The Core Concept and its applicability to cucurbits was further discussed and it was decided to construct core collections for each species based on geographic origin and proven value for one or more specific characters.

The National Seed Storage Laboratory (NSSL) is now accepting samples for long-term storage. NSSL has specific requirements for numbers of seeds and germination percentage that must be met. There were discussions on various aspects of deleting duplicates of accessions; ensuring entry of valuable breeding lines and varieties from scientists either

prior to or upon retirement when their program will not be continued; obtaining correct species identification for USDA taxonomists in a timely manner, conditioning acceptance of HortScience Cultivar and Germplasm Release manuscripts upon submission of samples to the National Plant Germplasm System (NPGS); germplasm exchange with Russia, The Netherlands and India; availability of accessions in NPGS; and increase of accessions outside of the US on a contract basis. The CAC Report is approximately four years old and needs to be updated for various reasons including the transfer of the melon collection from Georgia to Iowa, retirements, and progress in evaluating germplasm collections.

Two germplasm evaluation proposals were recommended for funding in FY 1993; Evaluation of the U.S. plant introduction collection of melon (*Cucumis melo*) and squash (*Cucurbita moschata* and *C. pepo*) for resistance to gummy stem blight (*Didymella bryoniae* Auersw.) Rehm (M.A. Kyle and T.A. Zitter), and Genetic diversity in cucumber (*Cucumis sativus* L.) and melon (*Cucumis melo* L.) accessions as measured by morphological and biochemical genetic markers (J.E. Staub and J.D. McCreight).

The 1993 meeting is scheduled for July 24 in Nashville, Tennessee, in conjunction with the Annual Meeting of the American Society for Horticultural Science.

Upcoming Meetings of Interest to Cucurbit Researchers

Groups meeting in conjunction with the **1993 Annual Meeting of the American Society for Horticultural Science** in Nashville, Tennessee (24-29 July 1993):

- **Cucurbit Crop Advisory Committee**
 - Saturday, 24 July, 8:00 a.m. - 12:00 p.m.
 - James D. McCreight
 - USDA-ARS, Salinas, CA USA
 - Tel: 408/755-2864; Fax: 408/755-2866
- **National Melon Research Group**
 - Saturday, 24 July, 1:00 p.m. - 3:00 p.m.
 - James D. McCreight
 - USDA-ARS, Salinas, CA USA
 - Tel: 408/755-2864; Fax: 408/755-2866
- **Squash Breeders**
 - Saturday, 24 July, 3:30 p.m. - 5:00 p.m.
 - Laura Merrick
 - University of Maine, Orono, ME USA
 - Tel: 207/581-2950; Fax: 207/581-2999
- **Cucurbit Genetics Cooperative**
 - (dates to be announced)
 - Timothy J. Ng
 - University of Maryland, College Park, MD USA
 - Tel: 301/405-4345; Fax: 301/314-9308

Other Meetings

The 13th annual meeting of the Watermelon Research Group will be held in conjunction with the 1994 annual meeting of the Southern Association of Agricultural Scientists (SAAS) and the Southern Region of the American Society for Horticultural Science (SR-ASHS) during the week of 5-9 February 1994 in Nashville, Tennessee.

Cucurbitaceae '94

Cucurbitaceae '94 will be held in the Lower Rio Grande Valley of Texas in either October or early November and will be hosted by the Texas Agricultural Experiment Station, Weslaco, and the USDA-ARS Subtropical Agricultural Research Laboratory, Weslaco. Jim Dunlap (TAES) and Gene Lester (USDA, ARS) will be the hosts for this meeting. The first announcement and call for papers is forthcoming.

Correlations Between Years for Foliar Gummy Stem Blight Disease Ratings on Field Grown Cucumbers

Paul C. St. Amand and Todd C. Wehner

Department of Horticultural Science, North Carolina State University, Raleigh, NC 27695-7609

It is often desirable to know if resistance data from one test is positively and highly correlated with other tests. A high positive correlation indicates that testing once in any year is a good measure of resistance and a low correlation indicates that multi-year tests are necessary to determine the resistance of a cultigen (breeding lines, cultivars, and plant introduction accessions). The objective of this study was to calculate correlations among years for foliar ratings of gummy stem blight in field-grown cucumbers.

Field tests were conducted in 1981, 1982, 1983, and 1986 to determine the resistance of diverse cucumbers to gummy stem blight. Each test was conducted at the Horticultural Crops Research Station at Clinton, N.C. with 3 (1982) or 6 (1983, 1986) replications, except for the 1981 test, which was conducted at the Horticultural Crops Research Station at Castle Hayne, N.C. without replication. Field plots were rated for foliar lesions using a 0 to 9 scale (0 = no foliar symptoms, 9 = plant dead). The rating system was modeled after the categories developed by Thompson and Jenkins (2). Plots were inoculated with equal numbers of spores from 12 isolates of *Didymella bryoniae*. Plants were sprayed at the vine-tip-over stage (4 to 6 true leaves) to run-off using a Solo back-pack sprayer at 15 to 20 psi. Overhead irrigation was used (25 to 38 mm/week) to spread the inoculum and encourage uniform disease development. Every 3rd row (4th row for 1986) was planted with susceptible 'Wisconsin SMR 18' to enhance the uniformity of disease spread. Plots were 6 m long (1981) with 40 plants each, or 3 m long (1982, 1983, 1986) with 30 plants each and were planted on raised, shaped beds 1.5 m apart (center to center) separated at each end by 1.5 m alleys. Standard cultural practices were used for crop production (1). A randomized complete block design was used for all tests. One rating was given for each plot 7, 14 and 21 days after inoculation, except in 1981 which was rated only 7 days after inoculation. Data were analyzed using PROC REG of SAS (SAS Institute, Cary, N.C.).

Disease ratings were greatest in 1982 and the mean rating was very similar for years in 1981, 1983, and 1986 (Table 1). The range of disease ratings was greatest in 1981 and was smallest in 1983 and 1986, Pearson product-moment correlations and Spearman rank correlations between years were moderate to high (Table 2); however, the highest correlations are based on small numbers of cultigens and the correlations tended to decrease as cultigen number increased. Rankings of cultigens were usually less correlated than were actual ratings, indicating that changes in rank often occurred. Wyszogrodzka et al. (3) calculated the Pearson correlation coefficient between gummy stem blight field tests conducted in Florida and Wisconsin. That correlation ($r=0.42$) was moderate and similar to those reported here for tests involving larger numbers of cultigens.

Table 1 The mean, standard deviation, median, minimum and maximum of foliar disease ratings of cucumber for gummy stem blight for four field tests conducted in different years. Also shown is the number of cultigens rated in each test.^Z

Year	Number of cultigens	Mean	Standard deviation	Median	Minimum	Maximum
1981	31	5.6	3.1	8.0	1.5	9.0
1982	46	6.3	2.0	6.6	1.7	9.0
1983	45	5.1	0.8	5.1	2.9	8.2
1986	36	5.5	0.9	5.5	3.6	7.2

^Z Foliar symptoms were rated from no symptoms (0) to complete necrosis (9).

Table 2. Correlations between years for foliar disease ratings of cucumber cultigens inoculated with gummy stem blight.

Pearson product-moment correlation coefficients are above the diagonal and Spearman rank correlation coefficients are below the diagonal.^z

	Year			
Year	1981	1982	1983	1986
1981	-	0.57 (31) ***	0.68 (10) *	0.93 (6) **
1982	0.56 (31) ***	-	0.73 (14) ***	0.73 (8) *
1983	0.82 (10) **	0.84 (14) ***	-	0.66 (32) ***
1986	0.70 (6) NS	0.64 (8) NS	0.48 (32) **	-

^z The symbols NS, *, **, and *** indicate not significant, or significant at the 0.05, 0.01, or 0.001 levels, respectively. Numbers in parenthesis indicate the number of cultigens shared between tests.

Literature Cited

1. Hughes, G.R., C.W. Averre, and K.A. Sorensen. 1983. Growing pickling cucumbers in North Carolina. N.C. Agric. Ext. Serv. AG-315.
2. Thompson, D.C. and S.F. Jenkins. 1985. Pictorial assessment key to determine fungicide concentrations that control anthracnose development on cucumber cultivars with varying resistance levels. Plant Dis. 69: 833-836.
3. Wyszogrodzka, A.J., P.H. Williams, and C.E. Peterson. 1986. Search for resistance to gummy stem blight (*Didymella bryoniae*) in cucumber (*Cucumis sativus* L.) Euphytica 35: 603-613.

Turgid flowers are essential for good fruit and seed set in cucumber

R. Szegedi, I. Cserni and P. Milotay

Vegetable Crops Research Institute, 6000 Kecskemet, P.O.B. 116, Hungary

Fruit number per unit area and shape is generally a good predictor of seed cucumber yield. In short-fruited varieties most of the available ovules are fertilized under optimum conditions resulting in high seed set throughout the ovary (3). However, late setting fruits, or fruits developed under dry field conditions often remain small and deformed containing a reduced number of seeds when compared to fruits grown under optimal conditions.

Pollination is successful up to 24 hours after anthesis under glasshouse conditions (5), and a minimum two-week irrigation period is sufficient to produce viable seeds (4). However, since drip irrigation significantly affects fruit size, shape and seed yield in a large containers study (1), an experiment was designed to evaluate the influence of withholding water during pollination on fruit and seed set.

Plants of the monoecious pickling line K 4599 were trellised during the summer of 1990 under an isolating net. The planting was drip irrigated every 6 hours with fixed doses. Before pollination and after pollination irrigation was withheld in one of two rows (water stress treatment) resulting in flagging during the noon hours, while plants in the second row remained turgid (control). 30-30 flowers were hand-pollinated using two turgid male flowers on plants without previous fruit set on both rows between 12:00 and 13:00 hours. Daily temperature during pollination and the days of early fruit development ranged from 18 to 31°C. After harvest the seed number, seed weight, germination %, radicle and hypocotyl elongation (4 days at 25°C) were recorded to compare treatment and control.

Pollinated flowers set fruit on 93% of the control plants, while only 57% of the pollinations were successful on tagging plants. Fruits of the control plants were regular in shape with an average length of 17.6 cm, while 12 of the 17 fruits developed from non-turgid flowers were pear shaped with an average length of 16.3 cm. Seed number from fruit harvested from control and treatment (water withheld) plants ranged from 124 to 289 and 45 to 168 respectively. Averages calculated from 10-10 randomly chosen fruits show that fruits from water withheld plants had a significantly reduced seed number and seed weight per fruit, but withholding water did not affect thousand seed weight and germination % (Table 1). Radicle and hypocotyl elongations did not show treatment differences (data not shown).

Because pollen germinates on the stigma within 30 minutes, pollen treatment tubes reach the ovary in 12 hours and fertilize ovules enlarge within 30-36 hours (2). Our results indicate that water status of the pistillate flower during this period appears to be a significant factor in determining the quantity and quality of fruit and seed set.

Table 1. Seed yield of cucumber fruits developed from turgid and non-turgid flowers at pollination.

Pollination of	Number of fruit set	Seed number per fruit	Seed weight per fruit (g)	Thousand seed weight (g)	Germination at 25°C (%)
Turgid flowers	28	213.2	4.31	21.3	99.0
Non-turgid flowers	17	90.3	1.94	21.6	100.1
LSD 5%	-	30.8	0.48	NS	NS

NS = not significant.

Literature Cited

- Cserni, I., P. Milotay, A.S. Hodossy and A-ne Trefas. 1990. The effect of fertilization and water supply on cucumber

- seed yield and quality on different soil types. Internat. Seed Testing and Growing Conf. 25-28th June 1990. Godollo, Hungary
2. Fuller, G.L. and A.C. Leopold. 1975. Pollination and the timing of fruit set in cucumbers. HortScience 10:617-619.
 3. Nijs, A.P.M. den and P. Milotay. 1991. Fruit and seed set in the cucumber (*Cucumis sativus* L.) in relation to pollen tube growth, sex type and parthenocarpy. Gartenbauwissenschaft 56.2:46-69.
 4. Wehner, T.C. and R.R. Horton Jr. 1986, Effect of pot size on growth and flowering of cucumbers in the greenhouse. Cucurbit Genet. Coop. Rpt. 9:47-49.
 5. Wehner, T.C. and R.R. Horton Jr. 1989. Delayed pollination successful for cucumbers in North Carolina greenhouse. Cucurbit Genet Coop Rpt. 12:15-16.

Vine Rolling vs. Conventional Multiple Harvest of Cucumbers in North Carolina

Todd C. Wehner and Conrad H. Miller

Department of Horticultural Science, North Carolina State University, Raleigh, NC 27695-7609

Most cucumbers (*Cucumis sativus* L.) in North Carolina are harvested 2 to 3 times per week for 2 to 4 weeks in two production seasons (spring and summer). Fruits are harvested by searching through the vines for marketable sizes. Since it is difficult to see fruits in the canopy, some fruits are not harvested each time. Those becoming oversized before they are picked in the next harvests are unmarketable, resulting in lost yield.

Ease of harvest has been improved using machines, new cultivar types, and more efficient techniques. Harvest machines reduce the labor input where they can be adapted to the production system (*Haffar and Van Ed, 1984). New cultivar types such as gynoeocious flowering permit fewer harvests for the same yield as monoecious types (Wehner and Miller, 1985). Techniques for improved harvest efficiency such as vine training can reduce vine damage, and provide a longer production period for the crop. We were interested in vine rolling for improved recovery of fruits hidden near the base of the plant.

The objective of this study was to determine if harvest was faster and easier if the vines were rolled over as they were picked.

Methods. The experiment was run at the Horticultural Crops Research Station near Clinton, NC. Seeds were planted in rows 1.5 m apart on raised beds. Plots were 9 m long with 1.5 m alleys at each end. Fertilizer was incorporated before planting at a rate of 90-39-84 kg-ha⁻¹ (N-P-K) and a side dressing of N (34 kg/ha) was applied at vine tip-over stage. Irrigation was applied as needed up to three times a week for a minimum of 25 mm of water per week. Recommended cultural practices were used (Hughes et al., 1983).

Plots were planted 22 July, 1982, and thinned 2 August to a density of 86,450 plants*ha⁻¹. Harvests were made Mondays and Thursdays 26 August to 7 September for pickling and 30 August to 10 September for slicing cucumbers. Cultivars tested were 'Calypso' pickle and 'Slicemaster' slicer, both gynoeocious hybrids. Pollen was supplied with border rows of 'Sumter' pickle or 'Poinsett 76' slicer, both monoecious inbreds.

The experiment design was a randomized complete block with 5 replications and 2 harvest methods (conventional vs vine rolling). The pickling and slicing crops were kept separate in harvest and grading operations. Plots were harvested 4 times (twice weekly), and the fruits graded and weighed. Grades were the NC grades for pickles and USDA grades for slicers. NC grades were based on fruit diameter as follows: No. 1=0-26 mm, No. 2=27-38 mm, No. 3=39-50 mm, No. 4=50 mm. USDA grades were based on quality as specified by federal guidelines (USDA 1958). Harvests were timed for each of the methods, conventional and vine rolling. Vines were rolled over on one side as a worker harvested that side, then rolled back the other way by the second worker. Vines were left in their rolled-up state after harvest, and were not laid back out over the soil. That resulted in the exposure of the base of the plant to sun and wind, and some stems and leaves were upside down.

Data were collected on fruit weight by grade, and summarized using analysis of variance. Treatment comparisons were essentially t-tests for conventional vs vine rolling harvest methods.

Results. Differences in yield between conventional and vine rolling were significant for fruit weight in pickling cucumbers but not for fruit value of pickling (Table 1) or for yield of slicing cucumbers (Table 2). Where the differences were significant, the vine rolling treatment resulted in lower yields (68% as much for pickles), and took longer (18% more for pickles, 2% more for slicers) than conventional plots. Vine rolling did, however, result in a significantly higher percentage of No. 1, and lower percentage No. 3 and 4 grade pickles than the conventional method (Table 1). In slicers, the vine rolling treatment produced fewer Fancy and No. 1 grade fruits than conventional, indicating more stress or less effective pollination (Table 2).

Vine rolling required more time to perform than conventional harvest, and resulted in lower yield. In the case of pickling cucumbers, vine rolling did shift the distribution of fruit sizes toward grade 1 and 2, and away from 3 and 4. That may be advantageous for producing small pickling cucumber sizes, but more frequent harvest would probably have the same effect without reducing the yield. Although there was less damage to the vines in the vine rolling treatment compared with conventional (data not shown), the effect was not translated into increased yield in the harvest crew, and improved the recovery of small grade pickles in the same way that vine rolling did. However, vine training would avoid the increased exposure of the base of the vine without leaving some vines turned upside down.

Table 1. Comparison of two multiple-harvest methods for yield and harvest time in pickling cucumbers.^z

Harvest method	Fruit yield		Harvest time (sec/plot/harvest)	Mean size grade ^y	% size grades			
	Value (\$/ha)	Weight (Mg/ha)			1	2	3	4
Conventional	2477	23.9	128	2.7	9	34	38	19
Rolled ^x	2186	16.1	151	2.4	17	39	33	10
LSD (5%)	NS	2.2	-	0.2	4	6	1	6
CV (%)	15	6	-	3	15	9	2	24

^z Data are means over 5 replications and 4 harvests. NS indicates F ratio not significant (5%).

^y Grades are 1 to 4 for the NC sizes.

^x Vines were flipped over the row while harvesting each side.

Table 2. Comparison of two multiple-harvest methods for yield and harvest time in slicing cucumbers.^z

Harvest method	Yield (Mg/ha)		Harvest time (sec/plot/harvest)	Mean size grade ^y	Fancy+ No. 1 (%)	Culls (%)
	Fancy & No. 1	Marketable				
Conventional	20.4	27.7	120	1.1	66	10
Rolled ^x	11.3	18.7	122	1.5	50	19
LSD (5%)	NS	NS	-	0.3	12	NS
CV (%)	50	41	-	14	12	50

^z Data are means over 5 replications and 4 harvests. NS indicates F ratio not significant (5%).

^y Grades are 0 to 3 for the 4 USDA sizes, were 0=Fancy, 1=No. 1, 2=No.2, 3=cull.

^x Vines were flipped over the row while harvesting each side.

Literature Cited

1. Haffar, I. and G. Van Ee. 1984. A comparison of hand and machine harvesting systems for pickling cucumbers in Michigan. Amer. Soc. Agric. Eng. 84-1575.
2. Hughes, G.R., C.W. Averre, and K.A. Sorensen. 1983. Growing pickling cucumbers in North Carolina. N.C. Agric. Ext. Serv. AG-315.
3. U.S.D.A. 1958. United States standards for cucumbers (effective 1 March, 1958). Agric. Marketing Service, U.S. Dept. Agric., Wash. D.C.
4. Wehner, T.C. and C.H. Miller. 1985. Effect of gynoecious expression on yield and earliness of a fresh-market cucumber hybrid. J. Amer. Soc. Hort. Sci. 110:464-466.

Observations on Fruit Netting in Cucumber

Michael S. Uchneat and Todd C. Wehner

Department of Horticultural Science, North Carolina State University, Raleigh, NC 27695-7609

Several fruit surface (skin) types have been identified in cucumber (*Cucumis sativus* L.), including warty (American), smooth (middle-eastern), ridged (greenhouse), and netted (wild). Hutchins (1940) identified the netted trait as a single gene (*H*) linked with black spines and red mature fruit color, and also mentioned the occurrence of intermediate netting. That trait was assigned to linkage group III by Pierce and Wehner (1990). More recently, Peterson and Pike (1992) suggested the netted skin characteristic was quantitatively inherited.

Cucumber accessions PI 197086 and PI 165509 were reported to be resistant to belly rot caused by *Rhizoctonia solani* Kuhn (AG-4) by Clark and Block (1984) and also have netted skin. Since that skin type is not horticulturally acceptable, belly rot resistance must be controlled by separate genes for those cultigens to be useful parents in a program aimed at breeding belly rot resistance. In muskmelon, it has been suggested that fruit netting provides enhanced disease resistance by acting as a physical barrier (Webster and Craig, 1976). In cucumber, it is not known whether netted skin provides a physical barrier, or whether some other mechanism is responsible for resistance.

During the summer of 1992, a study was conducted to determine whether netted skin was responsible for belly rot resistance. Both PI 197086 and PI 165509 have highly netted skin on the mature fruits. Young fruits (less than five days after pollination) do not have netted skin. Netting develops over the next few days, often starting at the ends of the fruit. At about that time, it appears that the epidermal layer stops expanding with the rest of the fruit. That phenomenon has been reported in *Cucumis* fruits (Webster and Craig, 1976). In the plant introduction accessions we observed, the result was corky tissue developing between pieces of the epidermis. The amount of netting increased slowly, and fruits became completely netted as they matured. In muskmelon, the skin netting was a combination of lenticellar tissue derived from a subepidermal periderm, and cork cells which were complementary tissue of lenticels (Webster and Craig, 1976). It appeared that the netting phenomenon in cucumber may be a result of the same mechanisms operating in muskmelon.

The two plant introduction accessions studied each had slightly different netting. In the case of PI 165509, the netting left the fruit with a red color, with cream colored veins (netting) over the surface. In the case of PI 197086, the netting was an obvious result of the dark, reddish-black epidermis becoming cracked, and then flaking to reveal a cream colored second layer. If the fruits were rubbed, most of the outer cracked skin layer was easily removed. It appeared that "netting" was the cracking the epidermal layer, which was red to dark maroon in color. That trait developed slowly, and resulted in what appeared to be an intermediate amount of netting on immature fruit.

The importance of this finding is that care must be taken when studying netted skin. Immature fruits may appear to be without netting, or intermediately netted, when in fact those fruits will develop complete netting if left in the field until maturity. This is a problem when ratings are made on immature fruit, and correlations to the netted skin phenotype are desired.

Figure 1. Four fruits of different ages from the same plant. Maturity and netting increase from left to right. The plant is from the F₂ generation of a cross between Wis. SMR 18 x PI 197086.




Figure 1. Four fruits of different ages from the same plant. Maturity and netting increase from left to right. The plant is from the F_2 generation of a cross between Wis. SMR 18 x PI 197086.

Literature Cited

1. Clark, R.L. and C.C. Block. 1984. Belly rot resistance in *Cucumis sativus*. *Phytopathology* 74:819 (abstract).
2. Hutchins, A.E. 1940. Inheritance in the cucumber. *J. Agr. res.* 60: 117-128.
3. Peterson, G.C. and L.M. Pike. 1992/ Inheritance of green mature seed-stage fruit color in *Cucumis sativus*. *J. Amer. Soc. Hort. Sci.* 117:643-645.
4. Pierce, L.K. and T.C. Wehner. 1990. Review of genes and linkage groups in cucumber. *HortScience* 25: 605-615.
5. Webster, B.D. and M.E. Craig. 1976. Net morphogenesis and characteristics of the surface of muskmelon fruit. *J. Amer. Soc. Hort. Sci.* 101(4): 412-415.




Figure 1. Four fruits of different ages from the same plant. Maturity and netting increase from left to right. The plant is from the F₂ generation of a cross between Wis. SMR 18 x PI 197086.

Leaf Structure and Photosynthetic Relationships in *Cucumis sativus* var. *sativus* and *Cucumis sativus* var. *hardwickii*

J.E. Staub, B.E. Struckmeyer

Vegetable Crops Research, USDA/ARS, Department of Horticulture, University of Wisconsin-Madison, WI 53076 U.S.A.

Department of Horticulture, University of Wisconsin, Madison 53076

Cucumis sativus var. *sativus* L. (cucumber) is widely cultivated in many parts of the world. *Cucumis sativus* var. *hardwickii* (r.) Alef. is a fully cross-compatible botanical variety of var. *sativus*. Variety *hardwickii* is of potential economic importance since it possesses a multiple lateral branching and sequential fruiting habit not found in var. *sativus*.

We have been using var. *hardwickii* as a source of genes for increasing yield potential in commercial cucumber. We have reported that there are morphological differences between these botanical varieties (Schuman et al., 1985). Three morphological plant characteristics and leaf anatomy were evaluated in two different seasons in the greenhouse for each of five var. *sativus* and five var. *hardwickii* cultigents. We found inter- and intravarietal differences which were, in the main, consistent across growing environments.

Because leaf anatomy and morphology could be important factors associated with productivity differences between var. *hardwickii* and var. *sativus*, we designed a series of studies to more completely define varietal leaf variation and photosynthetic response.

We used the var. *hardwickii* accessions PI 215589, PI 462369, PI 486336, PI 183967, PI 273648. and :K 91176, and the var. *sativus* breeding lines WI 1983 and WI 2238. In addition, we also evaluated the var. *sativus* x var. *hardwickii* derived (F7) line WI 2863. Accessions and lines were grown in a field nursery and leaf thickness, maximum photosynthetic potential and chlorophyll content measurements were taken from each of four plants arranged in a randomized complete block design grown on a m² spacing. Samples were taken during first flowering, but before the fruits had developed beyond approximately 1 cm in diameter (approximately 45 to 50 days after planting). Maximum photosynthetic rate was estimated from leaf slices (discs) prepared from the fourth leaf from the terminal whorl using the technique of Jones and Osmond (1973).

Mating were also made between the var. *sativus* (P2) breeding line WI 1606 and the var. *hardwickii* (P1) PI 215589 to produce reciprocal F1 and BC1 progeny. Studies were conducted in a greenhouse and field nursery. Maximum photosynthetic potential and chlorophyll content measurements of these cultigents and their cross-progeny were taken from each of four plants arranged in a randomized complete block design on a m² spacing. Parents, F1 and BC1 plants were flowering and fruits were enlarging (approximately 2 to 2.5 cm in diameter) on the sampling days.

The results of comparisons between var. *hardwickii* accessions, var. *sativus* lines and a derived line are given in Table 1 and Figure 1. Microscopic measurements indicate that leaves of the var. *sativus* lines examined were thicker than var. *hardwickii*, and the cells of var. *sativus* were larger with a more open arrangement. Maximum photosynthetic rates was greater in var. *hardwickii* when compared to var. *sativus*. Likewise, the dry weight of chlorophyll (mg/g) of var. *hardwickii* was often higher (three of five cases) than that of var. *sativus*. The derived line was usually closer to var. *sativus* for the traits examined.

Cross-progeny comparisons indicated that measurements taken in the field were higher than those in the greenhouse (Table 2). Reciprocal F1 differences (P1 X P2 . P2 X P1) were also observed in the BC1 indicating the cytoplasmic nature of this trait. There appears to be a relationship between the compressed leaf structure of var. *hardwickii* and its high maximum photosynthetic rate.


Table 1. Means (*means*) and standard errors (Sx) of leaf thickness, maximum photosynthetic rate (μ mO₂ evolved/mg chlorophyll/hr), and chlorophyll concentration of several *Cucumis sativus* var. *sativus* and var. *hardwickii* cultigens and a var. *sativus* x var. *hardwickii* derived line [(WI 1606 x PI 215589) F7].

Botanical variety	Line or PI	Chlorophyll (mg/g dry wt.)									
		Thickness (mm)		Photosyn. rate		a		b		a + b	
		Mean	Sx	Mean	Sx	Mean	Sx	Mean	Sx	Mean	Sx
<i>hardwickii</i>	215589	8.0	0.17	56.50	4.50	8.71	1.36	1.56	0.23	10.27	0.38
	462369	9.3	0.26	51.30	4.27	8.37	0.67	3.11	0.30	11.48	0.21
	LJ 91176	11.9	0.26	48.94	4.54	8.50	0.36	2.16	0.61	10.66	0.07
	486336	10.6	0.06	45.65	2.10	7.20	1.33	1.61	0.77	8.81	0.42
	183967	11.2	0.46	58.80	5.29	10.7	0.04	3.44	0.60	14.14	0.17
	273648	9.6	0.14	46.65	4.14	---	---	---	---	---	---
<i>sativus</i>	WI 1983	16.2	0.55	36.56	3.60	8.60	0.14	1.86	0.02	10.46	0.05
	WI 2238	13.5	0.32	28.30	0.75	7.59	0.65	2.75	0.14	10.36	0.19
Derivative	WI 2963	18.2	0.40	37.96	4.00	8.27	0.83	2.10	0.11	10.37	0.72

Table 2. field and greenhouse mean maximum photosynthetic rates (μ mO₂ evolved/mg chlorophyll/hr) of *Cucumis sativus* var. *hardwickii* (PI 215589) and var. *sativus* line WI 1606 and their F1 and BC1 progeny.




	Maximum photosynthetic rate							
	Parents				Cross-progeny			
					BC1	BC1	BC1	BC1
Days								
from	P1	P2	F1a	F1b	(P1 x F1a)	(F1a x P2)	(F1b x P1)	(F1b x P2)
plant	215589	1606	P1 x P2	P2 x P1				
Field								
55	54.24	39.10	105.41	56.28	35.63	31.41	38.45	54.24
56	59.00	36.86	80.77	41.84	32.46	29.30	27.49	30.00
Greenhouse								
42	97.14	34.62	---	---	98.26	162.59	86.75	123.50
43	62.17	42.59	91.64	71.52	112.78	143.39	47.04	75.57
50	68.42	39.26	83.78	31.27	126.36	141.07	---	---

Figure 1. A comparison of the leaf structure of *Cucumis sativus* var. *hardwickii* (A; PI 486336), *cucumis sativus* var. *sativus* (B; WI 2238), and a derivative (C; F7) of the cross between var. *hardwickii* (PI 215589) and var. *sativus* (WI 1606) (10 X).



Literature Cited

1. Jones, H.G. and C.B. Osmond. 1973. Photosynthesis by thin leaf slices in solution. I. Properties of leaf slices and comparison with whole leaves. *Aust. J. Biol. Sci.* 26: 15-24.
2. Schuman, D.A., J.E. Staub and B.E. Struckmeyer. 1985. Morphological and anatomical comparisons between two botanical varieties of *Cucumis sativus*: *hardwickii* and *sativus*. *CGC Rpt.* 8:15-17.



Cucumber (*Cucumis sativus* L.) induced mutations: A *Phaseolus* leaf mutant

M. Rucinska, E. Bergier, K. Niemirowicz-Szczytt and A. Korzeniewska

Department of Genetics and Horticultural Plant Breeding, Warsaw Agricultural University, SGGW, 02-766 Warsaw, Poland

A *Phaseolus* leaf mutation, along with other previously described cucumber mutants produced in our laboratory (1, 2, 3), was obtained by chemical mutagenesis. As far as we know no similar mutation has been described or listed in the CGC cucumber gene list.

This mutant has been generatively reproduced in our laboratory for several years and since it was derived from inbred Borszczagowski line, this was the line used to obtain subsequent generations. While the mutant's morphological description was based on 25 plants, genetic analysis of its inheritance utilized as several hundred individuals. The results of the genetic analysis were analyzed by chi-square contingency tests.

The mutant is detectable at the cotyledons and the first true leaf stage. Cotyledons are small, dark green with violet endings and frequently have a characteristic light spot which appears in the middle. The development of cotyledons is slightly retarded, because the seed coat is not easily separated from the emerging hypocotyl. Mutant plants are smaller than the standard (mutant - 83.5 to 152 cm and standard - 225 to 366 cm) with a smaller number of internodes (mutant = 9-26 and standard = 32-40) (Fig. 1). Mutant leaves are delicate, wrinkled at the end of the main vine, and the leaf margins are undulating and curled (Fig. 2). In most cases leaf blades are not serrated and as such deformations make them appear heart-shaped. The size of the mutant leaf is slightly smaller than that of the standard (mutant leaves = 14 to 16 cm long, and 13 to 17 cm wide with leaf petiole 8 to 12 cm long, while the respective standard dimensions = 19 to 26, 20 to 25 and 12 to 15 cm).

Mutant flowers are remarkably different from those of standard cucumber types (Fig. 3). Mutant plants bear either small and deformed or completely reduced petals. Mutant flowers are reminiscent of flowers in the Labiateae family. Most female flowers are sterile. As common in standard types, more male than female flowers are developed on the plant. The mutant usually sets fruits but they are often deformed and seedless (Fig. 4). As in the standard, mature fruits are brown in color, however the epidermis is not netted.

Following the results of the genetic analysis, it is possible to state that a single recessive gene (*phl* - phaseolus leaf) condition the morphological phenotype described above (Table 1).

Table 1. Inheritance of phaseolus-type leaf (*phl*)

Generation	No. observed		No. expected		Ratio tested ratio	χ^2	P
	Normal	Mutated	Normal	Mutated			
P1 (normal)	30	0	30	0	1:0	---	---
P2 (mutated)	0	25	0	25	0:1	---	---
F1	40	0	40	0	1:0	---	---
F2	335	101	327	109	3:1	0.78	0.50-0/80
F1 x P1	83	0	83	0	1:0	---	---
F1 x P2	124	116	120	120	1:1	0.26	0.50-0.80

Figure 1. Cucumber plant possessing *Phaseolus* leaf mutant.

Figure 2. Comparison of a *Phaseolus* mutant and standard leaf.




Figure 3. Comparison of female and male flowers from *Phaseolus* mutant (left) and standard (right) cucumber plants.

Figure 4. A deformed fruit from a *Phaseolus* mutant (left) compared to the typical fruit from Borszczagowski.




Figure 3. Morphology of flowers from normal and mutant plants.




Figure 4. Morphology of fruits from normal and mutant plants.

Literature Cited

1. Rucinska, M. K. Niemirowica-Szctt and A. Korzeniewska. 1991. A cucumber (*Cucumis sativus* L.) mutant with yellow stem and leaf petioles. *Cucurbit Genet. Coop. Rpt.* 14:8-9.
2. Rucinska, M., K. Niemirowica-Szcytt and A. Korzeniewska. 1992. Cucumber (*Cucumis sativus* L.) induced mutations. II. Short petioles mutations. *Cucurbit Genet. Coop. Rpt.* 15: this volume.
3. Rucinska, M., K. Niemirowica-Szcytt and A. Korzeniewska. 1992. Cucumber (*Cucumis sativus* L.) induced mutations. III and IV. divided and ginko leaves. Fifth Eucarpia Cucurbitaceae Symposium, Poland. July 27-31, 1992. pp. 66-69.




Figure 1. Chemically-induced *Phaseolus* leaf mutant of cucumber.




Figure 2. Morphology of leaves from normal and mutant plants.




Figure 3. Morphology of flowers from normal and mutant plants.




Figure 4. Morphology of fruits from normal and mutant plants.

Independence of fruit length and 10 other characters in cucumber.

Nick E., Fanourakis

Technical Educational Institute, Heraklion Crete 71500, Greece

Fruit length of cucumber exhibits great variability. Although the inheritance of fruit length has been investigated, data are not comparable because of the varying germplasm sources used and the collection and analysis measure applied (2, 3, 4, 5). Limited agreement exists regarding the inheritance of fruit length, however most researchers agree that it is a quantitative character. The quantitative nature of the inheritance of fruit length and its association with the fruit neck has been studied. (!). We report here the linkage relations of fruit length with 10 other qualitative characters of cucumber.

The long fruited line 791 from our cucumber breeding program was crossed with the non-necked American variety SMR-18 and F2 and BC progeny were produced. Parents, F1, F2, BCP1 and BCP2 progeny were evaluated for fruit length and the following characters: bitterness (**bi**), powdery mildew resistance (**pm**), female sex expression (**E**), spine size (**ss**), warted fruit (**Tu**), uniform color of immature fruit (**u**), mature fruit color (**R**), glossy fruit (**D**), structure of epidermis (**pe**), and spine color (**B**). Evaluation and classification was made as previously described (1).

The symbols "+" and "-" were used to denote contrasting phenotypic classes for the segregating characters. Thus "+" stands for bitterness, femaleness, rough spines, warted fruit, nonuniform color of immature fruit, red color of mature fruit, dull fruit epidermis, palisade epidermal structure, and black spines. The "-" stands for nonbitterness, monoecious flowers, fine spines, nonwarted fruit, uniform color of immature fruit, cream color of mature fruit, glossy epidermis, flat epidermal structure, and white spines respectively. For powdery mildew "R" designates resistance, "I" intermediate resistance and "S" susceptibility.

The continuous variation of fruit length allowed for separation of F2 and BC generations in classes differing by 8 cm. Plants were classified for other characteristics within each class of fruit length. Observed ratios within each class were compared with the total ratio of the characteristic over the classes by X^2 analysis. Thus the deviation of the partial ratios from the total ratio of the characteristic indicates linkage associations of any characteristic with the fruit length.

Partial phenotypic segregations and X^2 analysis for each particular character are generally indicated in Tables 1 and 2. Observed ratios within each class of fruit length do not indicate significant deviations from the total phenotypic character ratios. Significant deviations were, however, found in one F2 class for female sex expression, and in some classes of the BC1P2 generation for spine size, warted fruit, uniform color of immature fruit, dull fruit epidermis, and structure of epidermis. The rest of the partial BC ratios for these characters follow the expected segregation. Moreover, the respective partial ratios in the F2 generation segregate according to the expectation. Thus, it is hard to accept the observed deviation as indicative of any linkage relationships. Therefore, we propose that the segregation of the fruit length and the characteristics evaluated is independent.

Table 1. Phenotypic F2 segregation and X^2 values of the 10 characters within each class of fruit length of cucumber.

Trait		Total ratio	Fruit length (cm)					
			12-19, 99		20-27, 99		>28.00	
			Ratio	X^2	Ratio	X^2	Ratio	X^2
bi	+:	151	12	0.10	94	1.64	47	3.34
	-:	49	3		37		7	
	R:	14	0	1.54	11	0.49	3	1.85
pm	I:	22	1		13		9	

	S:	164	14		108		42	
F	+:	81	9	2.36	43	3.20	29	3.91
	-:	119	6		88		25	
SS	+:	157	12	0.02	95	2.33	40	1.09
	-:	43	3		35		15	
Tu	+:	157	12	0.02	106	0.43	39	1.88
	-:	43	3		24		16	
u	+:	157	12	0.02	106	0.43	39	1.88
	-:	43	3		24		16	
R	+:	156	13	0.59	102	0.01	41	0.51
	-	44	2		28		14	
D	+:	157	11	0.24	106	0.43	40	1.09
	-:	43	4		24		15	
pe	+:	156	12	0.02	106	0.43	38	2.89
	-:	44	3		24		17	
B	+:	155	13	0.59	101	0.06	41	0.51
	-	45	2		29		14	

* Significant difference at 0.05 level.

Table 2. Ssegregation and X^2 values of the BCX1P2 generation for the 10 characters of cucumber within each class of fruit length.

Trait		Total ratio	Fruit length (cm)					
			0-27.99		28-35.99		>36.00	
			Ratio	X^2	Ratio	X^2	Ratio	X^2
bi	+:	81	4	0.25	60	0.02	17	0.35
	-:	88	6		67		15	
pm	R:	33	2	0.28	27	0.80	4	2.40
	I:	46	4		30		12	
	S:	97	7		74		16	
F	+:	125	7	1.86	94	0.03	24	0.24
	-:	51	6		37		8	
ss	+:	55	9	8.74**	43	0.15	3	7.13**
	-:	121	4		88		29	
Tu	+:	72	9	4.31*	59	0.92	4	10.68**
	-:	104	4		72		28	
u	+	73	8	2.16	61	1.40	4	11.06**
	-	103	5		70		28	
R	+:	92	7	0.01	69	0.01	16	0.03
	-:	81	6		60		15	
	+:	74	9	3.93*	61	1.14	4	11.45**

D	-:	102	4		70		28	
	+:	72	9	4.15*	59	1.01	4	11.00**
pe	-:	102	4		70		28	
	+:	92	7	0.01	69	0.01	16	0.07
B	-:	84	6		62		16	

*, **: significant differences at 0.05 and 0.01 levels, respectively.

Literature Cited

1. Fanourakis N.E. and E.E. Tzifaki. 19193. Correlated inheritance of fruit neck with fruit length and linkage relations with 10 other characteristics of cucumber. *Euphytica* (in press).
2. Imam, M.K., M.A. Abobakr and H.M. Yacoub. 1977. Inheritance of some characters in cucumber, *Cucumis sativus* L. II. Some quantitative characters. *The Libyan J. Agric.*
3. Kingston, B.D. 1973. A genetic study of some fruit characteristics of the cucumber, *Cucumis sativus* L. Ph.D. Thesis, Texas A and M Univ., College Station.
4. Owens, K.O. 1982. Analysis of generation means and components of variance for fruit size in two cucumber populations, and genetic and breeding studies on cucumber fruit size utilizing inbred backcross lines. Ph.D. Thesis, Univ. of Wisconsin-Madison.
5. Pike, L.M. and B.D. Kingston. 1975. Inheritance of fruit length and shape in cucumber, *Cucumis sativus* L. *HortSci.* 10:318.
6. Smith, O.S., R.L. Lower, and R.H. Moll. 1978. Estimates of heritability and variance components in pickling cucumber, *J. Amer. Soc. Hort. Sci.* 103: 222-225.

Diallel analysis of cucumber germination at optimum and suboptimal temperatures

P. Milotay

Vegetable Crops Research Institute, 6000 Kecskemet, P.O.B. 116, Hungary

Germination of cucumber seeds under cool weather conditions is frequently erratic resulting in poor, or non-uniform emergence. cucumber seeds germinate rapidly between 20-30C, however there is a sharp reduction in germination percentage and initial seedling development below 15C (3). Although laboratory selection of a genetically broad-base cucumber population for improved germinability at suboptimal temperature resulted in better and faster emergence in the field (4), heritability for this character is low (6). To study the nature of genetic effects controlling germination under optimum and suboptimal temperature a laboratory experiment was conducted utilizing seeds produced from a set of diallel crosses.

Five F5 to F7 generation parthenocarpic pickling cucumber lines of different origin and their hybrids were used. Seeds were produced during the spring in a plastic house. One hundred seeds (four replications of 25 seeds) of parents and crosses were placed into the folding of moistened filter papers. Each row of seeds (one replication) were placed into the cut foldings to allow for unconstrained radicle and hypocotyl growth. The filter papers were then rolled up, and placed in plastic bags, and held in incubators in a vertical position at 17 and 25C. After 4 and 7 days at 17C and 4 days at 25C percentages of radicle protrusion, radicle and hypocotyl elongations were recorded/

Statistical analysis was carried out according to the model 2, , method 1 of Griffing (2). Before the analysis percentage data were angularly transformed. Maternal effects were estimated by the method of Topham (5).

At 25C all seed lots germinated over 95% (Table 1), and the average heterosis of hybrids was 1.02. At 17C germination proceeded slower and the average performance of hybrids over the mean of the parents was 0.89 at day four and 0.92 at day seven. At seven days the poorest seed lots germinated near 50%, while the best parents and hybrids showed maximum germination. Keeping lagging lots seeds at 17C to day 12 resulted in only a slight improvement.

Radicle elongations showed relatively small, but significant differences at optimum temperature (Table 2). More marked differences were detected at 17C resulting in an average heterosis of 1.22 and 1.06 at day four and day seven respectively. Variability increased within the seed lots at suboptimal temperature also. Hypocotyl development followed radicle protrusion by 1.5 days at optimum temperature and about 3 days at suboptimal temperature (data not shown).

The Griffing analysis resulted in significant general combining ability (gca) and specific combining ability (sca) effects for all characters at both temperatures. Reciprocal effects were also significant except for germination percentage at 25C. According to the variances (Table 3), dominance and non-allelic effects were much more important than additive effects at suboptimal temperature both for germination percentage and seedling growth. The average degree of dominance for these characters refers to over-dominance. At 25C, over-dominance was found only for germination percentage and radicle elongation was conditioned by additive genetic effects.

When analyzing gca effects, line KP exhibited significantly positive values, and line 17K significantly negative values for percent germination at 17C. In case of radicle growth, lines Ac18 and KP produced significantly positive gca values at the same temperature (data not shown).

Narrow-sense heritability values obtained for percent germination were not significant at both temperature and radicle elongation gave only higher heritability at 25C ($h^2 = 0.62$). Significant maternal effects in this study were obtained at sub-optimal temperature for percent germination (0.68) and radicle length growth (0.77). At optimum temperature only for radicle length (0.61) was significant. The nature of this effect should be studied further (1).

The parthenocarpic pickling cucumber lines involved in this study do not provide sufficient genetic base for improving germinating ability at suboptimal temperature.

Table 1. Germination of cucumber seeds at 35C after 4 days, at 17C after 7 days, and at 17C after 4 days (transformed data).

Parents	Temp/days	S750	Ac18	M72	KP	17K	Mean
S750	25/4	87.1	90.0	90.0	85.9	90.0	88.6
	17/7	85.9	82.0	70.7	47.3	52.4	67.6
	17/4	83.3	75.4	56.690.0	31.4	42.3	57.8
Ac18	25/4	90.0	90.0	82.6	87.1	90.0	89.4
	17/7	87.1	72.6	70.7	80.0	51.7	74.9
	17/4	83.3	66.5	77.1	72.6	42.4	67.1
M72	25/4	90.0	87.1	60.5	87.1	81.8	84.8
	17/7	87.1	75.6	53.8	81.3	47.9	70.5
	17/4	82.0	67.1	90.0	78.5	35.9	63.5
KP	25/4	90.0	87.1	87.1	90.0	90.0	89.4
	17/7	90.0	90.0	83.3	90.0	56.9	82.8
	17/4	90.0	85.9	90.0	80.4	35.0	74.9
17K	25/4	90.0	87.1	90.0	90.0	90.0	89.4
	17/7	83.0	84.1	66.9	86.4	90.0	78.1
	17/4	55.6	75.7	62.8	82.1	84.2	72.1
Mean	25/4	89.4	88.3	87.4	88.0	88.4	88.3
	17/7	82.6	80.9	73.5	77.0	59.8	74.8
	17/4	78.8	74.1	65.4	69.0	48.0	67.1

Critical differences at 25C after 4 days, 17C after 7 days and 4 days = 3.9, 8.8 and 10.7, respectively

Table 2. Radicle length (mm) of cucumber seedlings at 25C after 4 days, at 17C after 7 days and at 17C after 4 days.

Parents	Temp/days	S750	Ac18	M72	KP	17K	Mean
S750	25/4	93.8	103.6	100.3	90.2	93.5	96.3
	17/7	48.7	58.4	54.5	35.1	35.6	46.5
	17/4	9.1	12.4	11.4	2/9	6.3	8.4
Ac18	25/4	105.2	105.1	106.9	102.1	93.8	103.8
	17/7	81.3	49.9	56.0	64.0	38.6	54.0
	17/4	12.3	5.9	7.2	10.6	4.4	8.1
M72	25/4	99.2	105.1	93.1	99.4	80.5	95.5
	17/7	60.8	53.0	44.9	63.1	43.4	53.0
	17/4	15.1	10.2	4.1	11.5	5.8	9.3
KP	25/4	104.7	106.2	106.3	97.4	85.8	101.1
	17/7	65.2	68.6	64.8	53.2	34.1	57.2
	17/4	16.3	17.3	15.6	11.8	7.0	13.7
17K	25/4	89.6	95.6	97.5	96.9	85.3	95.0
	17/7	42.2	51.9	53.8	50.4	53.6	50.4

	17/4	11.0	11.0	12.4	10.4	12.3	11.4
Mean	25/4	98.5	103.1	100.8	87.2	91.0	98.1
	17/7	55.8	56.4	54.8	53.2	41.4	52.2
	17/4	12.9	11.4	10.1	9.4	7.2	10.2

Critical differences at 25C after 4 days, 17C after 7 days and 4 days = 3.4 5.5 and 2.6, respectively.

Table 3. Variances and genetic components of germination and seedling growth analyzed by Griffing's method 1.

Character	Mg	Ms	Mr	Me	O ² _g	O ² _s	h ²	a
OPTIMUM TEMPERATURE								
Germination %	16.25**	10.79*	5.44 ns	4.46	0.58	3.77	0.12	1.81
Radicle length	139.70***	21.50***	36.30***	4.85	23.80	9.33	0.62	0.65
SUBOPTIMUM TEMPERATURE								
Germination % day 4	236.93***	260.96***	428.47***	33.58	0.00	135.00	0.02	---
Germination % day 7	194.22***	204.21***	235.63***	24.87	2.15	106.76	0.02	6.81
Radicle length day 7	159.16***	98.12***	78.99***	6.61	6.54	54.43	0.18	2.04

*, **, *** = significant at 5, 1 and 0.1% level, respectively.

Literature Cited

1. Edwards, M.D. and R.L. Lower. 1982. The genetic regulation of several seed traits in compact (cpcp) cucumbers - Maternal vs. embryonic control. Cucurbit Genet. Coop. Rpt. 5:8-9.
2. Griffing, B. 1956. Concept of general and specific combining ability in relation to diallel crossing systems. Austr. J. Biol. Sci. 9:463-493.
3. Simon, E.W., A. Minchin, M.M. McMenamain, and J.M. Smith. 1976. The low temperature limit for seed germination. new Phytol. 77:301-311.
4. Staub, J.E., R.L. Lower and J. Nienhuis. 1988. Correlated responses to selection for low temperature germination in cucumber. HortScience 23.4:745-746.
5. Topham, P.B. 1966. Diallel analysis involving maternal and maternal-intraunion effects. Heredity 21:665-674.
6. Wehner, T.C. 1982. Genetic variation for low-temperature germination ability in cucumber,. Cucurbit Genetic Coop. Rpt. 5:16-17.

Application of factor analysis to cucumber breeding

Meng Zhang and Hongwen Cui

Department of Horticulture, Northwestern Agricultural University, Yangling Shanxi, 712100, P.R. China

Cucumber breeding objectives suggested by the "Chinese Cucumber Breeding Cooperation Team" include the development of an early maturing, high yielding cucumber which is resistant to anthracnose, fusarium wilt and downy mildew disease, and which possesses good commodity characters for Chinese markets. These objectives require the incorporation of many quantitatively inherited traits.

It is necessary to develop a comprehensive selective method for reaching these objectives. Studies which identify genetic relationships among traits (correlations) is a prerequisite for any comprehensive selection method. There are certain limitations inherent in traditional correlation analysis. Factor analysis, however, can be used to overcome some problems which are inherent in correlation analysis. The aim of this study was to investigate genetic relationships among several traits in cucumber using factor analysis.

An experiment was conducted at the Horticulture Station of the Northwestern Agricultural University. Twenty-four varieties and inbreds were evaluated in a randomized block design with 3 replications. Ten plants of each cultivar were randomly chosen to evaluate 26 quantitative traits during the growth period. Traits included: 1) the node position of the first pistillate flower (x_1); 2) the days from sowing to the first pistillate flowering plant in the population (x_2); 3) the days from sowing to first male flowering plant in the population (x_3); 4) the days from sowing to staminate flowering in 50% of the plants (x_4); 5) the days from sowing to staminate flowering in 50% of the plants (x_5); 6) leaf area per plant in the early stage (x_6); 7) fruit length (x_7); 8) fruit diameter (x_8); 9) leaf number in the early stage (x_9); 10) pistillate flower density (main vine) in the early stage (x_{10}); 11) number of pistillate flowers (main vine) in the early stage (x_{11}); 12) number of staminate flowers (main vine) in the early stage (x_{12}); 13) number of harvested fruit per plant in the early stage (x_{13}); 14) average fruit weight in the early stage (x_{14}); 15) early yield per plant (x_{15}); 16) total number of branches (x_{16}); 17) average length between two close nodes (x_{17}); 18) total number of leaves (x_{18}); 19) the highest fruit setting node (x_{19}); 20) number of harvested fruits per plant (x_{20}); 21) total fruit weight per fruit (x_{21}); 22) total yield per plant (x_{22}); 23) downy mildew resistance in the early stage (x_{23}); 24) downy mildew resistance in the late stage (x_{24}); 25) fusarium wilt disease incidence (percentage; x_{25}) and; 26) anthracnose disease incidence severity (x_{26}). Resistance to anthracnose was identified by an *in vitro* leaf method.

Original data of x_9 , x_{23} and x_{24} underwent anti-sine transformation and the variance was analyzed in a randomized block design. Genotype values (g_{ij}) of traits were estimated according to Chui-Yu Liu (1981) and then genotype correlation coefficients were calculated on the basis of genotypic value.

The main factor solution (factor analysis) was calculated using the genotype correlation matrix. The orthogonal factor load matrix was calculated by orthogonal rotation and transformation of the initial maximum variance using the BLQMIN oblique rotation method. The oblique factor load matrix and oblique factor correlation matrix were also analyzed.

Variance analysis detected significant differences among all observed traits, except for fruit diameter (x_8). Results show that the difference between model parameters was caused by genetic factors. Therefore, the genetic analysis could be evaluated in 25 x 25 matrix (25 traits).

Most of the trait loads centralized on a factor (F_{1-5} , Table 1). A biological explanation of all factors can be given: in factor F_1 , x_6 , x_8 , x_7 , x_{13} , x_{14} , x_{16} , x_{19} , x_{21} , x_{23} , x_{24} and x_{25} traits occupied the higher load. These traits centralized all yields and component factors, except the x_{12} trait. Therefore, F_1 factor is named the "Yield factor". Yhtrr kinds of disease (anthracnose, fusarium wilt and downy mildew) had a higher load, which indicated that there was close association between plant resistance disease level and yield. All of the susceptible parameters were negatively loaded, which indicated that they are negative factors for yield.

Higher traits loads in factor F_2 included x_1 , x_3 , x_9 , x_{10} , x_{11} , x_{12} , and x_{14} which were associated with traits of earliness, and F_2 was named the "early - mature factor". The node of the first female flower, pistillate flower density (main vine) in the early stage, and number of pistillate flowers (main vine) in early stage had very high loads. Results indicate that these traits had the greatest effect on earliness.

Higher trait loads in factor F_3 included x_6 , x_8 , x_{17} and x_{18} which reflected plant nutrient level and plant growth vigor. Therefore, this factor was named the "Yield physiology factor". The load of total average fruit weight was high and average fruit weight at early stage also had a positive load. These data indicate physiological factors had a direct positive effect on fruit weight.

The load of x_2 , x_3 , x_4 , and x_5 was higher than other traits in factor F_4 and was named the "flowering season factor". In addition, x_6 , x_8 , x_7 and x_{12} traits also had a certain load. The number of harvested fruit was only positive value. Results show that components of this factor could result in delayed growth and development, while influencing yield by increasing fruit weight and reducing the number of fruit.

High trait loads were observed for x_4 , x_5 , x_{11} , x_{15} , x_{22} and x_{23} in factor F_5 . This factor was named the "male flower development factor". The time of male flowering and the number of male flowers (increase) during the early part of the season was closely associated with downy mildew susceptibility.

Table 1. Factor loading matrix after orthogonal rotation transformation (genotype).

Traits	F_1	F_2	F_3	F_4	F_5
X_1	0.0909	-0.7137	0.3977	0.2239	0.1267
X_2	0.3323	-0.3775	-0.0021	0.7427	0.2136
X_3	0.1331	-0.5830	0.1883	0.5687	0.1845
X_4	0.2033	0.0274	-0.0219	0.8166	-0.3887
X_5	0.3820	-0.0692	-0.1084	0.7453	-0.3660
X_6	0.7670	-0.0303	0.5123	0.3280	-0.1242
X_7	0.8639	-0.0504	0.1894	0.3222	0.1484
X_8	0.5668	-0.0169	0.6095	0.3367	-0.1954
X_9	0.0943	0.9708	-0.0675	0.0824	0.1457
X_{10}	0.1077	0.8622	0.0986	-0.0330	0.1400
X_{11}	0.0516	-0.4363	0.3967	-0.0462	0.5504
X_{12}	0.3366	0.7722	-0.1883	-0.3558	0.0500
X_{13}	0.8306	-0.0427	0.2732	0.3145	0.1383
X_{14}	0.8109	0.5093	-0.0292	0.0283	0.0666
X_{15}	0.0012	-0.3291	0.1474	0.1302	-0.695

X ₁₆	0.7747	-0.1202	0.2234	0.0804	0.0307
X ₁₇	0.2910	-0.2044	0.8847	-0.136	0.0929
X ₁₈	0.4830	-0.1631	0.8038	-0.0436	0.1051
X ₁₉	0.8934	0.2664	0.1526	0.0530	-0.2160
X ₂₀	0.8191	-0.0834	0.4010	0.1978	-0.1020
X ₂₁	0.9173	0.0603	0.1689	0.1056	-0.2768
X ₂₂	-0.4571	0.2414	0.3065	-0.2842	0.5540
X ₂₃	-0.7594	-0.0988	0.1045	-0.1685	0.4936
X ₂₄	-0.8134	-0.1482	-0.2374	-0.2374	-0.1266
X ₂₅	-0.5227	-0.1296	0.1039	0.2561	0.3375

Factor analysis may be useful for describing and clarifying interrelationships among traits. As such this technique may be an important guide for selection during cucumber breeding.

Literature Cited

1. Liu Chio-Yu. 1981. The genetic correlation and its addibility of crop quantitative characters. *Anhui Journal of Agricultural Sciences*. Special Edit for Quantity Genetics, 84-88.

Application of narrow sense canonical characters to cucumber breeding

Hongwen Cui, Yongtao Qi

Department of Horticulture, Northwestern Agricultural university, Yangling, Shaanxi, 712100, P.R. China; Beijing Vegetable Research Center, Beijing, 1000081, P.R. China

Previous studies have shown that the correlation between heterosis of early yield in F_1 hybrids and its associated component (ACT) traits was near 1.0. Early maturity of F_1 hybrids is closely related to the trait constitution of parental lines. In order to develop a hybrid which possesses early maturity heterosis, parental lines must have both early maturity and ACT. Thus, the establishment of a reliable and effective selection scheme is essential. Such a scheme should possess the following: 1) a means for overcoming complicate correlations among important agronomic characteristics and unfavourable traits; 2) a means for overcoming low heritabilities often associated with quantitative characters, and, 3) a means for overcoming interference of non-additive effects in the early generations. A selection scheme which incorporates independent multiple characters with maximum narrow sense heritabilities may be reliable and effective during the course of pedigree breeding. Such a scheme is described below.

The additive variance and covariance of traits were separated from the genetic variance and covariance using analysis of combining ability covariance with eight parental inbred lines and 16 F_1 hybrids. This produced a narrow sense correlation heritability matrix (H_N). The H_N was used to construct two narrow sense canonical multiple characters with maximum narrow sense heritabilities. Four inbred lines, Changmi-1, Zanqiu-m, Pinli-m, and Hei235 were used as male parents and four inbred lines, Yue82, Jing4-3-1, Xilong58-5, and 7742 were used as female parents to produce an incomplete diallel.

The resulting F_1 hybrids were evaluated in a randomized complete block design with three replications. the narrow sense correlation heritability for two traits X_i and X_j (h^2_{Nij}) was estimated by means of variance and covariance of combining ability according to a random model.

The narrow sense correlation heritability matrix:

$H_N =$	h^2_{N1}	h^2_{N12}	h^2_{N1m}
	h^2_{N21}	h^2_{N2}	h^2_{N2m}
	h^2_{Nm1}	h^2_{Nm2}	h^2_{Nm}

The concepts of narrow sense canonical character were expressed as:

$$P^*_{Ni} = b_i'p \quad (i = 1, 2, \dots, S, S < M);$$

Such that p^*_N must satisfy the following two terms: 1) $p^2(b'p, b'A) = \max$ and 2) $D(b'p) = 1$,

The phenotypic random vector $P = (p_1 \ p_2 \ \dots \ p_m)'$ and

The additive random vector $A = (a_1, a_2, \dots, a_m)'$ consisting P^*_{Ni} is calculated by solving matrix equation $(H_N - F^2R_p)\beta = 0$.

The following traits were chosen as objective traits for the comprehensive selection scheme: 1) early yield per plant (S1); 2) number of early fruit (S2); 3) number of early female flowers on the main stem (S3); 4) total number of branches (S4); 5) number of leaves at the time of first flowering (S5); 6) the node position at which the first female flower appeared (S6).

H_N was estimated by means of combining ability using covariance analysis. Six eigenvalues and six eigenvectors of H_N for R_p were obtained by solving the matrix equation $(H_N - \lambda R_p) \beta = 0$ (Table 1).

Table 1. The broad sense eigenvalues and eigenvectors of H_N for R_p .

Eigenvalues	0.998 f_1	0.773 f_2	0.258 f_3	-0.895 f_4	-2.520 f_5	-5.306 f_6
Adjusted eigenvalues	0.998	0.773	0.258	0.000	0.000	0.000
S_1	0.449	0.492	-1.187	0.356	0.081	-2.382
S_2	-0.315	0.074	-0.094	0.328	0.099	4.179
Eigenvectors S_3	-0.353	-0.251	-0.385	0.482	-2.259	-1.538
S_4	0.062	0.640	1.106	1.452	-0.164	1.025
S_5	0.192	-0.904	-0.610	0.788	0.765	-1.482
S_6	0.490	0.077	-0.963	-0.902	-2.457	0.818

Eigenvalues (f) reflect the narrow sense heritabilities of corresponding canonical characters in biological sense (Table 2). For example, $f^2_3 = 0.2583$, it is very low as a heritability, f^2_4, f^2_5, f^2_6 are negative numbers and can be regarded as zero.

Table 2. The heritabilities (f^2) and their corresponding eigenvectors for the broad sense and narrow sense canonical characters f^2_1, f^2_2, f^2_3 used to structure canonical characters.

Heritabilities			Bigenvectors					
			s1	s2	s3	s4	s5	s6
f^2_1	Broad sense	0.999	-1.640	2.184	0.194	0.836	-1.984	1.283
	Narrow sense	0.998	0.449	-0.315	-0.353	0.062	0.192	0.490
f^2_2	Broad sense	0.999	0.517	-1.290	1.450	0.092	0.782	-0.823
	Narrow sense	0.773	0.492	0.074	-0.251	0.640	-0.904	0.077
f^2_3	Broad sense	0.928	0.382	-0.743	-1.508	-0.906	0.113	-1.542
	Narrow sense	0.258	-1.187	-0.094	0.385	1.106	-0.610	-0.963
f^2_4	Broad sense	0.897	1.118	-1.364	0.473	1.109	0.167	-0.990
	Narrow sense	0.000	0.356	0.328	0.482	1.452	0.788	-0.902
f^2_5	Broad sense	0.785	1.600	-2.837	1.800	-1.416	0.143	1.515
	Narrow sense	0.000	0.081	0.099	-2.259	-0.164	0.765	-2.457
f^2_6	Broad sense	0.724	-0.863	-0.896	0.405	0.244	-0.143	0.847
	Narrow sense	0.000	-2.382	4.179	-1.538	1.025	-1.482	0.818

The broad sense canonical heritabilities is much larger than those of the narrow sense canonical characters. These results indicate that genes controlling each objective trait have many non-additive effects. There is a large difference between broad sense and narrow sense canonical characters in eigenvalues and weighting coefficients for objective traits.

*The first canonical character (P^*_{N1}).* In accordance with the quantity and direction of weighting coefficient of each target trait, we used a reverse P^*_{N1} selection. That is, selecting the lineage with small P^*_{N1} value (i.e., the lineage with low node position at which the first female flower appeared, more female flowers in main stem and early fruits harvested, fewer branches and leaves at first flowering). These characteristic are all important in early maturity.

*The second canonical character (P^*_{N2}).* We employed a positive P^*_{N2} selection. that is, the lineage with fewer leaves at first flowering and high early fruit yield were selected. Nevertheless, P^*_{N2} selection was ineffective in reducing the number of branches.

Table 3. The narrow sense canonical character values of 8 parental and 16 hybrids (F1)

	Narrow sense canonical character 1 (P^*_{N1})	Narrow sense canonical character 2 (P^*_{N2})
Yue 82 (M1)	1.0757	-3.7898
Jing 4-3 (M2)	1.6548	-0.8218
Xilong 58-5 (M3)	1.6158	-1.9314
7742 (M4)	3.8848	-3.5500
Changmi-1 (F1)	1.3303	-1.4092
Zhangqiu-m (F2)	7.2461	-6.9080
Pinli-m (F3)	1.9358	-1.9746
Hei235 (F4)	3.4756	-1.5404
H ₁	0.3051	-2.2553
H ₂	2.2231	-2.7390
H ₃	2.3907	-1.7190
H ₄	1.6023	01.4923
H ₅	1.2318	-0.3444
H ₆	2.5815	-1.1398
H ₇	2.0961	-0.5351
H ₈	2.8573	-0.0409
H ₉	2.2798	-0.9479
H ₁₀	3.1833	-1.6005
H ₁₁	2.4834	0.0571
H ₁₂	3.1118	-1.1334
H ₁₃	2.7619	-2.2661
H ₁₄	3.5662	-2.1424

H ₁₅	3.1481	-1.3824
H ₁₆	3.0969	-1.3155

F₁, M₂, and M₃ (Table 3) were selected in P*_{N1} reverse selection and F₁, F₄, M₂, and M₃ were selected in P*_{N2} positive selection. F₁, M₂ and M₃ were all chosen in the selection of P*_{N1} and P*_{N2}. Data indicate that all inbred lines F₁ (Changmi-1), M₂ (Jing4-3-1) and M₃(Xilung 58-5) possess better characteristics for early maturity. However, in the course of P*_{N2} selection, none of the inbred lines were outstanding and thus all lines tested have shortcomings in this regard.

Maximum heritabilities and independence of canonical characters embody the advantages of this breeding method ⁽¹⁾ ⁽²⁾ . Yang De and DaiJunti (1983) reported on canonical character selection in the offspring of wheat populations ⁽³⁾ . Canonical analysis of early maturing cucumber lines overcame serious interferences with non-additive effects and thus is viewed as a reliable comprehensive selection scheme for early generations. As an overall appraisal, it can also define the best combinations from which unique early-mature diphyletic lines can be synthesized during the course of pedigree selection. Nevertheless, only the use of ines with maximum broad sense heritabilities of canonical characters can avoid the serious interferences with broad sense canonical selection in early generations. In order to overcome such defeats, new breeding methods should be developed which allow for the establishment of narrow sense canonical characters with maximum narrow sense heritabilities. Such methods will increase the accuracy and reliability of canonical selection and thus overall appraisal of parental lines and F₁ hybrids.

Literature Cited

1. Yongtao, Q. and C. Hongwen. 1991. High effective multiple selection of parental lines of cucumber hybrid with strong early mature heterosis. *Scientia Agriculture Sinica* 24:51-58.
2. De, Y. and D. Junti. 1983. Approaches to canonical correlations of multiple quantitative traits. (I). canonical character and its heritability. *J. Huanan Agr. college*, No. (1): 27-36.
3. De, Y. and D. Junti. 1983. Approaches to canonical of multiple quantitative traits. (I) canonical character selection for the offspring populations of crossing combination of wheat. *J. Hunan Agr. college*, No. (1):37-51.

Cucumis Germplasm: 1992 Collection Expedition in India

J.D. McCreight, J.E. Staub, N.M. Koppar and U. Ch. Srivastava

U.S. Department of Agriculture, Agricultural Research Service, U.S. Agricultural Research Station, 1636 East Alisal Street, Salinas, CA 93905, USA; U.S. Department of Agriculture, Agricultural Research Service, Department of Horticulture, University of Wisconsin, Madison, WI 53706, USA; National Board for Plant Genetic resources, Pusa, New Delhi-110012, INDIA

The government of India and the United States of America recently joined in a cooperative effort for: 1) the construction of modern germplasm storage facilities and establishment of a computerized germplasm database in India; 2) joint germplasm explorations in the two countries; and 3) the training of Indian scientists in the U.S. Most of the U.S. portion of the funding of this cooperative effort is from the U.S. Agency for International Development (USAID) through the USDA, Office of International Cooperative Development (OICD), a sister agency of USDA, ARS. The first cooperative effort of this program was a sunflower germplasm collection trip in the U.S. in 1992. The second cooperative effort was a *Cucumis* exploration trip in India which we undertook in October and November, 1992.

The primary objective of the *Cucumis* expedition was the collection of landraces of cucumber (*Cucumis sativus* L.) and melon (*Cucumis melo* L.) Other cucurbits were also collected during the expedition. Collections were made in three states (Rajasthan, Madhya Pradesh, and Uttar Pradesh) immediately following the rainy season (July through September). Seeds were collected from cultivated and non-cultivated areas, vegetable markets (subji mundi) and from seed dealers. Though many samples were collected as fruit we were not always able to observe the plants or the growing areas. Whenever possible, notes were taken at the collection site about the origin, description, and use of the collections.

The expedition provided several surprises. Cucumber landraces were difficult to find due to 1) the timing of the expedition (i.e. too late in the year), 2) a five-year drought that had completely eliminated the stocks of landraces in some areas (e.g., Sri Ganganagar), and 3), the slow adoption of open-pollinated varieties available through local seed dealers. An overwhelming number of melons were found in fields and markets. There were differences in botanical nomenclature of melon between the Indian and U.S. scientists (1). All melons including *Cucumis melo agrestis* are used as fresh (salad), cooked (vegetable) or dried preparations in India.

Approximately 677 samples were collected. Of these, 665 were cucurbits (Table 1) and 12 were non-cucurbits. There were approximately 186 cucumber and 447 melon collections, and they fell into several sub-groups (Table 2). Exact numbers of each will be known after discrepancies in the records have been resolved, and plants of some accessions have been observed.

These seeds will be available after they have been increased and properly documented (phenotypic and genotypic characterization) in the U.S. Germplasm Resources Information Network (GRIN) database. To that end, tentative 1993 plans call for an increase of a portion of these cucumber and melon collections in the U.S. and a substantial parallel increase of the collections in India. Based on the success and cooperation of the 1992 exploration, a proposal has been submitted to USAID through ARS, OICD, for a second joint *Cucumis* collection exploration in southern India and the Himalayan foothills in the 1994 summer season (May-June).

Table 1. Number of cucurbit and non-cucurbit species collected in India during October and November, 1992.

Species	By State			
	Rajasthan	Madhya Pradesh	Uttar Pradesh	Total
Cucurbit				
<i>Citrullus</i> sp.				

	7			7
<i>Cucumis melo</i>	301 ^{zy}	146 ^x		447
<i>Cucumis sativus</i>	104	63 ^{yw}	18	185
<i>Cucurbita maxima</i>	1			1
<i>Cucurbita moschata</i>	4			4
<i>Cucurbita pepo</i>	3	2	1	6
<i>Langenaria siceraria</i>	4			4
<i>Luffa acutangula</i>	1	2		3
<i>Luffa cylindrica</i>	1			1
<i>Momordica charantia</i>	2	1		3
<i>Momordica dioca</i>		1		1
<i>Praecitrullus</i> sp.	2			2
<i>Triconsanthea bracteata</i>	1			1
Non-Cucurbit				
<i>Abelmoschus esculentus</i> , okra		6		6
<i>Cyamopsis tetragonoloba</i> , cluster bean	1			1
<i>Raphanus sativus</i> , radish	1			1
<i>Vigna unguiculata</i> , cowpea	1			3
<i>Zea mays</i> , maize		3		

^zFive samples obtained in Rajasthan were collected by D.C. Bhandar, Central Arid Zone Research Institute, Jodhpur, Rajasthan in Gujarat.

^yOne sample could be a mixture of *C. melo* and *C. sativus*.

^xFive samples could be mixtures of *C. melo* and *C. sativus*.

^wFour samples grown in Madhya Pradesh were purchased at a market in Rajasthan.

Species	By State			Total
	Rajasthan	Madhya Pradesh	Uttar Pradesh	
<i>Cucumis melo</i>				
<i>agrestis</i>	88 ^z	73		161

<i>flexuosus</i>	7	12		19
<i>momordica</i>	194 ^z	53 ^y		247
not designated	12	8 ^x		20
<i>Cucumis sativus</i>				
land races	80	43 ^{yw}	14	138
open pollinated varieties	23	21	4	48

^zFive samples obtained in Rajasthan were collected by D.C. Bhandar, Central Arid Zone Research Institute, Jodhpur, Rajasthan in Gujarat.

^yOne sample could be a mixture of *C. melo* and *C. sativus*.

^xFour samples could be mixtures of *C. melo* and *C. sativus*.

^wFour samples grown in Madhya Pradesh were purchased at a market in Rajasthan.

Literature Cited

1. Munger, H.M. and R.W. Robinson. 1991. Nomenclature of *Cucumis melo* L. Cucurbit Genet. Coop. Rpt. 14:43-44.

Resistance to Melon Dieback in Spanish Landraces of Melon

F. Nuez, J. Esteva

Depto. Biotecnología. Universidad Politécnica de Valencia (UPV), SPAIN, Dpto. Ingeniería Aplicada. Universidad de Murcia, SPAIN

Melon dieback is a serious vine decline disease of melon in Spain (2, 3). The primary origin of this disease has been attributed to several ground fungus [*Rhizoctonia solani* (1), *Acremonium* sp. (4) and *Monosporascus cannonballus* (6)]. Melon dieback etiology associated with *Acremonium* sp. has been smartly by the Department of Plant Production of UPV (5). Inoculation with this fungus in hydroponic culture conditions produces root rot on melon seedlings (4). Nevertheless this technique seems not to be effective to carry out a screening of resistant genotypes to melon dieback. While these difficulties persist, a possible way to localize sources for resistance is to work in the field conditions.

We have evaluated the response of 47 melon accessions to melon dieback in field conditions during spring-summer season of 1992. The trials were carried out in two locations (Puzol and Romani) in Valencia (Spain). In both places plants were grown in plots where the disease had previously been noted during 1991. The majority of evaluated accessions are Spanish landraces belonging to 'Amarillo', 'Rochet', and 'Piel de Sapo' types. The number of replicate was four per each accession of the 'Amarillo' type and three for the rest of the accessions (Table 1). The number of plants per replicate was variable. It was sown in May. Affected plants during July and August were registered. The end of the trial was in the middle of August.

The following data summarize the incidence of melon dieback in this trial:

- Number of accessions with an incidence higher than 50% in both locations: 21.
- Number of accessions with an incidence higher than 50% in one location: 19.
- Number of accessions with an incidence lower than 50% in both locations: 7.

The accession No. 6 ("Cantaloupe" type) stand out among the accessions with an incidence lower than 50% in both locations (Table 1). This accession showed a consistent response although it did not escape completely to the disease. This could indicate that this accession has a notable degree of resistance to melon dieback and it could have interest as source for resistance. This should be confirmed in further trials. The accessions No. 12 and No. 18 ("Amarillo" type) could be employed as direct use materials. On the other hand the use of plots where the disease had previously been noted could be feasible as previous trial for the screening of sources for resistance, according to the results of this work.

Table 1. Accessions with an incidence lower than 50% in both locations.

		Location*							
		Puzol				Romani			
Accession	Incidence**	R1	R2	R3	R4	R1	R2	R3	R4
6	a	4	4	4	4	4	3	2	3
6	b	0	0	0	0	0	0	2	0
7	a	3	4	4	4	4	3	3	4
7	b	1	0	1	0	1	2	0	4
12	a	4	4	4	4	3	3	4	3
12	b	1	3	1	0	0	1	3	0

18	a	1	4	4	6	3	4	4	4
18	b	0	0	0	4	2	0	1	0
43	a	4	5	4	--	4	4	2	--
43	b	2	1	0	--	2	3	0	--
28	a	4	5	3	--	4	5	6	--
28	b	3	1	0	--	2	2	2	--
30	a	5	2	1	--	1	2	2	--
30	b	0	1	0	--	0	2	0	--

*R: replicate

**a: tested plants

b: affected plants

Literature Cited

1. Cebolla, V., T. Campos, and M. Garzia. 1990. *Rhizoctonia solani* Kuhn, causante del colapso del melon en el Pais Valenciano. *Actas de Horticultura* 2:150-155.
2. Esteva, J. F. Nuez, and J. Garcia-Jimenez. 1992 . Search for sources of resistance to a melon dieback disease in Spain. *Cucurbit Genetics Cooperative* 15:55-56.
3. Garcia-Jimenez, J., A. Alfaro, J. Esteva, F. Nuez, and M.T. Velazquez. 1991. Resistance to *Acremonium* sp. in Spanish landraces of melon. *Cucurbit Genetics Cooperative* 14:49-50.
4. Garcia-Jimenez, J., M.T. Velaquez and A. Alfaro. 1991. *Acremonium* sp., agente causal del copapso del melon en el Levante Espanol. *In* Moral de la Vega (Ed.), *Estudios de Fitopatologia. Actas del V Congreso Nacional de Fitopatologia*. Octubre de 1989. Badajoz (Spain): 68-72.
5. Jimenez, R. M. 1991. Enfermedades causadas por hongos en la horticultura espanola. *In* Rallo, L., F. Nuez (Eds.). *La horticultura espanola en la C.E. Sociedad Espanola de Ciencias Hortícolas*: 378-383.
6. Lobo, M. 1991. Las graves alteraciones de melonares y sandiares. *Bol. San. Veg. Plagas* 17:133-163.

Club Mildew: Working Group on Resistance of Melon to Powdery Mildew

James D. McCreight, Michel Pitrat

U.S. Department of Agriculture, Agricultural Research Service, U.S. Agricultural Research Station, 1636 East Alisal Street, Salinas, CA 93905, USA; Station d'Amelioration des Plantes Maraicheres, I.N.R.A., BP 94, 84143 Montfavet Cedex, FRANCE

Determinations of races of powdery mildew and resistance genes in melon are confusing. For instance, plants resistance to "race 2" and susceptible to "race 1" have been observed in the F₂ families crfom crosses of PI 12411F or PI 124112 with a susceptible line inoculated with Israeli isolates. This characterization was not observed using a French "race 2" isolate on the same families or on a collection of more than 400 accessions. Apparently these two "race 2" isolates do not have the same virulence genes.

At the 5th EUCARPIA Symposium on Cucurbitaceae in Skierniewice and Warsaw, Poland, in July 1992, a group of melon breeders and pathologists (Table 1) formed a working group to clarify the relationships between the genes for resistance and the powdery mildew races. This cooperative research includes three objectives: First, obtain melon lines possessing only one gene for powdery mildew resistance. Second, test these lines with different isolates of powdery mildew at different locations around the world. Third, conduct allelism tests among the powdery mildew resistance genes.

Table 1. Club Mildew members, affiliation, and country.

J.M. Alvarez INIA Zaragosa (Spain)	D. Gabillard Royal Sluis France Nimes (France)	M. Pitrat INRA Montfavet (France)
F. Bertrand VRED Nimes (France)	M.L. Gomez-Guillamon CSIC Malaga (Spain)	C. Robledo VRED Nimes (France)
J.L. Bourguet Graines Gutier Eyuragues (France)	J.D. McCreight USDA, ARS Salinas, Calif. (USA)	C.E. Thomas USDA, ARS Charleston, S.C. (USA)
E. Floris Univ. Politecnica Huesca (Spain)	P. Mas INRA MONTfavet (France)	J.A. Tores CSIC Malaga (Spain)

Downy Mildew (*Pseudoperonospora cubensis* B & C) Resistance in Melon (*Cucumis melo* L.)

R.G. Somkuwar and T. A. More

Division of Vegetable Crops, Indian Agricultural Research Institute, New Delhi - 110012, INDIA

Downy mildew has become a major threat to commercial cucurbits including melon and cucumber in India (5). In the sub-tropical parts during the premonsoon rains and in other humid and high rainfall areas, it is a devastating disease. In several trials conducted during rainy season by Dr. T. A. More at Delhi he noted almost 100 percent losses in cucumber and melon (4). He also observed that 'Phoot' or snapmelon (*Cucumis melo* var. *momordica*), a non-dessert type of Indian origin and FM 1 of U.S. origin hold resistance to downy mildew. Hence, efforts were made to screen advanced progenies (F₆ and F₇) of 'Phoot' (R) x Monoecious-4 (S) and FM 1 (R) Pusa Madhuras (S) crosses for resistance to downy mildew thrice under the natural epiphytotic conditions (Table 1) as per the procedure of Bonnet and Blancard (1) and also to study their performance in two replicated yield trials.

Results indicated that lines VRM 1-3, 5-10, 7-12, and 16-5 # B-1 of FM 1 x Pusa Madhuras cross and only one line VRM 31-1-2 of 'Phoot' x Monoecious-4 cross were highly resistant to downy mildew (Table 1). Combining these results with two replicated yield trials, two lines namely VRM 5-10 and 31-1-2 showed promise for downy mildew resistance and high productivity. These lines have attained homozygosity for major horticulturally important characters and now are in F₉ generation.

Detailed study on inheritance of downy mildew resistance under artificial inoculation conditions in three resistant x susceptible crosses using six generations - P₁, P₂, F₁, F₂, BC 1 and BC 2 (2, 3) was conducted. Inheritance of downy mildew resistance is governed by two dominant genes in 'Phoot' x Monoecious-3 and 'Phoot' x Pusa Madhuras crosses and two recessive genes are responsible for the same in 'Phoot' x Lucknow Safeda cross. Duplicate type of epistasis was evidenced in all the crosses.

Breeding lines	F ₆	F ₇		Average
	Rainy season - 1990	Summer season - 1991	Rainy season - 1991	
VRM-1-3	NT	12.5	20.8	16.7
VRM-5-10	8.3	31.3	42.9	27.5
VRM-7-12	12.5	33.9	21.9	22.8
VRM-16-5 # B-1	5.6	17.5	NT	11.6
VRM-31-1-2	7.1	22.3	21.9	17.8
Phoot (R)-Check	3.1	9.4	17.9	10.1
Monoecious-4(S)-Check	71	62.5	NT	67.1
Total no. of lines tested	33	18	10	---

R = Resistant, S = Susceptible, NT = Not tested

*All these lines are in F₉ generation at the present

Literature Cited

- Bonnet, A. and D. Blancard. 1987. Resistance of radish (*Raphanus sativus* L. to downy mildew *Peronospora*

- parasitica*. Cruciferae Newsletter 12:98-99.
2. Hayman, B.I. 1958. The separation of epistatic from additive and dominance variation in generation means. II. *Genetica* 31:133-146.
 3. Hayman, B.I. 1960. The separation of epistatic from additive and dominance variation in generation means. II. *Genetica* 31:133-146.
 4. More, T.A. 1989. Phoot (*Cucumis melo* var. *momordica*) resistant to downy mildew. Personal communication.
 5. Palti, J. 1974. The significance of pronounced divergence in the distribution of *Pseudoperonospora cubensis* on its crop hosts. *Phytoparasitica* 2:109-115.
 6. Sambandam, C. N., V.A. Parthasarthy, and S. Anthony Raj. 1979. Reaction of certain forms of *Cucumis* to downy mildew. *AUARA* 7/8:149-150.

Gene for Resistance to Fusarium Wilt Race 1 in Oriental Pickling Melon

Harry S. Paris, Ron Cohen, Yael Danin-Poleg, and Shoshanna Schreiber

Department of Vegetable Crops, Agricultural Research Organization, Neve Ya'ar Experiment Station, P.O. Haifa, ISRAEL

Several races of *Fusarium oxysporum* f. *melonis* afflict melons. The dominant genes *Fom-1*, carried by 'Doublon', and *Fom-3*, carried by 'Perlita', confer resistance to races 0 and 2, whereas the dominant gene *Fom-2*, carried by 'CM 17187', confers resistance to races 0 and 1. 'Charentais T' is susceptible to races 0, 1, and 2 (2, 3).

Resistance to Fusarium wilt races 0 and 1 was found in a number of Oriental pickling melon (*Cucumis melo* L. var. *conomon* Makino) accessions and breeding lines, including 'Freeman's Cucumber' (R. Cohen and H. Herson, unpublished observations). The objective of this investigation was to determine the mode of inheritance of resistance and identify the gene(s) responsible. For this purpose, 'Freeman's Cucumber' (seed sample kindly provided by H. M. Munger, Cornell Univ., U.S.A.) was crossed with 'Charentais T.' and 'CM 17187' (seed samples kindly provided by G. Risser, Station d'Amelioration des Plantes Maraicheres, France) to obtain filial, backcross, and testcross generations. The plants used for crossing had resulted from one or two generations of self-pollination of the original samples. Inoculation of melon seedlings with Fusarium wilt race 1 was described previously (1); identification of the pathogen as race 1 was confirmed by using 'Hemed' (seed sample obtained from Hazera' Co., Israel), which is resistant to races 0 and 2, as a control. The results presented in Table 1 are based on inoculations made in May 1991 and in November of the following year.

All plants of 'Charentais T' and 'Hemed' were, as expected, susceptible to Fusarium wilt race 1 (Table 1). All plants of 'CM 17187' and 'Freeman's Cucumber' were resistant. All plants of all possible F₁s between 'Charentais T', 'CM 17187', and 'Freeman's Cucumber' were also resistant. Therefore, the resistance to race 1 in 'Freeman's Cucumber' is dominant.

The F₂ of 'Charentais T' and 'Freeman's Cucumber', with the former the female parent, fit the expected 3:1 ratio of a single dominant gene conferring resistance. Although the reciprocal cross did not fit this ratio, having an excess of susceptible individuals, the overall F₂ population of this cross had a reasonable fit. In addition, the backcross to the susceptible 'Charentais T' fit reasonably well the expected 1:1 ratio for a single dominant gene conferring resistance. Therefore, it would appear that 'Freeman's Cucumber', like 'CM 17187', carries a single gene for resistance to Fusarium wilt race 1. All plants of the backcross to 'Freeman's Cucumber' were, as expected, resistant.

No susceptible individuals out of 176 were found in the F₂ of the cross of the two resistant accessions. Likewise, no susceptible individuals were found in the testcross for allelism: 'Charentais T' crossed with the F₁ of the two resistant accessions. As stated above, 'Freeman's Cucumber' was found to be resistant to race 0. These results and observations suggest that 'Freeman's Cucumber' carries the same gene for resistance as 'CM 17187', *Fom-2*.

Generation	Description	Number of plants Expected			ratio	X ²	P
		Total	R	S			
P ₁	Charentais T	48	0	48			
P ₂	CM 17187	17	17	0			
P ₃	Freeman's Cucumber	52	52	0			
P ₄	Hemed	81	0	81			

F ₁	P ₁ x P ₂	32	32	0			
F ₁	P ₂ x P ₁	40	40	0			
F ₁	P ₁ x P ₃	56	56	0			
F ₁	P ₃ x P ₁	49	49	0			
F ₂	(P ₁ x P ₃) ^(X)	127	96	31	3:1	0.024	0.87
F ₂	(P ₃ x P ₁) ^(X)	87	53	34	3:1	9.199	<0.01
F ₂	Total	214	149	65	3:1	2.472	0.12
BC ₁	P ₁ x (P ₁ x P ₃)	220	100	120	1:1	1.818	0.18
BC ₁	P ₃ x (P ₃ x P ₁)	104	104	0			
F ₁	P ₂ x P ₃	16	16	0			
F ₁	P ₃ x P ₂	13	13	0			
F ₂	(P ₂ x P ₃) ^(X)	40	40	0			
F ₂	(P ₃ x P ₂) ^(X)	136	136	0			
Test	P ₁ x (P ₃ x P ₂)	224	224	0			

Literature Cited

1. Cohen, R. T. Katan, J. Katan, and R. Cohn. 1989. Occurrence of *Fusarium oxysporum* F. sp. *melonis* race 1, 2 on muskmelon in Israel. *Phytoparasitica* 17:319-322
2. Risser, G., Z. Banihashemi, and D.W. Davis. 1976. A proposed nomenclature of *Fusarium oxysporum* f. sp. *melonis*. *Phytopathology* 66:1105-1106.
3. Zink, F.W. and Gubler. 1985. Inheritance of resistance in muskmelon to Fusarium wilt. *J. Amer. Soc. Hort. Sci.* 110:600-604.

Contribution No. 3701-E from the Agricultural Research Organization, Bet Dagan, Israel.

Breeding and Development of Cucumber Green Mottle Mosaic Virus (CGMMV) Resistant Lines in Melon (*Cucumis melo* L.)

T.A. More, A. Varma, V.S. Seshadri, R.G. Somkuwar, and L. Rajamony

Division of Vegetable Crops, Indian Agricultural Research Institute, New Delhi - 110012, INDIA

In sub-tropical parts of India around Delhi, CGMMV has been found more common and dangerous than any other virus on melon (5, 6). A research project on breeding for CGMMV resistance in melon was started in 1984. A extensive research carried out enabled to identify the 'Phoot' or snap-melon (*C. melo* var. *momordica*), 'Kachri' (both non-dessert Indian types) and FM 1 as resistant to CGMMV (1, 3, 4). Back inoculation and transmission electron microscopic studies confirmed that they possess symptomless carrier nature of CGMMV resistance. 'Phoot' and FM 1 were utilized to incorporate the CGMMV resistance into Monoecious 4 (M4) and Pusa Modjuras (PM) respectively (Fig. 1).

The crosses 'Phoot' x M4 and FM 1 x PM were advanced to F2 by selfing. Progeny of the two crosses were studied independently. In F2 and F3 generations, sibbing between the selected CGMMV resistant plants was done in order to pool the resistant genes, keeping in view the polygenic recessive nature of inheritance of CGMMV resistance (2). From F4 to F8 generations selected plants were selfed to advance the generations. From F2 to F5 screening for CGMMV resistance was done in two stages. The seedling at cotyledonary stage was inoculated with pure isolate of CGMMV under artificial inoculation conditions in insect-proof nethouse (in January-February) and selected resistant plants were transplanted in the field in the first week of March. They were again screened for CGMMV resistance under natural epiphytotic conditions and simultaneously selection was made for better horticultural characters and selected plants were sibbed (in F2 and F3) or selfed (in F4 to F5). From F6 to F8, generations were advanced by selfing. In F6 and F7 generations screening for CGMMV resistance was done under natural epiphytotic conditions. During F6 and F7 generations, performance for better horticultural characters was evaluated in the replicated yield trials. In F8, the level of CGMMV resistance was ascertained by employing the direct antigen coating ELISA technique after artificial inoculation with pure isolate of CGMMV.

Finally lines VRM 5-10, 29-1, 31-1, 31-2, 42-4, and 43-6 were selected from F5 to F10 which are highly resistant or resistant to CGMMV and possess better horticultural characters of economic important (Table 1).

Fig. 1. Breeding procedure, screening technique for CGMMV resistance and yield trials.

'PHOOT' x M4			FM1 x PM	
(R)	S)		(R)	(S)
F1			F1	
(X)			(X)	
F2		Artificial screening - field screening - selection for better horticultural traits.	F2	
#			#	
F3		- do -	F3	
#			#	
F4		- do -	F4	
(X)			(X)	
F5		- do -	F5	
(X)			(X)	

F6	Field screening and yield trial	F6
(X)		(X)
F7	- do -	F7
(X)		(X)
F8	Artificial inoculation with CGMMV-screening by direct antigen coating ELISA technique. Selection of lines having CGMMV resistance and better horticultural traits.	F8

(X) = selfed, # - Sibbed

Artificial inoculation was done with pure isolate of CGMMV.

Table 1. Evaluation of melon cultigens to CGMMV.

Cultigens ^z	CGMMV reaction to PDI (%) ^y		Reaction ^{z, x}	Yield/plant (kg) ^w	T.S.S. (%) ^v
	Field	Inoculation			
VRM-5-10	11.5	16.1	HR	1.165	9.2
VRM-29-1	12.3	20.9	HR	3.020	6.9
VRM-31-1-2	14.7	8.8	R	3.475	6.7
VRM-42-4	21.1	26.3	HR	1.880	7.7
VRM-43-6	33.5	2.5	R	1.626	7.4
Phoot (R)-Check	21.9	13.5	R	2.135	5.2
FM 1 (R)-Check	0.0	16.0	NT	1.090	8.9
Pusa Madhuras (S)-Check	67.5	62.6	HS	1.085	9.5
Monoecious-4 (S)-Check	80.1	81.6	HS	1.155	7.1

^zR = resistant, S = susceptible, HS = highly susceptible, NT = not tested.

^yPDI - present disease index. Field results of avg. of 1 yr and inoculation results are from avg. of 3 separate screenings.

^xDirect antigen coating ELISA technique was employed after artificial inoculation with pure isolate of CGMMV.

^wAverage of 2 replicated yield trials.

^vT.S.S. = total soluble solids.

Literature Cited

1. More, T.A. and A. Varma. 1991. Breeding for virus resistance in muskmelon - (*Cucumis melo* L.). Golden Jubilee Symp. on Genetic Research and Education - Current Trends and the Next Fifty Years, New Delhi, India. February 12 - 15, 1991. Abstract I : 184-185.
2. Rajamony, L., T.A. More, and V.S. Seshadri. 1990. Inheritance of resistance to cucumber green mottle mosaic virus in muskmelon (*Cucumis melo* L.). *euphytica* 47:93-97.
3. Rajamony, L., T.A. More, V.S. Seshadri, and A. Varma. 1987/ resistance to cucumber green mottle mosaic virus (CGMMV) in muskmelon. *Cucurbit Genet. Coop. Rpt.* 10:58-59.
4. Rajamony, L., T.A. More, V.S. Seshadri and A. Varma. 1990. Reaction of muskmelon collections to cucumber green mottle mosaic virus. *Phytopath. Z.* 129:237-244.
5. Raychaudhuri, M. and A. Varma. 1975. Virus disease of cucurbits in Delhi. *Proc. 62nd Indian Sci. Congr., Part III*, p. 74.
6. Raychaudhuri, M. and A. Varma. 1978. Mosaic disease of muskmelon caused by a minor variant of cucumber green mottle mosaic virus. *Phytopath. Z.* 93:120-125.

The Role of Viruses in Sudden Wilt of Melons in New York

Henry M. Munger

Department of Plant Breeding, Cornell University, Ithica, NY 14853, USA

It is now 50 years since I saw melons near Geneva, N.Y. wilt and die just as the first fruits were ripening. This was in a field in which W.D. Enzie was breeding for improved horticultural features and was typical of what came to be called "sudden wilt". Over the years since then, horticulturists and plant pathologists studying the problem in New York have implicated low soil temperatures, fusarium, pythium, and cucumber mosaic virus (CMV). However, the disorder has been so difficult to duplicate experimentally that there has been no consensus about the cause except that it is probably complex and can involve more than one agent.

In occasional years when sudden wilt is severe, differences have appeared in variety trials and breeding pots which offer some clues. The most frequent observation has been the superior survival of breeding lines and hybrids selected for cucumber mosaic resistance (CMR). On the other hand, there have so often failed to show superiority that other factors must have been more important or the level of the CMR has been inadequate.

One contributing factor may be powdery mildew. Over the past 30 years there have been three occasions when susceptible lines wilted while their PMR counterparts remained turgid. Even though mildew did not appear to be severe, temperature likewise reduces water uptake but by itself has not been shown to cause the wilting and death so often observed.

There have been two recent incidents in which viral resistance made a great difference in the survival of melons late in the season. In 1990, progenies segregating for WMV resistance (WMR), and with three backcrosses to Topmark, Honeydew, TAM Uvalde, and Cornell CPM339, were inoculated as seedlings with WMV and set in the field along with the susceptible recurrent parents. During pollination, the superiority of resistant segregates in all four backcrosses over the corresponding recurrent parents was evident, but it was also clear that CMV had come into the planting because CPM330, which has a moderate level of resistance, was the only recurrent parent to set fruit. Then in early September, as fruit was about half developed, the CPM330 plants collapsed and died, as did most of the WMR segregates except for those with backcrosses to CPM339. In short with both CMV and WMV present, resistance to either virus kept plants alive long enough to set fruit, and in most cases viable seed, but only those with resistance to both produced edible fruit.

In 1992, resistance to papaya ringspot virus (PRR) made a large difference in survival in a planting of diverse melons which we hoped would remain disease-free. One purpose was to compare fruit type of PR resistant Cornell PPM339 and its hybrids with its recurrent parent PM339 and its hybrids. PM339 has a low level of cucumber mosaic resistance and is also the recurrent parent of CPM339 mentioned above. Soon after transplanting, a few plants with virus symptoms were removed from the field, and PRV and CMV were both found to be present. Spread did not appear to be rapid through the flowering period, but plants began to die as fruit developed slowly in the cold wet season. In September the superior survival of PPM339 and its hybrids as compared with their near-isogenic counterparts became evident and Table 1 gives the proportion of live plants just before frost.

Table 1. Proportion of live plants on September 22.

Row. No.	Identity	-PPR	+PPR
68, 69	PM339	2/24	
70,71	PPM339		15/24
607	PM339 X MR324	0/6	
608	PPM339 X MR325		3/6

615	PM339 X TAM Uvalde		1/6	
616	PM339 X TAM Uvalde			2/6
638	PM339 X Gulfcoast		1/6	
639	PPM339 X Gulfcoast			3/6
		Total	4/42	23/42

It is doubtful that anyone would have suspected PRV as a cause of death had it not been for the isogenic comparisons. While we did not follow symptoms closely in this case, we have often seen in the past that when CMV infection occurs late in the season, there may be only a few days between the appearance of typical mosaic symptoms on the vine tips and their death, leaving little or no indication of the cause.

Screening of Melons for Sweetpotato Whitefly Resistance: 1992

James D. McCreight

U.S. Department of Agriculture, Agricultural Research Service, U.S. Agricultural Research Station, 1636 East Alisal Street Salinas, CA 93905, USA

Sweetpotato whitefly, *Bemisia tabaci* Genn. (SPWF) B strain virtually destroyed the Fall 1991 melon crop in the lower desert valleys of Arizona and California (4). SPWF-B does not appear to be an important vector of lettuce infectious yellows virus (LIYV, 1). In 1991, 17 of 150 PIs from India appeared to have some level of resistance to SPWF-B (3). In 1992, these 17 PIs were retested for SPWF-B resistance along with 108 previously untested PIs from India plus 27 standard varieties, breeding lines, and F₁, F₂ and backcross families from crosses among LIYV and SPWF-B resistant and susceptible parents.

A field test to evaluate SPWF-B resistance was planted on August 5 at the USDA, Irrigated Desert Research Station, Brawley, California. Plots were planted on 80 inch centers and consisted of five two-plant hills spaced 30 inches apart; there were two replications. Plots were treated once with Thiodan shortly after emergence for flea beetle control. The test was evaluated on a plot basis for four weeks (September 2 and 3) and eight weeks (September 30) later for number of live plants, plant size, plant condition, yellowing, leaf necrosis (burn) and flowering.

The SPWF-B population at Brawley during the test period was high. Four weeks after planting (September 2-3), mean number of plants per plot, and plant size and condition varied among the entries (Table 1). Because of little foliar yellowing was observed, yellowing data are not presented. A few entries had begun to flower, and as expected none of the entries was large enough to completely cover the bed. Plant condition was in general very good at this time. After an additional four weeks (8 weeks after planting) the plots had deteriorated. Many plants died; the proportion of plants surviving ranged from zero for 41 entries to 85% for one entry (F₁ Freeman Cucumber x Snakemelon); nine other entries had survival rates \geq 50% (PI 271329, PI 370021, PI 381775, PI 277281, BC Snakemelon (Freeman Cucumber x Snakemelon), BC Freeman Cucumber (Freeman Cucumber x Snakemelon), F₂ Freeman Cucumber x Snakemelon, PI 145594, PI 210542). Despite the higher survival rates, these lines were comparable to lines with lower (<50%) survival rates for plant size and condition. It is, however, significant to note that Snakemelon had a 42.1% survival rate and that F₁, F₂ and BC families from crosses of Snakemelon with Freeman Cucumber comprised four of the 10 entries with survival rate \leq 50%. Snakemelon, and BC Snakemelon (Freeman Cucumber x Snakemelon) were among the best five entries for plant condition after eight weeks. Five entries had a mean plant condition. 3.0 (PI 212895, Snakemelon, BC Snakemelon (Freeman Cucumber x Snakemelon), F₂ (90625 x Top Mark), PI 370021). Snakemelon was previously identified as a potential source of LIYV resistance and has very large vines in the presence of high levels of LIYV (2).

In 1991, a single plant of PI 164825 appeared to be resistant to SPWF-B feeding. Self-pollinated seed from that individual was not obtained, but crosses were made with WMR 29 and I5. PI 164825 was retested in this test but none of the plants were as outstanding as the individual observed in 1991. The two F₁ families made with PI 164825 were not remarkable (Table 1). PI 145594 was the only PI identified in 1991 as a potential source of SPWF-B resistance that performed well in this test (Table 1). Five PIs tested for the first time in 1992 were among the best entries for survival, and plant size and condition after eight weeks. These will be retested.

None of the entries in this test was superior for survival, and plant size and condition. Snakemelon from the Middle East and families from crosses of Snakemelon were among the best entries in 1992 for plant survival, size and condition after eight weeks exposure in the field to SPWF-B. Five previously untested wild melon introductions from India (PI 210542, PI 271329, 277281, 370021, PI 38775) were identified in 1992 as having some potential as sources for SPWF-B resistance. The better lines for these characteristics will be examined in subsequent field tests exposed to natural populations of SPWF-B. They should also be compared with Top Mark and other varieties in controlled greenhouse tests for response to SPWF-B feeding and reproduction.

Table 1. Mean number per plot^Z, plant size^Y and condition^X on September 2-3, 1992 and changes in these parameters on September 30 in response to sweetpotato whitefly feeding.

Group and Entry	September 2-3			September 30		
	Live	Size	Cond	% Surv	? Size	? Cond.
Varieties and Parental Lines						
90625	9	2	9	16.7	1	-8
AR 5	8	1.5	8	37.5	-0.5	-7
Freeman Cucumber	9	2	8.5	16.7	-1	-7.5
PMR Honeydew	8	1.5	7	06.2	-0.5	-6
Snakemelon	9.5	2.5	6.5	42.1	2	-3
Top Mark	7.5	1.5	7	40	-0.5	-6
WMR 29	9.5	1	6	21.0	0	-4
Snakemelon x Freeman cucumber						
F ₁ FC x Snake	10	2	7.5	85	1	-4.5
F ₂ FC x Snake	20	2.5	6	75	1.5	-4.5
BC FC (FC x Snake)	10	2	7	70	0.5	-5
BC Snake (FC x Snake)	8.5	3	6.5	58.8	2	-3
90625 x AR 5						
F ₁ 90625 x AR 5	10.5	2.5	7.5	0	--	--
F ₂ 90625 x AR 5	9	1.8	6.8	27.8	-0.5	-4.2
BC (90625 x AR 5) x AR	8	2	7	31.2	-0.5	-6
90625 x Top Mark						
F ₁ 90625 x Top Mark	9.8	2	7.5	12.8	-0.3	-6.5
F ₁ Top Mark x 90625	9.5	2.2	7.8	32.2	0	-6.2
F ₂ 90625 x Top Mark	9.5	1.8	7	5.3	3.2	-3
F ₂ Top Mark x 90625	9.5	2.5	7.5	31.6	2.5	-5
BC Top Mark x 28377	9.5	2	6.5	13.2	-1	-5.5

BC 28380 x Top Mark	10.2	2	7
BC 28380 x 90625	9.2	2.2	7.8
BC Top Mark x 28380	10	2	8
90625 x PMR Honeydew			
F ₁ PMR ND x 90625	9.5	2.5	8
F ₂ PMR ND x 90625	7.8	1.8	7.5
PI 164825 crosses			
F ₁ PI 164825 x WMR 29	1	2	7.5
F ₁ PI 164825 x 15	7.5	2.5	6.5
Plant Introductions tested in 1991			
123682	7	1	5.5
124092	8.5	2	5.5
124103	10	1.5	5.5
124106	6.5	1.5	5
124109	9.5	2.5	6.5
124431	7	2	5
124440	10	2	5
124447	6.5	2	5
124550	9.5	2.5	7
145594	9	2	9
164749	8	2	7.5
164825	8.5	2	5.5
165515	9	2	5.5
165525	7.5	1	6.5
179669	9.5	2	7
179671	6	2	6

9.8	-1	-6
13.5	0.8	-6.4
17.5	-1	-5.8
15.8	0.5	-7
25.8	-0.5	-5.5
0	--	-7
26.7	-0.5	-5.5
7.1	0	-4.5
0	--	--
5	-0.5	-4.5
15.4	1.5	-3
0	--	--
0	--	--
0	--	--
0	--	--
10.5	1.5	-6
77.8	1	-6.5
12.5	0	-6.5
11.8	1	-4.5
5.6	3	-4.5
13.3	0	-5.5
26.3	1	-6
0	--	--

179890	10	2	6.5	0	--	--
Plant Introductions tested in 1992						
179901	9.5	2	6.5	10.5	0	-5.5
179902	8.5	2	6.5	17.6	1	-5
179903	10	3	6.5	45	0	-5.5
179904	8	2	6.5	0	--	--
179905	9.5	2	6.5	10.5	0	-4.5
179906	8.5	2	6.5	11.8	0	-5.5
179909	8	3	8	25	0	-6.1
179911	6	2	6.5	25	0.5	-5.5
179913	10	2.5	6.5	0	--	--
179915	10	2.5	7.5	35	-1.5	-6
179916	10	2	5.5	5	-1	-4.5
179916	10	3	7.5	5.5	1.5	-6
179917	10	3	7.5	5.5	1.5	-6
179919	10.5	2	6.5	9.5	1	-5.5
179920	9.5	2.5	6.5	15.9	0.5	-5.5
179922	6.5	1	8.5	15.4	0	-6.5
179923	6	1	9	0	--	--
180280	6	1	6.5	8.3	0	-3.5
180281	7.5	1.5	7	13.3	-0.5	-5
180283	5	1	7.5	30	0	-5.5
181051	9.5	2	6	10.5	2	-3
182937	6.5	2	8.5	0	--	--
182938	9.5	2	6.5	0	--	--

182943	9	1.5	5	5.6	-0.5	-4
182949	8.5	2	6	17.6	0	5
182952	10	2	6	5	-1	-5
182959	9	2	6.5	22.2	-0.5	-5
182964	5	1.5	9	20	0.5	-8
183025	7.5	2	5.5	6.7	-1	-4.5
183029	9.5	2	6	0	--	--
183031	10	2	5.5	0	--	--
183032	8	1.5	5	0	--	--
183036	10	1.5	5	0	--	
183040	10	1	6	11.8	0	-5
183045	8.5	2	6	0	--	--
183048	7	1	4	8.3	0	-3
183049	6	2.5	6	5.9	0.5	-5
183051	8.5	2	6.5	15.4	-1	-5.5
183052	6.5	2	8	9.1	1	-6
183054	9	2	7	27.8	2	-4.5
183055	9.5	2.5	7.5	0	--	--
183128	10	2.5	6	10	-1.5	-5
183303	10	3.5	7	30	-0.5	-6
183305	9.5	2	6	10.5	2	-3
183307	3.5	1	8.5	0	--	--
183311	9.5	2.5	6	5.3	-1.5	-4
183397	10	2.5	7	15	0.5	
183444	9.5	1.5	6.5	26.3	1	

210077	10.5	1.5	7		14.3	0.5	-6
210541	8	12	8.5		81	1	-5
210542	8.5	2.5	7		2	0.5	-6
212803	8.5	2	7		25	1.5	-6
212895	8.5	2.5	6		29.4	1	-3.5
213247	8.5	2	6		5.9	1	-5
214048	8.5	2.5	7		0	--	--
214154	9.5	2	6.5		0	--	--
214318	9.5	2	7		0	--	--
216030	9	2	6		22.2	-0.5	-4.5
217599	8.5	2	6.5		11.8	0	-5
271329	10	2	8		50	1.5	-7
271335	10	2	6.5		10	0	-5.5
275633	8.5	2	6.5		17.6	-1	-5
277281	9.5	2.5	8		57.9	2	-6.5
277282	8	2	6.5		6.2	-1	-5.5
277283	9	2	6		0	--	--
277284	9	2	5.5		0	--	--
279367	9	2	6		0	--	--
288330	9	2	6.5		16.7	0	-5
288333	10	2	7		20	1	-5.5
302445	10	3	6.5		25	1	-5
302446	9	1.5	7		0	--	--
323316	8.5	2.5	7		5.9	-1.5	-5
358942	8.5	2	7		17.6	0	-5.5

370020	9	2	6	0	--	--
370021	10	3	7.5	50	0.5	-3.5
381758	9.5	2.5	6.5	0	--	--
381760	7.5	2	7	26.7	-1	-6
381761	9.5	1.5	6	10.6	-0.5	-4
381762	8.5	1.5	6	11.8	0	-5
381765	7	1.5	6.5	21.4	0	-5.5
381766	8.5	2	7	0	--	--
381770	7.5	2	6.5	0	--	--
381771	7.5	1.5	5.5	6.7	-0.5	-4.5
381772	9.5	1.5	6	10.5	-0.5	-4
381773	10	1	5.5	10	0	-4.5
381774	9.5	2	7	5.3	-1	-6
381775	8.5	2	9	52.9	1.5	-7
381773	10.5	2	7	19.0	0.5	-6
381781	10	2.5	7.5	5	-0.5	-6.5
381782	10	1.5	6.5	10	-0.5	-3.5
381783	9	2	7.5	38.9	0.5	-6.5
381784	8.5	2	6	5.9	0	-4
381786	10	1.5	6.5	15	-0.5	-5
381787	10	2	7.5	20	-0.5	-6.5
381788	9.5	2	7	21.0	-1	-6
381790	7.5	2	8	6.7	--	-7
381792	10	2	7	0	--	--
381794	10	1.5	6	10	-0.5	-5

381797	7	2	8		42.8	0.5	-6.5
381800	9.5	2	8		0	--	--
381802	10	2	6		30	1	-4.5
381803	8	1.5	5.5		6.2	-0.5	-2.5
401731	8	1.5	7		12.5	-0.5	-6
431581	10	1.5	6.5		0	--	--
504523	5.5	1.5	6.5		0	--	--
504524	8.5	1.5	4.5		0	--	--
504525	10	2.5	6.5		5	1.5	-5.5
504526	7.5	2.5	6.5		0	--	--
504527	9.5	2	5.5		0	--	--

^zNumber of live plants is the mean; Surv is the Percentage plants remaining on September 30.

^y Size was rated on 1 (very small, only a few true leaves) to 9 (completely covering the bed) scale.

^x Condition was rated on a 1 (dead) to 9 (vigorous, flowers) scale.

Literature Cited

1. Cohen, S. J.E. Duffus, and H.Y. Liu. 1991. A new *Bemisia tabaci* biotype in the Southwestern United States and its role in silverleaf of squash and transmission of lettuce infectious yellows virus. *Phytopathology* 82:86-90.
2. McCreight, J.D. 1991. Potential sources of resistance to lettuce infectious yellows in melon. *Cucurbit Genet. Coop. Rpt.* 14:51-52.
3. McCreight, J.D. 1992. Preliminary screening of melons for sweetpotato whitefly resistance. *Cucurbit Genet. Coop. Rpt.* 15:59-61.
4. Perring, T.M., A. Cooper, D.J. Kazmer, C. Shields, and J. Shields. 1991. New strain of sweetpotato whitefly invades California vegetables. *California Agriculture* 45 (6):10-12.

Genotypic Control of Regeneration Potential in *Cucumis melo*

J. Jain and T.A. More

Division of Vegetable Crops, Indian Agricultural Research Institute, New Delhi - 110012, INDIA

Present study was undertaken to outline the genotypic control of regeneration potential in *C. melo*. Three sex forms viz. andromonoecious (Pusa Madhuras), gynoeceious (GM-5D, GM-7 and GM-6E-7) and monoecious M_4 were utilized in regeneration studies to find out the genotypic control of regeneration potential of different sex forms.

Materials and Methods. In vitro regeneration studies were carried out in Pusa Madhuras, GM-5D, GM-6C-4, GM-7, GM-6E-7, and M_4 . Both epicotyl and cotyledonary explants were used. Pre-standardized de-differentiation medium of MS + 0.5 mg/l (MB) and differentiation medium of MS + 1.0 mg/l IAA and 5.0 mg/l Kinetin (MIK) were used following the usual procedure of regeneration (1). Callus obtained on MB medium was transferred to MIK medium after 2-3 weeks, for differentiation of shoot buds.

Results. Among the various sex forms of *C. melo*, andromonoecious sex form (Pusa Madhuras) has been found to be highly responsive to regeneraiton on pre-standardized de-differentiation and differentiation medium (Table 1). The four gynoeceious forms showed no response to callus formation on MB medium. Similarly neither any direct regeneration nor callus induced regeneraiton was observed on MIK medium. In monoecious sex form (M_4) cotyledonary leaf explants were found to be equally responsive to differentiation medium than explants of Pusa Madhuras. In epicotyl explants of M_4 , 23.82 (± 10.80) per cent callus induced regeneration was observed in comparison to 66.66 (± 7.21) per cent callus induced regeneration observed in Pusa Madhuras epicotyl explants.

Discussion. Various sex forms of *Cucumis melo* differ in twow genes *A* & *G*. The andromonoecious form (Pusa Madhuras) is homozygous recessive for *a* and homozygous dominant for *G* whereas the monoecious form M_4 has a genotype of *A-G* (3, 4, 5). The gynoeceious sex form is designated as *A-g g* plus modifiers at *g* locus (5). The high regeneration potential observed in Pusa Madhuras (1) and M_4 (2) may be governed by dominant *G*. Dominant *A* has a repressing effect on regeneration.

Table 1. Regeneration response of various sex forms of *C. melo*¹

Various sex forms	Explant	Explants forming callus (%)	Explants with shoots (%)	No response (%)
Andromonoecious	Epicot.	33.33 \pm 7.21	66.66 \pm 7.21	0
(Pusa Madhuras)	Cot. leaf	83.33 \pm 28.86	16.66 \pm 8.86	0
Monoecious (M_4)	Epicot.	75.18 \pm 12.30	23.82 \pm 10.80	0.98 \pm 0.67
	Cot leaf	56.10 \pm 20.30	30.49 \pm 6.76	13.39 \pm 4.20
Gynoeceious				
GM-5D	Epico.	0	0	100
	Cot. leaf	0	0	100
GM-6C-4	Epicot.	0	0	100

	Cot. leaf	0	0	100
GM-7	Epicot.	0	0	100
	Cot. leaf	0	0	100
GM-6E-7	Epicot.	0	0	100
	Cot. leaf	0	0	100

¹Epicotyl and cotyledonary explants were incubated on de-differentiation medium (MS + 0.5 mg/1 BA) for 2-3 weeks before transfer to shoot bud differentiation medium (MS + 1.0 mg/1 IAA and 5.0 mg/1 Kinetin).

Literature Cited

1. Jain, J. and T.A. More. 1992. In vitro regeneration in *Cucumis melo* cv. Pusa Madhuras. Cucurbit Genetics Cooperative Report No. 15, pp.62-64.
2. Jain, J. and T.A. More. 1992. Regeneration potential of various accessions of muskmelon, *Cucumis melo* L. Fifth Eucarpia Cucurbitaceae Symposium, held at Poland.
3. Rosa, J.T. 1928. The inheritance of flower types in *Cucumis* and *Citrullus*. Hilgardia 3:233.
4. Vegetable Breeding. 1988. Vol. I, pp. 3-11. Ed., Dr. Kalloo. CRC Press Inc., Florida.
5. Vegetable Research - A resume (1956-90). 1991. pp 17-25. Eds., H.S. Gill, T.A. re, N. Singh, and A. S. Kataria. Ankur Publishers; New Delhi.

Genetic Parthenocarpy in *Cucurbita Pepo* L.

R.W. Robinson

Horticultural Sciences Department, New York State Agricultural Experiment Station, Geneva, New York 14456

Fruit set of summer squash can be very poor when conditions are unfavorable for pollination. Many F₁ hybrid cultivars have been bred to have a high ratio of female to male flowers and they may not have sufficient male flowers for good pollination at all times. The problem of poor fruit set is especially acute early in the season when prices are highest, but production may be limited because the low temperature, long day length, and high light intensity of that season promotes female sex expression and reduces male flower formation. Genetic parthenocarpy would be of value in situations where male flower production is insufficient and could also increase yield when there are insufficient bees for good pollination, when bee activity is restricted by wet weather, and when pollinating insects are kept away from squash blossoms by plastic tunnels or other row coverings. If row covers did not need to be opened or removed for pollination, they could be left on squash plants longer to improve growth and early yield by increasing the temperature around the squash plants, and also reduce losses from insects and insect-transmitted virus diseases.

Genetic differences in *Cucurbita pepo* for ability to set parthenocarpic fruit were reported by den Nijs and Veldhuyzen den Zanten (1982). They found that early production of parthenocarpic fruit ranged from none for some cultivars to 42% for 'DG-4'. Previously, Rylski (1972) noted that 'Zucchini Elite' had a stronger parthenocarpic characteristic than 'Bushy White'.

For the past three years we have been selecting for genetic parthenocarpy in the summer squash breeding program at Geneva, NY. Several CMV resistant breeding lines, including 82-138 and 82-141, compared favorably with 'DG-4' for parthenocarpic fruit set. Preliminary tests this past season, however, indicated that there may already be commercially available cultivars with even better parthenocarpic fruit set than the best of our breeding lines selected for parthenocarpy.

Thirty-three *C. pepo* cultivars and breeding lines were grown in the field and the first female flowers to develop on each plant were enclosed in paper bags before anthesis to prevent insect pollination. The number of female flowers closed for each cultivar or line is given in Table 1. Only fruit that developed to a marketable stage were recorded as being set parthenocarpically, and fruit that had some ovary enlargement, then turned brown and ceased development were not regarded as set. The fruit that were considered parthenocarpic had normal size and shape, and at maturity contained no seeds.

Two thirds of the entries in the trial set parthenocarpic fruit, the best being 'Chefini Hybrid' with 82% fruit set (Table 1). It is not known if the unusually cool and wet season of 1992 in New York was a factor in the extraordinary set of parthenocarpic fruit by some cultivars.

Many summer squash cultivars with the same fruit color have a similar gene background and may have similar incidence of parthenocarpy. It was previously reported (1, 2) that zucchini-type cultivars have better parthenocarpic fruit set than yellow- or white-fruited cultivars. That was generally true in our test, but there were exceptions. Green-fruited 'Ambassador', for example, was relatively poor in parthenocarpic fruit set in our trial and also in Holland (1), and none of the five female flowers we tested for 'Dark Green Zucchini' set fruit, but the yellow-fruited cultivar Gold Strike had excellent set of parthenocarpic fruit. Results from our breeding program and cultivar tests indicate that it should be possible to breed parthenocarpic squash of different fruit colors and types.

Table 1. Parthenocarpic fruit set of squash cultivars and lines.

Cultivar or Line	Type	No. female flowers tested	% Parthenocarpy
Chefini Hybrid	zucchini	11	82
Gold Strike	yellow straightneck	8	75
Black Beauty	zucchini	7	71

Black Magic	zucchini	9	67
NY-82-138	zucchini	2	50
NY-92-728	zucchini	7	43
Green Magic	zucchini	12	42
NY-82-141	zucchini	8	38
Gold Slice	yellow straightneck	6	33
Cocozelle	striped straightneck	7	29
Goldie Hybrid	yellow crookneck	17	29
President	zucchini	14	29
Black Jack	zucchini	16	25
Gold Rush	precocious yellow straightneck	10	20
Caserta	striped straightneck	17	18
Golden Girl	yellow straightneck	12	17
Onyx	zucchini	7	14
Senator	zucchini	14	14
NY-92-727	zucchini	14	11
Hyrific	yellow straightneck	9	11
White Scallop	white scallop	9	10
Ambassador	zucchini	10	0
Yellow Crookneck	yellow crookneck	8	0
Gold Bar	yellow straightneck	10	0
Honey Boat	Delicata winter squash	4	0
Multi Pik	precocious yellow straightneck	12	0
Peter Pan	green scalop	11	0
Royal Acorn	green acorn winter squash	5	0
Scallopini	gren scalop	9	0
Slendegold	yellow straightneck	6	0
Early Prolific Straightneck	yellow straightneck	5	0
Dark Green Zucchini	zucchini	5	0
NY-92-730	zucchini	6	0

Literature Cited

1. Nijs, A.P.M. den and N.J.D. Veldhuyzen van Zanten. 1982. Parthenocarpic fruit set in glasshouse grown zucchini squash. *Cucurbit Genet. Coop. Rpt.* 5:44-45.
2. Rylski, I. 1974. Effects of season on parthenocarpic and fertilized summer squash (*Cucumis pepo* L.). *Expl. Agric.* 10:39-44.

Productivity of Naked Seed Squash, *Cucurbita pepo* L.

Warid A. Warid, Jaime J. Martinex, and Juan M. Loaiza

Department of Agriculture - DAG, University of Sonora, Hermosillo, Sonora, Mexico

The naked-seed squash has been reported by Curtis (2, 3). The seed lacks the lignified seed coat. Whitaker (5) referred to studies on this mutatin by several austrian and German investigators. Abak et al. (1) started a breeding project in 1985 for seed production in Turkey. Percent oil in naked seed strains of the present study is 35.4\$ as reported by Farag (4).

Seeds of naked seed Kurbis squash were kindly provided by Dr. L. Tanasch of University of Austria in 1988. S₀ plants were grown in Hermosillo at the University of Sonoro Experiment Station from seeds planted in April 1991 at a 0.5 X 3.0 m spacing. Fourteen plants were self pollinated in June 1991 and the fruits were harvested in August. S₁ seeds were extracted and segregation for seed type, normal or naked, was observed. All but two of the selfed plants produced fruits having naked seeds. S₁ naked seeds were planted in September 1991 under isolation, and harvested in December for observation. The plants exhibited normal growth, and produced open-pollinated fruits having naked seeds, confirming homozygosity of the naked trait.

Another planting of S₁ seed was made in March 1991 under isolationl. The growth of S₁ plants was normal and they partially recovered from viral infection by spraying with Dithane M-45. Vigorous, spiny vines with profuse male and female flowering, and normal set of fruits by open pollinationw ere observed. Mature fruits were harvested in July 1992. Data were recorded on 74 fruit from 31 plants. These were cut cross-wise and seeds were extracted and dried at ambient temperature. Fruits were morphologically described, and data were recorded on number and weight of fruits per plant. Fruit and seed traits were correlated on a single fruit basis. Simple and partial correlations were computed for fruit weight, number and weight of seeds, and seed size (mg/seed).

Results. There was no segregation for seed type. Fruits of all S₁ plants contained naked seeds. However, the green color of the exposed cotyledons (inner seed coat) varied in intensity from medium green to dark green. The mature fruit was round, oblate in shape. The pedicel end was spiny, solid, rather soft and ridged with 8 angles and inconspicuous flaring. The color of fruit skin (rind) was orange and splashed green. The rind was smooth and not hard. The flesh had a creamy color and fibrous texture. The placenta was gelatinous and orange.

The number of fruits per plants varied from 1 to 9 and averaged 3.3. Total fruit weight per plant ranged from 0.47 to 12.67 kg with a mean weight of 4.64 kg. Among the 74 individual fruits, the range of fruit weight was 0.47 to 3.17 kg, and the average was 1.55 kg. Number of seeds per fruit ranged from 16 to 393 and averaged 136. The average seed weight per fruit was 18.2 g and ranged from 1.4 to 64.1 g. Seed size averaged 134 mg and varied from 46 to 223 mg.

A strong association existed between number and weight of seeds per fruit, as evidenced from simple and partial correlation values (0.921 - 0.925) that were positive and highly significant (Tables 1 and 2). The values of the coefficient of determination indicated that 85 to 86 percent of the variation in seed weight can be ascribed to number of seeds. Seed size showed a strong, positive association with fruit weight ($r = 0.488$). Twenty-four percent of the variation in seed size can be attributed to fruit size.

Table 1. Simple correlation between fruit characters of naked seed squash (n = 74 fruits).

Characters correlated	Coefficient of	
	Correlation r	Determination r ²
Fruit wt. vs Number of seeds	0.207 ns	0.043

Fruit wt. vs Weight of seeds	0.346 **	0.120
Number of seeds vs Weight of seeds	0.921 **	0.848
Fruit wt. vs Seed size	0.488 **	0.238

Table 2. Partial correlation between fruit characters of naked seed squash (n = 74 fruits).

Characters correlated	Coefficient of	
	Correlation r	Determination r ²
Fruit wt. vs Number of seeds (fixing Wt. of seeds)	-0.306 *	0.094
Fruit wt. vs Wt. of seeds (fixing Number of seeds)	0.408 **	0.166
Number of seeds vs Wt. of seeds (fixing Fruit wt.)	0.925 **	0.856

Literature Cited

1. ABak, K. M. Sakin and S. Sakin-karakullukcu. 1990. Improvement of pumpkin for naked seeds. XXIII Intern. Hort. Congr. Abstr. 3074.
2. Curtis, L.C. 1946. The possibilities of using species of perennial cucurbits as source of vegetable fats and proteins. Chemurgic Digest 5:221-224
3. Curtis, L.C. 1948. The use of naked seed in *Cucurbita pepo* as a source of high quality liquid vegetable fat, as a high analysis protein, as a new confection, and as a sandwich spread. Proc. Amer. Soc. Hort. Sci. 52:403-406.
4. Farag, R.S. 1992. Personal communication. Lab. Chemical Analyses, College of Agriculture, University of Cairo, Egype.
5. Whitaker, T.W. 1960. Breeding squash and pumpkins. Handbuch der Pflanzenzuchtung 6:331-350.

The Chromoplast Ultrastructure of Two Isogenic Lines of *Cucurbita pepo* fruits at Different Developmental Stages

H.T. Lim, C.D. Boyer

Department of Horticulture, Kangwon National University, Chuncheon, 200-701, The Republic of Korea; Department of Horticulture, The Pennsylvania State University, University Park, PA 16802.

Introduction: Many agriculturally important processes take place in the various developmental states of plastids. Chloroplasts are extensively studied due to their photosynthetic process. Chromoplasts, the principal carotenoid-bearing organelles in higher plants, are the major sites of vitamin A precursors for fruits and vegetables. In most of higher plants, chloroplasts are much alike in morphology, but chromoplasts are found to be in a great diversity of shapes and sizes, especially during plastid differentiation.

The understanding of plastid differentiation and interconversion is of fundamental biological interest. We chose *Cucurbita pepo*, for they display a fascinating array of colors during fruit development. The biochemical aspects of carotenoid accumulation in the chromoplast of different squash varieties were studied by Schafer et al. (6). We reported restriction site and genetic maps of chloroplast DNA of *Cucurbita pepo*, which corresponds to 153 kb in size (3).

Changes in plastid ultrastructure from two genotypes affecting fruit pigment ($YY B^+B^+$: fruits is initially green, then yellow) and $YY BB$: precociously yellow) of squash with isogenic backgrounds were compared at different stages of fruit development.

Materials and Methods. Fruits of two isogenic lines of squash (c.v. Early Prolific) were harvested at five developmental stages: 2 days postpollination, at pollination, 3 days postpollination, 10 days postpollination, and 20 days postpollination. We also used other squash varieties which display distinct green and yellow colors representing chloroplasts and chromoplasts, respectively. Small pieces of squash fruits were cut from the pericarp of fruits at five progressive stages, fixed in 3% glutaraldehyde in a 0.06 M phosphate buffer (pH 7.3) in ice for 3 hrs, and post-fixed it in 2% OsO₄ for 2 hrs. The samples were dehydrated through a graded series of acetone and propylene oxide and then embedded in Epon resin. Thin sections were stained with 6% uranyl acetate for 90 min and lead citrate for 6 min (4) and observed with an electron microscope (Phillips) at 80 V.

Results and Discussions: Longitudinal section of chloroplast of green zucchini fruits showed the granal and stomatal thylakoid network similar to that observed in leaf tissue of Early Prolific squash.

Appearance of plastoglobuli and internal membrane structure in prochromoplasts paralleled the disappearance of thylakoid materials in chloroplasts in $YY B^+B^+$ fruits. Therefore, we assume that thylakoid materials in chloroplasts may be used for the formation of internal membranes and plastoglobuli during chromoplast differentiation (Fig. 1. a-e).

According to their structure and pigment composition of $YY B^+B^+$ fruits and of tomato fruits (2), we suggest that chromoplasts undergo structural changes from the membranous chromoplasts to the globular chromoplasts (Fig. 1. d-e).

Chromoplasts in fruits were indeed derived directly from proplastids in precocious ($YY BB$) fruits (Fig. 1.f-h). It was reported that chromoplasts of the precociously pigmented yellow portion of bicolor ornamental gourd were differentiated directly from proplastids (1).

Very similar line structure of plastids was observed in mature fruits of both genotypes. Based on previous hypothesis (6) and our results, we suggest that *B* and *Y* genes control the timing of chromoplast or chloroplast appearance rather than structure or content of plastids.

Fig. 1.a. Proplastid of $YY B^+B^+$ ovary 2 days before anthesis. 1.b. Chloroplast of $YY B^+B^+$ fruit at anthesis. 1.c. Prochromoplast from $YY B^+B^+$ fruit 3 days postanthesis. 1.d. Cross section of membraneous chromoplast from $YY B^+B^+$ fruits 10 days postanthesis. 1.e. Longitudinal sectioned globular chromoplast of $YY B^+B^+$ fruit 20 days postanthesis. 1.f. Prochromoplast of $YY BB$ fruit at anthesis. 1.g. Membraneous chromoplast of $YY BB$ fruit 10 days postanthesis. 1.h. Globular chromoplast of $YY BB$ fruit 20 days postanthesis.




Fig.1.a. Proplastid of $YY B^+B^+$ ovary 2 days before anthesis. 1.b. Chloroplast of $YY B^+B^+$ fruit at anthesis. 1.c. Prochromoplast of $YY B^+B^+$ fruit 3 days postanthesis. 1.d. Cross section of membraneous chromoplast from $YY B^+B^+$ fruits 10 days postanthesis. 1.e. Longitudinal sectioned globular chromoplast of $YY B^+B^+$ fruit 20 days postanthesis. 1.f. Prochromoplast of $YY BB$ fruit at anthesis. 1.g. Membraneous chromoplast of $YY BB$ fruit 10 days postanthesis. 1.h. Globular chromoplast of $YY BB$ fruit 20 days postanthesis.

Literature Cited

1. Devide, Z. 1970. Ultrastructurll changes of plastid in ripe fruits of *Cucurbita pepo* cv. ovofera. Acta Bot. Croat. 29:57-62.
2. Harris, W.M. and A.R. Spurr. 1969 Chromoplasts of tomato fruits. 1. Ultrastructure of low-pigment and high-beta mutants. Carotene analyses. Amer J. Bot. 56 (4) : 369-379.
3. Lim, H-T., I. Gounaris, R.C. Hardison, and C.D. Boyer. 1990/ Restriction site and Genetic Map of *Cucurbita pepo* chloroplast DNA. Chrent Genetics. 18:273-275.
4. Reynolds, E.S. 1963. The use of lead citrate at high pH as an electron opaque stain in electron microscopy. J. Cell Biol. 17: 208-212.
5. Schaffer, A.A., C.D. Boyer, and T. Gianfagna. 1984. Genetic control of plastid carotenoids and transformation in the skin of *Cucurbita pepo* L. fruit. Theor. Appl. Genet. 68: 493-501.




Fig.1.a. Proplastid of $YY B^+B^+$ ovary 2 days before anthesis. 1.b. Chloroplast of $YY B^+B^+$ fruit at anthesis. 1.c. Prochromoplast of $YY B^+B^+$ fruit 3 days postanthesis. 1.d. Cross section of membraneous chromoplast from $YY B^+B^+$ fruits 10 days postanthesis. 1.e. Longitudinal sectioned globular chromoplast of $YY B^+B^+$ fruit 20 days postanthesis. 1.f. Prochromoplast of $YY BB$ fruit at anthesis. 1.g. Membraneous chromoplast of $YY BB$ fruit 10 days postanthesis. 1.h. Globular chromoplast of $YY BB$ fruit 20 days postanthesis.

A Case of Extremely Low Expressivity of Gene *B2* in *C. moschata*.

Oved Shifriss

21 Walter Avenue, Highland Park, NJ 08904

An F_2 population of the cross IL-B x NJ-B, *C. moschata*, was grown in the fall of 1989 in Naples, Florida. According to present interpretation (1, 2), IL-B is $B1 + B1 + B2B2$ and NJ-B, $B1B1 B2 + B2 +$. Each of the *B* genes conditions chlorophyll deficiency.

Both parents exhibit precocious depletion of chlorophyll in ovaries, prior to anthesis, and their fruits are uniformly pigmented. But the stems of IL-B are persistently golden, devoid of chlorophyll, and the stems of NJ-B are persistently green.

One of the F_2 segregates, plant 807-34, produced bicolor fruits and persistently green stems. Although its ovaries were uniformly green, at pre-anthesis stages, they later on turned into bicolor fruits at some point during the post-anthesis stages. The fact that the bicolor design was first visible sometime during the post-anthesis stages indicated, according to present interpretation, that 807-34 carried gene *B2*.

An F_3 progeny was obtained from 807-34 and this progeny was grown in spring of 1990. It consisted of 110 plants, 106 of which were classified with a high degree of confidence. An analysis of this F_3 progeny suggested that 807-34 was $B1 + B1 + B2B2+$ (see test 5, table 2, in reference 1).

Of the 106 F_3 plants, 56 were bicolor. All the 56 plants produced fruits that turned bicolor sometime during the post-anthesis stages, and were therefore considered to be $B2B2+$. However, these bicolor individuals greatly varied in size of the chlorophyll deficient area in their fruits and in extent of chlorophyll deficiency in their stems.

A program of inbreeding was initiated in order to clarify the basis for the wide range of variation in the above F_3 progeny. As a result, two phenotypically distinct lines evolved through selection and inbreeding two different bicolor-fruited individuals: (a) plants that produced fruits in which chlorophyll deficiency affected about one-half of their surface, and (b) plants that produced fruits in which chlorophyll deficiency was confined to a few golden spots or a relatively small area (about 1.0 cm²) at the proximal end. The data for case (b) are presented and interpreted here. Three phenotypic classes were identified and the symbols for these phenotypes are described in the following.

Symbols for three phenotypic classes (Table 1):

I. PDC-UP, PDC-SL = precocious depletion of chlorophyll uniformly affects ovaries prior to anthesis; fruits are uniformly golden; chlorophyll deficiency clearly affects the stems early in plant development, the basal portion of the plant being golden, but later on the expressivity level of chlorophyll deficiency is *low*, fluctuating between golden and green.

Unlike the PDC-UO, PDC-S phenotype of IL-B whose petioles are golden (tests 10-11, Table 1, in reference 1), the petioles of PDC-UO, PDC-SL are green.

II. GOT-VL, GS = ovaries are uniformly green at pre-anthesis stages; chlorophyll deficiency becomes visible late in post-anthesis stages and is confined to a small area, often inconspicuous, at the proximal end of the fruit (late *turning*); stems are persistently green.

This phenotype represents a *very low* expressivity level of chlorophyll deficiency. and significantly, some fruits of a given plant may be uniformly green.

III. GO, GS = ovaries, fruits and stems are persistently green.

It should be emphasized that unless a large number of fruits is carefully examined in each plant, a potentially phenotype of class II may be mistakenly in class III. Furthermore, there is evidence suggesting that the environment can transform some or all plants of class II into class III (e.g., see the offspring of plant 3 in the present Table 1).

Interpretation

The analysis of the data in Table 2 does not disagree with either the 3:1 or the 13:3 hypothesis. However, the 3:1 hypothesis is favored with some elaboration. The present contention is (i) that the PDU-UO, PDC-SL phenotype of this line is $B2B2$; (ii) that the GOT-0VL, GS phenotype is $B2B2+$; (iii) that the GO, GS class consists of either $B2+B2$ exclusively or a combination of $B2+B2+$ and $B2B2+$ in various proportions; (iv) that the genetic background of this line is homozygous for certain regulators of $B2$ (v) that said regulators attenuate or partially suppress the action of $B2$; (vi) that the degree of suppression is especially high in the heterozygotes; and (vii) that the environment can further intensity this suppression. Consequently, in some environments, $B2$ can operate as recessive rather than as dominant or codominant gene.

If the 13:3 hypothesis were applicable, one would have expected (a) that the GOT-VL, GS phenotype was heterozygous for both $B2$ and its partial suppressor, and (b) that the offspring of such double heterozygote would have consisted of some bicolor individuals in which chlorophyll deficiency effected a large portion of the fruit. No such individuals were observed in the present experiment.

Table 1. Offspring of self-pollinated plants that manifested extremely low expressivity of gene $B2$ (see text)

Parents ^z			Offspring					
			Phenotypic Classes			I	II	III
Plants ^y	Pedigree	Growing Season	Growing Season	PDC-UO	PDC-SL			
1	807-34-110 ^x	Spring, '90	Fall '90	10		18	12	40
			Fall '92	8		11	13	32
2	807-34-110-2	Fall, '90	Spring '91	4		8	8	20
			Fall, '92	2		12	18	32
3	807-34-110-2-10	Spring, '91	Fall, '91	12		11	8	31
			Spring, '92	4		0	27	31
4	807-34-110-2-10-13	Fall, '91	Spring '92	7		0	25	32
			Fall '92	4		8	11	23
5	807-34-110-2-10-30	Fall '91	Spring, '92	4		0	28	32
			Fall, '92	3		0	4	7
Grand Total				58		68	154	280

^z All parental plants exhibited extremely low expressivity (phenotype GOT-VL, GS) of gene $B2$ (see text).

^y Two samples of each parental plant were tested. the two samples were drawn from the same seed packet.

^x F₃ segregate of the cross IL-B (phenotype PDC-UO, PDC-S) X NJ-B (phenotype PDC-UO, GS). See reference 1.

Table 2. Analysis of the data presented in Table 1.

Parental Plants		Offspring			Hypothesis #1	Hypothesis #2
		II & III	I	Total		
					3:1	13:3

				χ^2	χ^2
1	54	18	72	0.0000	1.8462
2	46	6	52	5.0256	1.7751
3	46	16	62	0.0699	2.0265
4	44	11	55	0.7333	0.0564
5	32	7	39	1.0341	0.0442
	222	58	280	6.8629	5.7484
				2.7428	0.7092
				4.1201	5.0392

		χ^2	df	P
Hypothesis #1	Deviation	2.74	1	0.05-0.10
	Heterogeneity	4.12	4	0.30-0.50
Hypothesis #2	Deviation	0.71	1	0.30-0.50
	Heterogeneity	5.04	4	0.20-0.30

Acknowledgement: I thank Rogers NK SEed Co. for enabling me to conduct this study.

Literature Cited

1. Shifriss, O. 1991. Expression of the *B* Genes in *Cucurbita*. Cucurbit Genetics Coop. 14:113-115.
2. Shifriss,). 1991. The Two *B* Genes in *Cucurbita* are Unlinked. Cucurbit Genetics Coop. 14:116-122.

Inheritance of Immature Fruit Color in *C. Moschata*

Antonio I.I. Cardoso and Paulo T. Della Vecchia, Norberto Silva

Agroflora S/A, Caixa Postal 427, 12900-000 Braganca Paulista - SP, Brazil; Departamento de Agricultura e Melhoramento Vegetal, FAC-Campus Botucatu-UNESP, 186-000 Botucatu - SP, Brazil

Bush types of *C. moschata* with field resistance to Papaya Ring Spot Virus-W were developed in Brazil, as an alternative to *C. pepo* for the production of immature fruits (1). Lines were obtained for different immature fruit color such as: *Mottled Light and Dark Green*, *Uniform Dark Green* and *Precocious Yellow*. Several studies have been reported on the inheritance of immature fruit color in *C. pepo* (2, 3, 4, 5, 6). However, no report was found in the literature on the inheritance of this trait in *C. moschata*. The present paper reports on the inheritance of immature fruit color in a cross between two *C. moschata* lines which produce Mottled Light and Dark Green (P1) and Uniform Dark Green (P2) immature fruits, respectively.

Parental lines were selfed for seven generations before being crossed to produce the F1, F2, BCP1 and BCP2 populations. Parents F1, F1, BCP1 and BCP2 populations were grown under field conditions at Sao Manuel-SP, Brazil, during the Summer season of 1992. Plants were scored for immature fruit color 3 to 5 days after the anthesis of the female flowers.

The results are presented in Table 1. Chi-square analysis revealed a good fit for a single completely dominant gene hypothesis for the Mottled Light and dark Green immature fruit color. It is proposed the use of the symbol *Mldg* for this gene in *C. moschata*.

Table 1. Inheritance of immature fruit color in *C. moschata*

Population	Number of Plants			Expected Ratio	χ^2	P
	Mottled Light and Dark Green	Uniform Dark Green	Total Plants			
P1	90	0	90	1:0		
P2	0	89	89	0:1		
F1	0	0	83	1:0		
F2	283	96	379	3:1	0.014.>	0.99
BCP1	189	0	189	1:0		
BCP2	90	95	185	1:1	0.135>	0.99

Literature Cited

1. Della Vecchia, P.T., P. Terenciano S. and A. Terenciano. 1993. Breeding bush types of *C. moschata* with field resistance to PRSV-W Cucurbit Genetics Coop. Rpt. 16:
2. Globerson, D. 1969. The inheritance of white fruit and stem colour in summer squash. *Euphytica* 18: 249-255.
3. Paris, H.S. and H. Nerson. 1986. Genes for intense pigmentation of squash. *J. Heredity* 77: 403-409.
4. Paris, H.S. 1989. List, description and interactions of the genes affecting fruit color in *C. pepo*. Cucurbit Genetics Coop. Rpt. 12:72-74.
5. Scharhuk, J. 1954. Fruit and leaf characters in summer squash. *J. Heredity* 45: 295-297.

Breeding Bush Types of *C. moschata* with Field Resistance to PRSV-W.

Paulo T. Della Vecchia, Petro Terenciano S. and A,adeu Terenciano

Agroflora S/A, Caixa Postal 427, 12900-000 Braganca Paulista-SP, Brazil

Immature fruits of *C. pepo* and *C. moschata* are used in Brazil for raw or steamed salads. the market preference is towards *C. pepo* cultivars which bear slightly tapered, cylindrical fruits. However, the available *C. pepo* cultivars in Brazil are very susceptible to Papaya Ring Spot Virus-W which makes difficult their production, particularly during the late Spring and Summer periods. Some cultivars of *C. moschata* show good field resistance to PRSV-W, but they are late, produce plants with long vines and fruits that resemble the straightneck Butternut types. The present paper relates to the development of novel types of *C. moschata* with bush grown habit, good PRSV-W field resistance, earliness and which produce slightly tapered cylindrical fruits, similar to those of *C. pepo*.

Method. The lines were obtained through pedigree and backcross selection as shown in Fig. 1 and Fig. 2. Selections for field resistance to PRSV-W were practiced in all generations, under natural field infection conditions, using spread rows and plants of *C. pepo* cv. Caserta as a standard immature squash type. Cv. Piramoita contributed the bush growth habit and PRSV-W resistance (1,2), F1 hybrid Seaulmadi the slightly tapered cylindrical fruit shape and PPI 165561 the precocious yellow fruit color (3).

Results. The main characteristics of the novel *C. moschata* breeding lines developed by the program are presented in Table 1.

Conclusions. Preliminary commercial field trials have indicated a good acceptance of the novel breeding lines among farmers in main production areas of Brazil. We believe that these novel types of *C. moschata* will be interesting to other tropical countries besides Brazil.

Table 1. Characteristics of some *C. moschata* breeding lines with bush growth habit and PRSV-W field resistance.

Characteristics	Breeding Lines/Cultivar			
	AF1075 L	AF1094 L	AF1351 L	Piramoita
Plant diameter (m)	1,50	1,20	1,50	1.80
Days for flowering	55	50	55	55
Sex of the 1st flower	female	male	female	female
Fruit length/diameter (cm)	15/7	15/5	15/7	---
Fruit colour	mottled light and dark green	uniform dark green	precocious yellow	mottled light and dark green
Fruit shape	tapered cylindr.	tapered cylindr	tapered cylindr.	long straightneck
Fruit growth ¹ (days)	5	5	6	5
Productivity ² (fruits/plant)	20-30	20-30	20-30	15-20

¹ Fruit growth - days from anthesis to the commercial harvesting size.

² Productivity - fruit/plant for a 40 days harvest period.




Fig. 1 Pedigree of the bush type, PRSV-W field resistant *C. moschata* cv. Piramoita.




Fig. 2 Pedigrees of the bush type PRSV-W field resistant *C. moschata* breeding lines AF 1075 L, AF 1094 L and AF PY1351 L.

Literature Cited

1. Costa, C.P. da 1974. Obtencao de aborinha (*Cucurbita moschata* Duch ex Poir) Menina Brasileira com habito de crescimento tipo moita e com tolerancia ao Mosaico da Melancia. Revista de Olericultura XIV: 21-22.
2. Munger, H.M, 1990. Availability and use of interspecific populations involving *Cucurbita moschata* and *C. pepo*. Cucurbit Genetics Coop. 13:49.
3. Paris, H.S., Nerson, H. and Burger, Y. 1985. Precocious PI 165561, and Precocious PI 165561 R pumpkin breeding lines. Hort Science 20 (4): 778-779.




Fig. 1 Pedigree of the bush type, PRSV-W field resistant *C. moschata* cv. Piramoita.




Fig. 2 Pedigrees of the bush type PRSV-W field resistant *C. moschata* breeding lines AF 1075 L, AF 1094 L and AF PY1351 L.

PI Num.	Origin	Field	Greenhouse		PI Num.	Origin	Field	Greenhouse
193499	Ethiopia ¹	1.0	3.0		379295	Yugoslavia ¹	0.0	0.0
201254	Mexico ¹	1.5	---		414906	India ¹	1.0	5.0
234251	Japan	1.5	4.0		438811	Mexico	1.5	2.0
249565	Thailand	1.5	5.0		482490	Zimbabwe	1.0	4.0
298036	Australia ¹	0.0	0.0		282523	Zimbabwe ¹	1.5	2.5
357916	Yugoslavia	1.5	3.0		540906	Unknown	1.5	4.0
2693456	Costa Rica	1.5	3.5		----	----	---	---

¹ Accession appears to be misclassified as *C. moschata*

Table 2. *C. moschata* accessions with a mean field rating of 0 for downy mildew.

PI Number	Origin		PI Number	Origin		PI Number	Origin
168547	Mexico		190185	Mexico ¹		196923	Mexico ¹
200736	El Salvador		201254	Mexico ¹		201471	Mexico
201473	Mexico		326184	Mexico ¹		281810	India
381815	India		438577	Guatemala		438578	Guatemala
438723	Mexico		438726	Mexico		438731	Mexico
438747	Mexico		438748	Mexico		438756	Mexico
438760	Mexico		438772	Mexico		438775	Mexico
438776	Mexico		438781	Mexico		438784	Mexico
438787	Mexico		438790	Mexico		438792	Mexico ¹
438794	Mexico		438824	Mexico		442248	Mexico
442249	Mexico		442250	Mexico		442251	Mexico
442253	Mexico		442256	Mexico		442257	Mexico
442258	Mexico		442272	Mexico		442274	Mexico
452276	Mexico		442281	Mexico		442284	Mexico
451836	Guatemala		451837	Guatemala		451845	Guatemala

¹ Accession appears to be misclassified as *C. moschata*.

Literature Cited

1. Cienfuegos-Agreda, R.E. 1991. Metodos de inoculacion para cuantificar resistencia poligenica a anublo polvoriento *Cucurbita moschata* [(Duch) ex. Poir.]. M.S. Thesis, University of Puerto Rico, Mayaguez Campus, Mayaguez, PR. 62 pp.
2. Hiiata, K. 1955. On the shape of germ tubes of Erysiphae (II). Bul. Faculty Agric Niigata Univ. 7:24-26 (In Japanese with English summary).
3. Siterly, W.R. 1978. Powdery mildews of cucurbits, p. 359-379. In: D.M. Spencer (ed). The powdery mildews, Academic Press Inc., New York.

This research is a contribution from the Agricultural Experiment Station, UIniversioty of Puerto Rico and was supported in part by the U.S. Dept. of Agriculture under CSRS Special Grant 88-34135-4661 managed by the Caribbean Basin Advisory Group (CBAG).

Leaf Silvering of Squash: A Brief Review

Harry S. Paris

Department of Vegetable Crops, Agricultural Research Organization, Neve Ya'ar Experiment Station, P.O. Haifa, Israel

Leaf silvering is an important malady of squash and pumpkins in the Middle East, Puerto Rico, the southern United States, and possibly other regions (1, 2, 3, 9, 11, 12, 13). Leaf silvering was first recognized and reported as a serious disorder of squash in Israel (1, 2, 3, 4). The symptoms were illustrated and described by Paris et al. (10) in Israel and by Simons et al. (13) and Maynard and Cantliffe (9) in Florida. The symptoms, in mild cases, are silvering in and parallel to the veins in the upper surface of the leaves; in severe cases, the entire upper leaf surface is silvered and the petioles, stems, flowers, and fruits are pale in color (10). The rate of photosynthesis is about 30% lower in completely silvered than in green leaves (5).

Leaf silvering is induced by the sweetpotato whitefly, *Bemisia tabaci* Genn., especially by the nymphs (11, 14) of what is sometimes referred to as the poinsettia or IV-90 strain (6). Efforts to find a pathogenic organism such as a virus or viroid, associated with silvering have failed, leading to the conclusion that silvering is a systematic phytotoxemia (6, 14). Silvering is exacerbated by drought stress in the broad sense (i.e. plant water deficit) and its components (low soil moisture, high temperatures, high light intensity, long days, etc.) (3, 4, 10). Chemical control of the whitefly (8) and cultural practices which reduce plant water deficit (3, 4) have been reported to reduce the severity of the disorder. Silvering was reduced in a cultivar that was less susceptible to silvering when grown on reflective mulch with full irrigation (7).

Differential susceptibility to silvering occurs among cultivar groups, cultivars, and even among different strains of the same cultivar in *Cucurbita pepo* (H.S. Paris, P.J. Stoffella, and C.A. Powell, manuscript in preparation). Whilst genetic material immune to silvering has not been found, the cocozelle and vegetable groups of *C. pepo* have been found to contain some less susceptible cultivars.

Literature Cited

1. Ayyalon, A. 1969. Growing green summer squash for export. Publ. 68/110/19, Agric. Exten. Serv., Israel Ministry of Agriculture, Tel Aviv (in Hebrew).
2. Be'eri, Y. and B. Kapuler. 1963 Growing of vegetables, Part B. Summer squash, pp. 235-240. Y.M. Margalit, ed. Sifriyyat Hassadeh Publ., Tel Aviv (in Hebrew).
3. Burger, Y., H.S. Paris., H. Nerson, Z. Karchi, and M. Edelstein. 1983. Overcoming the silvering disorder of *Cucurbita*. Cucurbit Genet. Coop. Rep. 6: 70-71.
4. Burger, Y., H.S. Paris, H. Nerson, Z. Karchi, Y. Roter, D. Lozner, M. Edelstein, and A. Govers. 1983. On the leaf silvering problem of summer squash. Hassadeh 63: 2580-2581 (in Hebrew, English abstract).
5. Burger, Y., A. Schwartz, and H.S. Paris. 1988. Physiological and anatomical features of the silvering disorder of *Cucurbita*. J. Hort. Sci. 63:6350640.
6. Cohen, S., J.E. Duffus, and H.Y. Liu. 1992. A new *Bemisia tabaci* biotype in the southwestern United States and its role in silverleaf of squash and transmission of leatuce infectious yellows virus. Phytopathology 82: 86-90.
7. Edelstein, M., H.S. Paris, R. Shir, G. Leibowitz, H. Nerson, and Y. Burger. 1991/ Effects of reflective plastic mulch on fall-grown summer squash. Hassadeh 71:868-869 (in Hebrew, English abstract).
8. Gornick, A. and N. Belizer. 1991. Prevention of silvering and virus in summer squash. Gan, sade, UMesheq: Dec. pp. 79-80 (in Hebrew).
9. Maynard, N.D. and D.J. Cantliffe. 1989. Squash silverleaf and tomato irregular ripening: new vegetable disorders in Florida. Veg. Crops Fact Sheet VC-37, Coop. Exten. Serv. Univ. Florida, Gainesville, 4 pp.
10. Paris, H.S., H. Nerson, and Y. Burger. 1987. Leaf silvering of *Cucurbita*. Can. J. Plant Sci. 67: 593-598.
11. Schuster, D.J., J.B. Kring, and J.F. Price. 1991. Association of the sweetpotato whitefly with a silverleaf disorder of squash. HortScience 26: 155-156.
12. Segarra-Carmona, A.E., J. Bird, J. Escudero, G. Fornaris-Rullan, and R.A. Franqui. 1990. Silvering of *Cucurbita moschata* (Duchesne) Poir associated with *Bemisia Tabaci* Genn. (Homoptera: Aleyrodidae) in Puerto Rico. J. Agric.

Univ. Puerto Rico 74: 477-478.

13. Simons, J.N., P. J. Stoffella, K.D. Shuler, and R.N. Raid. 1989. Silverleaf of squash in south Florida. Proc. Fla. St. Hort. Soc. 101 (1988): 397-399.
14. Yokomi, R.K., K.A. Hoelmer, and L.S. Osborne. 1990. Relationships between the sweetpotato whitefly and the squash silverleaf disorder. Phytopathology 80: 895-900.

Contribution No. 3668-E from the Agricultural Research Organization, Bet Dagan, Israel.

Inheritance of Resistance to Races 0, 1, and 2 of *Fusarium oxysporum* f. sp. *niveum* in Watermelon (*Citrullus* sp. PI 296341)

Xingping Zhang and Bill Rhodes

Horticulture Department, Clemson University, Clemson, SC 29634-0375

Martyn and Netzer (1) reported *Citrullus* sp. PI 296341-FR to be 100% resistant to races 0 and 1 of *Fusarium oxysporum* f. sp. *niveum* but segregating for resistance to race 2. We report here inheritance studies with this PI.

Materials and Methods. Seeds of parents, PI 296341-FR, New Hampshire Midget (NHM) and F₁, F₂ and BC₁ generations were surface sterilized and germinated. Eight-day seedlings with fully expanded cotyledons were used for inoculation. Roots of the seedlings were rinsed with water and then dipped in a spore suspension for 10 min. Inoculation concentration of each race of fusarium was 10 spores/ml. Inoculated plants were single planted in styrofoam trays and moved into the greenhouse. Plants were watered daily and fertilized weekly. The temperature in the greenhouse was 26 ± 1-2°C. Wilting began 5 to 7 days after inoculation. Data collected before transplanting (18 days post inoculation) and after planting (50 days after inoculation) are shown in Table 1. A set of crosses were made in the summer using resistant PI 296341 and F₁ individuals. Resistant F₁ plants were selfed or backcrossed in each case to produce F₂ and backcross progeny.

Results. The data support Martyn and Netzer's observation that the resistance genes in the resistant parent PI 296341-FR were not fixed. Inheritance to each race in this PI will be discussed separately.

Race 0 Resistance: F₁ progeny from reciprocal crosses with NHM are segregating 1:1 resistant to susceptible, indicating heterozygosity of one or more dominant genes for resistance in PI 296341. Segregation in the F₂ population from a selfed resistant F₁ plant fits a 9:7 ratio, suggesting an interaction between nonallelic genes. Segregation of the backcross population with NHM also indicates modifier(s) which overcome the dominant gene for resistance.

Race 1 Resistance: Dominance for resistance to race 1 in PI 296341 is indicated. This result is in agreement with Netzer and Weintall (2). However, the presence of a susceptible class in the F₁ indicates other modifier gene(s). The susceptible class from the testcross to the recessive parent is much higher than expected, also indicating segregation of modifier gene(s). At 18 days, the F₂ progeny segregate 3:1 R:S as expected, but the susceptible class has increased by 50 days.

Race 2 Resistance: Both resistant and susceptible individuals exist in the parent PI 296341. The cross of a resistant individual with the susceptible NUM parent results in predominant fly, but not exclusively, susceptible individuals. Thus, it is clear that resistance to race 2 is governed by at least one recessive pair of genes. If a 13:3 model is hypothesized for the F₂ generation, i.e., a dominant gene from NHM is epistatic over a recessive gene for resistance in PI 296341, then the data fit the model perfectly. If the backcross to the susceptible parent is assumed to be AaBb x aaBB, then all the backcross progeny will carry the B gene that is epistatic over a. If the few progeny scored as resistant in the F₁ and backcross populations were actually susceptible, the hypothesis that one or more recessive genes is interacting with a dominant gene is consistent with the data.

Our data suggest epistasis between resistance genes in PI 296341 and the susceptible cultivar New Hampshire Midget.

Table 1. Resistance/susceptible distribution in different families inoculated with races 0, 1 and 2 of *Fusarium oxysporum* f. sp. *niveum*.

Races of the	Families	Resistant:Susceptible			p
		18 days	50 days	Hypothesis	

Pathogen					
Race 0	PI 296341	15:0	15:0	1:0	1.00
	NHM	0:15	0:15	0:1	1.00
	296341xNHM	9:11	9:11	1:1	0.70-0.50
	NHMx296341	13:7	10:10	1:1	0.99-0.95
	(296341xNHM)NHM	9:27	4:32	0:1	0
	(296341xNHM)F ₂	70:24	52:42	9:7	0.90-0.75
Race 1	PI 296341	15:0	15:0	1:0	1.00
	NHM	0:15	0:15	0:1	1.00
	296341xNHM	16:4	15:5	3:1	0.99-0.95
	NHMx296341	17:3	16:4	3:1	0.75-0.50
	(296341xNHM)NHM	12:23	6:29	1:3	0.25-0.10
	(296341x296341)F ₂	67:28	61:34	2:1	0.75-0.50
Race 2	PI 296341	10:5	8:7	1:1	0.75-0.50
	NHM	2:13	0:15	0:1	1.00
	296341xNHM	4:14	1:17	0:1	0
	NHMx296341	3:17	2:18	0:1	0
	(NHMx296341)NHM	1:39	1:39	0:1	0
	(NHMx296341)F ₂	20:72	18:77	3:13	>0.90

Literature Cited

1. Martyn, R.D. 1991. Resistance to races 0, 1, and 2 of *Fusarium* wilt of watermelon in *Citrullus* sp. PI 296341-FR. HortScience 26(4):429-432.
2. Netzer, D. and C. Weintall. 1980. Inheritance of resistance in watermelon to race 1 of *Fusarium oxysporum* L. sp. *niveum*. Plant Dis. 64:853-854.

Male Sterile, *ms*, in Watermelon not Linked to Delayed Green, *dg* and *I-dg*

Xingping Zhang and Bill Rhodes

Department of Horticulture, Clemson University, Clemson, SC 29634-0375

The genetic male-sterile *ms*, from China (1) has been reported to be unlinked to *gms* (1) and *Sp* (2). Data are reported here on its relationship to the two genes that determine delayed green - *dg* and *I-dg*.

To transfer *dg* (and its epistatic gene, *I-dg*) into the male-sterile line, the following cross was made in 1990: *ms ms Dg Dg i-dg i-dg* and *Ms Ms dg dg I-dg I-dg*.

The fertile, normal green F_1 (*Ms ms Dg dg I-dg i-dg*) was selfed in the spring of 1991. The F_2 population was tested in the greenhouse in the fall of 1991. An F_2 population of 111 individuals was selected first for delayed green and then male sterility. The F_2 seedling population consisted of 88 normal: 23 delayed green (13:3 ratio, $X^2=0.283$). The 23 delayed green plants were grown out for fertility testing and seed. Seven (7) male-sterile recombinants were found among the 23 delayed green progeny, consistent with a 61:3 segregation of male sterile, delayed green ($X^2=0.651$).

Finally, the delayed green, male fertile class was obtained by subtracting the 7 delayed green, male sterile from the 23 delayed green total in the population. Thus, there were 16 delayed green, male fertile individuals in the F_2 , consistent with a 55:9 segregation of delayed green, male sterile ($X^2=0.011$).

These preliminary data suggest that *ms* is assorting independently.

Literature Cited

1. Murdock, B.A., N.H. Ferguson and B.B. Rhodes. 1991. Segregation data suggest male sterility genes *gms* and *ms* in watermelon are not in the same linkage group. Cucurbit Genetics Coop. Rpt. 14:90-91.
2. Murdock, B.A. and N.H. Ferguson. 1992. Segregation data suggest male sterility gene *ms* and the spotted gene *Sp* in watermelon are not in the same linkage group. Cucurbit Genetics Coop Rpt. 15:72-73.
3. Rhodes, B.B. 1986. Genes affecting foliage color in watermelon. J Hered. 77:134-135.
4. Zhang, Xingping and Ming Wang. 1990. A genetic male-sterile (*ms*) in watermelon from China. Cucurbit Genetics Coop. Rpt. 13:45-46.

Fertility Shift in an *In Vitro* Regenerated Male-sterile Line in Watermelon

Xingping Zhang and Bill Rhodes

Department of Horticulture, Box 340375, Clemson University, Clemson, SC 29634-0375

Regeneration of a male-sterile line was begun in November of 1991. Cotyledons from 20-day old fruit (*ms ms* x *Ms ms*) were inoculated on 10 μ M BA regeneration medium (Adelberg and Rhodes (1)). Regenerated shoot buds were subcultured on 10 μ M BA during January and early February, 1992. The shoot buds were inadvertently subjected to a severe heat shock (over 37 ° C) for a period of 2 days during the subculture when an air conditioner motor failed. The subcultured shoot buds were transferred to 5 μ M IBA for rooting in late February, 1992. Rooted plantlets were transplanted into the greenhouse in early March, 1992. A total of 249 plantlets were transplanted into the greenhouse, and 230 plantlets survived (92.4% survival).

Two hundred regenerated plants were transplanted into the field at Edisto Research and Education Center, Blackville. A great range of pollen sizes were observed in this population - from 1673 μ M² to 5419 μ M². Only two putative tetraploids, identified by morphology, were observed. No visible somaclonal variation was obtained in the first generation other than the fertility segregation ratio. A 1:1 fertile:sterile ratio was expected, but 146 fertile:54 sterile were actually found.

Why did the fertile:sterile ratio change so drastically? One possibility was that the *Ms ms* tissue generated more plants than the *ms ms* tissue without any effect of temperature. The other possibility is that, under the conditions of this regeneration trial, the *ms* gene reverted to *Ms*. Perhaps the heat shock was responsible for this reversion. The mechanism responsible for this effect is being investigated. Nearly one hundred individuals were either selfed or sib crossed. If some of the fertile plants are transiently fertile because of temperature, their selfed progeny may be sterile.

Literature Cited

1. Adelberg, J.W. and B.B. Rhodes. 1989. Micropropagation from zygotic tissues of watermelon. Proc. Cucurbitaceae 89: Evaluation and Enhancement of Cucurbit Germplas,. pp.110-112.

Protoplast Isolation and Culture of Watermelon Cotyledons

Caiping Ma, CX.P. Zhang, B.B. Rhodes and J.W. Adelberg

Horticulture Dept., Box 340375, Clemson University, Clemson, SC 29634

Protoplast culture and subsequent regeneration could lead to more rapid genetic progress with watermelon. Experiments were conducted to define conditions for protoplast culture of cotyledons.

Effects of Mannitol, Cellulase and Pectinase on Protoplast Isolation from Watermelon Cotyledons.

A factorial experiment was performed to determine effects of mannitol cellulase and pectinase on protoplast isolation. Three levels of mannitol (0.3, 0.4 and 0.5 M), three levels (0.5, 1.0, 1.5, and 2.0%) of cellulase (E.C. 3.2.1.4, C-9422, Sigma) and two levels (0 and 0.4%) of pectinase (E.C. 3.2.1.15, P-4625, Sigma) were dissolved in the following buffer: 0.1 M glycine, 10 mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 0.7 mM KH_2PO_4 at pH 5.7. Enzyme solutions were sterilized by filtration. Cotyledons of 1 week old seedlings grown in the greenhouse were used as the explant source. Cotyledons were first washed with tap water one hour, then surface-disinfected with 10% Clorox for 10 min., then rinsed four times in sterile distilled waer. Lower epidermis of the cotyledon was peeled off with fine curved forceps, cut into 0.4 x 0.4 mm segments and put into a sterile enzyme solution. The tissue was incubated in darkness at 28 ° C for 4-6 hr. At the end of digestion, the cotyledon tissue and enzyme solution was shaken gently by hand to release the protoplasts. Enzyme solution and protoplasts were filtered through a nylon mesh (45 μ M pore size), and filtrate was transferred to a centrifuge tube and spun at 200 RCF (Microcentrifuge Model 5-9A, Fisher Scientific) for 3-5 min. Protoplasts were resuspended in 0.5 ml of buffer solution with different concentrations of mannitol but without cellulase and pectinase. Density of protoplasts was determined with a hemocytometer, and viability of the protoplasts was examined after staining with 0.1% phenosafranin.

The optimal concentration of enzyme solution for isolating watermelon cotyledon protoplasts was 0.4 M mannitol, 1.5% cellulase and 0.4% pectinase (Table 1). Large standard errors indicate that lal three factors are important for watermelon protoplast isolation, Sorbitol at 0.4 M was an equally effective osmoticum for isolation of osmoticum in trials not presented here.

Effect of Temperature and Time during the Enzyme Digestion on Protoplast Denisty and Viability.

Materials and procedures were the same as in the first trial. Enzyme solution consisted of 0.4 M mannitol, 1% cellulase, 0.4% pectinase, 0.1 M glycine, 10 mM $\text{CaCl}_2 \cdot \text{H}_2\text{O}$ and 0.7 mM $\text{K}_2\text{H}_2\text{PO}_4$ at pH 5.7. Three temperatures and digestion times were tested. Every treatment had three replications. In order to obtain a high intact protoplast density in a short time, we selected a suitable digestion period of 4-6 h at 29 °C (Table 2).

Effect of Calcium Concentration in Enzyme Solution

It is known that Ca^{+2} can stabilize the protoplast membrane and enhance the survival of protoplasts. Four levels of Ca^{+2} in enzyme solution were evaluated: 0, 10, 15 and 20 mM. Enzyme solution consisted of 1% cellulase, 0.4% pectinase and 0.4 M mannitol at pH 5.7. In this experiment, immature cotyledons were used to isolate protoplasts. First, proximal portion of cotyledons were dissected from the embryo, cut in half lengthwise and transferred to MS medium with 10 mM BA, 3% sucrose, and 0.7% agar at pH 5.7. Tjese were maintained at 25 ± 2 ° C, with a 16 h light period produced by cool white fluorescent lamps at $30 \mu \text{mol m}^{-1} \text{sec}^{-2}$. After 10-14 days of culture, green, expanding cotyledons were cut into 0.1 cm strips and put into sterile enzyme solution kept in darkness at 26 ° C for 4h. Every treatment had three replicates. The results are shown in Table 3.

Protoplast viability was highest at 20 mM Ca^{+2} although the protoplast density was lower at 15 mM Ca^{+2} . Unless a high

density is desired, 20 mM Ca is suitable for protoplast isolation (Table 3).⁷

Effect of Cotyledon Source on Protoplast Isolation

Because harvested watermelon seeds are often difficult to decontaminate prior to tissue culture, protoplast isolation for cotyledons of harvested (dried) seed and cotyledons of seed from the initial fruit were compared.

1. Fresh seed from mature fruit. Mature fruit harvested in the greenhouse were surface disinfected with 95% ethanol, and seeds were removed aseptically. Embryos were removed from seed coats. Cotyledon explants were dissected from embryos and transferred to MS salts medium with 0.7% agar, 3% sucrose, 10 mM BA, 100 mg l⁻¹ myo-inositol, 2 mg l⁻¹ glycine, 0.2 mg l⁻¹ nicotinic acid at pH 5.7. Tissue was maintained at 25 ± 2 ° C under light. After 8-15 days the cotyledons were cut into 0.1 cm strips and put into the enzyme solution.
2. Aseptic seed from mature fruit were dried in the laminar flow hood and kept in sterile bottles. Seeds were soaked in sterile distilled water before use. After 24-48 h. embryos were removed from seed coats and cultured as above.

In a second test with dried seeds, aseptic mature seeds were soaked in sterile distilled water and shaken on a platform shaker at 150 rpm for 24 hours. Seeds were transferred to a Magenta GA₇ vessel with a layer of wet filter paper and kept in the dark at 30 ° C. After 2 days, embryos with ca. 1 cm radicle were removed from seed coats and transferred to the same medium used in test 1, but with out BA. Cotyledons 8-14 days old were cut into 0.1 cm strips and put into the enzyme solution.

Cotyledons of fresh seeds from mature fruit were the best explant source for isolation and culture of protoplasts. The density of the protoplasts was 1.0-1.3 x 10⁵ ml⁻¹ and protoplast viability was 68-74%.

Protoplasts were cultured on liquid and agar gelled B5 medium (Gambourg et al. 1968) with different levels of growth regulators. Protoplasts became more oval after 2-3 days, indicating the synthesis cell wall. The protoplasts survived 15 days. However, protoplasts from dried seeds died in 2-3 days although the density and viability was not significantly different from protoplasts of fresh seeds. Results indicate that cotyledons from fresh seeds of mature fruit should be explants for protoplast isolation.

Table 1. Effect of 3 levels of mannitol, 2 levels of pectinase and 4 levels of cellulase on watermelon cotyledon protoplast density and variability.

	Ingredient	Density (x 10⁴)	Variability (%)
Mannitol			
	0.3 M	14.5 ± 12.4 ^a	46.9 ± 28.3
	0.4 M	14.7 ± 13.8	58.6 ± 15.6
	0.5 M	13.7 ± 18.7	51.6 ± 22.3
Pectinase^b			
	none	2.6 ± 2.2	55.2 ± 15.2
	0.4	16.0 ± 8.7	47.0 ± 25.6
Cellulase^c			
	0.5%	15.2 ± 14.90	48.5 ± 16.0
	1.0%	14.5 ± 15.3	47.1 ± 21.0
	1.5%	16.6 ± 14.1	60.2 ± 10.4
	2.0%	10.9 ± 6.2	48.7 ± 30.4

^aStandard errors.

^{b,c}Pectinase was E.C. 3.2.1.15,P4625; Cellulase, E.C.3.2.1.4,C-9422, both Sigma.

Table 2. Effect of temperature and digestion time on watermelon cotyledon protoplast density and variability.

	20		25.5		29	
	D(x 10 ⁴) ^a	V	D(x10 ⁴)	V	D(x10 ⁴)	V
digestion (h)						
4	8.2	85.4	12.3	92.4	19.4	88.2
6	12.0	87.8	16.0	83.8	20.0	82.2
8	12.8	78.9	22.3	75.8	27.4	72.5
10	19.0	69.0	26.2	77.9	23.8	61.0

^aD = density; V = viability

Table 3. Effect of Ca⁺² in enzyme solution on density and viability of protoplasts from immature watermelon cotyledons.

	Ca ⁺² (mM) ^a			
	0	10	15	20
Density (x10 ⁴)	1.9	3.9	2.5	2.2
Viability (5)	64.7	80.8	88.7	94.1

^aCaCl₂* 2H₂O

Literature Cited

1. Gambourg, O.L., r.A. Miller, and K. Ojima. 1968. Plant cell cultures. I. Nutrient requirements of suspension cultures of soybean root cells. Experimental Cell Research 50:151-158.

Crossability between *Momordica charantia* and *Momordica dioica*

M. Abdul Vahab and K.V. Peter

College of Horticulture, Vellanikkara, India - 680 654

Bittergourd (*Momordica charantia*) is an important fruit vegetable in the warm humid tropics, grown throughout the year. Preference for this vegetable varies highly with the extent of bitterness. In the present study an attempt was made to cross *M. charantia* with *M. dioica*, the bitterless, small fruited, tuberous perennial to seek possibilities of transferring desirable attributes of the latter to the former.

Both direct and reciprocal crosses were made between *M. charantia* and *M. dioica*. Anthesis in *M. charantia* is between 4:00 and 10:00 a.m. while that in *M. dioica* is between 5:30 a.m. and 12:00 p.m. Therefore, pollen from both species were collected at anthesis time, stored at 10 ° C and hand pollinated at the corresponding time of anthesis in both species.

The results (Table 1) indicated that the cross *M. charantia* x *M. dioica* or its reciprocal failed to set fruit when normal pollens were used as reported earlier (Trivedi and Roy, 1972; Dutt and Pandey, 1982). The ovary did not grow further; it shriveled and dried in three days. However, when crosses were made using pollens collected and stored at 10 ° C, the percentage of success was above 90. The fruits and F₁ seeds resembled their respective female parents in both direct and reciprocal crosses (Fig. 1 and 2). The number of chaffy seeds was more in crossed fruits than in selfed fruits. Further studies are in progress.

This is the first report of success of crossing between *M. charantia* and *M. dioica*. The study indicated the possibility of utilizing the bitterless, perennial tuberous *M. dioica* in transferring the desirable attributes to the commercially cultivated large fruited bittergourd. The study also shows the possibility of identifying the progenitor of bittergourd through genome analysis. The failure in earlier studies could be attributed to nonsynchronization of anthesis in two species, which is solved through storing pollens at low temperature and pollinating at anthesis in both species.

Table 1. Crossability between *M. charantia* and *M. dioica*.

Crosses/selfs	Number of				
	Crosses or self	Fruit set	Seeds/fruit	Bold seeds	Chaffy seeds
<i>M. charantia</i> x <i>M. dioica</i> (using pollens stored at 10 ° C)	50	46	23	14	9
<i>M. dioica</i> x <i>M. charantia</i> (using pollens stored at 10 ° C)	50	47	21	11	10
<i>M. charantia</i> (selfed)	25	24	22	19	3
<i>M. dioica</i> (selfed)	25	23	20	18	2
<i>M. charantia</i> x <i>M. dioica</i>	50	0	---	---	---

(using normally collected pollens)					
<i>M. dioica</i> x <i>M. charantia</i>	50	0	---	---	---
(using normally collected pollens)					

Figure 1. Fruits of selfs and reciprocal crosses between *Momordica charantia* and *M. dioica*.

Figure 2. Seeds from selfs and reciprocal crosses between *Momordica charantia* and *M. dioica*.




Figure 1. Fruits of selfs and reciprocal crosses between *Momordica charantia* and *M. dioica*.




Figure 2. Seeds from selfs and reciprocal crosses between *Momordica charantia* and *M. dioica*.

Literature Cited

1. Dutt, B. and C. Pandey. 1982. Callose deposition in relation to incompatibility in *Momordica* In: Roy, R.P. and U. Sinha (Ed.) Cytogenetics in India.
2. Trivedi, R.N. and P.P. Roy. 1972. Cytological studies in some species of *Momordica*. *Genetica* 43:282-291.




Figure 1. Fruits of selfs and reciprocal crosses between *Momordica charantia* and *M. dioica*.




Figure 2. Seeds from selfs and reciprocal crosses between *Momordica charantia* and *M. dioica*.

Preliminary screening of cucurbits species for *Bemisia tabaci* Genn. whitefly resistance

V. Moreno, J.L. Gomez Agulera, C. Guerau de Arellano and L.A. Roig

Plant Cell & Tissue Laboratory, Biotechnology Department, Universidad Politecnica de Valencia. C.Camino de Vera, 14.46022-Valencia (Spain)

The majority of the Spanish production of early cucurbits dedicated to the foreign markets is being produced in greenhouses in Almeria, Southern Spain. In the last few years, one of the most important problems in the culture of melon in this area was the yellowing virus disease (1,2,3) transmitted by the greenhouse whitefly *Trialeurodes vaporariorum* W. (4). A systematic investigation of sources for resistance to this disease led to the discovery of resistance in several wild species of *Cucumis* and *Citrullus* (1,5) as well as in *cucumis melo* var *agrestis* (6).

More recently, a progressive substitution of *Trialeurodes* by another whitefly *Bemisia tabaci* Genn. was detected in that geographic area. In 1991 this was responsible for the severe losses in production of melon cultured in the greenhouses of Almeria. This fact seems to be similar to those previously described in the lower desert valleys of California, Arizona and Mexico fall melon production between 1978 and 1991 (7), and in the United Arab Emirates with both melon (8) and watermelon (9).

The acquisition of new sources of resistance to whitefly could represent the first step in the fight against the direct as well as indirect damages which whitefly produces. Interestingly, preliminary screening of accessions in *agrestis*-type melon for resistance to SPWF-P opened up hopeful results (7). In this paper we present the response versus *Bemisia* of a series of 30 entries belonging to genera *Cucumis* (including melon, cucumber and wild relatives), *Citrullus* (watermelon and the wild relative *C. colocynthis*), *Lagenaria*, *Momordica*, *Trichosanthes*, *Benincasa*, *Cucurbita*, *Echballium* and *Luffa*, grown in greenhouse in Almeria, Spain.

The enormous population of whitefly in that area permitted us to evaluate the resistance to *Bemisia* feeding and reproduction in natural conditions. The 30 entries (12-15 plants per accession) of the 18 species were cultivated in the same greenhouse in sandy soil with drip irrigation in the period July-October 1992. Adult population as well as eggs which settled on the leaves were scored and evaluated in a range of 0-5. The aspect of the whole plant and production of fruits were also scored. Table 1 shows the results of whitefly attack on the different entries studied.

All the melon cultivars and the cucumber line were severely affected by whitefly while the *agrestis*-type melon used in this experiment showed low susceptibility. This observation confirms the results reported by MacCreight (7) in a study made in another geographic area with a series of *agrestis*-type melons in which he found certain degree of resistance to *Bemisia* in some entries.

We have also observed a different response within the entries of the wild relatives *Cucumis africanus*, *C. Anguria*, *C. myriocarpus*, *C. dipsaceus* and *C. zeyheri*, but all of them showed several degrees of susceptibility to whitefly. On the contrary, the three accessions of the other wild relative, *Cucumis metuliferus*, were completely resistant to the pest. Since resistance to other plagues and diseases have been described in this species (see a revision in 10) its behavior versus *Bemisia* is very interesting and we will try to evaluate and corroborate these results in subsequent experiments.

Citrullus lanatus (watermelon) plants were completely devastated as a consequence of *Bemisia* attack. The three accessions of their wild relative *Citrullus colocynthis* were susceptible but not as seriously damaged.

Benincasa hispida was as susceptible as melon, cucumber and watermelon. The two entries of *Lagenaria siceraria* were affected by whitefly although at a different level. However, the two species of *Luffa* showed completely different response: *L. acutangula* was susceptible while *L. cylindrica* appeared resistant and free of whitefly. Similarly, all the entries of *Echballium elaterium*, *Momordica Balsamina* and *Trichosanthes cucumerina* appeared permanently free of insects throughout the culture cycle in spite of being cultivated in the same greenhouse together with the other starved species and with a very high

population of *Bemisia* in the environment.

Although the real resistance to *Bemisia* should be confirmed in additional cycles of culture and under different culture conditions, these results can be useful for successive breeding programs. The lesser susceptibility of some lines of *C. Melo* var. *agrestis* (7) (see also table 1) could be relevant since, in this case, there is not any cross ability barriers with the cultivars of melon. On the other hand, to profit from the sources of resistance present in the wild relative *Cucumis metuliferus* or in the genera *Echballium*, *Luffa*, *Momordica* or *Trichosanthes*, it would be necessary to apply a program of somatic hybridization by protoplast fusion. At present, our group is carrying out a wide program of this kind doing symmetric as well as asymmetric hybridizations between melon as recipient species and several of the wild species as donors of desirable genes.

Acknowledgments: The authors express their appreciation to CICYT (Comision Interministerial de Ciencia y Tecnologia, Ministry of Education and Science, Spanish Government for financial support (Project BIO89-0446) and to Dr. F. Garcia-Mari (Dept. of Entomology, U.P.V.) for their technical assistance in the analysis of whitefly population.

Table 1. Susceptibility of different cucurbits to the whitefly *Bemisia tabacci*.

Species and line	Source and accession	Whitefly Population	
		Adults	Eggs
<i>Cucumis melo</i> 'Cant. Charentais'	Clause	+++++	+++++
<i>Cucumis melo</i> 'Valenciano Tardio'	Intersemillas	+++++	+++++
<i>Cucumis melo</i> 'Tokio Early'	J. Abadia (CEBAS, Murcia)	+++++	+++++
<i>Cucumis sativus</i> 'Marketer'	Clause	+++++	+++++
<i>Cucumis melo</i> var. <i>agrestis</i>	Gatersleben CuM 190/1982	+	+
<i>Cucumis africanus</i> L5	IVT Gbnr. 1984	++	-
<i>Cucumis anguria</i> var. <i>longipes</i> L1	Pretoria, 71113	+	+
<i>Cucumis anguria</i> var. <i>longipes</i> L3	IVT Gbnr 1790	+++	+++
<i>Cucumis dipsaceus</i>	Gatersleben Ha 408/1981	++	++
<i>Cucumis metuliferus</i> L1	IVT Gbnr. 1802	-	-
<i>Cucumis metuliferus</i> L4	Gatersleben CuC 16/1981	-	-
<i>Cucumis metuliferus</i> L3	Pretoria, 78263	-	-
<i>Cucumis myriocarpus</i> L1	IVT Gbnr. 1079	+++++	+++++
<i>Cucumis myriocarpus</i> L2	IVT. Gbnr. 1051	++	++
<i>Cucumis zeyheri</i> L1	IVT Gbnr 1786	++++	+++
<i>Cucumis zeyheri</i> L2	IVT Gbnr. 1785	++	+
<i>Cucumis zeyheri</i> L3	Pretoria, 77048	+++	++
<i>Citrullus lanatus</i> 'Dulce Maravilla'	F1 Hybrid Sluis&Groot	+++++	+++++
<i>Citrullus colocynthis</i> 'Rhodes'	Edisto R-309	++	+++
<i>Citrullus colocynthis</i> 'Argelia'	UVP-87 (Argelia)	+	+++
<i>Citrullus colocynthis</i> 'Canarias'	UVP-85 (Gran Canaria)	++	+++
<i>Benincasa hispida</i>	Gatersleben BEN 14/1982	+++++	+++++
<i>Cucurbita martinezii</i>	INRA, Montfavet, 1981	++	-
<i>Echballium elaterium</i>	UPV-EE87	-	-
<i>Lagenaria siceraria</i> 'Murcia'	UPV.ETSIA, 87	++++	++++
<i>Lagenaria siceraria</i> 'Gatersleben'	Gatersleben LAG 41/1986	++	+++
<i>Luffa cylindrica</i>	Gatersleben LUF 25/1985	-	-

<i>Luffa acutangula</i>	Gatersleben LUF 19/1982	+++	+++
<i>Momordica balsamina</i>	Gatersleben MOM 16/1985	-	-
<i>Trichosanthes cucumerina</i>	Gatersleben TCH 4/1982	-	-

EDISTO: Clemson University, College of Agricultural Sciences, S.C. Agricultural Experiment Station/EDISTO Branch Station, P.O. Box 247, Blackville, South Carolina 29817, USA

GATERSLEBEN: Zentralinstitut für Genetik und Kulturpflanzenforschung (ZIGuK), DDR-4325 Gatersleben.

INRA-Montfavet: Institut National de la Recherche Agronomique, Station d'Amélioration des Plantes Maraichères, Domaine Sainte Maurice, 8410, Montfavet, France.

INTERSEMILLAS: Intersemillas, S.A. Quart de Poblet, Valencia, Spain.

CLAUSE: Clause Graines d'Elite.91221 Bretigny-sur-orge. Cedex. France.

CEBAS: Centro de Edafología y Biología Aplicada del Segura, Murcia, Spain.

SLUIS & GROOT: Sluis & Groot, Almeria, Spain.

UPV: Universidad Politécnica de Valencia, 46022-Valencia, Spain.

I.V.T: Institute for Horticultural Plant Breeding, P.O. Box 16, 6700 AA, Wageningen, The Netherlands

PRETORIA: Department of Agricultural Economics and Marketing, Directorate of plant and Seed Control (DPSC), Private Bag X179, Pretoria 0001, South Africa

Literature Cited

1. Esteva, J., F. Nuez and J. Cuatero. 1988. Resistance to yellowing disease in wild relatives of muskmelon. *Cucurbit Genet. Coop. Rpt.* 11:52-53.
2. Esteva, J., F. Nuez, and M.L. Gomez-Guillamon. 1989. Resistance to yellowing disease in muskmelon. *Cucurbit Genet. Coop. Rpt.* 12:44-45.
3. Jorda, C. and A. Alfar. 1989. El amarilleamiento del melon: su diagnosis. Documentos de trabajo del V Congreso Nacional de Fitopatología. Sección: Etiología y Epidemiología (comunicaciones). Badajoz, 17-20 Octubre 1989.
4. Soria, C. and M.L. Gomez-Guillamon. 1989. Transmission of the causal agent of muskmelon yellowing disease. *Cucurbit Genet. Coop. Rpt.* 12:40-41.
5. Soria, C. M.L. Gomez-Guillamon, J. Esteva, and F. Nuez. 1989. Search for sources of resistance to yellowing disease in *Cucumis* spp. *Cucurbit Genet. Coop. Rpt.* 14:59-60.
6. Nuez, F. J. Esteva, C. Soria, and M.L. Gomez-Guillamon. 1991/ Search for sources of resistance to a whitefly transmitted yellowing disease in melon. *Cucurbit Genet. Coop. Rpt.* 14:59-60.
7. McCreight, J.D. 1992. Preliminary screening of melons for sweetpotato whitefly resistance, *Cucurbit Genet. Coop. Rpt.* 15:59-61.
8. Hassan, A.A., H.H. Al-Masri, U.A. Obaji, M.S. Wafi, N.E. Quronfilah, and M.A. Al-Rays. 1991a. Screening of domestic and wild *Cucumis melo* germplasm for resistance to the yellowing-stunting disorder in the United Arab Emirates. *Cucurbit Genet. Coop. Rpt.* 14: 56-58.
9. Hassan, A.A., N.E. Quronfilah, U.A. Obaji, M.A. Al-Rays and M.S. Wafi. 1991b. Screening of Domestic and wild *Citrullus* germplasm for resistance to the yellow-stunting disorder in the United Arab Emirates. *Cucurbit Genet. Coop. Rpt.* 14: 98-101.
10. Moreno V. and L.A. Roig. 1990. Somaclonal variation in Cucurbits. In: "Biotechnology in Agriculture and Forestry". Bajaj YPS (ed.). Vol 11. Springer-Verlag, Berlin, pp. 435-464.

Cucumber, Melon and Squash Germplasm from the Cornell Collection

Molly Kyle, George Moriarty and Henry M. Munger

Department of Plant Breeding and Biometry, Cornell University, Ithica, NY 14853

A number of accessions and breeding lines representing H. M. Munger's *Cucumis* and *Cucurbita* germplasm collection have been increased in recent years and are available to interested workers. This work was supported in part by Hortinova and Petoseed Co.

Cucumis

Cucumber
Tokyo Long Green
Yomaki - Source of high level PMR
Hawaii 60G-10 - Source of high level PMR bred by Gilbert
Denna's Dwarf - Tablegreen ype
Marketmore 80 <i>B_w</i> (Bacterial Wilt Resistance)
Albion (PMR551) <i>B_w</i> (Bacterial Wilt Resistance)
additional cucumber germplasm listed in Munger (1985)
Melon
Iroquois
PMR Iroquois
PMR Monoecious Iroquois
Delicious 51 - (typical Delicious 51)
PD 23 - selection 23, PMR Delicious 51, male parent of 'Progress' hybrid
PB 13 - Minn 90-36, Fusarium resistant, similar to PB12 (Minn 99-36), the resistant parent of Iroquois
MR 324, 335 - Eastern type with low level cucumber mosaic (CM) resistance
M (+MMR) 324, 328, 339 Eastern type with low level CM resistance
PM (= PMM) 324, 328, 339 Eastern type with low level CM resistance + monoecious + PMR
PPM 339 - Monoecious, PMR + papaya ringspot resistance
ZPPM 339 - PPM 339 + ZYM resistance
PMR Charentais
Siberian Honeydew
<i>C. melo conomon</i> 'Freeman cucumber' - source of CM resistance (Enzie 1943)
<i>C. melo dudaim</i> - Source of monoecious (Wall 1967) and WM resistance
PI 414723-4 S ₃ - Source of WMV and ZYM resistance
<i>Cucurbita</i>

Genic male sterility in Eskandarany Egyptian squash (*C. pepo*)

Male sterile EPS type derived from Eskandarany

Male Sterile Caserta type derived from Eskandarany

CMR 469 - CMV-resistant yellow *C. pepo* derived from *C. martinezii*

W225 - CMV-resistant yellow *C. pepo* derived from *C. martinezii* (viny)

PI 174186 selected for scab resistance

Early Prolific Straightneck x PI 174186 F₂ and backcrosses

PI 174183 selected for scab resistance

C. martinezii Source of powdery mildew resistance (Contin and Munger 1977), CMV resistance and gummy stem blight resistance

C. pepo x *C. moschata* bush butternut populations (Munger 1990).

C. mixta Variant, silver skin gourd

Literature Cited

1. Contin, M. 1978. Interspecific transfer of powdery mildew resistance in the genus *Cucurbita*. Ph.D. Thesis. Cornell Univ.
2. Contin, M. and H.M. Minger. 1977. Inheritance of powdery mildew resistance in interspecific crosses with *Cucurbita martinezii*. 12:29 (Abstr.)
3. Enzie, W.D. 1943. A source of muskmelon mosaic resistance found in the oriental pickling melon, *Cucumis melo* var. conomon. Proc. Amer. Soc. Hort. Sci.
4. Munger, H.M. 1985. Near-isogenic lines of several cucumber varieties. CGC Rept. 8:4-6.
5. Munger, H.M. 1990. Availability and use of interspecific populations involving *Cucurbita moschata* and *C. pepo*. CGC Rept. 13:49.
6. Wall, J.R. 1967. Correlated inheritance of sex-expression and fruit shape in *Cucumis*. Euphytica 16:199-208.

Gene List for Cucumber

Todd C. Wehner

Department of Horticultural Science, North Carolina State University, Raleigh, NC 27695-7609

Lists of the known genes for the Cucurbitaceae have been published previously (12, 16). However, in the interest of updating the information, following is a list of the 43 genes introduced or modified since the last report (13). That makes a total (105 plus 43 minus 2 renamed) 146 reported gene mutants.

A wild type cultivar has not been proposed for cucumber as the reference for all "+" alleles. 'Marglobe' tomato has been chosen as the normal type for genetic studies in that crop. Perhaps 'Wisconsin SMR 18' should be the choice for cucumber, since it is an inbred that is used widely in genetic studies. In addition, it has many dominant and wild-type alleles.

The genes on the following list are of four categories, seedling markers, virus resistance, isozymes, and fruit mutants (Table 1). The seedling markers include a reclassification of *gc* as *ls* (25, 26) and a new group produced using gamma rays on pollen (4, 5). The fruit mutants are green mature fruit, *gn* (11), and palisade epidermis, *Pe* (2). Isozyme mutants now include 14 markers, many of which are linked (6). One virus resistance gene should be renamed according to the revised taxonomy of watermelon mosaic virus. Watermelon mosaic virus-1 was renamed papaya ringspot virus-waermelon strain (PRSV-W), so *wmv-1-1* should be renamed *prsv*, retaining *wmv-1-1* as a synonym. Watermelon mosaic virus-2 is renamed watermelon mosaic virus (WMV). Thus, the dominant gene for resistance to WMV does not change its symbol, remaining *Wmv*.

Isozyme variant nomenclature for this gene list follows the form according to Staub et al. (22), such that loci coding for enzymes (e.g. glutamine dehydrogenase, G2DH) are designated as abbreviations, where the first letter is capitalized (e.g. G2dh). If an enzyme system is conditioned by multiple loci, then those are designated by hyphenated numbers, which are numbered from most cathodal to most anodal and enclosed in parentheses. The most common allele of any particular isozyme is designated 100, and all other alleles for that enzyme are assigned a value based on their mobility relative to that allele.

Researchers are encouraged to send reports of new genes, as well as seed samples to the cucumber gene curators (Todd C. Wehner, Jack E. Staub and Richard W. Robinson). Please let me know of any omissions or errors in the following list.

Table 1. The 43 new or revised genes of cucumber*.

Gene symbol			
Preferred	Synonym	Character	References
<i>al</i>		<i>albino cotyledons</i> .. White cotyledoms and slightly light green hypocotyl; dying before the first true leaf stage. Wild type <i>al+</i> from 'Nishiki-suyo'; <i>al</i> from M ₂ line from pollen irradiation.	4, 5
<i>by</i>	<i>bu</i>	<i>bushy</i> . Short internodes; normal seed viability. Wild type <i>by+</i> from 'Borszczagowski'; <i>by</i> from induced mutation of 'Borszczagowski'. . Linked with <i>F</i> and <i>gy</i> , not with <i>B</i> or <i>bi</i> ..	9
<i>chp</i>		<i>choripetalous</i> . Small first true leaf' choripetalous flowers; glossy ovary; small fruits; few seeds. Wild type <i>chp+</i> from 'Borszczagowski'; <i>chp</i> from chemically induced mutation.	7
<i>cp-2</i>		<i>compact-2</i> . Short internodes; small seeds, similar to <i>cp</i> , but allelism not checked. Wild type <i>cp-2+</i> from 'Borszczagowski'; <i>cp-2</i> from induced mutation of 'Borszczagowski' called W97. Not linked with <i>B</i> or <i>F</i> ; interacts with <i>by</i> to produce super dwarf.	10
<i>de-2</i>		<i>determinate-2</i> . Main stem growth ceases after 3 to 10 nodes, producing flowers	21

		at the apex; smooth, fragile, dark-green leaves; similar to <i>de</i> , but not checked for allelism. Wild type <i>de-2+</i> from 'Borszczagowski'; <i>de-2</i> from W-sk mutant induced by ethylene-imine from 'Borszczagowski'.	
<i>dm-1</i>	<i>dm?</i>	<i>downy mildew resistance</i> . One of three genes for resistance to downy mildew caused by <i>Pseudoperonospora cubensis</i> (Berk & Curt). Wild type <i>dm-1+</i> from Wisconsin SMR 19; <i>dm-1</i> from WI 4783. Not checked for allelism with <i>dm</i> .	1
<i>dm-2</i>		<i>downy mildew resistance-2</i> . One of three genes for resistance to downy mildew caused by <i>Pseudoperonospora cubensis</i> Berk & Curt). Wild type <i>dm-2+</i> from Wisconsin SMR 18; <i>dm-2</i> from WI 4783. Not checked for allelism with <i>dm</i> .	1
<i>dm-3</i>		<i>downy mildew resistance-3</i> . One of three genes for resistance to downy mildew caused by <i>Pseudoperonospora cubensis</i> (Berk & Curt). Wild type <i>dm-3+</i> from Wisconsin SMR 18; <i>dm-3</i> from WI 4783. Not checked for allelism with <i>dm</i> .	1
<i>dvl-2</i>	<i>dl-2</i>	<i>divided leaf-2</i> . Divided leaves after the 2nd true leaf; flower petals free; similar to <i>dvl</i> , but allelism not checked. Wild type <i>dvl-2+</i> from 'Borszczagowski'; <i>dvl-2</i> from mutant induced by ethylene-imine from 'Borszczagowski'.	19
<i>dwc-1</i>		<i>dwarf cotyledons-1</i> . Small cotyledons; late germination; small first true leaf; died after 3rd true leaf. Wild type <i>dwc-1+</i> from 'Nishiki-suyo'; <i>dwc-1</i> from M ₂ line from pollen irradiation.	4, 5
<i>dwc-2</i>		<i>dwarf cotyledons-2</i> . Small cotyledons, later germination; small first true leaf. Wild type <i>dwc-2+</i> from 'Nishiki-suyo'; <i>dwc-2</i> from M ₂ line from pollen irradiation.	4, 5
<i>G2dh</i>		<i>Glutamine dehydrogenase</i> (E.C.# 1.1.1.29). Isozyme variant found segregating in PI 285606; 5 alleles observed.	6
<i>gi-2</i>		<i>ginko-2</i> . Spatulate leaf blade with reduced lobing and altered veins; recognizable at the 2nd true leaf stage; similar to <i>gi</i> , fertile instead of sterile. wild type <i>gi-2+</i> from 'Borszczagowski'; <i>gi-2</i> from mutant in the Kubicki collection.	19
<i>gig</i>		<i>gigantism</i> . First leaf larger than normal. Wild type <i>gig</i> from 'Borszczagowski'; <i>gig</i> from chemically induced mutation.	8
<i>gn</i>		<i>green mature fruit</i> . Green mature fruits when <i>R+R+ gngn</i> ; cream colored when <i>R+R+ gn+gn+</i> orange when <i>R_</i> . Wild type <i>gn+</i> from 'Chipper', SMR 58 and PI 165509; <i>gn</i> from TAMU 830397.	11
<i>Gpi-1</i>		<i>Glucose phosphate isomerase</i> . (E.C. # 5.3.1.9). Isozyme variant found segregating (1 and 2) in PI 176524, 200815, 249561, 422192, 432854, 436608; 3 alleles observed.	6
<i>Gr-1</i>		<i>Glutathione reductase-1</i> (E.C. # 1.6.4.2). Isozyme variant found segregating in PI 109275; 5 alleles observed.	6
<i>hl</i>		<i>heart leaf</i> . Heart shaped leaves. Wild type <i>hl+</i> from Wisconsin SMR 18; <i>hl</i> from WI 2757. Linked with <i>ns</i> and <i>ss</i> in the linkage group with <i>Tu-u-D-pm</i> .	23
<i>hn</i>		<i>horn like cotyledons</i> . Cotyledons shaped like bull horns; true leaves with round shape rather than normal lobes; circular rather than ribbed stem cross section; divided petals; spineless fruits; pollen fertile, but seed sterile. Wild type <i>hn+</i> from 'Nishiki-suyo'; <i>hn</i> from M ₂ line from pollen irradiation.	4, 5
<i>hsl</i>		<i>heart shaped leaves</i> . Leaves heart shaped rather than lobed; tendrils branched. Wild type <i>hsl+</i> from 'Nishiki-suyo' <i>hsl</i> from M ₂ line from pollen irradiation.	4, 5
<i>Idh</i>		<i>Isocitrate dehydrogenase</i> (E.C. # 1.1.1.42). Isozyme variant found segregating in PI 183967, 21589; 2 alleles observed.	6

<i>lg-1</i>		<i>light green cotyledons-1</i> . Light green cotyledons, turning dark green; light green true leaves, turning dark green; poorly developed stamens. Wild type <i>lg-1+</i> from 'Nishiki-suyo'; <i>lg-1</i> from M ₂ line from pollen irradiation.	4, 5
<i>lg-2</i>		<i>light green cotyledons-2</i> . Light green cotyledons, turning dark green (faster than <i>lg-1</i>); light green true leaves, turning dark green; normal stamens. Wild type <i>lg-2+</i> from 'Nishiki-suyo'; <i>lg-2</i> from M ₂ line from pollen irradiation.	4, 5
<i>ls</i>	<i>gc</i>	<i>light sensitive</i> . Yellow cotyledons, lethal in high light. Abstract gave <i>cg</i> as symbol; article that followed gave <i>ls</i> as symbol. Mutant <i>ls</i> from a selection of 'Burpless Hybrid'.	25, 26
<i>Mdh-1</i>		<i>Malate dehydrogenase-1</i> (E.C. # 1.1.1.37.). Isozyme variant found segregating in PI 171613, 209064, 326594; 3 alleles observed.	6
<i>Mdh-2</i>		<i>Malate dehydrogenase-2</i> (E.C.# 1.1.1.37). Isozyme variant found segregating in PI 174164, 185690, 357835, 419214; 2 alleles observed.	6
<i>Mdh-3</i>		<i>Maleate dehydrogenase-3</i> (E.C.# 1.1.1.27). Isozyme variant found segregating in PI 255236, 267942, 432854, 432887; 2 alleles observed.	6
<i>Mpi-2</i>		<i>Mannose phosphate isomerase</i> (E.C. # 5.3.1.8). Isozyme variant found segregating in PI 109275, 175692, 200815, 209064, 263049, 354952; 2 alleles observed.	6
<i>mpy</i>	<i>mpi</i>	<i>male pygmy</i> . Dwarf plant with only staminate flowers. Wild type <i>mpy+</i> from Wisconsin SMR 12' <i>mpy</i> from Gnome 1, a selection of 'Rochford's Improved'.	14
<i>Pe</i>		<i>Palisade epidermis</i> . Epidermal cells arranged perpendicular to the fruit surface. Wild type <i>Pe</i> from 'Wisconsin SMR 18', 'Spartan Salad' and Gy 2 compact; <i>pe</i> from WI 2757.	2
<i>Pep-la</i>		<i>Peptidase with leucyl-leucine</i> (E.C. # 3.4.13.11). Isozyme variant found segregating in PI 169380, 175692, 263049, 289698, 354952, 5 alleles observed.	6
<i>Pep-pap</i>		<i>Peptidase with phenylalanyl-L-proline</i> (E.C. # 3.4.13.11). Isozyme variant found segregating in PI 163213, 188749, 432861; 2 alleles observed.	6
<i>Per-4</i>		<i>Peroxidase</i> (E.C. # 1.11.1.7). Isozyme variant found segregating in PI 215589; 2 alleles observed.	6
<i>Pgd-1</i>		<i>Phosphogluconate dehydrogenase-1</i> (E.C. # 1.1.1.43). Isozyme variant found segregating in PI 169380, 175692, 222782; 2 alleles observed.	6
<i>Pgd-2</i>		<i>Phosphogluconate dehydrogenase-2</i> (E.C. # 1.1.1.43). Isozyme variant found segregating in PI 171613, 177364, 188749, 263049, 285606, 289698, 354952, 419214, 432858; 2 alleles observed.	6
<i>Pgm-1</i>		<i>Phosphoglucomutase</i> (E.C. # 5.4.2.2.). Isozyme variant found segregating in PI 171613, 177364, 188749, 263049, 264229, 285606, 289698, 354952; 2 alleles observed.	6
<i>prsv</i>	<i>wmv-1-1</i>	<i>watermelon mosaic virus 1 resistance</i> . Resistance to papaya ringspot virus (formerly watermelon mosaic virus 1). Wild type <i>prsv+</i> from WI2757; <i>prsv</i> from 'Surinam'.	24
<i>sh</i>		<i>short hypocotyl</i> . Hypocotyl of seedlings 2/3 the length of normal. Wild type <i>sh+</i> from 'Borszczagowski'; <i>sh</i> from khp, an induced mutant from 'Borszczagowski'.	20
<i>shl</i>		<i>shrunk leaves</i> . First and 2nd true leaves smaller than normal; later leaves becoming normal; slow growth; often dying before fruit set. Wild type <i>shl+</i> from 'Nishiki-suyo'; <i>shl</i> from M ₂ line from pollen irradiation.	4, 5
<i>sp-2</i>		<i>short petiole-2</i> . Leaf petioles shorter, darker green than normal at 2-leaf stage; crinkled leaves with slow development; short hypocotyl and stem; little branching. Not tested for allelism with <i>sp</i> . Wild type <i>sp-2+</i> from 'Borszczagowski'; <i>sp-2</i> from	18

		chemically induced mutation.	
<i>wi</i>		<i>wilty leaves</i> . Leaves wilting in the field, but not in shaded greenhouse; weak growth; no fruiting. Wild type <i>wi+</i> from greenhouse; weak growth; no fruiting. <i>wi</i> from M ₂ line from pollen irradiation.	4, 5
<i>wy</i>		<i>wavy rimed cotyledons</i> . Wavy rimed cotyledons, with white centers; true leaves normal. Wild type <i>wy+</i> from 'Nishiki-suyo'; <i>wy</i> from M ₂ line from pollen irradiation.	4, 5
<i>ys</i>		<i>yellow stem</i> . Yellow cotyledons, becoming cream-colored; cream-colored stem, petiole and leaf veins; short petiole; short internode. Wild type <i>ys+</i> from 'Borszczagowski'; <i>ys</i> from chemically induced mutation.	17

*Isozyme nomenclature follows a modified form (22) previously described by Richmond (15) and Gottlieb (3).

Literature Cited

- Dorochowski, R.W. and E. Lakowska-Ryk. Inheritance of resistance to downy mildew (*Pseudoperonospora cubensis* Berk & Curt) in *cucumis sativus*. Proc. 5th EUCARPIA, Cucurbitaceae Symp. p. 66-69, Warsaw, Poland.
- Fanourakis, N.E. and P.W. Simon. 1987. Inheritance and linkage studies of the fruit epidermal structure in cucumber. J. Hered. 78: 369-371.
- Gottlieb, T.D. 1977. Evidence for duplication and divergence of the structural gene for phosphoglucose isomerase in diploid species of *Clarkia*. Genetics 86: 289-307.
- Iida, S. and E. Amano. 1990. Pollen irradiation to obtain mutants in monoecious cucumber. Ga., a Field Symp. 29: 95-111.
- Iida, S. and E. Amano. 1991. Mutants induced by pollen irradiation in cucumber. Cucurbit Genet. Coop. Rpt. 14:32-33.
- Knerr, L.D. and J.E. Staub. 1992. Inheritance and linkage relationships of isozyme loci in Cucumber (*Cucumis sativus* L.). Theor. Appl. Genet. 84:217-224.
- Kubicki, B. and A. Korzeniewska. 1984. Induced mutations in cucumber (*Cucumis sativus* L.) III. A mutant with choripetalous flowers. Genetica polonica 25: 53-60.
- Kubicki, B., I Goszczycka and A. Korzeniewska. 1984. Induced mutations in cucumber (*Cucumis sativus* L.) II. Mutant of gigantism. Genetica Polonica 25: 41-52.
- Kubicki, B., U. Soltysiak and A. Korzeniewska. 1986a. Induced mutations in cucumber (*Cucumis sativus* L.). IV. A mutant of the bush type of growth. Genetica Polonica 27:273-287.
- Kubicki, B., U. Soltysiak and A. Korzeniewska. 1986b. Induced mutations in cucumber (*Cucumis sativus* L.). V. Compact type of growth. Genetica Polonica 27: 289-298.
- Peterson, G. C. and L. M. Pike. 1992. Inheritance of green mature seed-stage fruit color in *Cucumis sativus* L. J. Amer. Soc. Hort. Sci. 117:643-645.
- Pierce, L.K. and T.C. Wehner. 1989. Gene list for cucumber. Cucurbit Genet. Coop. Rpt. 12:91-103.
- Pierce, L.K. and T.C. Wehner. 1990. Review of genes and linkage groups in cucumber. HortScience 25: 605-615.
- Pyzhenkov, V. and G.A. Kosareva. 1981. Description of features and their inheritance pattern in a new cucumber mutant. Biull. Vses. Inst. Rasteniievod 109: 5-8 (Plant Breeding Abstracts 1983, 53: 478).
- Richmond, R.C. 1972. Enzyme variability in the *Drosophila williston* group. 3. Amounts of variability in the super species *D. Paulistorum*. Genetics 70: 87-112.
- Robinson, R.W., H.M. Munger, T.W. Whitaker and G.W. Bohn. 1976. Genes of the Cucurbitaceae. HortScience 11:554-568.
- Rucinska, M., K. Niemirowicz-Szczytt and A. Korzeniewska. 1991. A cucumber (*Cucumis sativus* L.) mutant with yellow stem and leaf petioles. Cucurbit Genet. Coop. Rpt. 14: 8-9.
- Rucinska, M. K. Niemirowicz-Szczytt and A. Korzeniewska. 1992a. Cucumber (*Cucumis sativus* L.) induced mutations. II. A second short petiole mutant. Cucurbit Genet. Coop. Rpt. 15:33-34.
- Rucinska, M., K. Niemirowicz-Szczytt and A. Korzeniewska. 1992b. Cucumber (*Cucumis sativus* L.) induced mutations. III and IV. Divided and ginko leaves. Proc. 5th EUCARPIA Cucurbitaceae Symp. p. 66-69, Warsaw, Poland.
- Soltysiak, U. and B. Kubicki. 1988. Induced mutations in the cucumber (*Cucumis sativus* L.). VII. Short hypocotyl mutant. Genetica Polonica 29: 315-321.
- Soltysiak, U., B. Kubicki and A. Korzeniewska. 1986. Induced mutations in cucumber (*Cucumis sativus* L.). VI. Determinate type of growth. Genetica Polonica 27: 229-308.

22. Staub, J.E., R.S. Kupper, D. Schuman, T.C. wehner and B. May. 1985. Electrophoretic variation and enzyme storage stability in cucumber. *J. Amer. Soc. Hort. Sci.* 110:426-431.
23. Vakalounakis, D.J. 1992. Heart leaf, a recessive leaf shape marker in cucumber: linkage with disease resistance and other traits. *J. Heredity* 83: 217-221.
24. Wang, W.J., R. Provvidenti and R.W. Robinson. 1984. Inheritance of resistance in cucumber to watermelon mosaic virus. *Phytopathology* 51: 423-428.
25. Whelan, E. D.P. 1971. Golden cotyledon: a radiation-induced light sensitive mutant in cucumber, *HortScience* 6:343.
26. Whelan, E.D.P. 1972. Inheritance of a radiation-induced light sensitive mutant of cucumber. *J. Amer. Soc. Hort. Sci.* 97-765-767.

Scientists should consult the above list as well as the rules of gene nomenclature for the Cucurbitaceae (16) before choosing a gene name and symbol. That will avoid duplication of gene names and symbols. The rules of gene nomenclature were adopted in order to provide guidelines for naming and symbolizing genes. Scientists are urged to contact members of the Gene List committee regarding rules and gene symbols.

Gene List Committee:

Cucumber	T.C. Wehner
Cucurbita spp.	R.W. Robinson
	M.G. Hutton
Melon	M. Pitrat
Watermelon	W.R. Henderson
Other genera	R.W. Robinson

Gene Nomenclature for the Cucurbitaceae

[From: Robinson, R. W., H.M. Munger, T.W. Whitaker and G.W. Bohn 1976.]

Genes of the Cucurbitaceae, HortScience 11:554-568.]

1. Names of genes should describe a characteristic feature of the mutant type in a minimum of adjectives and/or nouns in English or Latin.
2. Genes are symbolized by italicized Roman letters, the first letter of the symbol being the same as that for the name. A minimum number of additional letters are added to distinguish each symbol.
3. The first letter of the symbol and name is capitalized if the mutant gene is dominant, and all letters of the symbol and name are in lower case if the mutant gene is recessive to the normal type. The normal allele of a mutant gene is represented by the symbol "+", where it is needed for clarity, the symbol of the mutant gene followed by the superscript "+". The primitive form of each species shall represent the + allele for each gene, except where long usage has established a symbol named for the allele possessed by the normal type rather than the mutant.
4. A gene symbol shall not be assigned to a character unless supported by statistically valid segregation data for the gene.
5. Mimics, i.e. different mutants having similar phenotypes, may either have distinctive names and symbols or be assigned the same gene symbol, followed by a hyphen and distinguishing Arabic numeral or Roman letter printed at the same level as the symbol. The suffix-1 is used, or may be understood and not used, for the original gene in a mimic series. It is recommended that allelism tests be made with a mimic before a new gene symbol is assigned to it.
6. Multiple alleles have the same symbol, followed by a Roman letter or Arabic number superscript. Similarities in phenotype are insufficient to establish multiple alleles; the allelism test must be made.
7. Indistinguishable alleles, i.e., alleles at the same locus with identical phenotypes, preferably should be given the same symbol. If distinctive symbols are assigned to alleles that are apparent recurrences of the same mutation, however, they shall have the same symbol with distinguishing numbers or letters in parentheses as superscripts.
8. Modifying genes may have a symbol for an appropriate name, such as intensifier, suppressor, or inhibitor, followed by a hyphen and the symbol of the allele affected. Alternatively, they may be given a distinctive name unaccompanied by the symbol of the gene modified.
9. In cases of the same symbol being assigned to different genes, or more than one symbol designated for the same gene, priority in publication will be the primary criterion for establishing the preferred symbol. Incorrectly assigned symbols will be enclosed in parentheses on the gene lists.

[From: CGC Gene List Committee, 1982. Update of cucurbit gene list and nomenclature rules. CGC 5:62-66.]

The same symbol shall not be used for nonallelic genes of different *Cucurbita* species. Allelic genes of compatible species are designated with the same symbol for the locus.

Cucurbit Genetics Cooperative 1993 Membership Directory

Cucurbit Genetics Cooperative

Individual Members

1. **Al-Sulaiman, Abdul Mohsen I.** National Agricultural & Water Research Center, P.O. Box 54763, Ridyah 11524, Saudi Arabia
2. **Ayuso, Ma Cruz** Petoseed Iberica, S.A. Carretera de Malaga, 34,04710 Santa Maria del Aguila, Almeria, Spain.
3. **Barbercheck, Mary** Dept. Entomology, Box 7634, North Carolina State Univ. Raleigh, NC, 27695-7634. Tel: (919) 737-2638
4. **Beekman, A.G.B.** ROYAL SLUIS, P.O. Box 22, 1600 AA Enkhuizen, The Netherlands
5. **Ben Tahar, Sofia** BIOSEM, Campus Universitaire des Cezeaux, 24, Avenue des Landais, 63170 Aubiere, France
6. **Bohn, G. W.** 1094 Klish Way, Del Mar, CA 92014 Tel: (619) 755-4780. Breeding and genetics of resistance to fungi, viruses and insects.
7. **Boorsma, P.A.** Vegetable Research, Sluis & Groot, P.O. Box 26, 1600 AA Enkhuizen, The Netherlands
8. **Bradbury, Gary K.** Saphir International, The Oast, Perry Court, London Road, Faversham, Kent, ME13 8RY, Great Britain. Tel: (44) 795 530700. FAX: (44) 795-530797. Squash and pumpkin breeding for virus disease resistance and horticultural traits.
9. **Burkett, Al** Petoseed Woodland Research Station, 37437 State Highway 16, Woodland, CA 95695.
10. **Callaway, M. Brett** Atwood Research Facility, Kentucky State University, Frankfort, KY, 40601. Tel: (502) 227-7291.
11. **Canos Rios, Pedro** Apartado Postal No. 247, Tereon, Coah., Mexico
12. **Carey, Edward E.** IIE-CIP, 809 UN Plaza, New York, NY. 10017-3580. E-mail: CGI801%NSFMAIL@INTER
13. **Carle, R. Bruce** Univ. New Hampshire, Dept. Plant Science - Newmith Hall, Durham, NH, 03824. Tel: (603) 862-3205. Cucurbit breeding, gametophytic selection, herbicide resistance.
14. **Chambonnet, Daniel** Station d'Amelioration des Plantes Maraicheres, B.P. 94 , 84140 Montfavet, France.
15. **Charpiot, Bernard** TEZIER - Centre de Recherche, Domaine de Maninet, Route de Beaumont, 26000 Valence, France.
16. **Chen, Fure-Chyi** Univ. Hawaii, Dept. of Horticulture, 3190 Maile Way # 102, Honolulu, HI, 96822. Tel: (808) 956-7992. Gene transfer, breeding, tissue culture.
17. **Chung, Paul** Petoseed Woodland Research Station, 37437 State Highway 16, Woodland, CA 95695. Tel: (916) 666-0931.
18. **Clayberg, C. D.** Department of Horticulture, Waters Hall, Kansas State University, Manhattan, KS 66502.
19. **Close, Timothy J.** University of California, Riverside, Dept. Botany & Plant Sciences, Roverside, CA, 92521-0124. Tel: (714) 787-3318. FAX: (714) 787-4437. E-mail: timclose@ucrac1.ucr.edu.
20. **Cohen, Yigal** Department of Life Sciences, Bar-Ilan University, Ramat-Gan 52 100, Israel .
21. **Corella, Pilar** Asgrow SEed Co. Apdo. 175, 04700 El Ejido (Almeria), Spain.
22. **Coyne, Dermot P.** Department of Horticulture, University of Nebraska, Lincoln, NE 68583-0724 Tel: (402) 472-1126. FAX: (402) 472-2853. Breeding, genetics, physiology, plant development and disease resistance of squash.
23. **Croom, Edward M. Jr.** 716 University Avenue, Oxford, MS, 38655. Tel: (601) 232-5324. FAX: (601) 232-5118. Medical & pharmaceutical uses for cucurbit secondary plant products; cultural practices to enhance yield.
24. **Cui, Hongwen** Department of Horticulture, Northwestern Agricultural Univ., Yangling, Shaanxi 712100, P.R. China
25. **Dane, Fenny** Dept. Horticulture, Auburn University, Auburn, AL 36849. Tel: (205) 844-3047.
26. **de Groot Erik** Breeding, SEmenti Nunhems S.R.L., Via Ghiarone, 2, 40019S. Agata Bolognese, Italy.
27. **de Ruiter, A.C.** de Ruiter Zonen CV, Postbus 4, 2665 ZG Bleiswijk, The Netherlands. Tel: 18G2-2741. Breeding and seed production of cucumbers.
28. **Della Vecchia, Paulo T.** Av Das Nacoes 68 Jardim Europa, CP 12900 Braganca, Paulista SP, Brazil. Tel: 011-433-7447. Breeding & genetics.
29. **Denlinger, Phil** Mt. Olive Pickle Co. Inc., P.O. Box 609, Mount Olive, NC 28365. Tel: (919) 296-1996.

30. **Dhaliwal, Major Singh** Dept. of Vegetable Crops, L.S. & F. Punjab Agriculture University, Ludhiana-141001, Punjab, India
31. **DiNitto, Louis** Sunseeds, 8850 59th Ave., N.E., Brooks, OR, 97305
32. **Drowns, Glenn** Sand Hill Preservation Center, 1878 230th Street, Calamus, IA 52729. Tel: (515) 246-2299.
33. **Dumlao, Rosa** Harris Moran Seed Co.4331 C ockroach Bay Road, Ruskin, FL 33570-2612
34. **Dunlap, James R.** Texas Agric. Expt. Ststion, 2415 E. Highway 83, Weslaco, TX 78596. Tel: (512) 968-5585. FAX: (512) 968-0641. Stress physiology and senescence.
35. **Eigsti, Orié J.** 1602 Winsted, College Green, Goshen, ID 46526. Tel: (219) 533-4632.
36. **El Jack, Ali Elamin** Dept. Horticulture, Fac. Agric. Sciences, University of Gezira, Wad-Medani, P.O. Box 20, Sudan.
37. **El-Downey, Hamdy Hassan Ali** c/o Ms. A. El-Menshawy, Foreign Agric. Relations, Ministry of Agriculture, Nady El-Seid Street, Dokki, Cairo,. Egypt. Cucurbit breeding program, including: diseases (virus, fungal), salinity, greenhouses, hybrids.
38. **Elmstrom, Gary.** University of Florida, Central Florida Res & Educ Center, 5336 University Avenue, Leesburg, FL 34748-8232. Tel: (904) 78-3423. FAX: (904) 392-8940
39. **Fanourakis, Nicholas E.** Technological Educational Institute, Heraklion Crete, 71500, Greece.
40. **Funakushi, Hisashi** Mikado Seed Growers Co., Ltd., 1203 Hoshikuki, Chuo-Ku, Chiba City 260, Japan. Tel: 81-265-4847. FAX: 81-43-266-6444.
41. **Gaba, Victor** Dept. Virology, The Volcani Center, P.O.B. 6, Bet Dagan 50250, Israel. Tel: 972-3-9683568/9. FAX: 972-3-9604180. E-mail: VPGABA&VOLCANI.Tissue Culture & Transformation of melon.
42. **Gabert, August C.** Sunseeds Genetics, Inc. 8850 59th Avenue NE, Brooks, OR 97305-9625. Tel: (503) 393-3243
43. **Gaggero, James M.** Micro Flo, 719 2nd Street #12, Davis, CA 95616
44. **Gautier, Granes B.P.** No. 1, 13630, Eyragues, Franceeurs Grainiers, B. P No. 1, 13630 Eyragues, France. Tel: 90.94.13.44. FAX: 90.92.83.96
45. **Goblet, J.P.** Phytotec SA, Chausse Romaine, 77, 5030 Gemblous, Belgium.
46. **Gonon, Yves.** Mas de Rouzel, Route de Generac, 3000 Nimes, France. Tel: 66.84.21.32, FAX: 66.29.87.81.
47. **Groenewal, Irma** ENZA ZADEN, De Enkhuizer Zaadh, b.v., Postbox 7, 1600 AA Enkhuizen, The Netherlands. Tel: 02280-1-58-44. FAX: 02280-1-59-60.
48. **Groff, David.** Asgrow Seed Company, R. R. #1, Tifton, GA, 31794. Tel: (912) 386-8701. Cucumber and squash breeding.
49. **Grumet, Rebecca** Department of Horticulture, Plant and Soils Building, Michigan State University, East Lansing, MI 48824-1325. Tel: (517) 353-5568. Genetic engineering of disease resistance; vegetable crop improvement; plant transformation.
50. **Hagihara, Toshitsugu** Hagihara-Noujou 984, Hokgi, Tawaramotocho, Siki-gun, Nara-ken, Japan.
51. **Han, Sang Joo** Seoul Seed Co., Chongil Bldg., 736-17 Yeoksam-Dong, Kangnam-gu, Seoul, Korea. Tel: (2) 569-7147. FAX: (2) 552-9439. Vegetable breeding and seed production.
52. **Hassan, Ahmed Abdel-Moneim** Department of Horticulture, Faculty of Agriculture, Cairo University, Giza, Egypt.
53. **Havey, Michael J.** USDA/ARS, Department of Horticulture, University of Wisconsin, Madison, WI, 53706. Tel: (608) 262-1830.
54. **Henderson, W. R.** Department of Horticultural Science, Box 5216, North Carolina State University, Raleigh, NC. 27650-5216. Tel: (919) 737-3167. Breeding and genetics of watermelon.
55. **Herman, Ran** "Zeraim" Seed Growers Company Ltd., Department of Breeding, Gedera 70 700, Israel. Tel: 08-59 27 60. FAX: 08-59 43 76.
56. **Herrington, Mark Edward.** 21 Warner Street, Welling Point, Queensland 4160, Australia. Tel: 07 2861488.
57. **Hertogh, K.** Nickerson-Zwaan b.v., Postbus 19, 2990 AA Barendrecht, The Netherlands.
58. **Himmel, Phyllis** Asgrow Seed Company, 500 Lucy Brown Lane, San Juan Bautista, CA 95045
59. **Hirabayashi, Tetsuo** Nihon Horticultural Production Institute, 207 Kamishki, Matsudo-shi, Chiba-ken, Japan. Tel: 0473-87-3827. FAX: 0473-86-1455. Varietal improvementof cucurbit crops, especiallymelon, cucumber and pumpkin.
60. **Hollar, Larry A.** Hollar & Co., Inc., P. O. Bos 106, Rocky Ford, CO 81067. Tel: (719) 254-7411. FAX: (719) 254-3539. Cucurbit breeding and seed production.
61. **Holle, Miguel** Choquehuanca 851, Lima 27, Peru. Tel: 220474. Plant genetic resources.
62. **Humaydan, Hasib** Ag Consulting International, 317 Red Maple Drive, Danville, CA. 94506
63. **Hung, Lih** National Taiwan Univ. College Agric., Dept. Horticulture, Vegetable Crops Lab., Taipei, Taiwan 107, Republic of China.
64. **Hutton, Mark** Alf Christianson, P.O. Box 304, Woodstown, NJ 08098. Tel: (609) 769-0393. FAX: (609) 769-0393.
65. **Ibrahim, Aly M.** USREP/JECOR/AGWAT, Unit 61306, Box 43, APO, AE, 09308-1306.
66. **Ignart, Frederic** Eta TEZIER Centre de Recherche, Domaine de Maninet, Route de Beaumont, 26000 Valence, France.

67. **Iida, Akira** Minowa Noen, 63-1 Ichieda-cho, Yamato-Kohriyama City, Nara Pref., Japan, T639-11.
68. **Iida, Shuichi** Inst. Radiation Breeding, NIAR, MAFF, P.O. Box 3, Ohmiya-Machi, Naka-Gun, Ibaraki-Ken, 319-22, Japan.
69. **Ikeda, Satoru** Sakata Seed America, Research Station, 105 Boronda Road, Salinas, CA 93907. Tel: (408) 758-0585. FAX: (408) 758-0908
70. **Jain, Jaagrati** Division of Vegetable Crops, Indian Agricultural Research Institute, New Delhi - 110012, India. Tel: 583628. Muskmelon genetics.
71. **Jaramillo-Vasquez, Juan Jefe** Regional Seccion Hortalizas, Instituto Colombiano Agropecuario, AA 233, Palmira, Colombia.
72. **Kaminimura, Shoji** 421-19 Furuichi-machi, Macbashi City, Gunma-ken 371, Japan.
73. **Kampmann, Hans Henrick** Breeding Station Danefield, Odensevej 82, 5290, Marslev, Denmark.
74. **Kanno, Tsuguo** Kurume Branch, NRI Vegetables, Orn. Plants & Tea, Min. Agric. Fishery & For., 1823, Mii-cho, Kurume, Fukuoka, 830, Japan.
75. **Karchi, Zvi** Div. Vegetable Crops, Agr. Research Org., Newe Ya'ar Experiment Station, P.O. Haifa, Israel. Tel: 04-833186. FAX: 972-3-9665327.
76. **Kaswari, Mahmoud** Department of Plant Production, University of Jordan, Amman, Jordan. Tel: 962-6-842167. FAX: 962-6-832318
77. **Katsiotis, Andreas** Dept. Agronomy, Univ. Wisconsin-Madison, 1575 Linden Drive, Madison, WI 53706-1597. Tel: (608) 262-7452.
78. **Kirkbride, Joseph H. Jr.** USDA-ARS-PSI, Systemic Botany & Mycology Lab, Bldg. 265, BARC-East, Beltsville, MD 20705. Tel: (301) 504-9447, E-mail: JKIRKBRIDE&ASRR.ARSUSDA.GOV.
79. **Knerr, Larry D.** Ferry Morse Seed Co., 1570 County Road, 951, W. Naples, FL, 33999. Tel: (813) 455-1817. Protein level biochemical diversity and genetics of herbicide resistance.
80. **Kraakman, Peter B** Torre Verde 7-2, Aguadulce, Roquetas De Mar, Spain
81. **Kuginuki, Uashuhisa** Ntl. Research Inst. Vegetables, Orn. Plants & Tea, Ano. Age-Gun, Mic. Japan 514-23.
82. **Kuti, Joseph O.** College of Agriculture, Horticulture Res. Lab., Texas A&L University, Kingsville, TX, 78363. Tel: (512) 595-3711. FAX: (512) 595-3713. Breeding and genetics; host-parasite interrelationships; postharvest physiology.
83. **Kwack, Soo-Nyeon** Department of Horticultural Breeding, Mokpo Natl. Univ., Dorimri, Chonggyemyun, Muangun, Chonnam 534-729, Korea
84. **Kyle, Molly** Cornell Univ., Dept. Plant Breeding & Biometry, 306 Bradfield Hall, Ithaca, NY, 14853-1902. Tel: (607) 255-8147. Breeding for disease resistance.
85. **Ladd, Krystyna M.** 8420 Kelton Drive, Gilroy, CA 95020
86. **Lari, Andrea** ORIS S.p.A., Via Vittorio Veneto, 81, 20090 Salerano Sul Lambro, Milano, Italy, Melon, squash and watermelon breeding.
87. **Layton, Jeanne G.** Monsanto Co., Mail Zone: GG4H, 700 Chesterfield Village Parkway, St. Louis, MO, 63198. Tel: (317) 537-7158.
88. **Leaver, Chris** Dept. Plant Science, Univ. Oxford, South Parks Road, Oxford OX1 3RB, England
89. **Lecouvior, Michel** Clause Semences Professionelles, 24, boulevard P. Brossolette, 91221 Bretainy-sur-Orge, France
90. **Lehmann, Louis Carl** Svalof AB, S-268 00 Svalov, Sweden.
91. **Levy, Roni** Rua General Antonio Fernandes, Dantes No. 221, Bell, Horizonte, Assu, R/N 59-650-000, Brazil. Fertigation of cucurbits in sandy soils (and effects on shelf life); pollination effects on fruit deformities.
92. **Lim, Haktae** Department of Horticulture, Kangnung National University, Kangnung, Kangwon-Do, South Korea, 210-702. Plastid gene regulation; organelle genetics; RFLP mapping; somatic hybridization.
93. **Lin, Depei** Xinjing August 1st Agricultural College, Department of Horticulture, Urumqi, 830052, People's Rep. China.
94. **Love, Stephen L.** University of Idaho, Research and Extension Center, Aberdeen, ID 83210. Tel: (208) 397-4181. Breeding, culture and management; mutation breeding.
95. **Lower, Richard. L.** College of Agriculture, Univ. of Wisconsin, 1450 Linden Drive, Room 136, Madison, WI 53706. Tel: (608) 262-2349, FAX: (608) 262-4556. E-mail: richard.lower@mail.admin.wisc.edu. Effects of plant type genes on yield, sex-expression, growth.
96. **Loy, J, Brent** Plant Biology Department, Nesmith Hall, University of New Hampshire, Durham NH 03824. Tel: (603) 862-3216. FAX: (603) 862-4757. Squash, melon, pumpkin. Genetics, breeding, plasticulture, mulch, row-covers.
97. **Ma, Dewei** Department of Horticulture, Hebei Agricultural University, Bao Ding, Hebei 071000, P.R. China.
98. **Mackay, Wayne A.** 1380 A&M Circle, El Paso, TX, 79927. Tel: (915) 859-9111.
99. **Maluf, Wilson Roberto** Escola Superior de Agricultura de Lavras, Departamento de Agricultura, Caixa Postal 37, CEP 27300 - Lavras - MG, Brazil. Tel: (035) 821-3700.
100. **Maneesinthu Likhit** Chia Tai Company Limited, 299-301 Soingsawad Road, Bangkok 10100, Thailand. Tel: 02-2338191-9. FAX: 662-2371540. Breeding and seed production of open-pollinated and hybrid cucurbits.

101. **McCright, J. D.** USDA-ARS, 1636 E. Alisal St., Salinas, CA 93915. Tel: (408) 755-2864, FAX: (408) 753-2866. Melon breeding and genetics.
102. **McFerson, James R.** Germplasm Resources Unit, New York St. Agric. Experiment Sta., Geneva, NY. 14456-0462. Tel: (315) 787-2393. FAX: (315) 787-2397. Conservation and utilization of genetic resources.
103. **McGrath, D.J.** Dept. Primary Industries, Hortic. Research Sta., P.O. Box 538, Bowen. 4805. Queensland, Australia .
104. **Meadows, Mike** Rogers NK Seed Co., 10290 Greensway Road, Naples, FL 33961.
105. **Merrick, Laura C.** Department of Plant & Soil Sciences, 105 Deering Hall, University of Maine, Orono, ME 04469. Tel: (207) 581-2950. FAX: (207) 581-2199. Crop evolution, germplasm evaluation and conservation, genetic resources of cucurbits.
106. **Milotay, Peter** Vegetable Crops Research Institute, P. O. Box 116, Kecskemet, 6000, Hungary.
107. **Miranda, Baldwin** Rogers NK Seed Co., 10290 Greenway Road, Naples, FL 33961. Tel: (813) 775-4090. FAX: (813) 774-6852.
108. **Mochizucki, Tatsuya** Kurume Br, Natl Res Inst Veg Ornament Plant & Tea, 1823 Mii-machi, Kurume, Fukuoka 830, Japan.
109. **Montiero, Antonio A.** Section of Horticulture, Inst. Superior de Agronomia, Tech. Univ. Lisbon, Portugal. Tel: 351-1-3638161. FAX: 351-1-3635031. Melon protection & cultivation in adverse environments.
110. **Moraghan, Brian J.** Asgrow Seed Co. P.O. Box 667, Arvin, CA 93203. Tel: (805) 854-2360. Melon and watermelon breeding and disease resistance.
111. **More, T.A.** Dept. Vegetable Crops, Indian Agricultural Research Institute, New Delhi, 110012, India.
112. **Morelock, Ted** Dept. Horticulture & Forestry, University of Arkansas, Fayetteville, AR, 72701. Tel: (501) 575-2603.
113. **Muhyi, Rejah I.** Petoseed Woodland Research Station, 37437 State Highway 61, Woodland, CA 95695. Tel: (916) 666-0931.
114. **Munger, H. M.** Cornell University, 410 Bradford Hall, Ithaca, NY 14853. Tel: (607) 255-1661. FAX: (607) 255-6683. Cucurbit breeding and disease resistance.
115. **Murdock, Brent A.** Clemson University, Department of Horticulture, Clemson, SC 29634-0375.
116. Tel: (803) 296-1871. FAX: (803) 656-4960. Watermelon breeding: genetic improvement of neglected tropical vegetables.
117. **Nagai, Hiroshi** Instituto Agronomico, Cx. Postal 28, 13.100-Campinas, Sp., Brazil.
118. **Navazio, John** Dept. Hort., 1575 Linden Dr., University of Wisconsin, Madison, WI 53706. Tel: (608) 238-8567. Breeding for stress tolerance, postharvest physiology, improved cultivars and inbreds.
119. **Nechama Shulamit** Breeding Department, Mivhor Farm, Post Sde Gat 79570, Israel .
120. **Ng, Timothy J.** Department of Horticulture, University of Maryland, College Park, MD 20742 . Tel: (301) 405-4345. FAX: (301) 314-9308. E-mail: tng@grad.umd.edu. Melon breeding and genetics; postharvest physiology; seed germination.
121. **Niemirowicz-Szczytt, Katarzyna** UL. Nowoursynowska 166, Dept. Genetics and Plant Breeding, 02-766 Warsaw, Poland. Tel: 430982. Breeding of cucumber, melon, watermelon & squash. Downy mildew res., wide crosses, tissue culture, haploids.
122. **Norton, J.D.** Department of Horticulture, Auburn University, Auburn, AL, 36830. Tel: (205) 844-3031. FAX: (205) 844-3131. Multiple disease resistant melon and watermelon.
123. **Nourizadeh, Saeid** Purdue University, 1165 Horticulture Bldg., West Lafayette, IN 47907-1165. Tel: (317)494-1300. FAX: (317) 494-0391.
124. **Nuez, Fernando** Catedra de Genetica, E.T.S. Ingenieros Agronomos, Universidad Politecnica, Camino de Vera,14, Valencia-22, Spain.
125. **Nugent, Perry** USDA-ARS, U.S. Vegetable Laboratory, 2875 Savannah Highway, Charleston, SC, 29414. Tel: (803) 556-0840. Melon and watermelon inheritance studies, pest resistance, stress resistance, and fruit quality.
126. **Om, Young-Hyuan** Horticulture Experiment Station, 475 Imok Dong, Suwon 440-310, Republic of Korea. Tel: 82-331-41-5931. FAX: 82-331-46-7148.
127. **Omara, Sadig Khdir** Dept. Horticulture, Fc. Agric. Sciences, University of Gezira, Wad-Medani, P.O. Box 20, Sudan.
128. **Oridate, Toshiroh** 15 Karasawa, Minami-ku, Yokohama-shi, Kanagawa-ken, Japan.
129. **Ortea, Sergio Garza** Universidad de Sonora, Dept. de Agricultura y Ganaderia, A.P. Postal 1853, Hermosillo, Sonora, Mexico. Breeding of Cucurbita spp.; testing of new muskmelon lines.
130. **Ouyang, Wei** Petoseed Woodland Research Station, 37437 State Highway 16, Woodland, CA, 95695.
131. **Owens, Ken.** PetoSeed Woodland Research Station, 37437 State Highway 16, Woodland, CA 95695. Tel: (916) 666-0931. Cucumber breeding.
132. **Palomares, Gloria.** Dept. de Biotecnologia, Univ. Politecnica, Camino e Vera, 14, 46022 Valencia, Spain.
133. **Paris, Harry** Division of Vegetable Crops, Agr. Research Org., Neve Ya'ar Expt. Station, P. O. Haifa, Israel . Tel: 972-4-894516. Breeding and genetics of cucurbits.

134. **Peter, K.V.** National Ressearch Centre for Spices, ICAR, Post Bag No. 1701, Marikunnu P.O. Calicut - 673 012, Kerala, India.
135. **Picard, Florence** Vilmorin, La Menitre, 49250 Beaufort-en-Vallee, France.
136. **Pierce, Lawrence** 3091 Lynview Drive, San Jose, CA, 95148. Tel: (408) 258-0307.
137. **Pierce, Vicki** 1583 Endicott Drive, San Jose, CA 95122 Tel: (408) 258-0307.
138. **Pitrat, Michel** Centre de Recherches Agronomiques de Avignon, Stat d'Amelior des Plantes Mar, Domaine St-Maurice, 841430 Montfavet, France.
139. **Pootstchi, Iraj** 97 St. Marks Road, Henley-on-Thames RG9 1LP, England.
140. **Price, E. Glen** American Sunmelon Research Center, P.O. Box 153, Hinton, OK 73047. Tel: (405) 542-3456. FAX: (405) 542-3457. Seedless watermelon; polyploidy, genetics, breeding, cytogenetics.
141. **Provvidenti, Rosario** Department of Plant Pathology, NYAES, Cornell University, Geneva, NY 14456-0462
142. **Punja, Zamir K.** Dept. Biological Sciences, Simon Fraser University, Burnaby, B.C. V5A 1S6, Canada.
143. **Quemada, Hector** The Upjohn Company, 9612-50-1, Kalamazoo, MI 49001. Tel: (616) 384-2642. FAX: (616) 384-2725.
144. **Raharjo, Simon H.T.** Dept. Biological Sciences, Simon Fraser University, Burnaby, B.C. V5A 1S6, Canada. Tel: (604) 291-3090. FAX: (604) 291-3496. Tissue culture and genetic transformation of cucumber.
145. **Ray, Dennis** Department of Plant Sciences, University of Arizona, Tucson, AZ 85721. Tel: (602) 621-7612. FAX: (602) 621-7186. New crops; cytogenetics.
146. **Reitsma, Kathy** NC Regional Plant Introduction Station, Iowa State University, Ames, IA, 50011. Tel: (515) 292-6502. FAX: (515) 294-4880.
147. **Reuling, G.** Nunhens Zaden B.V., P.O. Box 4005, 6080 Haelen, The Netherlands, Tel: 04759-9222. FAX: 04759-9223/5104.
148. **Rhodes, Billy B.** Clemson Univ./Horticulture, Poole Agricultural Center, Clemson, SC 29634-0375. Tel: (803) 656-0410. Watermelon genetics, breeding, micropropagation, disease resistance, male sterility, triploids.
149. **Risser, Georgette** Centre de Recherches Agronomiques de Avignon, Stat d'Amelior des Plantes Mar, Domaine St Maurice, 84140 Montfavet, France.
150. **Robinson, R. W.** Department of Horticultural Sciences, New York State AES, Geneva, NY 14456-0462 . Tel: (315) 787-2237. FAX: (315) 787-2397. Breeding and genetics of cucurbits.
151. **Robledo, C.** V.R.E.D., Centre de Recherches, Mas d'Aptel, 30510 Generac, France. Tel: 66 01 89 07.
152. **Rodriguez, Jose Pablo** 25 De Mayo 75, 2930-San Pedro, Buenos Aires, Argentina.
153. **Roig, Luis A.** Departamental Biotechnology, E.T.S.Ingenieros Politecnica, Camino de Vera 14, 46022-Valencia, Spain. Tel: 34(6) 3877424. FAX: 34(6) 3877429.
154. **Rovelo, Claudia** Horticulture Research Laboratory, Coll. Agriculture, Box 156, Texas A&I Univ., Kingsville, TX, 78363.
155. **Rumsey, Anthony E.** New World Seeds Pty Ltd., P.O. Box 18, Dural 2158, 22-24 Crosslands Road, Galston, N.S.W., Australia.
156. **Scheirer, Douglas M.** Nestle USA/Libby Division, P. O. Box 198, Morton, IL 61550. Tel: (309) 263-2651. Processing pumpkin; breeding and cultural practices.
157. **Schnock, Martin G.** Norsingen, Fridolin-Mayer-Strasse 5, D-7801 Ehrenkirchen, Fed. Rep. Germany. Tel: 07633-13095.
158. **Schroeder, Robert H.** Harris Moran Seed Co., R.R. 1, Box 1243, Davis, CA. 95616. Tel: (916) 756-1382. FAX: (916) 756-1016. Cucurbit genetics and breeding; germplasm evaluation and utilization.
159. **Sekioka, Terry T.** Kauai Branch Station, University of Hawaii, Kapaa, HI 96746. Tel: (808) 822-4984. Cucumber and bitter melon breeding; disease resistance.
160. **Semillas Fito, S.A.** c/. Selva de Mar, 111,08019, Barcelona, Spain.
161. **Shann, David A.** Arthur Yates & Co., Pty. Limited, Research Farm, Burroway Road, Naromine, N.S.W. 2821, Australia. Tel: (68) 89-1144.
162. **Shiffris, Oved** 21 Walter Avenue, Highland Park, NJ 08904.
163. **Shiga, Toshio** Plant Biotech. Ctr., Sakata Seed Corp., 358 Uchikoshi, Sodegaura, Chiba, 299-02 Japan. Tel: 0438-75-7276.
164. **Shintaku, Yurie** 2-10-2, Shimizu, Suginami-ku, Tokyo, 167, Japan.
165. **Simon, Philipp W.** 5125 Lake Mendota Drive, Madison, WI 53705. Tel: (608) 264-5406. FAX: (608) 262-4743. Breeding and genetics.
166. **Skirvin, Robert M.** Univ. Illinois, Dept. Horticulture, 1707 S. Orchard St. Urbana, IL 61801. Tel: (217) 333-1530. Micropropagation, somaclonal variation.
167. **Snyder, Jim** PetoSeed Co., Inc. RR2, Box 80A, Bridgeton, NJ, 08302-8723. Tel: (609) 451-6231.
168. **Song, Jin-Soo** Plant Breeding & Res. Inst., Nong-Woo Seeds, 387-2 Sasa-2Ri, Panwol, Whasong, Kyonggi, 445-820, Republic of Korea. Tel: 0345-0466.

169. **Staub, Jack E.** USDA, ARS, Dept. Horticulture, Univ. Wisconsin, Madison, WI 53706-1590. Tel: (608) 262-0028. FAX: (608) 262-4743. Cucumber breeding & genetics, physiology, biochemical genetic markers, evolution, environmental stress.
170. **Stern, Joseph** Royal Sluis Inc., 910 Duncal Road, San Juan Bautista, CA 95045.
171. **Stevens, M. Allen** Petoseed Woodland Research Station, 37437 State Highway 16, Woodland, CA, 95695. Tel: (916) 666-0931.
172. **Tasaki, Seikoh** AOS/Q-O2, Bloco-E, Apt. 603, CEP-70660, Brasilia, DF, Brazil.
173. **Tatlioglu, Turan** Institut of Applied Genetics, Univ. Hannover, Herrenhauser Str. 2, 3000 Hannover, Germany.
174. **Taurick, Gary** Ferry Morse Seed Company, P.O. Box 392, Sun Praris, WI 53590. TEL: (608) 8370-6573. FAX: (608) 837-3758. Population improvement and hybrid development for cucumber and summer squash.
175. **Teppner, Herwig** Karl-Franzens-Universitat Graz, Institut fur Botanik, Holteigasse 6, A-8010 GRAZ, Austria.
176. **Thomas, Claude E.** USDA-ARS, U.S. Vegetable Labaratory, 2875 Savannah Highway, Charleston, SC 29407.
177. **Thomas, Paul** PetoSeed Woodland Research Station, 37437 State Highway 16, Woodland, CA 95695. Tel: (916) 666-0931.
178. **Tolla, Greg** Campbell Research & Development. Napoleon, OH, 43545. Tel: (419) 592-8015. Development of pickling cucumber varieties.
179. **Trulson, Anna** Petoseed Woodmand Research Station, 37437 State Highway 16, Woodland, CA 95695.
180. **van Deursen, S.** Sluis & Groot Research, Blaker 7, 2678 L W de Lier, The Netherlands
181. **van Kooten, Hank** Bruinsma Selectiebedrijven B.V., P.O. Box 24, 2670 AA Naaldwijk, The Netherlands.
182. **van Leeuwen, Loes** Sementi Nunhems, Via Ghiarone, 2, 40019 S. Agata Bolognese, Italy.
183. **Vecchio, Franco** Pioneer Hi-Bred Italia SpA, via Provinciale 42/44, 43018 Sissa (PR), Italy.
184. **Walters, Deena Decker** Fairchild Tropical Garden Science Center, 11935 Old Cutler Road, Miami, FL, 33156. Tel: (305) 665-2844. FAX: (301) 665-8032.
185. **Walters, Terrence** Fairchild Tropical Garden Science Center, 11935 Old Cutler Road, Miami, FL, 33156. Tel: (305) 665-2844. FAX: (301) 665-8032.
186. **Wann, E. Van** USDA-ARS, P.O. Box 159, Lane, OK, 74555. Tel: (405) 889-7395.
187. **Warid, Warid A.** Paseo de las Fuentes No. 18, Col. Valle Verde, 83200 Hermosillo, Sonora, Mexico.
188. **Wasilwa, Lusike** Dept. Horticulture & Forestry, University of Arkansas, Fayetteville, AR, 72701.
189. **Watterson, Jon** Petoseed Woodland Research Station, 37437 State Highway 16, Woodland, CA 95695.
190. **Wehner, Todd C.** Department of Horticultural Science, Box 7609, North Carolina State University, Raleigh, NC 27695-7609. Tel: (919) 515-5363. FAX: (919) 515-7747. E-mail: Todd_Wehner@ncsu.edu. Cucumber genetics and breeding; yield, earliness, quality, disease, cold tolerance.
191. **Wessel-Beaver, Linda** Department of Agronomy & Soils, College of Agriculture, Univ. Puerto Rico, Mayaguez, PR, 00708. Tel: (809) 832-4040. Pumpkin & squash breeding; disease resistance; insect resistance.
192. **Whiteaker, Gary** Nunhems Seed Corporation, 221 East Main Street, Lewisville, ID 83431. Tel: (208) 754-8666. FAX: (208) 754-8669.
193. **Williams, Tom V.** Rogers NK Seed Co., 10290 Greenway Road, Naples, FL 33961. Tel: (813) 775-4090. FAX: (831) 774-6852. Watermelon breeding.
194. **Winkler, Johanna** Saatzzucht Gleisdorf, Ges. m.b.g. & Co. KG, Am Tioeberhof 33, 8200 Gleisdorf, Austria.
195. **Wolff, David W.** Texas A&M Experiment Station, 2415 East Hwy. 83, Weslaco, TX 78596-8399. Tel: (512) 968-5585.
196. **Wu, Mingzhu** Hort. Inst. Xinjiang Acad. Agric. Sciences. Nanchang Rod NO. 38, Urumqi, Xinjiang, People's Rep. China.
197. **Wunderlin, Richard P.** Dept. Biology, University of South Florida, 4202 East Fowler Ave., LIF 169, Tampa, FL, 33620-5150. Tel: (813) 974-2359. FAX: (813) 874-3557. Systematics of neotropical species: Zanonioideae.
198. **Wyatt, Colen** Petoseed Woodland Research Station, 37437 State Highway 16, Woodland, CA 95695.
199. **Yamanaka, Hisako** Yamato-Noen Co., Ltd. 100, Byodobo-cho, Tenri-City NARA, Japan 632. Tel: 07436-2-1182.
200. **Yan, Yin** Institute of Vegetable & Flower, Chinese Academy of Agricultural Science, Beijing 100081, P.R. China.
201. **Yang, Si-Lin** Dept. Horticulture, Southwest Agric. Univ., Bei-bei, Chong-qing, Si-chuan 630716, P.R. China. Ethnobotany, crop evolution, genetic resources (wild & cultivated) of Asian Cucumis, Benincase, Momordica.
202. **Yorty, Paul** Rogers NK Seed Co., P.O. Box 104, Twin Falls, ID, 83303-0104. Tel: (208) 733-0077. Cucurbit breeding.
203. **Yukura, Yasou** 46-7, 3-Chome, Miyasaka, Setagaya-Ku, Tokyo, Japan.
204. **Zhang, Jiannong** Melon Research Institute, Gansu University of Agriculture, Lanzhou, Gansu, 730070, P.R. China.
205. **Zhang, Zingping** Department of Horticulture, Clemson University, Clemson, SC, 29634-0375.
206. **Zhao, Yanru** Beijing Vegetable Research Center, P.O. Box 2443, Beijing 100081, P.R. China.
207. **Zink, Frank** Department of Vegetable Crops, University of California, Davis, CA, 95616. Tel: (916) 79504967. Plant breeding and disease resistance.
208. **Zitter, Thomas** Cornell Univ., Dept. Plant Pathology, 334 Plant Science Building, Ithica, NY 14853-5908.

Geographical Distribution of CGC Members in the United States

Cucurbit Genetics Cooperative

- **Alabama**
 - Fenny Dane
 - J.D. Norton
- **Arizona**
 - Dennis Ray
- **Arkansas**
 - Ted Morelock
 - Lusike Wasilwa
- **California**
 - G.W. Bohn
 - Al Burkett
 - Paul Chung
 - Timothy J. Close
 - James M. Gaggero
 - Phyllis Himmel
 - James C. Hollar
 - Hasib Humaydan
 - Satoru Ikeda
 - Krystyna M. Ladd
 - J.D. McCreight
 - Brian J. Moraghan
 - Rejah I. Muhyl
 - Wei Ouyang
 - Ken Owens
 - Lawrence Pierce
 - Vicki Pierce
 - Robert .H. Schroeder
 - Joseph Stern
 - M. Allen Stevens
 - Paul Thomas
 - Anna Trulson
 - Jon Watterson
 - Colen Wyatt
 - Frank Zink
- **Colorado**
 - Larry A. Hollar
- **Florida**
 - Rosa Dumlao
 - Gary Elmstrom
 - Larry D. Knerr
 - Mike Meadows
 - Baldwin Miranda
 - Deena Decker Walters
 - Tom V. Williams
 - Richard P. Wunderlin

- **Georgia**
 - David Groff
- **Hawaii**
 - Fure-Chyi Chen
 - Terry T. Sekioka
- **Idaho**
 - Steven L. Love
 - Gary Whiteaker
 - Paul Yorty
- **Illinois**
 - Douglas M. Scheirer
 - Robert M. Skirvin
- **Indiana**
 - Orie J. Eigsti
 - Saeid Nourizadeh
- **Iowa**
 - Glenn Drowns
 - Kathy Reitsma
- **Kansas**
 - C.D. Clayberg
- **Kentucky**
 - M. Brett Callaway
- **Maine**
 - Laura C. Merrick
- **Maryland**
 - Joseph H. Kirkbride, Jr.
 - Timothy J. Ng
- **Michigan**
 - Rebecca Grumet
 - Hector Quemada
- **Missouri**
 - Jeanne G. Layton
- **Mississippi**
 - Edward M. Croom
- **Nebraska**
 - Dermot P. Coyne
- **New Hampshire**
 - R. Bruce Carle
 - J. Brent Loy
- **New Jersey**
 - Mark Hutton
 - Oved Shifriss
 - Jim Snyder
- **New York**
 - Edward E. Carey
 - Aly M. Ibrahim
 - Molly Kyle
 - James R. McFerson
 - H.M. Munger
 - Rosario Provvidenti
 - R.W. Robinson
 - Thomas Zitter
- **North Carolina**
 - Mary Barbercheck
 - Phil Denlinger
 - W.R. Henderson
 - Todd C. Wehner

- **Ohio**
 - Greg Tolla
- **Oklahoma**
 - E. Glen Price
 - E. Van Wann
- **Oregon**
 - Louis Di Nitto
 - August C. Gabert
- **Puerto Rico**
 - Linda Wessel-Beaver
- **South Carolina**
 - Brenta A. Murdock
 - Perry Nugent
 - Billy B. Rhodes
 - Claude E. Thomas
 - Xingping Zhang
- **Texas**
 - James R. Dunlap
 - Joseph O. Kuti
 - Wayne A. Mackay
 - Claudia Roveló
 - David W. Wolff
- **Wisconsin**
 - Michael J. Havey
 - Andreas Katsiotis
 - R.L. Lower
 - John Navazio
 - Philipp W. Simon
 - Jack E. Staubb
 - Gary Taurick

Geographical Distribution of CGC Members Outside the USA

- **Argentina**
 - Jose Pablo Rodriguez
- **Australia**
 - Mark Edward Herrington
 - D.J. McGrath
 - Anthony E. Rumsey
 - David A. Shann
- **Austria**
 - Herwig Teppner
 - Johanna Winkler
- **Belgium**
 - J.P. Goblet
- **Brazil**
 - Paulo T. Della Vecchia
 - Roni Levy
 - Wilson Roberto Maluf
 - Hiroshi Nagai
 - Seikoh Tasaki
- **Canada**
 - Zamir K. Punja
 - Simon H.T. Raharjo
- **China, People's Rep.**
 - Hongwen Cui
 - Dewei Ma
 - Si-Lin Yang
 - Jiannong Zhang
 - Yanru Zhao
 - Depei Lin
 - Mingzhu Wu
- **China, Republic of**
 - Lih Hung
- **Columbia**
 - Juan Jaramillo-Vasquez
- **Denmark**
 - Hans Henrik Kampmann
- **Egypt**
 - Hamdy Hassan Ali El-Doweny
 - Ahmed Abdek-Moneim Hassan
- **England**
 - Gary K. Bradbury
 - Chris Leaver
 - Iraj Poostchi
- **France**
 - Sofia Ben Tahar
 - Daniel Chambonnet..
 - Bernard Charpiot
 - Graines Gautier
 - Yves Gonon

- Frederic Ignart
- Michel Lecouviour
- Florence Picard
- Michael Pitrat
- Georgette Risser
- C. Robledo
- **Germany**
 - Martin G. Schnock.
 - Turan Tatliglu.
- **Greece**
 - Nicholas E. Fanourakis.
- **Hungary**
 - Peter Milotayr
- **India**
 - Major Singh Dhaliwal
 - Jaagrati Jain
 - T.A. More
 - K.V. Peter
- **Israel**
 - Yigal Cohan
 - Victor Gaba
 - Ran Herman
 - Zvi Karchi
 - Shulamit Nechama
 - Harry Paris
- **Italy**
 - Erik de Groot
 - Andrea Lari
 - Loes van Leeuwen
 - Franco Vecchio
- **Japan**
 - Hisasgu Funakushi
 - Toshitsugu Hagihara
 - Tetsuo Hirabayashi
 - Akira Iida
 - Shuichi Iida
 - Shoki Kamimura
 - Tsuguo Kanno
 - Yasuhisa Kuginuki
 - Tatsuya Mochizuki
 - Toshiroh Oridate
 - Toshio Shiga
 - Yurie Shintaku
 - Hisako Yamanaka
 - Yaso Yukura
- **Jordan**
 - Mahmoud Kaswari
- **Korea , Republic of**
 - Sang, Joo Han
 - Soo Nyeon Kwack
 - Haktae Lim
 - Young-Hyun Om
 - Jin-Soo Song
- **Mexico**
 - Pedro Cano Rios
 - Sergio Garza Ortega
 - Warid A.Warid

- **The Netherlands**
 - A.G.B Beekman
 - P.A Boorsma
 - A.C. de Ruyter
 - Irma Groenewal
 - K. Hertogh
 - G. Reuling
 - S. van Deursen
- **Peru**
 - Miguel Holle
- **Poland**
 - Karatzyna Niemirowicz-Szczytt
- **Portugal**
 - Antonio Monteiro
- **Saudi Arabia**
 - Abdul Mohsen I. Al-Sulaiman
- **Spain**
 - Ma Cruz Ayuso
 - Pilar Corella
 - Peter Kraakman
 - Fernando Nuez
 - Gloria Palomares
 - Luis A. Roig
 - Semillas Fito, S.A.
- **Sudan**
 - Ali Elamin El Jack
 - Sadig Khidir Omara
- **Sweden**
 - Luis Carl Lehmann
- **Thailand**
 - Likhit Maneesinthu

Covenant and By-Laws of the Cucurbit Genetics Cooperative

Article I. Organization and Purposes

The Cucurbit Genetics Cooperative is an informal, unincorporated scientific society (hereinafter designated "CGC") organized without capital stock and intended not for business or profit but for the advancement of science and education in the field of genetics of cucurbits (Family: Cucurbitaceae). Its purposes include the following: to serve as a clearing house for scientists of the world interested in the genetics and breeding of cucurbits, to serve as a medium of exchange for information and materials of mutual interest, to assist in the publication of studies in the aforementioned field, and to accept and administer funds for the purposes indicated.

Article II. Membership and Dues

The membership of the CGC shall consist solely of active members; an active member is defined as any person who is actively interested in genetics and breeding of cucurbits and who pays biennial dues. Memberships are arranged by correspondence with the Chairman of the Coordination Committee.

The amount of biennial dues shall be proposed by the Coordinating Committee and fixed, subject to approval at the Annual Meeting of the CGC. The amount of biennial dues shall remain constant until such time that the Coordinating Committee estimates that a change is necessary in order to compensate for a fund balance deemed excessive or inadequate to meet costs of the CGC.

Members who fail to pay their current biennial dues within the first six months of the biennium are dropped from active membership. Such members may be reinstated upon payment of the respective dues.

Article III. Committees

1. The Coordinating committee shall govern policies and activities of the CGC. It shall consist of six members elected in order to represent areas of interest and importance in the field. The Coordinating Committee shall select its Chairman, who shall serve as spokesman of the CGC, as well as its Secretary and Treasurer.

Approvals: W. Bemis; J.D. Norton; R.W. Robinson; W.R. Henderson; M.L. Robbins; R.L. Lower

2. The Gene List Committee, consisting of five members, shall be responsible for formulating rules regulating the naming and symbolizing of genes, chromosomal alterations, or other hereditary modifications of the cucurbits. It shall record all newly reported mutations and periodically report lists of them in the Report of the CGC. It shall keep a record of all information pertaining to cucurbit linkages and periodically issue revised linkage maps in the Report of the CGC. Each committee member shall be responsible for genes and linkages of one of the following groups: cucumber, *Cucurbita* spp., muskmelon, watermelon, and other genera and species.

3. Other committees may be selected by the Coordinating Committee as the need or fulfilling other functions arises.

Article IV. Election and Appointment of Committees

1. The Chairman will serve an indefinite term while other members of the Coordinating Committee shall be elected for ten-year terms, replacement of a single retiring member taking place every other year. Election of a new member shall take place as follows: A Nominating Committee of three members shall be appointed by the Coordinating Committee. The

aforesaid Nominating Committee shall nominate candidates for an anticipated opening on the Coordinating Committee, the number of nominees being at their discretion. The nominations shall be announced and election held by open ballot at the Annual Meeting of the CGC. The nominee receiving the highest number of votes shall be declared elected. The newly elected member shall take office immediately.

In the event of death or retirement of a member of the Coordinating Committee before the expiration of his/her term, he/she shall be replaced by an appointee of the Coordinating Committee.

Members of other committees shall be appointed by the Coordinating Committee.

Article V. Publications

1. One of the primary functions of the CGC shall be to issue an Annual Report each year. The Annual Report shall contain sections in which research results and information concerning the exchange of stocks can be published. It shall also contain the annual financial statement. Revised membership lists and other useful information shall be issued periodically. The Editor shall be appointed by the Coordinating Committee and shall retain office for as many years as the Coordinating Committee deems appropriate.

2. Payment of biennial dues shall entitle each member to a copy of the Annual Report, newsletters, and any other duplicated information intended for distribution to the membership. The aforementioned publications shall not be sent to members who are in arrears in the payment of dues. Back numbers of the Annual Report, available indefinitely, shall be sold to active members at a rate determined by the Coordinating Committee.

Article VI. Meetings

An Annual Meeting shall be held at such a time and place as determined by the Coordinating Committee. Members shall be notified of time and place of meetings by notices in the Annual Report or by notices mailed not less than one month prior to the meeting. A financial report and information on enrollment of members shall be presented at the Annual Meeting. Other business of the Annual Meeting may include topics of agenda selected by the Coordinating Committee or any items that members may wish to present.

Article VII. Fiscal Year

The fiscal year of the CGC shall end on December 31.

Article VIII. Amendments

These By-Laws may be amended by simple majority of members voting by mail ballot, provided a copy of the proposed amendments has been mailed to all the active members of the CGC at least one month previous to the balloting deadline.

Article IX. General Prohibitions

Notwithstanding any provision of the By-Laws or any other document that might be susceptible to a contrary interpretation:

1. The CGC shall be organized and operated exclusively for scientific and educational purpose.
2. No part of the net earnings of the CGC shall or may under any circumstances inure to the benefit of any individual.
3. No part of the activities of the CGC shall consist of carrying on propaganda or otherwise attempting to influence legislation of any political unit.
4. The CGC shall not participate in, or intervene in (including the publishing or distribution of statements), any political campaign on behalf of a candidate for public office.
5. The CGC shall not be organized or operated for profit.
6. The CGC shall not:

- (a) lend any part of its income or corpus without the receipt of adequate security and a reasonable rate of interest to;
- (b) pay any compensation in excess of a reasonable allowance for salaries or other compensation for personal services rendered to;
- (c) make any part of its services available on a preferential basis to;
- (d) make any purchase of securities or any other property, for more than adequate consideration in money's worth; or
- (e) sell any securities or other property for less than adequate consideration in money or money's worth; or
- (f) engage in any other transactions which result in substantial diversion of income or corpus to any officer, member of the Coordinating Committee, or substantial contributor to the CGC.

The prohibitions contained in this subsection (6) do not mean to imply that the CGC may make such loans, payments, sales, or purchases to anyone else, unless authority be given or implied by other provisions of the By-Laws.

Article X. Distribution on Dissolution

Upon dissolution of the CGC, the Coordinating Committee shall distribute the assets and accrued income to one or more scientific organizations as determined by the Committee, but which organization or organizations shall meet the limitations prescribed in sections 1-6 of Article IX.

Approvals: W. Bemis, J.L. Norton, R.W. Robinson, W.R. Henderson, M.L. Robbins, R.L. Lower

Cucurbit Genetics Cooperative Report 16:113 (article 41) 1993

Cucurbit Genetics Cooperative

Financial Statement

31 December 1992

Balance (31 December 1991)		\$4,026.69
Receipts		
Dues and back issue orders	\$2,971.00	
Interest on savings	\$164.28	
Total receipts		\$3,135.28
Expenditures		
CGC Report No. 15 (1992)		
Printing	\$1,874.17	
Mailing	\$542.46	
Call for papers (Report No. 16)	\$108.35	
Miscellaneous (envelopes, postage, etc.)	\$135.92	
U.S. FDIC bank fees	\$9.90	
Total Expenses		\$2,670.80
Balance (31 December 1992)		\$4,481.27