

GERB



United States
Department of
Agriculture

Human Nutrition
Information
Service

Home Economics
Research Report
Number 44

Foods Commonly Eaten by Individuals:

Amount Per Day and Per Eating Occasion

Eleanor M. Pao
Kathryn H. Fleming
Patricia M. Guenther
Sharon J. Mickle

FOODS COMMONLY EATEN BY INDIVIDUALS: AMOUNT PER DAY AND PER EATING OCCASION, Eleanor M. Pao, Kathryn H. Fleming, Patricia M. Guenther and Sharon J. Mickle
Consumer Nutrition Center, Human Nutrition Information Service,
U.S. Department of Agriculture,
Hyattsville, Maryland 20782.
Home Economics Research Report Number 44

ABSTRACT

Data on 3-day intakes of 200 commonly used foods and food groups by individuals in the Nationwide Food Consumption Survey 1977-78 are reported. The survey was conducted by the U.S. Department of Agriculture from April 1977 to March 1978 and included about 38,000 individuals in the 48 conterminous States. Data are tabulated by 16 sex-age categories and for total individuals. The 200 tables include the percentage of individuals using the food, the number of eating occasions at which the food was consumed, and the quantities (grams) of the food consumed by users per day and per eating occasion. Foods included in each table are explained briefly in the text. The number of eating occasions and quantities of food consumed by users are described by averages, standard deviations, and maximum values. Quantities consumed per day and per eating occasion are further described by values at seven specified percentiles.

Keywords: Dietary survey, eating occasion, food intake.

Trade names are used in this publication solely for the purpose of providing specific information. Mention of a trade name does not constitute a guarantee or warranty of the products by the U.S. Department of Agriculture or an endorsement of the Department over other products not mentioned.

Although data collection in the NFCS 77-78 was completed in mid-1978, this survey is the only source of national information on food intakes of individuals, by item and quantity, since 1965 when the first USDA nationwide survey of individuals was conducted. The results are still valid and will be used in coming years in the evaluation of many issues relating to food safety and product development, consumer demand, and nutrition education.

Supersedes Portion Sizes and Day's Intakes of Selected Foods, ARS-NE-67, issued December 1975.

CONTENTS

| | Page |
|---|------|
| LIST OF TABLES..... | iii |
| INTRODUCTION..... | 1 |
| METHODS AND PROCEDURES..... | 1 |
| Data collection..... | 1 |
| Sample..... | 2 |
| Data handling..... | 2 |
| Selection of foods for tables..... | 3 |
| Content of tables..... | 3 |
| DESCRIPTION OF FOODS..... | 4 |
| Beverages..... | 4 |
| Eggs..... | 5 |
| Fats and oils..... | 5 |
| Fruits and fruit juices..... | 6 |
| Citrus fruit..... | 6 |
| Noncitrus fruit..... | 6 |
| Grain products..... | 8 |
| Yeast breads..... | 8 |
| Quick breads and tortillas..... | 9 |
| Cakes, cookies, pies, doughnuts, and sweet rolls.. | 9 |
| Crackers and snacks..... | 10 |
| Pasta, pasta mixtures, and pizza..... | 10 |
| Rice and cooked cereals..... | 11 |
| Ready-to-eat cereals..... | 11 |
| Legumes and nuts..... | 12 |
| Meat, poultry, and fish..... | 12 |
| Meat, poultry, and fish..... | 12 |
| Processed meat..... | 14 |
| Mixtures..... | 15 |
| Milk and milk products..... | 16 |
| Fluid milk and milk beverages..... | 16 |
| Cheese..... | 16 |
| Milk products..... | 17 |
| Sugar and sweets..... | 17 |
| Vegetables..... | 18 |
| White potatoes..... | 18 |
| Raw vegetables..... | 19 |
| Cooked vegetables..... | 19 |
| Condiments..... | 20 |
| LITERATURE CITED..... | 21 |
| TABLES..... | 22 |
| APPENDIXES | |
| A -- System for classifying foods based on the first three digits of seven-digit food codes..... | 423 |
| B -- Table notes..... | 431 |

LIST OF TABLES

| | Table No. | Page |
|---|--------------|------|
| <u>Beverages</u> | | |
| Coffee..... | 1.01 | 22 |
| Decaffeinated coffee..... | 1.02 | 24 |
| Tea..... | 1.03 | 26 |
| Carbonated soft drinks..... | 1.04 | 28 |
| Cola-type carbonated soft drinks..... | 1.05 | 30 |
| Artificially sweetened carbonated soft drinks..... | 1.06 | 32 |
| Noncarbonated soft drinks..... | 1.07 | 34 |
| Fruit ades and drinks..... | 1.08 | 36 |
| Beer and ale..... | 1.09 | 38 |
| Wine..... | 1.10 | 40 |
| Distilled liquors..... | 1.11 | 42 |
| <u>Eggs</u> | | |
| Eggs..... | 2.01 | 44 |
| Boiled and poached eggs..... | 2.02 | 46 |
| Fried eggs..... | 2.03 | 48 |
| Scrambled eggs..... | 2.04 | 50 |
| <u>Fats and oils</u> | | |
| Butter..... | 3.01 | 52 |
| Margarine..... | 3.02 | 54 |
| Whipped butter and margarine..... | 3.03 | 56 |
| Salad dressings and mayonnaise..... | 3.04 | 58 |
| Mayonnaise..... | 3.05 | 60 |
| Salad dressings..... | 3.06 | 62 |
| Mayonnaise-type salad dressing..... | 3.07 | 64 |
| French dressing..... | 3.08 | 66 |
| Italian dressing..... | 3.09 | 68 |
| <u>Fruits</u> | | |
| Citrus fruit: | | |
| Fresh grapefruit..... | 4.01 | 70 |
| Grapefruit juice..... | 4.02 | 72 |
| Fresh oranges..... | 4.03 | 74 |
| Orange juice..... | 4.04 | 76 |
| Noncitrus fruit: | | |
| Raw apples..... | 4.05 | 78 |
| Applesauce and cooked apples..... | 4.06 | 80 |
| Applejuice..... | 4.07 | 82 |
| Bananas..... | 4.08 | 84 |
| Cantaloup..... | 4.09 | 86 |
| Fruit cocktail..... | 4.10 | 88 |

| | Table No. | Page |
|--|--------------|------|
| Grapejuice..... | 4.11 | 90 |
| Raw peaches..... | 4.12 | 92 |
| Cooked, canned, and frozen peaches..... | 4.13 | 94 |
| Raw pears..... | 4.14 | 96 |
| Cooked and canned pears..... | 4.15 | 98 |
| Cooked and canned pineapple..... | 4.16 | 100 |
| Raisins..... | 4.17 | 102 |
| Raw strawberries..... | 4.18 | 104 |
| <u>Grain products</u> | | |
| Yeast breads: | | |
| Yeast breads..... | 5.01 | 106 |
| White breads..... | 5.02 | 108 |
| White rolls..... | 5.03 | 110 |
| French, vienna, and italian breads..... | 5.04 | 112 |
| Whole and cracked wheat breads..... | 5.05 | 114 |
| Rye and pumpernickel breads..... | 5.06 | 116 |
| High fiber, wheat germ, bran, oatmeal, and multigrain breads..... | 5.07 | 118 |
| English muffins..... | 5.08 | 120 |
| Quick breads and tortillas: | | |
| Biscuits other than refrigerator type..... | 5.09 | 122 |
| Refrigerator-type biscuits..... | 5.10 | 124 |
| Cornbread..... | 5.11 | 126 |
| Pancakes..... | 5.12 | 128 |
| Waffles..... | 5.13 | 130 |
| Tortillas..... | 5.14 | 132 |
| Cakes, cookies, pies, doughnuts, and sweet rolls: | | |
| Cakes and cupcakes..... | 5.15 | 134 |
| Chocolate cakes and cupcakes..... | 5.16 | 136 |
| Cookies..... | 5.17 | 138 |
| Chocolate chip cookies..... | 5.18 | 140 |
| Oatmeal cookies..... | 5.19 | 142 |
| Peanut butter cookies..... | 5.20 | 144 |
| Sugar cookies..... | 5.21 | 146 |
| Pies..... | 5.22 | 148 |
| Fruit pies..... | 5.23 | 150 |
| Cream and custard pies..... | 5.24 | 152 |
| Doughnuts..... | 5.25 | 154 |
| Sweet rolls..... | 5.26 | 156 |
| Crackers and snacks: | | |
| Crackers..... | 5.27 | 158 |
| Graham crackers..... | 5.28 | 160 |
| Saltines and soda crackers..... | 5.29 | 162 |
| Butter crackers..... | 5.30 | 164 |
| Popcorn..... | 5.31 | 166 |

| | Table No. | Page |
|--|--------------|------|
| Pretzels..... | 5.32 | 168 |
| Potato chips..... | 5.33 | 170 |
| Corn-based salty snacks..... | 5.34 | 172 |
| Pasta, pasta mixtures, and pizza: | | |
| Pasta..... | 5.35 | 174 |
| Spaghetti..... | 5.36 | 176 |
| Noodles..... | 5.37 | 178 |
| Macaroni..... | 5.38 | 180 |
| Spaghetti with tomato sauce..... | 5.39 | 182 |
| Macaroni and cheese..... | 5.40 | 184 |
| Lasagne..... | 5.41 | 186 |
| Pizza..... | 5.42 | 188 |
| Rice and cooked cereals: | | |
| Rice..... | 5.43 | 190 |
| Cooked cereals..... | 5.44 | 192 |
| Grits..... | 5.45 | 194 |
| Oatmeal..... | 5.46 | 196 |
| Wheat cereals..... | 5.47 | 198 |
| Ready-to-eat cereals: | | |
| Ready-to-eat cereals..... | 5.48 | 200 |
| Ready-to-eat cereals, 15-21 grams per cup, all sugar levels..... | 5.49 | 202 |
| Ready-to-eat cereals, 25-32 grams per cup, 30 percent or more sugar..... | 5.50 | 204 |
| Ready-to-eat cereals, 25-32 grams per cup, less than 15 percent sugar.... | 5.51 | 206 |
| Ready-to-eat cereals, 35-45 grams per cup, 30 percent or more sugar..... | 5.52 | 208 |
| Ready-to-eat cereals, 35-45 grams per cup, less than 20 percent sugar.... | 5.53 | 210 |
| Ready-to-eat cereals, 50-75 grams per cup, all sugar levels..... | 5.54 | 212 |
| Ready-to-eat cereals, 80 grams or more per cup, all sugar levels..... | 5.55 | 214 |
| <u>Legumes and nuts</u> | | |
| Cooked dried beans..... | 6.01 | 216 |
| Baked beans..... | 6.02 | 218 |
| Nuts..... | 6.03 | 220 |
| Peanuts..... | 6.04 | 222 |
| Peanut butter..... | 6.05 | 224 |
| <u>Meat, poultry, and fish</u> | | |
| Meat, poultry, and fish: | | |
| Meat, poultry, fish, and their mixtures... | 7.01 | 226 |
| Meat, poultry, and fish..... | 7.02 | 228 |

| | Table No. | Page |
|---|--------------|------|
| Beef, pork, lamb, and veal..... | 7.03 | 230 |
| Beef..... | 7.04 | 232 |
| Beef steaks..... | 7.05 | 234 |
| Beef roasts..... | 7.06 | 236 |
| Ground beef..... | 7.07 | 238 |
| Pork..... | 7.08 | 240 |
| Pork chops and steaks..... | 7.09 | 242 |
| Ham..... | 7.10 | 244 |
| Pork roasts..... | 7.11 | 246 |
| Bacon..... | 7.12 | 248 |
| Lamb..... | 7.13 | 250 |
| Veal..... | 7.14 | 252 |
| Poultry..... | 7.15 | 254 |
| Chicken..... | 7.16 | 256 |
| Turkey..... | 7.17 | 258 |
| Liver..... | 7.18 | 260 |
| Fish and shellfish..... | 7.19 | 262 |
| Finfish other than canned, dried, and raw..... | 7.20 | 264 |
| Canned tuna..... | 7.21 | 266 |
| Shrimp..... | 7.22 | 268 |
| Processed meat: | | |
| Frankfurters..... | 7.23 | 270 |
| Bologna..... | 7.24 | 272 |
| Salami..... | 7.25 | 274 |
| Pork breakfast sausage..... | 7.26 | 276 |
| Sausage other than bologna, frank- furters, salami, and pork breakfast sausage..... | 7.27 | 278 |
| Mixtures: | | |
| Beef stew with vegetables..... | 7.28 | 280 |
| Chili con carne..... | 7.29 | 282 |
| Meat loaf..... | 7.30 | 284 |
| Tuna salad..... | 7.31 | 286 |
| Pot pies..... | 7.32 | 288 |
| Frozen plate meals..... | 7.33 | 290 |
| Hamburger and cheeseburger sandwiches..... | 7.34 | 292 |
| Soups..... | 7.35 | 294 |
| <u>Milk and milk products</u> | | |
| Fluid milk and milk beverages: | | |
| Fluid milk, milk beverages, and milk- based infant formulas..... | 8.01 | 296 |
| Fluid milk..... | 8.02 | 298 |

| | Table No. | Page |
|--|--------------|------|
| Fluid whole milk..... | 8.03 | 300 |
| Fluid lowfat milk..... | 8.04 | 302 |
| Fluid skim milk..... | 8.05 | 304 |
| Reconstituted dry milk..... | 8.06 | 306 |
| Chocolate milk..... | 8.07 | 308 |
| Cocoa and hot chocolate..... | 8.08 | 310 |
| Cheese: | | |
| Natural and processed cheese..... | 8.09 | 312 |
| Cheddar cheese..... | 8.10 | 314 |
| Swiss cheese..... | 8.11 | 316 |
| Natural cheese other than cheddar, swiss, and parmesan..... | 8.12 | 318 |
| Processed cheese, cheese food, and cheese spread..... | 8.13 | 320 |
| Cottage cheese..... | 8.14 | 322 |
| Milk products: | | |
| Fluid cream..... | 8.15 | 324 |
| Liquid and frozen cream substitutes..... | 8.16 | 326 |
| Powdered cream substitutes..... | 8.17 | 328 |
| Ice cream and ice milk..... | 8.18 | 330 |
| Nondairy whipped topping..... | 8.19 | 332 |
| Sour cream..... | 8.20 | 334 |
| Yogurt..... | 8.21 | 336 |
| <u>Sugar and sweets</u> | | |
| White sugar..... | 9.01 | 338 |
| Sirup..... | 9.02 | 340 |
| Honey..... | 9.03 | 342 |
| Jellies..... | 9.04 | 344 |
| Jams, preserves, and marmalades..... | 9.05 | 346 |
| Plain gelatin desserts..... | 9.06 | 348 |
| Gelatin desserts with fruit..... | 9.07 | 350 |
| Candy..... | 9.08 | 352 |
| <u>Vegetables</u> | | |
| White potatoes: | | |
| White potatoes..... | 10.01 | 354 |
| Baked potatoes..... | 10.02 | 356 |
| Boiled potatoes..... | 10.03 | 358 |
| Creamed, scalloped, and au gratin potatoes..... | 10.04 | 360 |
| Mashed potatoes..... | 10.05 | 362 |
| Home-fried and hash-browned potatoes..... | 10.06 | 364 |

| | Table No. | Page |
|---|--------------|------|
| French fried potatoes..... | 10.07 | 366 |
| Potato salad..... | 10.08 | 368 |
| Raw vegetables: | | |
| Cabbage salad and coleslaw..... | 10.09 | 370 |
| Raw carrots..... | 10.10 | 372 |
| Raw celery..... | 10.11 | 374 |
| Raw cucumbers..... | 10.12 | 376 |
| Lettuce and tossed salads..... | 10.13 | 378 |
| Lettuce..... | 10.14 | 380 |
| Tossed salads..... | 10.15 | 382 |
| Raw mature onions..... | 10.16 | 384 |
| Raw tomatoes..... | 10.17 | 386 |
| Cooked vegetables: | | |
| Cooked broccoli..... | 10.18 | 388 |
| Cooked cabbage..... | 10.19 | 390 |
| Cooked carrots..... | 10.20 | 392 |
| Cooked whole kernel corn..... | 10.21 | 394 |
| Cream-style corn..... | 10.22 | 396 |
| Cooked lima beans..... | 10.23 | 398 |
| Cooked mixed vegetables..... | 10.24 | 400 |
| Cooked cowpeas, field peas, and blackeye peas..... | 10.25 | 402 |
| Cooked green peas..... | 10.26 | 404 |
| Cooked spinach..... | 10.27 | 406 |
| Cooked string beans..... | 10.28 | 408 |
| Cooked summer squash..... | 10.29 | 410 |
| Cooked sweetpotatoes..... | 10.30 | 412 |
| Tomato juices..... | 10.31 | 414 |
| Condiments: | | |
| Catsup..... | 10.32 | 416 |
| Mustard..... | 10.33 | 418 |
| Cucumber pickles..... | 10.34 | 420 |

Foods Commonly Eaten by Individuals:

Amount Per Day and
Per Eating Occasion

Eleanor M. Pao
Kathryn H. Fleming
Patricia M. Guenther
Sharon J. Mickle¹

INTRODUCTION

This publication reports average intakes per day and per eating occasion for foods most commonly eaten by individuals who participated for 3 days in the Nationwide Food Consumption Survey (NFCS) conducted by the U.S. Department of Agriculture (USDA) in the 48 conterminous States April 1977 through March 1978. It updates and extends information for 1965 (10)² that has been widely used by business firms, trade associations, private researchers, journalists, Government agencies, and private individuals concerned with variations in the amounts of different kinds of foods consumed by individuals classified by sex and age. The data are also organized to meet the needs of nutrition specialists, consumer educators, and dietitians. The list of tables shows the commodity detail in which data are supplied to meet the requirements of market research and of educational and scientific applications.

This report provides information on many more foods than did the earlier publication and is derived from intakes of almost 38,000 individuals in 16 sex-age groups. Tables for 200 foods show the number and proportion of individuals who reported using each food, the number of occasions during the 3-day period on which they ate the food, and the quantities eaten per occasion and per day. Averages for actual users of foods are often quite different from the customary averages for whole population groups. An eating occasion was defined by the time at which one or more foods were consumed. Foods eaten at different times on the diet record were treated as separate eating occasions. The time of the eating occasion referred to the time eating began, and a few eating occasions, such as evening snacking, probably extended over a considerable time.

METHODS AND PROCEDURES

Data Collection

Information in the NFCS 1977-78 was collected during home visits by trained interviewers using a 1-day interviewer-recorded recall and a 2-day self-administered record. After the household respondent reported on food used in the household during the past week, the interviewer asked members of the household present to recall their intakes of food and beverage during the preceding day. Household members then were asked to record their intakes for the day of

¹The authors acknowledge the assistance of Bruce C. Gray, Jan L. Janiczek, and Michael S. Hamada, who developed the programs for data processing and table preparation.

²Underlined numbers in parentheses refer to Literature Cited, p. 21.

the interview and the next day, thus providing data for 3 consecutive days. Interviewers used a four-page leaflet of instructions to assist participants in describing foods and amounts consumed and left the leaflet for reference. A set of measuring cups and spoons and a ruler were given to each household for use in estimating quantities eaten. The household respondent usually provided information for children under 12 years and others unable to answer for themselves. If a household member was absent at the time of the interview but was expected to return within 2 days, forms were left to be completed. The interviewers returned to pick up the records and review them for clarity and completeness.

Sample

A stratified area probability sample of households was drawn in the 48 conterminous States in each of the four quarters from April 1977 through March 1978. About 15,000 households participated. All individuals in the households were asked to give food intake information in spring 1977. In the other three seasons all individuals under the age of 19 and one-half of those 19 years and over were asked to participate. In the three seasons other than spring, proportional representation was maintained among sex-age categories by treating each record for an individual 19 years and over as two records. In addition, a household weighting factor was applied to each individual to adjust for households included in the sample but providing no response. The unweighted number of individuals was 30,770. More than 90 percent of the individuals responding provided information for 3 days.

Data Handling

Interviewers mailed completed food intake forms to a central location for editing and coding. Coders assigned numerical codes to food items using a manual prepared at USDA (5). A seven-digit code was used to help classify foods into groups for tabulation. The first digit categorizes food items and beverages into 9 major food groups, the second digit identifies 47 major subgroups, and the third digit, 221 minor subgroups (see appendix A). The fourth digit further specifies minor subgroups. The remaining digits code information unique to the individual food. Foods with complete descriptions were assigned codes to preserve most of the detail specified. If the description of an item lacked detail, such as the type (e.g., type of cooky or meat), method of preparation (e.g., raw or cooked), or accompaniment (e.g., butter), the item was assigned a code for items lacking such specification.

Mixtures were usually classified in the food group of the major ingredient. Some mixtures such as salads, sandwiches, and frozen meals were reported either as a single item or by listing component parts. Some food items, such as peanut butter (table 6.05), are tabulated in this report only when they were reported separately, even though the food item might also have been a component of a sandwich or other mixture reported as a single unit. Therefore, the peanut butter in a sandwich reported as a unit is not captured, and peanut butter is underreported in the table.

Participants in the survey were asked to report amounts of foods they ate in measures most convenient to them, i.e., cups, tablespoons, dimensions, ounces. USDA nutritionists supplied the gram weight conversion factors in the coding manual (5) used by the coders. One ounce is equivalent to 28.35 grams. One cup, unless

otherwise noted in the text, refers to the standard 8-ounce measuring cup. The gram weights of common measures of food reported in the survey came mostly from Agricultural Handbook No. 456 (1), available revisions of Agricultural Handbook No. 8 (2-4), other USDA bulletins (7, 9), food industry, and actual determinations. If the amount eaten was inadequately specified, a standard portion size was assigned and identified. The weights in grams per food measure mentioned in the food descriptions in section III of this publication are those given in the coding manual. The number of times a food item is reported refers to the number of eating occasions in which it was consumed and not to the number of individual entries for that item. If a food item was entered more than once for a single eating occasion (such as milk on cereal and in coffee), the amounts on the dietary forms were summed and the total was the portion for that eating occasion. All maximum food consumption values were checked on the original dietary forms to be sure they were not errors caused by coding or transcription.

Selection of Foods for Tables

The food items and food groups selected for inclusion in this publication were those most frequently reported by survey participants. Most were mentioned on at least 1,000 occasions. The text provides a brief description of the foods included in each table. Several tables include all items in a major food group, for example, meat, poultry, and fish (table 7.02). Subsequent tables include items in a subgroup, such as beef, pork, lamb, and veal (table 7.03), which is followed by a table for each component--such as beef (table 7.04)--of the more cumulative table. Another level of subgrouping is provided in tables for beefsteaks (7.05), beef roasts (table 7.06), and ground beef (table 7.07). Unless specifically mentioned in the text, dietetic foods, such as artificially sweetened or low-sodium foods, were not included in the tables.

Content of Tables

For each table the numbers of individuals in total and in each of the 16 sex-age categories are identified. Five sets of intake data for users of the food or food group were computed. These intake data are based solely on persons who consumed the food item, and all averages reported are for the group of users only. The relevant formulas for the computations are given in appendix B. The first data set presents the percentages of individuals using the food on only 1 day, on only 2 days, and on all 3 of the 3 days for which data were collected. The second set includes the average number of eating occasions for the food, the standard deviation, and the maximum number of occasions at which the food was consumed during the 3 days.

In the third data set, the average quantity of food consumed per user per day and the standard deviation are given for individuals who used the food on only 1 day, on only 2 days, and on all 3 days reported. Also presented are the maximum quantity of the food item consumed on any 1 day and the maximum quantity consumed over 3 days. The average quantities of food per user per day are identified at specified percentiles in the distribution of the intakes of each food or food group. Percentile data for foods consumed by fewer than 50 individuals in a sex-age group are not reported. The fourth set of data consists of the average quantities, the standard

deviations, and the maximum quantities consumed by users per eating occasion. Also, the average quantities per eating occasion at specified percentiles in distributions of at least 50 individuals are reported. The last set of data has the number and the percentage of eating occasions for which portion sizes were assigned because amounts eaten were inadequately described. These assigned portion sizes are included in all calculations except in the fourth set of data, the average quantities consumed per eating occasion.

Additional information concerning foods consumed by individuals in the NFCS 1977-78 has been published (6, 11, 12).

DESCRIPTION OF FOODS

Beverages

Separate tables are included in the beverage section for coffee, tea, soft drinks, fruit ades and drinks, beer, wine, and distilled liquors. Milk and milk drinks are reported in the "Milk and Milk Products" section, fruit juices in "Fruits and Fruit Juices," and vegetable juices in the "Vegetables" section.

Coffee (table 1.01) includes liquid coffee made from ground roasted coffee, powdered instant coffee, and liquid coffee concentrate and coffee that was either acid-neutralized or decaffeinated. Usually sugar was reported separately, but some coffee included in this table had sugar added. Not included are turkish coffee; espresso; coffee made from instant cafe au lait; coffee combined with chicory, cereal, or cocoa; unreconstituted instant or liquid concentrate coffee; and coffee substitutes such as Postum. For all eating occasions with coffee, coffee prepared from ground roasted beans was specified for about 41 percent and coffee prepared from instant for 34 percent. Type of coffee was not specified for 25 percent of the eating occasions. About 10 percent of the eating occasions with coffee were for decaffeinated coffee (table 1.02). Of the occasions with decaffeinated coffee, 66 percent were for coffee prepared from powdered instant. One 6-fluid-ounce cup of coffee weighs 180-182 grams.

Tea (table 1.03) includes tea made from tea leaves (loose and in bags), powdered instant, and frozen concentrate and tea sold ready-to-drink in cans. On 46 percent of the occasions the type of tea was not specified. Tea for 34 percent of the occasions was identified as tea prepared from tea leaves and 20 percent as tea made from powdered instant. On less than 1 percent of the occasions tea was identified as ready-to-drink or prepared from frozen concentrate. Data in this table include tea with added cream or milk, sweeteners, or lemon but not the few reports of tea with added chocolate. Tea prepared from herbs, such as saffras or mint, is not included. A 6-fluid-ounce cup of tea weighs 180-182 grams.

About two-thirds of the carbonated soft drinks (table 1.04) were identified as cola-type carbonated soft drinks (table 1.05) and 14 percent as artificially sweetened carbonated soft drinks (table 1.06). Of the noncarbonated soft drinks (table 1.07), most were prepared from powdered mixes containing sugar and fruit

flavor, including some with added vitamin C. No artificially sweetened noncarbonated soft drinks are included with the noncarbonated soft drinks (table 1.07). Carbonated and non-carbonated soft drinks weigh 356-372 grams per 12-ounce can or 237-248 grams per cup. Fruit ades and drinks (table 1.08) include ready-to-drink beverages as well as beverages made from powdered mixes and frozen concentrates but exclude low-calorie or low-sugar drinks. Fruit ades and drinks weigh 247-256 grams per cup.

Alcoholic beverages tabulated herein are beer and ale, wine, and distilled liquors that are not part of mixtures such as cocktails reported as a unit. Cocktails were not tabulated because of the great diversity in their alcoholic content. Beer and ale (table 1.09) include beer, light beer, and ale. Wine (table 1.10) includes table and dry wines, such as burgundy, claret, chianti, sauterne, rhine, and champagne and sweet or dessert wines such as marsala, port, tokay, madeira, muscatel, angelica, sherry, and sweet vermouth. Cooking wines are not included in this table. Distilled liquors (table 1.11) include brandy, whiskey, gin, vodka, and rum when reported separately but not in cocktails reported as a single item. Beer, wine, and distilled liquors weigh 28-30 grams per fluid ounce. One 12-ounce can or bottle of beer weighs 360 grams, and one glass of wine (3.5 fluid ounces) weighs 100 grams.

Eggs

Eggs (table 2.01) include eggs cooked by all methods, raw eggs, baby-food egg yolks, and mixtures having egg as the main ingredient such as omelets, creamed and deviled eggs, and egg salads. Excluded are meringues, egg soups and sandwiches, and egg substitutes. Intakes of the most popular forms in which eggs are prepared are reported in the following three tables. One large boiled or poached egg (table 2.02) weighs 50 grams, one large fried egg (table 2.03) 46 grams, and one large egg scrambled or made into a plain omelet (table 2.04) 64 grams (2).

Fats and Oils

Fats and oils tabulated include butter, margarine, and salad dressings when these items were reported separately. Fats in mixtures and fats used in cooking, such as deep-fat frying, are not included in these tables nor is salad dressing when a salad with dressing was reported as a unit, i.e., the amount of salad dressing was not reported separately by the respondent.

Butter (table 3.01) does not include whipped butter or butter-margarine blends. Margarine (table 3.02) includes unsalted margarine but does not include liquid, diet, whipped, or imitation margarine or butter-margarine blends. Butter and margarine weigh 14 grams per tablespoon or 5 grams per pat (1 inch square, 1/3 inch high).

Whipped butter and margarine (table 3.03) weigh 9 grams per tablespoon.

Salad dressings and mayonnaise (table 3.04) include all salad dressings and mayonnaise except the low-calorie types. Salad dressings and mayonnaise weigh 14-16 grams per tablespoon. About 42 percent of the reports of salad dressing and mayonnaise were

mayonnaise, 13 percent italian dressing, 11 percent french dressing, 9 percent mayonnaise-type salad dressing, 7 percent thousand island dressing, and 11 percent other types of salad dressing. The remaining 7 percent was not specified as to type.

Mayonnaise (table 3.05) includes regular mayonnaise only and does not include mayonnaise-type salad dressing, mayonnaise made with yogurt, and imitation mayonnaise. These three items are included with salad dressings and mayonnaise (table 3.04). Salad dressings (table 3.06) include all the items in table 3.04 except regular mayonnaise. The most frequently consumed salad dressings were the mayonnaise type (table 3.07), french dressing (table 3.08), and italian dressing (table 3.09). French dressing includes catalina, ranch style, and homemade but does not include low-calorie french dressing. Italian dressing includes california onion and green onion dressings but does not include low-calorie italian dressing.

Fruit and Fruit Juices

Citrus fruit

Only the most frequently reported citrus fruits and citrus fruit juices were tabulated. Fruit juice combinations and mixtures containing citrus fruit, such as gelatin desserts, fruit cup, and salads, were excluded. Fresh grapefruit (table 4.01) includes halves and fresh sections with juice. Almost all reports of grapefruit were for fresh halves. One-half medium grapefruit was calculated as yielding 134 grams of edible portion (9), and 1 cup of sections with juice weighs 230 grams (1). Grapefruit juice (table 4.02) encompasses fresh, reconstituted from frozen, and canned. For 25 percent of the occasions canned grapefruit juice was specified, for 16 percent fresh or frozen was specified, and for 58 percent the form was not specified. On 44 percent of the occasions the juice was specified as unsweetened. Fresh oranges (table 4.03) include all varieties. One medium orange was calculated to yield 145 grams of edible portion and 1 cup of sections to weigh 195 grams. Orange juice (table 4.04) covers all forms--fresh, reconstituted from frozen, and canned, sweetened and unsweetened. Of the reported orange juices, nearly one-half (48 percent) were reconstituted frozen, 17 percent were fresh, 4 percent were canned, and the rest were not specified. Nearly half (49 percent) of the reports specified unsweetened orange juice. One cup of orange juice weighs 248-250 grams, depending upon whether sugar is added.

Noncitrus fruit

Noncitrus fruits and fruit juices reported most often were tabulated. Raw apples (table 4.05) include all varieties, both peeled and unpeeled, but do not include apples in such mixtures as waldorf salad. A medium apple was calculated to provide an edible portion of 138 grams. Applesauce and cooked apples (table 4.06) include apples that are baked, fried, pickled, stewed, and prepared for babies, but most of the items (85 percent) were applesauce. One cup of applesauce with sugar weighs 255 grams and 1 cup unsweetened weighs 244 grams. Applejuice (table 4.07) excludes apple cider and juices that are only part applejuice, such as cranberry-applejuice. About 7 percent of

the reports of applejuice specified baby-food applejuice. One cup weighs 248 grams.

Bananas (table 4.08) include only raw bananas. The yield from one medium banana was calculated as 119 grams (9), and 1 cup of sliced bananas weighs 150 grams (1). Cantaloup (table 4.09) refers to fresh only and excludes the few reports of frozen balls. One-half cantaloup was calculated to yield 272 grams of edible portion (9), and 1 cup of fresh cubes to weigh 160 grams (1). Fruit cocktail (table 4.10) is mainly the canned or cooked type and includes canned fruit salad and fruit compote. Raw fruit cup and raw fruit cocktail are excluded. One cup of canned fruit cocktail with heavy sirup weighs 255 grams. Grapejuice (table 4.11) includes all forms, canned and frozen concentrate reconstituted with water, sweetened and unsweetened, with and without added vitamin C. Grapeade and grape drink are reported with fruit ades and fruit drinks (table 1.09). One cup of grapejuice weighs 253 grams.

Peaches were tabulated for both the raw and the cooked, canned, and frozen forms. Raw peaches (table 4.12) accounted for 35 percent of the occasions on which peaches were consumed. One medium raw peach (about 2-3/4 inches in diameter) was calculated to yield 152 grams of edible portion, and 1 cup of sliced peaches weighs 178 grams. Cooked, canned, and frozen peaches (table 4.13) include peaches not specified as to form (about 13 percent of those mentioned) and commercially prepared baby food (8 percent). On about 75 percent of the occasions entries were identified as canned or cooked and on about 2 percent as frozen. Of the peaches reported as canned or cooked (excluding the baby-food items), 27 percent were in light sirup, 9 percent in heavy sirup, 4 percent unsweetened, and the type of liquid was not mentioned for the rest. One cup of canned or cooked peaches with heavy sirup weighs 256 grams, 1 cup with light sirup 250 grams, 1 cup unsweetened 244 grams, and 1 cup drained halves 224 grams. Peaches in cobblers, pies, and other mixtures are not included in these two tables.

Pears, like peaches, were consumed raw almost as often as cooked or canned. Raw pears (table 4.14) comprised 46 percent of the pear items in this and the next table combined. One medium pear was calculated to provide an edible portion of 164 grams. Cooked and canned pears (also baby-food pears) (table 4.15) include 5 percent of the pear items for which the form (raw, cooked, or canned) was not specified. Of the cooked or canned items (excluding baby food), 71 percent lacked a description of the sirup, 19 percent had light sirup, 5 percent had heavy sirup, and 5 percent were specified as unsweetened. Pears in combination with other fruits such as the baby-food pears and pineapple are not included in these tables. One cup of canned pear halves with heavy sirup weighs 255 grams.

Pineapple was specified as cooked or canned in 64 percent of the reports of this food. Pineapple items were not specified as to type in 20 percent of the reports and were specified as raw for the remainder. Cooked and canned pineapple (table 4.16) includes all pineapple items not reported as raw. Of these cooked and

canned items, 10 percent were reported as cooked or canned in fruit juice, slightly more (12 percent) as unsweetened, and the remainder did not provide such information. One cup of canned pineapple (chunk, tidbit, or crushed) with heavy sirup weighs 255 grams.

Raisins (table 4.17) were the most frequently reported dried fruit with dried prunes reported about one-half as often and dates and apricots each less than one-tenth as often. One cup of raisins weighs 145 grams. Raisins in mixtures are not included in the table.

Among berries, strawberries were reported most often, followed by cranberries, which were mentioned about one-half as often, and blueberries and raspberries each mentioned about one-tenth as often. Of strawberry items reported (excluding mixtures), 77 percent were specified as raw, 19 percent as frozen, and the remaining 4 percent were cooked, canned, or not specified. Quantities for raw strawberries (table 4.18) only are tabulated. One cup of raw strawberries weighs 149 grams.

Grain Products

Yeast breads

Yeast breads (table 5.01) include all toasted and not toasted breads, rolls, and sweet rolls. Bread and rolls used in sandwiches with ingredients reported separately by the respondent are included here. Bread and rolls used in sandwiches with amounts of ingredients not reported separately are excluded. Also excluded from table 5.01 are such mixtures as french toast. Tables 5.02-5.08 treat separately several of the products included in table 5.01.

White breads (table 5.02) are plain and toasted white breads including sourdough, french, italian, portuguese, and vienna breads, and bread that was not identified as to type. About 80 percent were reported as "white bread" and 13 percent as "bread," i.e., type was not specified. Excluded are flat breads, cheese breads, egg breads, and other specialty breads. Most breads and toasts weigh 20-25 grams per medium slice. White rolls (table 5.03) are plain and toasted white rolls, including hard, soft, sourdough, french, vienna, garlic, submarine, and sandwich rolls, and rolls that were not specified as to type. Most medium-sized rolls weigh 25-37 grams, and a submarine roll 100 grams. Excluded are rolls that were used in sandwiches reported as a single unit (ingredients and amounts not listed individually by the respondent). Also excluded are cheese, egg, and sweet rolls, bagels, and bread sticks. The bread group identified as french, vienna, and italian bread (table 5.04) also includes toast, rolls, and portuguese, cuban, and spanish varieties. These breads weigh 20-30 grams per slice, and a medium roll weighs 45 grams.

Whole and cracked wheat breads (table 5.05) include plain and toasted, whole and cracked wheat bread and rolls. Excluded are flat bread and bagels. Table 5.06 has plain and toasted rye, pumpernickel, and black bread and rolls, excluding bagels. High fiber, wheat germ, bran, oatmeal, and multigrain breads (table

5.07) include plain and toasted breads made with other types of flours and grains such as barley, buckwheat, oatmeal, triticale, and soy or with added parts of grains such as bran and wheat germ. Breads and toast in tables 5.05-5.07 weigh 23-28 grams per slice.

English muffins (table 5.08) are those made with white, whole wheat, or cracked wheat flour; 95 percent were white or not specified as to type. One english muffin weighs 57 grams, and one toasted english muffin 50 grams.

Quick breads and tortillas

The next five tables report data for quick breads. Biscuits (table 5.09) exclude whole wheat and cheese biscuits and those made from refrigerated dough. One medium biscuit was assigned a weight of 38 grams. Information on all types of biscuits made from refrigerated dough is found in table 5.10. The weight of a regular-sized biscuit is 20 grams. Cornbread (table 5.11) includes cornbread (86 percent of the occasions) and corn muffins, sticks, and rounds. Excluded are other cornmeal products such as cornbread stuffing, corn pone, hush puppies, and spoonbread. One piece of cornbread (2-1/2 inches by 2-1/2 inches by 1-1/2 inches) weighs 78 grams.

All varieties of pancakes except crepes and chinese pancakes are included in table 5.12, and waffles are tabulated in table 5.13. One medium pancake weighs 27 grams, and one medium waffle 50 grams.

Tortillas (table 5.14) are only those that were reported separately and include corn, wheat flour, and whole wheat flour tortillas. Excluded are tortillas that were a part of mixtures reported as units such as tacos and burritos. Tortillas weigh 30-35 grams each.

Cakes, cookies, pies, doughnuts, and sweet rolls

Cakes and cupcakes (table 5.15) include fruitcakes, poundcakes, and shortcakes with fruit. Excluded are coffeecakes. Of the items reported, 80 percent were servings of cake and 20 percent were cupcakes. Of the cakes and cupcakes, 42 percent were chocolate, 11 percent were yellow, 9 percent were white, 35 percent were other flavors, and 3 percent were not specified as to flavor. Chocolate cakes and cupcakes (table 5.16) include chocolate, devil's food, and fudge-flavored cakes and cupcakes. One piece of cake (one-twelfth of a two-layer frosted cake) weighs 92 grams, and one frosted cupcake (2-1/2 inches diameter) 36 grams.

All varieties of cookies are combined in table 5.17. Graham crackers are not included but appear with crackers--tables 5.27 and 5.28. The four most frequently reported kinds of cookies are tabulated separately (tables 5.18-5.21). All five tables exclude dietetic cookies. Chocolate chip cookies (table 5.18) include those with and without such additional ingredients as nuts and raisins. One medium chocolate chip cookie was given a weight of 7 grams. Oatmeal cookies (table 5.19) include those with such additional ingredients as raisins and chocolate chips,

but oatmeal sandwich and oatmeal peanut butter cookies are excluded. One oatmeal cookie was given a weight of 13 grams. Peanut butter cookies (table 5.20) include peanut sandwich cookies and wafers and all other varieties of peanut cookies except those with oatmeal. One peanut butter cookie was given a weight of 12 grams. Sugar cookies (table 5.21) include those with such added ingredients as nuts, fruit, spice, sour cream, and jelly topping. Excluded are frosted sugar cookies and sugar wafers. One sugar cookie was given a weight of 16 grams.

Pies (table 5.22) include all types of pies and tarts. Fruit pies (table 5.23) include tarts, fried pies, and pies that were not specified as to kind by the respondent (3 percent of the reports of pies). Of the fruit pies, 56 percent were apple and 16 percent were cherry. Of the cream and custard-type pies and tarts (table 5.24), 27 percent were pumpkin and 18 percent were chocolate cream. Neither table 5.23 nor 5.24 includes chiffon, meringue, or pecan pies. A tart weighs 100 grams, a slice of fruit pie 135 grams, and one slice of cream or custard pie 130 grams.

All varieties of doughnuts are included in table 5.25 and sweet rolls, excluding danish pastries, in table 5.26. Most doughnuts weigh 42-57 grams, and one medium sweet roll 50 grams.

Crackers and snacks

Crackers (table 5.27) are of all types--graham crackers (plain, cinnamon, and chocolate covered), saltines, butter crackers, cheese crackers, toast thins, matzo, flat bread, and peanut butter sandwich crackers. Graham crackers (table 5.28) include the honey and cinnamon varieties but not the chocolate covered. One graham cracker square (or two small rectangular pieces) weighs 14 grams. Saltine and soda crackers (table 5.29) include those with seeds but exclude low-sodium crackers. One square saltine or soda cracker weighs 3 grams. With butter crackers (table 5.30) are all butter-type crackers except cheese. One round cracker weighs 3 grams and a rectangular cracker 4 grams.

Popcorn (table 5.31) includes plain, "battered," salted, and popcorn with cheese, but not popcorn with sugar or caramel sirup. One cup weighs 6-9 grams. Pretzels (table 5.32) exclude soft pretzels and baby pretzels. One three-ring pretzel weighs 3 grams. Potato chips, flavored potato chips, and chips made from granulated potatoes are reported in table 5.33. One cup of potato chips weighs 18 grams. Corn-based salty snacks (table 5.34) include corn chips, tortilla chips, puffed corn, and toasted corn. One cup weighs 35-40 grams, depending upon the type.

Pasta, pasta mixtures, and pizza

Pasta (table 5.35) includes cooked macaroni, noodles (egg, whole wheat, and spinach), and spaghetti. Pasta may be enriched or unenriched and with or without added fat. Chow mein noodles, rice noodles, and such pasta mixtures as macaroni and cheese are not included. The pasta was spaghetti (table 5.36) on 48 percent of the occasions, noodles (table 5.37) on 30 percent, and

macaroni (table 5.38) on the other 21 percent. Cooked pastas weigh 140-160 grams per cup.

Three frequently reported pasta mixtures are tabulated. Spaghetti with tomato sauce (table 5.39) also includes macaroni and noodles mixed with tomatoes or tomato sauce with or without meat and cheese. One cup weighs 250 grams. Macaroni and cheese (table 5.40) also includes macaroni casserole and egg noodle with cottage cheese. One cup of homemade macaroni and cheese weighs 200 grams, and 1 cup from mix or canned 240 grams. Lasagne (table 5.41) weighs 250 grams per cup.

Pizza (table 5.42) includes all kinds except thick crust pizzas. A pizza 5 inches in diameter weighs 84-94 grams, and one 12 inches in diameter 486-543 grams.

Rice and cooked cereals

White rice (table 5.43) includes yellow rice but not sweet rice, brown rice, wild rice, oriental long rice (transparent noodles), or rice cooked with milk. Cream of rice and rice mixtures also are not part of this category. The majority of the reports of rice (84 percent) were not specified as to type, i.e., regular cooking, instant, or converted. Cooked rice weighs 165-175 grams per cup.

In the cooked cereals group (table 5.44) are grits, oatmeal, barley, cream of rice, bulgar, and wheat cereals. Cereals may be regular, quick cooking, or instant, made with water or milk, and with or without added fat. Not included are cornmeal mush, cooked cornstarch, and millet. Of the cooked cereals reported, 34 percent were grits or hominy (table 5.45), 43 percent were oatmeal (table 5.46), and 21 percent were wheat cereals (table 5.47) such as cream of wheat, whole wheat cereal, and cocoa wheat cereal. Wheat cereals do not include bulgar or wheat pilaf. The weights of 1 cup of cereal are as follows: grits, 220-245 grams; oatmeal, 240 grams; cream of rice, 245 grams; wheat cereals, 245 grams; and bulgar, 135 grams.

Ready-to-eat cereals

Ready-to-eat cereals are categorized by cup weight and by sugar content as a percent of dry weight. Data on sugar content were obtained from published reports (8) and information supplied by manufacturers. The general category of ready-to-eat cereals (table 5.48) includes all the cereals tabulated separately in tables 5.49 through 5.55 plus dry cereals not specified as to type and dry baby cereals. Baby cereals in jars and breakfast bars are not included.

Ready-to-eat cereals, 15-21 grams per cup, all sugar levels (table 5.49) encompass Corn Kix, Honey Combs, puffed corn, puffed oats, puffed rice, puffed wheat, Quisp, and Special K. In ready-to-eat cereals, 25-32 grams per cup, 30 percent or more sugar (table 5.50) are Alpha Bits, Apple Jacks, Boo Berry, Coco Pebbles, Cocoa Puffs, Cookie Crisp, Corny Snaps, Count Chocula, Crazy Cow, Frankenberry, Freakies, Frosty O's, Fruit Brute, Fruit Loops,

Fruity Pebbles, Golden Grahams, Kaboom, Lucky Charms, Sugar Pops, and Trix. Ready-to-eat cereals, 25-32 grams per cup, less than 15 percent sugar (table 5.51) include unprocessed bran, Cheerios, Corn Chex, corn flakes, Crispy Rice, Pep, Post Toasties, Rice Chex, Rice Krispies, Team, Total, and Wheaties.

Categorized as ready-to-eat cereals, 35-45 grams per cup, 30 percent or more sugar (table 5.52) are Captain Crunch, Captain Crunch with Peanut Butter, Cocoa Krispies, Country Crisp, Frosted Rice, Sugar Frosted Flakes, Sugar Snacks, and Super Sugar Crisp. Ready-to-eat cereals, 35-45 grams per cup, less than 20 percent sugar (table 5.53) include Bran Chex, Bran Flakes, Buc Wheats, 40% Bran, Grape Nut Flakes, Life, Product 19, and Wheat Chex. The group of ready-to-eat cereals, 50-75 grams per cup, all sugar levels (table 5.54) is comprised of All Bran, Bran Buds, Cinnamon Frosted Mini Wheats, Frosted Mini Wheats, 100% Bran, Raisin Bran, and Shredded Wheat. The group of ready-to-eat cereals, 80 grams or more per cup, all sugar levels (table 5.55) includes Alpen, Concentrate, Country Morning, cracked whole wheat, Cracklin' Bran, C. W. Post, Familia, Golden Harvest, Granola, Granola Puffs, Grape Nuts, Heartland, 100% Natural, Pernola, Proteinola, and wheat germ.

Legumes and Nuts

Of the cooked dried beans (table 6.01) reported, including canned, 46 percent were pinto beans, 19 percent kidney beans, 14 percent lima beans, and 13 percent white beans such as navy beans and great northern beans. The remaining beans reported were black, soy, mungo, or not specified as to type. This table does not include soy-derived products, soy-based infant formulas, or such bean mixtures as baked beans, stewed beans, and bean soups. Immature lima beans are reported with cooked vegetables (table 10.23). Cooked dried beans weigh 185-190 grams per cup.

Baked beans (table 6.02) are bean mixtures in which dried beans are cooked with sauce (tomato, chili, barbecue, sweet and sour), sugar, fat, or meat. Stewed beans and other bean mixtures are not included in this table. Baked beans weigh 255 grams per cup.

Nuts (table 6.03) include all nuts except coconut and nuts mixed with fruit or seeds, and peanuts accounted for 54 percent of the reports. Other frequently reported nuts were almonds (8 percent of the eating occasions), cashews (4 percent), pecans (6 percent), and walnuts (6 percent). The cup weight of nuts varies from 120 to 160 grams. Almonds, pecan halves, and peanuts weigh about 1 gram each, and cashews, butternuts, ginko nuts, and english walnut halves about 2 grams each.

Peanuts (table 6.04) include regular and dry roasted peanuts but not boiled peanuts. Peanut butter (table 6.05) does not include low-sodium or imitation peanut butter or the peanut butter portion of sandwiches when the peanut butter sandwich was reported as a unit. Peanut butter weighs 16 grams per tablespoon.

Meat, Poultry, and Fish

Meat, poultry, and fish

Among meat, poultry, fish, and their mixtures (table 7.01) are plain and breaded meat, poultry, and fish, and foods that are combinations of meat and grains or vegetables but have meat as

the main ingredient. Baby foods are included as are frozen plate meals when they were reported as a unit. Meat soups (table 7.35) and gravies are not in this group nor are mixtures having some meat but not as a major component. All foods tabulated in tables 7.02-7.27 are included in table 7.01 as are other meats and meat mixtures not separately tabulated.

Meat, poultry, and fish (table 7.02) include plain and breaded meat, poultry, and fish and baby-food meats. Food items that are combinations of meat and grain products or vegetables are omitted as are frozen plate meals (see table 7.33), soups (see table 7.35), and gravies.

Beef, pork, lamb, and veal (table 7.03) include all the foods listed in the individual beef, pork, lamb, and veal tables, which are described below. Excluded are organ meats, sausages, luncheon meats, spreads, and mixtures. Of the items reported in this table, 57 percent are beef and 41 percent are pork. Lamb and veal account for only 1 percent each.

In the beef category reported in table 7.04 are beef steaks, roasts, stew meat, ground beef, brisket, corned beef, pastrami, beef bacon, dried chipped beef, and other types of beef including oxtails, neckbones, shortribs, and baby-food and junior beef and beef sticks. Excluded are beef sausages, luncheon meats, meat spreads, organ meats, and mixtures. Separate tables are provided for beef steaks (table 7.05), beef roasts (table 7.06), and ground beef (table 7.07), which includes plain patties and plain meatballs and excludes meatloaf.

The cuts of pork in table 7.08 are chops, steaks, cutlets, and roasts; ham; bacon; spareribs; miscellaneous parts such as pigs' feet and ham hocks; ground, pickled, and dehydrated pork; and baby-food pork, ham, and pork sticks. The table does not include pork sausages, luncheon meats, meat spreads, organ meats, or mixtures.

Pork cutlets are included with pork chops and steaks in table 7.09. Of the pork items reported, 91 percent were pork chops. One medium pork chop was calculated to yield a cooked, edible portion of 80 grams. Ham (table 7.10) includes baby-food ham but excludes fresh ham, i.e., uncured. Most of the pork roasts reported in table 7.11 were fresh, but a few were cured. The few reports of fresh ham are not included in table 7.11. Bacon (table 7.12) includes pork breakfast strips but not fresh, uncured bacon, side pork, fatback, beef bacon, or canadian bacon. One medium slice of cooked bacon weighs 8 grams.

Data for lamb in table 7.13 include a few (less than 1 percent) reports of mutton. The item most frequently reported was lamb chops (62 percent). One medium lamb chop has a cooked, edible yield (lean and fat) of 109 grams. Tables of data on lamb and veal (table 7.14) include baby foods but exclude organ meats and mixtures.

Poultry (table 7.15) includes chicken, turkey, cornish game hen, duck, dove, squab, pigeon, quail, partridge, goose, and

pheasant. Excluded are poultry organ meats, luncheon meats, spreads, and mixtures. Of the reports of poultry, 88 percent were chicken and 11 percent were turkey. Chicken (table 7.16) includes all chicken parts, breaded and floured chicken, and baby-food chicken and chicken sticks, but not cornish game hen, chicken loaf, chicken liver, or chicken spread. Cooked edible yields of three parts were calculated as follows: one-half medium roasted breast, 90 grams; one fried whole leg, 112 grams; and one fried thigh, 61 grams.

Turkey (table 7.17) includes turkey roll, ground turkey, and baby-food turkey and turkey sticks. Excluded are turkey luncheon meats, organ meats, and mixtures.

Liver (table 7.18) includes beef, calf, pork, poultry, rabbit, venison, and baby-food livers. Excluded are fish liver, liver sausage, and such mixtures as chopped liver and liver pate. Of the reports of liver, 45 percent were beef, 18 percent poultry, and 5 percent pork, rabbit, or venison. The remaining 32 percent were not specified as to kind.

The fish and shellfish (table 7.19) include fresh, breaded, floured, canned, raw, and dried fish but not fish mixtures. Finfish (table 7.20) encompasses all varieties of finfish except canned, dried, and raw fish. Fish mixtures are also excluded. The most frequently reported item was fish sticks (13 percent of the finfish reports). Other popular finfish items were flounder (7 percent), haddock (6 percent), and perch (6 percent). Canned tuna (table 7.21) includes only plain tuna in water or in oil. One 7-ounce can of tuna, drained, weighs 169 grams. Tuna salad appears in table 7.31. Shrimp in table 7.22 is breaded and plain but not dried shrimp. A 1/2-cup serving of breaded, fried shrimp weighs 68 grams, and a 1/2-cup serving of steamed shrimp 64 grams.

Processed Meat

Frankfurters (table 7.23) are made from any type of meat or poultry and include a few that are breaded. Junior meat sticks are excluded. A medium frankfurter was assigned a weight of 44 grams. All types of bologna are included in table 7.24, and all types of salami in table 7.25. One slice of either weighs 28 grams. Meats that were part of sandwiches reported as a unit, i.e., the amount of meat was not reported separately by the respondent, are not included in these tables.

Included with pork breakfast sausage (table 7.26) are fresh bulk sausage, patties, and links (about 75 percent of the items reported); brown-and-serve sausage; and country-style and smoked sausage. Sausage that respondents did not identify as to type was assumed to be pork breakfast sausage and was also put in this table. Sausages other than frankfurters, bologna, salami, and pork breakfast sausage (table 7.27) include pork and beef sausage, beef sausage, smoked beef sausage, brown-and-serve beef sausage, smoked beef sausage sticks, blood sausage, bockwurst, bratwurst, capicola, cervelat, chorizos (spanish sausage), head cheese, knoblauch, knockwurst, mortadella, pepperoni, polish sausage (kielbasa), italian sausage, lup chong (chinese smoked pork sausage), mettwurst, souse, thuringer, and vienna sausage.

Polish and Italian were the most frequently reported of these sausages, representing 24 percent and 13 percent of the total number of items, respectively.

Mixtures

Of the many meat, poultry, and fish mixtures consumed by individuals, the most frequently reported were beef stew, chili con carne, meat loaf, tuna salad, pot pies, frozen plate meals, hamburger and cheeseburger sandwiches, and soups. The following weights were assigned: beef stew with vegetables (table 7.28), 245 grams per cup; chili con carne (table 7.29) with or without beans, 255 grams per cup; meatloaf (table 7.30), 11 grams per cubic inch; and tuna salad (table 7.31), 205 grams per cup.

Pot pies (table 7.32) include beef, ham, chicken, turkey, and tuna and weigh 227 grams per 8-ounce frozen pot pie. Of the pot pies reported, 69 percent were chicken or turkey and 28 percent were beef.

Frozen plate meals (table 7.33) include a wide variety of beef, pork, veal, poultry, fish, legume, and pasta entrees obtained frozen then heated to serve. Poultry entrees such as fried chicken or turkey slices with potatoes and mixed vegetables comprised 45 percent of the frozen plate meals reported. Beef entrees such as sliced or chopped beef with potatoes and/or other vegetables accounted for 32 percent of the reports, fish dinners 10 percent, and pasta-based frozen plate meals such as lasagne and macaroni and cheese 5 percent. Frozen plate meals were reported in a variety of package sizes from 7 to 18 ounces (198-510 grams).

Hamburger and cheeseburger sandwiches (table 7.34) include all hamburgers and cheeseburgers reported as a unit with or without condiments such as mustard, catsup, and special sauces and accompaniments such as lettuce, tomatoes, pickles, and bacon. Chiliburgers, tacoburgers, pizzaburgers, and sloppy joes are excluded. Of the total hamburger and cheeseburger sandwiches reported, 75 percent were hamburgers. One hamburger on a bun with regular condiments weighs 97 grams and with accompaniments (lettuce, tomato, pickles) 154 grams. A double hamburger (two patties, double-decker bun) with condiments was given a weight of 183 grams, and a hamburger with one-fourth pound (raw weight) of meat and accompaniments 266 grams. Addition of a slice of cheese increases the weight by 14 grams.

Of the soups reported (table 7.35) 37 percent were vegetable; 27 percent grain mixtures such as chicken noodle soup; 18 percent beef, poultry, or fish; 5 percent legume such as bean or pea soup; and 11 percent tomato, including tomato beef and tomato rice. The remaining 2 percent of soups include pork, lamb, potato, egg drop, cheese, and milk soups, and baby-food soups. Soups included were homemade or prepared from canned or dry mixes. Unreconstituted dry soup mixes and low-sodium soups are not included. One cup (8 fluid ounces) of soup weighs 227-250 grams.

Milk and Milk
Products

Fluid milk and milk beverages

Fluid milk, milk beverages, and milk-based infant formulas (table 8.01) include whole, lowfat, skim, evaporated, condensed, and reconstituted dry milks; flavored milks such as chocolate milk, cocoa, and malted milk; milk-based infant formulas; buttermilk; and goat's milk. Excluded are eggnog, milk shakes, imitation milks such as soy milk, and meal replacements with milk such as instant breakfast. Of the total reports of fluid milk, milk beverages, and milk-based infant formulas, about 45 percent were whole milk, 16 percent lowfat milk, 4 percent skim milk, 1 percent reconstituted dry milk, 5 percent chocolate milk or cocoa, and 2 percent milk-based infant formulas. The remaining reports were evaporated milk, condensed milk, buttermilk, goat's milk (2 percent), and milk unspecified as to type (24 percent).

Under fluid milk (table 8.02) are whole, lowfat, and skim milk as well as milk not specified as to type. These types of milk are individually reported in tables 8.03, 8.04, and 8.05, respectively. Milk unspecified as to type is included with whole milk (table 8.03). Data for acidophilus and low-sodium whole milks are in table 8.02 but not in table 8.03. Buttermilk, evaporated milk, reconstituted dry milk, flavored milks, milk-based infant formulas, and goat's milk are not included with fluid milk (table 8.02). One cup (8 fluid ounces) of whole fluid milk weighs 244 grams.

Reconstituted dry milk (table 8.06) includes milk reconstituted from whole dry milk (3 percent of the eating occasions), lowfat dry milk (11 percent), and nonfat dry milk (48 percent). Dry milk not specified as to fat content was reported for 38 percent of the occasions. The weight of 1 cup (8 fluid ounces) of reconstituted dry milk is 245 grams.

With chocolate milk (table 8.07) and cocoa and hot chocolate (table 8.08) are beverages made with whole milk (27 percent of the occasions), lowfat milk (7 percent), skim or dry milk (4 percent), and milk not specified as to form and fat content (52 percent). Malted milk beverages are omitted. Chocolate milk, cocoa, and hot chocolate weigh 250-256 grams per cup (8 fluid ounces).

Cheese

Natural and processed cheeses (table 8.09) include natural cheeses such as blue, brick, camembert, brie, cheddar, gouda, edam, limburger, mozzarella, parmesan, and swiss, and processed cheeses such as cream cheeses, processed cheeses, cheese foods, and cheese spreads. Excluded are imitation cheese, cottage cheese, and cheese mixtures such as cheese cakes, fondues, dips, souffles, and soups. Of the reports of natural and processed cheese, 10 percent were cheddar cheese, 6 percent swiss, 8 percent other natural cheeses, 3 percent cream cheese, 50 percent processed cheese, and 23 percent cheeses not specified as to type. Separate tables are provided for cheddar cheese (table 8.10), swiss cheese (table 8.11), natural cheeses other than

cheddar, swiss, and parmesan (table 8.12), and processed cheese, cheese food, and cheese spread (table 8.13). One cubic inch of most cheeses weighs 17 grams.

Of the reports of cottage cheese (table 8.14), 29 percent were for creamed cottage cheese, 10 percent lowfat, 4 percent cottage cheese with fruits and vegetables and baby-food cottage cheese, and 57 percent were not specified as to type. Uncreamed, dry curd cottage cheese is not included in this table. One-half cup of cottage cheese weighs 113 grams.

Milk products

Under fluid cream (table 8.15) are data for light cream (8 percent of the eating occasions), half and half (28 percent), and heavy cream (2 percent), but more than half of the items (62 percent) were not specified as to type. Excluded from this table are powdered cream, whipped cream, and cream substitutes. Of the reports of liquid and frozen cream substitutes (table 8.16), 14 percent were for liquid and 28 percent for frozen; 58 percent were not specified as to type. One tablespoon of fluid cream, liquid cream substitute, or frozen cream substitute weighs 15 grams. Powdered cream substitutes (table 8.17) weigh about 6 grams per tablespoon, 3 grams per packet.

Ice cream and ice milk (table 8.18) include the so-called "hard serve" ice cream and ice milk but exclude "soft serve" ice cream, dietetic and imitation ice cream, and ice cream bars, sandwiches, cones, and sundaes. Of the reports of ice cream and ice milk, 96 percent were for ice cream and 4 percent for ice milk. Data on flavors indicated 16 percent were chocolate ice cream or ice milk, 70 percent flavors other than chocolate, but 14 percent were not specified as to flavor. One cup of ice cream weighs 133 grams, and 1 cup of ice milk 131 grams.

Of reported nondairy whipped toppings (table 8.19), 60 percent were frozen, 15 percent were made from powdered mixes, 4 percent were products in pressurized cans, and 21 percent were not specified as to type. Nondairy whipped toppings weigh 70-80 grams per cup.

Sour cream (table 8.20) includes a few reports of sour half and half but does not include imitation (nondairy) sour cream, filled sour cream, or sour cream in mixtures such as dips. Sour cream weighs 12 grams per tablespoon, 230 grams per cup (2).

Data for all types of yogurt except frozen are in table 8.21. Of reported yogurts, 60 percent were fruit varieties, only 4 percent were vanilla or coffee flavors, and 36 percent were not specified as to flavor. An 8-ounce (avoirdupois) container of yogurt weighs 227 grams and 1 cup 245 grams.

Sugar and Sweets

The sugar and sweets tabulated are those that were most commonly reported: white sugar, sirup, honey, jellies, jams, gelatin desserts, and candies. Under white sugar (table 9.01) are both the granular and lump forms as well as sugar not further specified, rock sugar, and rock candy. Table 9.01 does not include

confectioner's or powdered sugar. One teaspoon of granular white sugar weighs 4 grams, one individual packet 6 grams.

Sirup (table 9.02) includes cane, corn, fruit, maple, and sorghum sirups singly or in combination, buttered blends, and sirups not further specified as to type. Chocolate sirup, caramelized sugar, liquid brown sugar, sugar and water sirup, and dietetic sirups are excluded. One tablespoon of sirup or honey (table 9.03) weighs 20 grams.

Jellies (table 9.04) and jams, preserves, and marmalades (table 9.05) include all fruit flavors but exclude dietetic jellies, jam, and preserves. Jellies weigh 19 grams per tablespoon; jams, preserves, and marmalades 20 grams per tablespoon.

Plain gelatin desserts (table 9.06) and gelatin desserts with fruit (table 9.07) do not include gelatin desserts with added whipped cream, cream cheese, or nuts when desserts with these ingredients were reported as a unit. Dietetic gelatin desserts also are excluded from these tables. One-half cup of gelatin dessert, plain or with fruit, weighs 120 grams.

Candy (table 9.08) includes all types of candy except those specified as dietetic. Of the reports of candy, 59 percent specified chocolate candies or candies containing or coated with chocolate; 14 percent specified hard candies; 6 percent caramels (not chocolate), gumdrops, or taffy; and 4 percent marshmallows. Among other reports of candy were a variety of other kinds and those not specified as to type. One cup of Hershey Kisses, chocolate covered peanuts, or M&M's was assigned a weight of 170 grams, 1 cup of jelly beans (75 pieces) 220 grams, and 1 roll of Life-savers 22 grams. Most medium candy bars weigh 34-51 grams.

Vegetables

Only the most frequently reported vegetables were tabulated. Vegetable mixtures were excluded from tabulations with the exception of white potato mixtures, tomato juices, traditional mixed vegetables, and cabbage, lettuce, and tossed salads.

White potatoes

White potatoes (table 10.01) include all reports of white potatoes except potato chips and sticks (table 5.33), which are classified under crackers and snacks, and potato soups. Among the reports of white potatoes, 23 percent each were for mashed potatoes and french fried potatoes. Baked potatoes accounted for 15 percent, boiled potatoes 11 percent, and potato salad 6 percent of the occasions. Creamed, scalloped, and au gratin potatoes; home-fried potatoes; and hash-browned potatoes accounted for 5, 8, and 2 percent, respectively. The data for baked potatoes (table 10.02) pertain to all white potatoes baked in skin except stuffed baked potatoes, which were mentioned only a few times. One medium baked potato yields 93-98 grams of edible weight, excluding peel. Under boiled potatoes (table 10.03) are those with and without added fat and with and without peel but not canned potatoes. A medium boiled potato yields 122-127 grams edible portion without the peel. Creamed, scalloped, and au gratin potatoes (table 10.04) include potatoes cooked in sauce

with and without cheese. One cup weighs 235-245 grams. Mashed potatoes (table 10.05) weigh 210 grams per cup. Information on home-fried and hash-browned potatoes (table 10.06) is for german fried, o'brien, cottage style, and fried from raw as well as hash browned from fresh, frozen, and dry mix. One cup was assigned a weight of 155 grams for hash-browned and 170 grams for home fries. A fast food order of french fried potatoes (table 10.07) weighs about 70 grams. Potato salad (table 10.08) includes german style, and 1 cup weighs 250 grams.

Raw vegetables

Raw vegetables (tables 10.09-10.17) reported separately most often were cabbage, carrots, celery, cucumbers, lettuce, onions, and tomatoes. Other vegetables also were eaten raw but on too few occasions to be tabulated individually.

Raw cabbage salad and coleslaw (table 10.09) include a few salads with apples, raisins, or pineapple, and 1 cup weighs 120-135 grams. Chinese and red cabbage salad or coleslaw and plain cabbage are not included although they were reported a few times. Raw carrots (table 10.10) exclude carrot salads. One medium carrot strip weighs 4 grams. One large outer stalk of raw celery (table 10.11) weighs 40 grams. Raw cucumbers (table 10.12) do not include the few cucumber salads that were reported. One-half cup of cucumber slices weighs 70 grams.

Lettuce and tossed salads (table 10.13) include all the types treated separately in tables 10.14 and 10.15 as well as tossed salads with avocado, cheese, and/or eggs added to lettuce or assorted vegetables. Only data for plain lettuce with and without dressing are in table 10.14. Tossed salads (table 10.15) include salads with lettuce and other assorted vegetables, with and without salad dressing. One cup of plain lettuce weighs 55 grams, lettuce with dressing 65 grams, and tossed salad 60-108 grams because of some heavier ingredients such as tomatoes and cucumbers.

Raw mature onions (table 10.16) include onions when they were reported separately such as slices of onion mentioned as part of sandwiches and as garnishes, but the onions in salads and mixtures often were not reported separately. Green onions were seldom reported and are not included. One slice of onion weighs 18 grams. Raw tomatoes (table 10.17) include cherry, italian, and plum tomatoes. One slice of tomato weighs 30 grams, one cherry tomato 15 grams. Undoubtedly, many of the reports are for slices of tomatoes eaten as part of a sandwich or a salad. For many salads and some sandwiches with tomatoes, however, ingredients were not reported separately; thus raw tomatoes were used more often than shown in the table.

Cooked vegetables

Data on cooked vegetables, including canned, appear in tables 10.18-10.30. Included in all of these tables are vegetables that were eaten with or without added fat, e.g., butter or margarine. Baby-food and junior vegetables are included also. Excluded are

vegetables served with a sauce; creamed, dried, and pickled vegetables; and vegetable souffles and casseroles.

Broccoli (table 10.18) weighs 85-88 grams per 1/2-cup serving. Cabbage (table 10.19) includes white cabbage and cabbage that was not specified as to type. A 1/2-cup serving weighs about 75 grams. Carrots (table 10.20) exclude glazed. One-half cup weighs about 78 grams. Whole kernel corn (table 10.21) includes white and yellow corn on and off the cob. A 1/2-cup serving weighs 83-85 grams. Cream-style corn (table 10.22) includes white and yellow corn, and a 1/2-cup serving weighs 128 grams. Immature lima beans (table 10.23) weigh 85-88 grams per 1/2 cup. Mixed vegetables (table 10.24) include only the traditional mixture of corn, lima beans, peas, green beans, and carrots. Other vegetable combinations are excluded. One-half cup of mixed vegetables weighs 91-94 grams. Crowder peas are included with cowpeas, field peas, and blackeye peas, not dried (table 10.25). A 1/2-cup serving weighs 85-88 grams. For green peas (table 10.26) the weight is 83-85 grams per 1/2 cup.

Spinach (table 10.27) weighs 93-103 grams per 1/2 cup. String beans (table 10.28) include green string beans and pole beans but exclude yellow (wax) and italian beans. A 1/2-cup serving weighs 67-70 grams. With plain cooked and canned sweetpotatoes (table 10.30) are candied sweetpotatoes. One medium baked sweetpotato weighs 114-119 grams, and one medium boiled 151-156 grams. A 1/2-cup serving of canned sweetpotatoes weighs 113-115 grams.

Along with tomato juices (table 10.31) are tomato juice cocktails and combination juices such as tomato with other vegetable juices. One-half cup weighs 121-122 grams.

Condiments

The last three tables cover the most frequently reported condiments: catsup (table 10.32), mustard (table 10.33), and cucumber pickles (table 10.34), both sweet and sour, but not pickle relish. One tablespoon of catsup or mustard weighs 15 grams, and one sour pickle spear 30 grams.

LITERATURE CITED

- (1) Adams, C. F.
1975 Nutritive value of American foods in common units. U.S. Dept. Agr. Agr. Handb. 456, 291 pp.
- (2) Consumer and Food Economics Institute.
1976 Composition of foods: Dairy and egg products; raw, processed, prepared. U.S. Dept. Agr. Agr. Handb. 8-1, 157 pp.
- (3) _____
1978 Composition of foods: Baby foods; raw, processed, prepared. U.S. Dept. Agr. Agr. Handb. 8-3, 231 pp.
- (4) _____
1979 Composition of foods: Poultry products; raw, processed, prepared. U.S. Dept. Agr. Agr. Handb. 8-5, 356 pp.
- (5) _____
1979 Manual of food codes and conversions of measures and gram-weight for use with individual food intake data from Nationwide Food Consumption Survey 1977-78, U.S. Dept. Agr. CFE (Adm.) 352, 569 pp.
- (6) Consumer Nutrition Center.
1980 Food and nutrient intakes of individuals in 1 day in the United States, spring 1977. Nationwide Food Consumption Survey 1977-78, U.S. Dept. Agr. Prelim. Rpt. No. 2, 121 pp.
- (7) Fulton, L., Matthews, E., and Davis, C.
1977 Average weight of a measured cup of various foods. U.S. Dept. Agr. Home Econ. Res. Rpt. No. 41, 26 pp.
- (8) Li, B.W., and Schuhmann, P. J.
1980 Gas-liquid chromatographic analysis of sugars in ready-to-eat breakfast cereals. J. Food Science. 45:138-141.
- (9) Matthews, R., and Garrison, Y. J.
1975 Food yield summarized by different stages of preparation. U.S. Dept. Agr. Agr. Handb. 102, 136 pp.
- (10) Pao, E. M., and Burk, M. C.
1975 Portion sizes and day's intakes of selected foods. U.S. Dept. Agr. ARS-NE-67, 70 pp.
- (11) Pao, E. M.
1981 Changes in American food consumption patterns and their nutritional significance. Food Technology. 35:43-53.
- (12) Rizek, R. L., and Jackson, E. M.
1980 Current food consumption practices and nutrient sources in the American diet. 35 pp. Consumer Nutrition Center, Science and Education Administration, U.S. Dept. Agr., Hyattsville, Md.

TABLE 1.01.--COFFEE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 51.1 | 0 | 1.0 | 1.2 | 2.0 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.3 | 0 | .3 | .7 | 1.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 6.4 | 0 | .4 | .5 | .4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 39.4 | 0 | .3 | .1 | .3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 4.9 | 0 | 1.9 | 1.7 | 1.5 |
| STANDARD DEVIATION.....DO..... | 3.4 | 0 | .8 | 1.0 | .8 |
| MAXIMUM.....DO..... | 64.0 | 0 | 3.0 | 6.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 521 | 0 | 106 | 86 | 107 |
| STANDARD DEVIATION.....GM..... | 436 | 0 | 73 | 54 | 98 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 80 | | | | |
| 25TH.....GM..... | 240 | | | | |
| 50TH.....GM..... | 400 | | | | |
| 75TH.....GM..... | 681 | | | | |
| 90TH.....GM..... | 1,021 | | | | |
| 95TH.....GM..... | 1,281 | | | | |
| 99TH.....GM..... | 2,083 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 311 | 0 | 73 | 161 | 180 |
| STANDARD DEVIATION.....GM..... | 199 | 0 | 78 | 70 | 90 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 396 | 0 | 177 | 168 | 208 |
| STANDARD DEVIATION.....GM..... | 260 | 0 | 46 | 45 | 127 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 619 | 0 | 195 | 173 | 246 |
| STANDARD DEVIATION.....GM..... | 445 | 0 | 26 | 100 | 128 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 11,580 | 0 | 360 | 360 | 600 |
| MAXIMUM OVER 3 DAYS.....GM..... | 27,900 | 0 | 720 | 840 | 1,440 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 317 | 0 | 166 | 155 | 209 |
| STANDARD DEVIATION.....GM..... | 171 | 0 | 75 | 61 | 118 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 180 | | | | |
| 25TH.....GM..... | 240 | | | | |
| 50TH.....GM..... | 240 | | | | |
| 75TH.....GM..... | 480 | | | | |
| 90TH.....GM..... | 480 | | | | |
| 95TH.....GM..... | 720 | | | | |
| 99TH.....GM..... | 961 | | | | |
| MAXIMUM.....GM..... | 2,880 | 0 | 360 | 240 | 600 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 654.1 | 0 | 0 | 1.0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .7 | 0 | 0 | 2.9 | 0 |

1/. SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|--------|--------|--------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 4.7 | 4.1 | 16.4 | 17.1 | 53.9 | 53.0 | 84.2 | 82.4 | 84.7 | 85.0 | 85.1 | 81.7 |
| 2.6 | 2.2 | 5.0 | 6.6 | 9.5 | 9.6 | 5.5 | 5.0 | 4.5 | 4.1 | 3.0 | 4.0 |
| 1.0 | 1.1 | 4.5 | 4.1 | 10.6 | 10.2 | 8.5 | 8.5 | 6.2 | 6.0 | 6.3 | 5.5 |
| 1.2 | .8 | 6.8 | 6.4 | 33.8 | 33.2 | 70.2 | 68.9 | 74.1 | 74.9 | 75.9 | 72.2 |
| 1.8 | 1.7 | 2.6 | 2.8 | 4.3 | 4.2 | 5.7 | 5.3 | 5.2 | 4.7 | 5.2 | 4.7 |
| 1.1 | .9 | 1.7 | 2.3 | 3.3 | 3.2 | 4.1 | 3.3 | 3.1 | 2.4 | 2.7 | 2.6 |
| 6.0 | 4.0 | 16.0 | 16.0 | 32.0 | 29.0 | 64.0 | 43.0 | 29.0 | 18.0 | 13.0 | 32.0 |
| 141 | 129 | 244 | 268 | 474 | 453 | 633 | 566 | 532 | 444 | 484 | 421 |
| 106 | 88 | 207 | 289 | 435 | 420 | 538 | 418 | 371 | 280 | 277 | 312 |
| 40 | 40 | 60 | 60 | 80 | 80 | 120 | 120 | 149 | 120 | 160 | 140 |
| 80 | 80 | 80 | 80 | 180 | 160 | 300 | 260 | 240 | 240 | 240 | 240 |
| 80 | 80 | 160 | 160 | 360 | 320 | 480 | 480 | 480 | 400 | 480 | 320 |
| 160 | 160 | 320 | 320 | 640 | 561 | 841 | 721 | 680 | 561 | 641 | 520 |
| 240 | 240 | 480 | 560 | 1,020 | 960 | 1,200 | 1,061 | 961 | 800 | 880 | 721 |
| 400 | 240 | 641 | 880 | 1,280 | 1,200 | 1,441 | 1,360 | 1,201 | 960 | 1,040 | 960 |
| 480 | 480 | 1,280 | 1,580 | 1,921 | 2,162 | 2,563 | 2,082 | 1,702 | 1,441 | 1,282 | 1,363 |
| 254 | 230 | 287 | 264 | 355 | 314 | 324 | 321 | 305 | 274 | 229 | 246 |
| 114 | 88 | 184 | 120 | 275 | 183 | 174 | 196 | 161 | 165 | 132 | 122 |
| 210 | 263 | 305 | 381 | 428 | 365 | 448 | 416 | 323 | 386 | 330 | 303 |
| 66 | 128 | 207 | 292 | 298 | 208 | 293 | 276 | 118 | 212 | 170 | 136 |
| 269 | 213 | 381 | 462 | 632 | 617 | 715 | 635 | 584 | 478 | 522 | 456 |
| 134 | 100 | 225 | 355 | 464 | 446 | 548 | 418 | 366 | 277 | 266 | 314 |
| 840 | 780 | 1,680 | 2,400 | 7,680 | 5,520 | 11,580 | 4,320 | 3,360 | 3,120 | 1,680 | 6,480 |
| 2,040 | 1,440 | 4,320 | 6,180 | 15,840 | 11,760 | 27,900 | 12,000 | 9,120 | 6,840 | 4,320 | 13,440 |
| 233 | 222 | 286 | 286 | 330 | 321 | 331 | 320 | 307 | 284 | 277 | 271 |
| 104 | 100 | 131 | 170 | 190 | 190 | 182 | 169 | 142 | 127 | 113 | 116 |
| 60 | 60 | 149 | 150 | 180 | 180 | 180 | 180 | 180 | 180 | 180 | 150 |
| 180 | 180 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 |
| 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 |
| 240 | 240 | 240 | 240 | 480 | 480 | 480 | 480 | 360 | 240 | 240 | 240 |
| 360 | 242 | 480 | 480 | 480 | 480 | 480 | 480 | 480 | 480 | 480 | 480 |
| 480 | 480 | 480 | 480 | 720 | 720 | 720 | 720 | 480 | 480 | 480 | 480 |
| 480 | 540 | 721 | 961 | 961 | 961 | 961 | 960 | 721 | 721 | 720 | 720 |
| 720 | 540 | 960 | 1,920 | 2,400 | 2,400 | 2,880 | 2,160 | 1,440 | 1,920 | 1,200 | 1,440 |
| .8 | 5.8 | 4.7 | 5.8 | 56.3 | 99.9 | 185.1 | 173.5 | 29.3 | 59.5 | 12.3 | 20.1 |
| .5 | 3.8 | .8 | .8 | .6 | .8 | .8 | .6 | .6 | .9 | .5 | .5 |

TABLE 1.02.--DECAFFEINATED COFFEE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 7.3 | 0 | 0 | .1 | .1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.8 | 0 | 0 | .1 | .1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.4 | 0 | 0 | 0 | .1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 4.1 | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 3.7 | 0 | 0 | 1.0 | 1.6 |
| STANDARD DEVIATION.....DO..... | 2.6 | 0 | 0 | .0 | .5 |
| MAXIMUM.....DO..... | 25.0 | 0 | 0 | 1.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 347 | 0 | 0 | 60 | 85 |
| STANDARD DEVIATION.....GM..... | 292 | 0 | 0 | 0 | 40 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 80 | | | | |
| 25TH.....GM..... | 160 | | | | |
| 50TH.....GM..... | 240 | | | | |
| 75TH.....GM..... | 480 | | | | |
| 90TH.....GM..... | 720 | | | | |
| 95TH.....GM..... | 911 | | | | |
| 99TH.....GM..... | 1,341 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 371 | 0 | 0 | 180 | 120 |
| STANDARD DEVIATION.....GM..... | 285 | 0 | 0 | 0 | 0 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 407 | 0 | 0 | 0 | 180 |
| STANDARD DEVIATION.....GM..... | 261 | 0 | 0 | 0 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 467 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 314 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 5,520 | 0 | 0 | 180 | 180 |
| MAXIMUM OVER 3 DAYS.....GM..... | 11,760 | 0 | 0 | 180 | 360 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 282 | 0 | 0 | 180 | 163 |
| STANDARD DEVIATION.....GM..... | 129 | 0 | 0 | 0 | 27 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 180 | | | | |
| 25TH.....GM..... | 240 | | | | |
| 50TH.....GM..... | 240 | | | | |
| 75TH.....GM..... | 240 | | | | |
| 90TH.....GM..... | 480 | | | | |
| 95TH.....GM..... | 480 | | | | |
| 99TH.....GM..... | 721 | | | | |
| MAXIMUM.....GM..... | 1,440 | 0 | 0 | 180 | 180 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 41.5 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| .2 | .3 | 1.2 | 1.3 | 2.9 | 4.4 | 11.0 | 13.6 | 15.1 | 20.1 | 20.7 | 21.3 |
| .1 | .2 | .5 | .7 | .9 | 1.9 | 2.9 | 3.2 | 2.5 | 2.5 | 3.2 | 3.9 |
| 0 | .1 | .3 | .5 | .9 | 1.0 | 2.2 | 2.6 | 2.3 | 3.8 | 2.4 | 2.3 |
| .1 | 0 | .4 | .2 | 1.1 | 1.5 | 6.0 | 7.7 | 10.4 | 13.8 | 15.1 | 15.1 |
| 2.0 | 1.3 | 2.1 | 2.2 | 2.6 | 2.7 | 3.8 | 3.9 | 4.1 | 3.8 | 4.1 | 3.8 |
| 1.0 | .5 | 1.2 | 1.9 | 1.6 | 2.4 | 2.8 | 2.8 | 2.4 | 2.0 | 2.5 | 2.4 |
| 3.0 | 2.0 | 5.0 | 9.0 | 7.0 | 18.0 | 24.0 | 25.0 | 12.0 | 13.0 | 11.0 | 13.0 |
| 132 | 94 | 189 | 171 | 267 | 270 | 372 | 380 | 381 | 323 | 350 | 316 |
| 72 | 36 | 105 | 159 | 210 | 423 | 304 | 310 | 247 | 188 | 218 | 215 |
| | | | | 60 | 60 | 80 | 80 | 80 | 80 | 80 | 80 |
| | | | | 120 | 80 | 160 | 160 | 240 | 180 | 240 | 180 |
| | | | | 240 | 160 | 240 | 320 | 320 | 240 | 320 | 240 |
| | | | | 320 | 280 | 500 | 480 | 540 | 460 | 420 | 400 |
| | | | | 561 | 560 | 800 | 800 | 641 | 561 | 720 | 561 |
| | | | | 721 | 641 | 961 | 1,000 | 721 | 680 | 800 | 721 |
| | | | | 881 | 1,922 | 1,521 | 1,361 | 1,364 | 880 | 1,081 | 1,280 |
| 206 | 222 | 345 | 214 | 352 | 330 | 407 | 392 | 383 | 324 | 308 | 401 |
| 54 | 43 | 133 | 105 | 297 | 207 | 330 | 330 | 225 | 142 | 149 | 212 |
| 0 | 208 | 230 | 295 | 324 | 329 | 439 | 457 | 418 | 374 | 376 | 378 |
| 0 | 43 | 54 | 93 | 132 | 159 | 244 | 343 | 176 | 185 | 119 | 177 |
| 201 | 0 | 313 | 440 | 438 | 517 | 516 | 509 | 464 | 382 | 418 | 373 |
| 36 | 0 | 87 | 225 | 230 | 654 | 325 | 319 | 245 | 183 | 211 | 222 |
| 242 | 240 | 480 | 720 | 1,440 | 5,520 | 2,400 | 3,360 | 1,920 | 1,920 | 1,620 | 1,680 |
| 724 | 480 | 1,200 | 2,160 | 2,640 | 11,760 | 5,760 | 7,440 | 4,095 | 4,320 | 3,240 | 3,840 |
| 202 | 215 | 265 | 234 | 305 | 301 | 295 | 291 | 279 | 258 | 255 | 252 |
| 44 | 44 | 89 | 63 | 156 | 181 | 132 | 141 | 107 | 84 | 74 | 94 |
| | | | | 180 | 180 | 180 | 180 | 180 | 180 | 180 | 180 |
| | | | | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 |
| | | | | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 |
| | | | | 360 | 240 | 300 | 240 | 240 | 240 | 240 | 240 |
| | | | | 480 | 480 | 480 | 480 | 480 | 360 | 360 | 360 |
| | | | | 720 | 720 | 480 | 480 | 480 | 480 | 480 | 480 |
| | | | | 721 | 961 | 721 | 900 | 720 | 480 | 480 | 540 |
| 242 | 240 | 480 | 480 | 960 | 1,440 | 1,200 | 1,440 | 720 | 720 | 480 | 1,080 |
| 0 | 0 | 0 | 0 | 0 | 1.0 | 12.4 | 17.0 | 8.6 | 0 | .8 | 1.8 |
| 0 | 0 | 0 | 0 | 0 | .2 | .6 | .5 | 1.2 | 0 | .2 | .2 |

TABLE 1.03.--TEA
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|------------|------------|------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE | MALE | MALE | MALE |
| | | AND FEMALE | AND FEMALE | AND FEMALE | AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 38.5 | 4.9 | 19.9 | 22.4 | 24.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 12.0 | 3.1 | 9.3 | 9.5 | 10.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 10.1 | .7 | 5.6 | 6.4 | 7.5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 16.4 | 1.1 | 5.0 | 6.5 | 6.8 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER.... | 3.1 | 3.1 | 2.4 | 2.5 | 2.3 |
| STANDARD DEVIATION.....DO..... | 2.3 | 3.0 | 1.8 | 2.1 | 1.5 |
| MAXIMUM.....DO..... | 36.0 | 13.0 | 12.0 | 18.0 | 11.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 315 | 118 | 136 | 175 | 190 |
| STANDARD DEVIATION.....GM..... | 278 | 127 | 109 | 161 | 141 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 80 | | 30 | 40 | 60 |
| 25TH.....GM..... | 121 | | 60 | 80 | 80 |
| 50TH.....GM..... | 240 | | 100 | 160 | 160 |
| 75TH.....GM..... | 401 | | 200 | 221 | 240 |
| 90TH.....GM..... | 640 | | 282 | 360 | 363 |
| 95TH.....GM..... | 801 | | 362 | 403 | 440 |
| 99TH.....GM..... | 1,362 | | 480 | 646 | 720 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 330 | 232 | 200 | 243 | 276 |
| STANDARD DEVIATION.....GM..... | 200 | 260 | 130 | 151 | 139 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 379 | 98 | 207 | 243 | 275 |
| STANDARD DEVIATION.....GM..... | 218 | 68 | 95 | 99 | 104 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 504 | 264 | 261 | 327 | 340 |
| STANDARD DEVIATION.....GM..... | 310 | 146 | 121 | 211 | 161 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 5,280 | 960 | 847 | 1,980 | 1,331 |
| MAXIMUM OVER 3 DAYS.....GM..... | 11,760 | 1,380 | 2,178 | 4,800 | 2,760 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 302 | 115 | 171 | 207 | 246 |
| STANDARD DEVIATION.....GM..... | 145 | 64 | 79 | 82 | 90 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 180 | | 60 | 120 | 120 |
| 25TH.....GM..... | 240 | | 120 | 180 | 182 |
| 50TH.....GM..... | 240 | | 151 | 240 | 240 |
| 75TH.....GM..... | 360 | | 240 | 240 | 242 |
| 90TH.....GM..... | 480 | | 242 | 242 | 360 |
| 95TH.....GM..... | 484 | | 242 | 303 | 480 |
| 99TH.....GM..... | 912 | | 480 | 480 | 541 |
| MAXIMUM.....GM..... | 3,840 | 240 | 726 | 1,200 | 726 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER.... | 678.5 | 4.8 | 0 | 5.2 | 16.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.5 | 6.4 | 0 | .5 | 1.6 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|--------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 27.4 | 29.3 | 30.1 | 33.5 | 38.3 | 46.8 | 41.0 | 48.4 | 39.5 | 50.5 | 32.6 | 47.3 |
| 11.2 | 11.2 | 10.7 | 12.1 | 13.2 | 14.9 | 11.9 | 13.1 | 9.4 | 11.7 | 9.2 | 11.5 |
| 7.5 | 8.9 | 8.4 | 10.5 | 11.0 | 11.5 | 10.4 | 11.7 | 9.7 | 13.2 | 7.6 | 11.7 |
| 8.7 | 9.2 | 10.9 | 10.9 | 14.1 | 20.4 | 18.7 | 23.6 | 20.5 | 25.6 | 15.8 | 24.2 |
| 2.5 | 2.4 | 2.8 | 2.7 | 3.0 | 3.3 | 3.2 | 3.4 | 3.2 | 3.4 | 3.1 | 3.4 |
| 1.6 | 1.6 | 1.9 | 2.1 | 2.4 | 2.6 | 2.3 | 2.4 | 2.1 | 2.3 | 2.0 | 2.3 |
| 10.0 | 10.0 | 13.0 | 24.0 | 36.0 | 27.0 | 16.0 | 22.0 | 11.0 | 17.0 | 10.0 | 20.0 |
| 258 | 230 | 332 | 272 | 359 | 332 | 359 | 336 | 329 | 308 | 284 | 296 |
| 209 | 169 | 287 | 257 | 346 | 294 | 302 | 281 | 277 | 222 | 202 | 214 |
| 80 | 80 | 80 | 80 | 80 | 80 | 80 | 80 | 80 | 80 | 60 | 80 |
| 101 | 100 | 160 | 110 | 160 | 121 | 160 | 160 | 160 | 160 | 140 | 160 |
| 200 | 161 | 240 | 210 | 280 | 240 | 250 | 240 | 240 | 240 | 240 | 240 |
| 320 | 302 | 440 | 360 | 480 | 441 | 480 | 441 | 440 | 400 | 400 | 400 |
| 540 | 443 | 645 | 521 | 720 | 661 | 721 | 641 | 641 | 564 | 600 | 561 |
| 721 | 560 | 921 | 681 | 968 | 840 | 960 | 803 | 841 | 721 | 721 | 641 |
| 969 | 887 | 1,450 | 1,200 | 1,762 | 1,442 | 1,441 | 1,442 | 1,441 | 1,120 | 960 | 1,041 |
| 360 | 315 | 395 | 309 | 384 | 331 | 354 | 323 | 328 | 307 | 272 | 325 |
| 272 | 148 | 254 | 139 | 244 | 198 | 253 | 151 | 129 | 143 | 100 | 161 |
| 355 | 341 | 434 | 375 | 473 | 403 | 416 | 371 | 313 | 342 | 329 | 330 |
| 147 | 163 | 236 | 185 | 330 | 231 | 218 | 193 | 119 | 143 | 115 | 164 |
| 452 | 383 | 561 | 480 | 610 | 530 | 558 | 506 | 487 | 444 | 427 | 421 |
| 237 | 186 | 328 | 332 | 408 | 323 | 322 | 301 | 300 | 224 | 191 | 218 |
| 2,520 | 2,040 | 3,360 | 3,600 | 4,620 | 5,280 | 4,356 | 3,540 | 2,400 | 2,400 | 1,200 | 1,920 |
| 4,080 | 3,540 | 6,120 | 10,320 | 11,160 | 11,760 | 6,292 | 9,600 | 5,760 | 4,860 | 3,120 | 5,040 |
| 310 | 282 | 362 | 298 | 364 | 298 | 335 | 295 | 305 | 269 | 275 | 262 |
| 140 | 112 | 188 | 117 | 212 | 131 | 168 | 124 | 118 | 101 | 101 | 83 |
| 180 | 180 | 180 | 180 | 180 | 180 | 180 | 180 | 180 | 180 | 180 | 180 |
| 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 |
| 240 | 240 | 300 | 240 | 300 | 240 | 240 | 240 | 240 | 240 | 240 | 240 |
| 360 | 300 | 480 | 360 | 480 | 360 | 480 | 360 | 360 | 300 | 242 | 240 |
| 480 | 480 | 484 | 480 | 484 | 480 | 480 | 480 | 480 | 480 | 480 | 360 |
| 484 | 480 | 721 | 480 | 721 | 484 | 720 | 480 | 480 | 480 | 480 | 480 |
| 960 | 720 | 968 | 721 | 1,080 | 726 | 961 | 721 | 720 | 600 | 541 | 480 |
| 1,440 | 1,452 | 1,920 | 960 | 3,840 | 1,920 | 1,694 | 1,920 | 720 | 960 | 1,080 | 968 |
| 9.0 | 11.7 | 17.2 | 20.8 | 55.5 | 114.3 | 100.5 | 183.9 | 39.1 | 68.8 | 4.6 | 26.4 |
| .6 | .8 | 1.5 | 1.5 | 1.2 | 1.4 | 1.5 | 1.6 | 2.7 | 2.3 | .8 | 1.7 |

TABLE 1.04.--CARBONATED SOFT DRINKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 50.8 | 4.5 | 40.1 | 51.4 | 50.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 20.3 | 2.7 | 20.1 | 25.4 | 25.9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 15.2 | .7 | 12.5 | 16.6 | 14.5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 15.2 | 1.1 | 7.6 | 9.4 | 9.7 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1</u> /NUMBER..... | 2.6 | 2.3 | 2.2 | 2.2 | 2.1 |
| STANDARD DEVIATION.....DO..... | 2.0 | 1.8 | 1.6 | 1.5 | 1.5 |
| MAXIMUM.....DO..... | 22.0 | 7.0 | 13.0 | 13.0 | 16.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1</u> /GM..... | 274 | 88 | 123 | 153 | 177 |
| STANDARD DEVIATION.....GM..... | 241 | 75 | 111 | 114 | 129 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> / | | | | | |
| 5TH.....GM..... | 62 | | 21 | 41 | 41 |
| 25TH.....GM..... | 119 | | 51 | 82 | 82 |
| 50TH.....GM..... | 205 | | 82 | 123 | 144 |
| 75TH.....GM..... | 359 | | 164 | 205 | 246 |
| 90TH.....GM..... | 556 | | 247 | 297 | 338 |
| 95TH.....GM..... | 733 | | 328 | 382 | 410 |
| 99TH.....GM..... | 1,193 | | 574 | 554 | 657 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 342 | 145 | 187 | 255 | 297 |
| STANDARD DEVIATION.....GM..... | 188 | 89 | 105 | 136 | 169 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 387 | 162 | 197 | 266 | 309 |
| STANDARD DEVIATION.....GM..... | 197 | 85 | 105 | 129 | 128 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 503 | 172 | 269 | 295 | 343 |
| STANDARD DEVIATION.....GM..... | 291 | 83 | 150 | 138 | 146 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 5,228 | 679 | 1,538 | 1,107 | 1,845 |
| MAXIMUM OVER 3 DAYS.....GM..... | 10,824 | 775 | 3,198 | 2,276 | 3,444 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1</u> /GM..... | 310 | 115 | 169 | 209 | 248 |
| STANDARD DEVIATION.....GM..... | 124 | 61 | 74 | 89 | 95 |
| PERCENTILES: <u>1</u> / | | | | | |
| 5TH.....GM..... | 124 | | 62 | 93 | 123 |
| 25TH.....GM..... | 246 | | 123 | 124 | 185 |
| 50TH.....GM..... | 296 | | 154 | 185 | 246 |
| 75TH.....GM..... | 369 | | 246 | 246 | 308 |
| 90TH.....GM..... | 492 | | 246 | 310 | 369 |
| 95TH.....GM..... | 492 | | 308 | 369 | 372 |
| 99TH.....GM..... | 739 | | 372 | 492 | 496 |
| MAXIMUM.....GM..... | 1,968 | 248 | 496 | 984 | 984 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> / | | | | | |
| EATING OCCASIONS.....NUMBER..... | 371.3 | 0 | 2.5 | 13.3 | 2.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .7 | 0 | .3 | .7 | .1 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 56.5 | 56.8 | 69.1 | 72.0 | 69.4 | 65.9 | 43.9 | 46.5 | 21.0 | 20.5 | 18.3 | 16.3 |
| 26.4 | 25.5 | 22.4 | 23.4 | 22.6 | 21.6 | 18.8 | 20.1 | 10.0 | 9.9 | 7.9 | 9.2 |
| 17.5 | 18.4 | 22.8 | 22.8 | 20.7 | 20.2 | 12.3 | 13.0 | 6.4 | 5.8 | 5.3 | 3.4 |
| 12.6 | 13.0 | 23.9 | 25.8 | 26.1 | 24.1 | 12.7 | 13.4 | 4.6 | 4.9 | 5.1 | 3.7 |
| 2.4 | 2.4 | 2.9 | 2.5 | 3.1 | 3.0 | 2.5 | 2.5 | 2.2 | 2.3 | 2.1 | 1.9 |
| 1.6 | 1.7 | 2.0 | 1.9 | 2.3 | 2.1 | 1.9 | 1.9 | 1.5 | 1.9 | 1.2 | 1.3 |
| 12.0 | 20.0 | 15.0 | 15.0 | 17.0 | 22.0 | 16.0 | 19.0 | 9.0 | 19.0 | 6.0 | 7.0 |
| 247 | 237 | 352 | 312 | 366 | 313 | 276 | 242 | 218 | 206 | 211 | 163 |
| 195 | 185 | 271 | 223 | 297 | 264 | 243 | 217 | 165 | 236 | 143 | 125 |
| 81 | 62 | 82 | 82 | 82 | 79 | 79 | 62 | 62 | 59 | 82 | 41 |
| 123 | 103 | 164 | 144 | 164 | 124 | 123 | 103 | 103 | 82 | 103 | 82 |
| 198 | 185 | 277 | 246 | 287 | 246 | 205 | 165 | 164 | 142 | 163 | 119 |
| 328 | 308 | 453 | 410 | 492 | 410 | 369 | 316 | 287 | 248 | 296 | 205 |
| 492 | 482 | 700 | 615 | 740 | 616 | 554 | 474 | 489 | 400 | 375 | 328 |
| 615 | 575 | 862 | 738 | 949 | 797 | 697 | 616 | 492 | 474 | 496 | 410 |
| 988 | 882 | 1,395 | 1,087 | 1,395 | 1,274 | 1,215 | 1,184 | 862 | 1,422 | 657 | 656 |
| 375 | 344 | 430 | 381 | 410 | 353 | 354 | 322 | 323 | 264 | 316 | 258 |
| 195 | 172 | 251 | 214 | 231 | 189 | 172 | 156 | 136 | 93 | 111 | 103 |
| 408 | 386 | 467 | 438 | 465 | 389 | 410 | 346 | 364 | 318 | 350 | 315 |
| 199 | 169 | 232 | 216 | 249 | 175 | 196 | 150 | 165 | 132 | 151 | 128 |
| 469 | 447 | 586 | 496 | 610 | 531 | 511 | 454 | 423 | 437 | 354 | 312 |
| 235 | 228 | 294 | 224 | 316 | 302 | 301 | 277 | 184 | 376 | 150 | 141 |
| 3,690 | 2,460 | 2,598 | 2,224 | 2,952 | 3,936 | 5,228 | 2,952 | 1,107 | 3,336 | 984 | 984 |
| 5,166 | 5,904 | 6,428 | 4,554 | 7,565 | 10,824 | 9,410 | 7,110 | 2,745 | 9,024 | 1,968 | 1,968 |
| 314 | 296 | 364 | 321 | 355 | 312 | 330 | 291 | 300 | 273 | 301 | 255 |
| 119 | 116 | 142 | 115 | 133 | 118 | 129 | 104 | 108 | 91 | 107 | 95 |
| 185 | 178 | 186 | 185 | 185 | 178 | 185 | 153 | 183 | 123 | 124 | 119 |
| 246 | 246 | 246 | 246 | 246 | 246 | 246 | 237 | 246 | 237 | 237 | 185 |
| 296 | 246 | 369 | 308 | 369 | 296 | 308 | 246 | 296 | 246 | 296 | 246 |
| 369 | 369 | 372 | 365 | 372 | 369 | 369 | 356 | 369 | 310 | 369 | 308 |
| 492 | 492 | 492 | 492 | 492 | 492 | 492 | 415 | 372 | 369 | 492 | 369 |
| 492 | 492 | 496 | 492 | 496 | 492 | 492 | 492 | 492 | 474 | 492 | 474 |
| 739 | 683 | 984 | 738 | 922 | 711 | 744 | 533 | 829 | 496 | 496 | 492 |
| 1,479 | 1,538 | 1,538 | 1,235 | 1,488 | 1,968 | 1,968 | 984 | 948 | 738 | 496 | 492 |
| 33.2 | 17.2 | 28.3 | 22.8 | 59.7 | 69.9 | 47.5 | 55.4 | 1.9 | 17.1 | 0 | 0 |
| 1.2 | .6 | 1.0 | .7 | .7 | .7 | .9 | .7 | .4 | 2.1 | 0 | 0 |

TABLE 1.05.--COLA-TYPE CARBONATED SOFT DRINKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 36.1 | 2.2 | 31.0 | 37.8 | 38.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 16.7 | 1.4 | 17.7 | 20.9 | 22.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 10.5 | .2 | 8.4 | 10.9 | 10.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 9.0 | .7 | 4.8 | 6.0 | 6.1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 2.4 | 2.4 | 2.0 | 2.0 | 2.0 |
| STANDARD DEVIATION.....DO..... | 1.9 | 1.5 | 1.5 | 1.4 | 1.4 |
| MAXIMUM.....DO..... | 22.0 | 5.0 | 13.0 | 13.0 | 16.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 251 | 74 | 113 | 144 | 164 |
| STANDARD DEVIATION.....GM..... | 230 | 42 | 106 | 110 | 126 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 62 | | 21 | 41 | 41 |
| 25TH.....GM..... | 103 | | 41 | 62 | 82 |
| 50TH.....GM..... | 164 | | 82 | 103 | 123 |
| 75TH.....GM..... | 328 | | 144 | 185 | 226 |
| 90TH.....GM..... | 492 | | 226 | 287 | 328 |
| 95TH.....GM..... | 666 | | 287 | 369 | 390 |
| 99TH.....GM..... | 1,107 | | 574 | 553 | 657 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 342 | 174 | 192 | 249 | 289 |
| STANDARD DEVIATION.....GM..... | 186 | 106 | 109 | 136 | 157 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 383 | 154 | 179 | 264 | 299 |
| STANDARD DEVIATION.....GM..... | 189 | 0 | 78 | 127 | 127 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 502 | 102 | 280 | 297 | 347 |
| STANDARD DEVIATION.....GM..... | 299 | 43 | 159 | 134 | 159 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 5,228 | 400 | 1,538 | 984 | 1,845 |
| MAXIMUM OVER 3 DAYS.....GM..... | 10,824 | 492 | 3,198 | 2,276 | 3,444 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 312 | 95 | 170 | 213 | 248 |
| STANDARD DEVIATION.....GM..... | 121 | 56 | 75 | 87 | 93 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 154 | | 62 | 123 | 123 |
| 25TH.....GM..... | 246 | | 123 | 123 | 185 |
| 50TH.....GM..... | 308 | | 153 | 185 | 246 |
| 75TH.....GM..... | 369 | | 246 | 246 | 308 |
| 90TH.....GM..... | 492 | | 246 | 308 | 369 |
| 95TH.....GM..... | 492 | | 308 | 369 | 369 |
| 99TH.....GM..... | 738 | | 369 | 492 | 492 |
| MAXIMUM.....GM..... | 1,968 | 246 | 492 | 984 | 984 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 222.8 | 0 | .8 | 11.8 | 1.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .7 | 0 | .1 | .9 | .1 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 44.9 | 44.4 | 54.8 | 54.8 | 57.3 | 46.1 | 30.3 | 26.4 | 12.8 | 10.4 | 9.2 | 8.2 |
| 24.0 | 22.4 | 21.6 | 21.1 | 21.5 | 19.5 | 14.4 | 13.2 | 7.1 | 5.4 | 4.8 | 5.3 |
| 12.3 | 13.9 | 18.1 | 17.7 | 16.6 | 13.5 | 8.6 | 7.2 | 3.5 | 2.6 | 2.8 | 1.5 |
| 8.6 | 8.1 | 15.1 | 16.0 | 19.2 | 13.1 | 7.3 | 6.0 | 2.3 | 2.4 | 1.6 | 1.5 |
| 2.1 | 2.2 | 2.6 | 2.6 | 2.9 | 2.7 | 2.3 | 2.2 | 2.0 | 2.0 | 2.0 | 1.8 |
| 1.5 | 1.5 | 1.9 | 1.7 | 2.2 | 2.1 | 1.7 | 1.7 | 1.4 | 1.2 | 1.3 | 1.1 |
| 12.0 | 20.0 | 12.0 | 11.0 | 17.0 | 22.0 | 15.0 | 19.0 | 7.0 | 6.0 | 6.0 | 6.0 |
| 220 | 213 | 314 | 278 | 339 | 278 | 255 | 219 | 204 | 170 | 218 | 153 |
| 169 | 163 | 243 | 203 | 288 | 264 | 229 | 203 | 149 | 115 | 169 | 128 |
| 82 | 62 | 82 | 82 | 82 | 62 | 82 | 62 | 62 | 41 | | 41 |
| 103 | 82 | 123 | 123 | 123 | 123 | 123 | 82 | 103 | 82 | | 67 |
| 164 | 164 | 246 | 246 | 246 | 205 | 164 | 164 | 164 | 164 | | 103 |
| 287 | 267 | 410 | 369 | 451 | 369 | 328 | 267 | 246 | 246 | | 205 |
| 431 | 410 | 656 | 534 | 738 | 574 | 492 | 410 | 492 | 328 | | 308 |
| 534 | 534 | 779 | 657 | 903 | 739 | 656 | 554 | 492 | 390 | | 410 |
| 821 | 780 | 1,232 | 985 | 1,395 | 1,209 | 985 | 985 | 616 | 512 | | 657 |
| 371 | 344 | 435 | 379 | 401 | 343 | 363 | 318 | 334 | 252 | 311 | 244 |
| 186 | 188 | 248 | 190 | 228 | 173 | 168 | 147 | 140 | 115 | 132 | 106 |
| 387 | 377 | 467 | 430 | 449 | 382 | 394 | 346 | 372 | 321 | 413 | 342 |
| 196 | 163 | 240 | 211 | 214 | 171 | 172 | 148 | 163 | 105 | 147 | 102 |
| 434 | 417 | 558 | 469 | 603 | 544 | 508 | 452 | 420 | 319 | 467 | 332 |
| 201 | 209 | 270 | 217 | 324 | 341 | 310 | 287 | 153 | 90 | 193 | 165 |
| 3,690 | 2,460 | 2,214 | 2,091 | 2,952 | 3,936 | 5,228 | 2,952 | 1,107 | 738 | 984 | 984 |
| 5,166 | 5,904 | 4,428 | 3,813 | 6,704 | 10,824 | 9,410 | 6,888 | 2,214 | 1,845 | 1,968 | 1,968 |
| 308 | 292 | 359 | 321 | 355 | 312 | 333 | 294 | 302 | 256 | 327 | 257 |
| 108 | 116 | 135 | 112 | 127 | 114 | 126 | 99 | 90 | 79 | 111 | 102 |
| 185 | 185 | 185 | 185 | 185 | 185 | 185 | 185 | 185 | 123 | | 123 |
| 246 | 246 | 246 | 246 | 246 | 246 | 246 | 246 | 246 | 200 | | 185 |
| 246 | 246 | 369 | 308 | 369 | 246 | 308 | 246 | 308 | 246 | | 246 |
| 369 | 369 | 431 | 369 | 369 | 369 | 369 | 369 | 369 | 308 | | 308 |
| 492 | 492 | 492 | 492 | 492 | 492 | 492 | 492 | 369 | 369 | | 369 |
| 492 | 492 | 492 | 492 | 492 | 492 | 492 | 492 | 492 | 369 | | 492 |
| 615 | 616 | 739 | 677 | 739 | 615 | 739 | 492 | 492 | 492 | | 492 |
| 1,107 | 1,538 | 1,538 | 1,230 | 1,476 | 1,968 | 1,968 | 984 | 677 | 738 | 492 | 492 |
| 19.5 | 10.6 | 16.5 | 14.0 | 42.6 | 44.7 | 25.8 | 32.7 | 0 | 2.2 | 0 | 0 |
| 1.0 | .5 | .8 | .7 | .7 | .7 | .8 | .8 | 0 | .6 | 0 | 0 |

TABLE 1.06.--ARTIFICIALLY SWEETENED SOFT DRINKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 8.1 | .4 | 2.5 | 3.8 | 3.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.0 | .4 | 1.6 | 2.6 | 2.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.1 | 0 | .6 | 1.0 | .8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.1 | 0 | .3 | .3 | .3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER.... | 2.4 | 1.0 | 1.6 | 1.7 | 1.6 |
| STANDARD DEVIATION.....DO..... | 2.0 | .0 | 1.0 | .9 | .9 |
| MAXIMUM.....DO..... | 19.0 | 1.0 | 5.0 | 5.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 245 | 17 | 82 | 97 | 109 |
| STANDARD DEVIATION.....GM..... | 232 | 17 | 68 | 64 | 67 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 59 | | | 20 | 40 |
| 25TH.....GM..... | 99 | | | 59 | 59 |
| 50TH.....GM..... | 158 | | | 79 | 79 |
| 75TH.....GM..... | 316 | | | 119 | 138 |
| 90TH.....GM..... | 474 | | | 217 | 237 |
| 95TH.....GM..... | 711 | | | 237 | 237 |
| 99TH.....GM..... | 1,185 | | | 277 | 287 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 322 | 52 | 174 | 207 | 226 |
| STANDARD DEVIATION.....GM..... | 160 | 50 | 72 | 111 | 100 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 383 | 0 | 123 | 206 | 228 |
| STANDARD DEVIATION.....GM..... | 192 | 0 | 29 | 84 | 84 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 499 | 0 | 218 | 208 | 213 |
| STANDARD DEVIATION.....GM..... | 298 | 0 | 118 | 83 | 76 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 2,844 | 119 | 533 | 711 | 474 |
| MAXIMUM OVER 3 DAYS.....GM..... | 8,532 | 119 | 1,244 | 830 | 860 |
| QUANTITY CONSUMED PER EATING OCCASION: - | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 306 | 52 | 150 | 168 | 200 |
| STANDARD DEVIATION.....GM..... | 117 | 50 | 56 | 75 | 81 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 178 | | | 59 | 59 |
| 25TH.....GM..... | 237 | | | 119 | 147 |
| 50TH.....GM..... | 296 | | | 178 | 178 |
| 75TH.....GM..... | 356 | | | 237 | 237 |
| 90TH.....GM..... | 474 | | | 237 | 356 |
| 95TH.....GM..... | 474 | | | 296 | 356 |
| 99TH.....GM..... | 712 | | | 356 | 474 |
| MAXIMUM.....GM..... | 1,896 | 119 | 356 | 356 | 474 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER.... | 52.8 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .7 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.0 | 4.7 | 3.4 | 11.4 | 5.2 | 15.7 | 7.0 | 14.2 | 2.2 | 5.1 | 2.2 | 4.1 |
| 2.1 | 2.7 | 2.2 | 6.6 | 2.8 | 6.2 | 3.4 | 6.8 | .7 | 3.0 | .7 | 2.7 |
| .4 | 1.3 | .6 | 2.5 | 1.1 | 4.4 | 1.8 | 3.7 | .8 | 1.0 | .5 | .8 |
| .5 | .7 | .5 | 2.2 | 1.2 | 5.1 | 1.7 | 3.8 | .7 | 1.1 | .9 | .6 |
| 1.8 | 1.9 | 1.8 | 2.2 | 2.5 | 2.7 | 2.4 | 2.4 | 2.5 | 2.4 | 2.7 | 1.6 |
| 1.2 | 1.2 | 1.1 | 1.9 | 2.4 | 2.0 | 1.9 | 2.0 | 1.4 | 2.8 | 1.6 | .9 |
| 6.0 | 7.0 | 5.0 | 12.0 | 15.0 | 12.0 | 12.0 | 19.0 | 6.0 | 18.0 | 6.0 | 4.0 |
| 172 | 189 | 185 | 222 | 273 | 287 | 258 | 243 | 267 | 249 | 193 | 141 |
| 130 | 141 | 125 | 201 | 257 | 243 | 239 | 228 | 187 | 375 | 87 | 99 |
| 40 | 59 | | 59 | 79 | 59 | 59 | 59 | | 79 | | |
| 79 | 79 | | 99 | 119 | 119 | 113 | 119 | | 79 | | |
| 119 | 138 | | 158 | 158 | 237 | 158 | 158 | | 119 | | |
| 237 | 237 | | 257 | 339 | 359 | 326 | 316 | | 237 | | |
| 435 | 375 | | 474 | 612 | 593 | 592 | 474 | | 415 | | |
| 464 | 514 | | 633 | 949 | 790 | 712 | 633 | | 474 | | |
| 513 | 633 | | 1,265 | 1,302 | 1,224 | 1,068 | 1,185 | | 2,844 | | |
| 298 | 315 | 388 | 340 | 356 | 322 | 315 | 341 | 294 | 292 | 341 | 273 |
| 140 | 146 | 242 | 165 | 116 | 172 | 111 | 176 | 65 | 102 | 25 | 113 |
| 449 | 379 | 332 | 375 | 529 | 397 | 416 | 369 | 383 | 357 | 237 | 242 |
| 182 | 160 | 140 | 131 | 342 | 168 | 190 | 178 | 221 | 144 | 0 | 92 |
| 370 | 408 | 392 | 520 | 561 | 524 | 537 | 473 | 439 | 652 | 272 | 327 |
| 70 | 150 | 93 | 260 | 293 | 267 | 294 | 307 | 158 | 630 | 74 | 92 |
| 948 | 1,067 | 1,422 | 2,133 | 2,133 | 2,074 | 2,489 | 2,844 | 948 | 2,844 | 533 | 474 |
| 1,541 | 2,489 | 1,896 | 3,911 | 3,911 | 5,036 | 5,333 | 7,110 | 2,489 | 8,532 | 1,126 | 1,422 |
| 292 | 304 | 312 | 304 | 330 | 317 | 327 | 298 | 326 | 309 | 218 | 264 |
| 107 | 109 | 106 | 103 | 107 | 126 | 123 | 108 | 161 | 90 | 70 | 101 |
| 119 | 178 | | 178 | 178 | 178 | 178 | 178 | | 178 | | |
| 237 | 237 | | 237 | 237 | 237 | 237 | 237 | | 237 | | |
| 296 | 296 | | 296 | 356 | 356 | 356 | 296 | | 296 | | |
| 356 | 356 | | 356 | 356 | 356 | 356 | 356 | | 356 | | |
| 356 | 356 | | 474 | 474 | 474 | 474 | 356 | | 474 | | |
| 474 | 474 | | 474 | 474 | 474 | 474 | 474 | | 474 | | |
| 474 | 711 | | 534 | 712 | 712 | 948 | 711 | | 533 | | |
| 948 | 1,067 | 474 | 948 | 1,067 | 1,896 | 1,067 | 948 | 948 | 533 | 356 | 474 |
| 0 | 2.2 | 0 | 3.1 | 3.1 | 12.6 | 3.5 | 14.3 | 0 | 13.9 | 0 | 0 |
| 0 | 1.2 | 0 | .9 | .6 | .6 | .4 | .6 | 0 | 6.4 | 0 | 0 |

TABLE 1.07.--NONCARBONATED SOFT DRINKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 10.9 | 4.2 | 25.4 | 30.4 | 27.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.5 | 1.8 | 11.1 | 13.3 | 12.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.1 | 1.8 | 7.3 | 9.4 | 8.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.3 | .6 | 7.0 | 7.6 | 6.6 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 2.2 | 2.2 | 2.5 | 2.6 | 2.4 |
| STANDARD DEVIATION.....DO..... | 1.6 | 1.2 | 1.9 | 1.8 | 1.6 |
| MAXIMUM.....DO..... | 12.0 | 5.0 | 12.0 | 12.0 | 10.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 216 | 95 | 160 | 192 | 200 |
| STANDARD DEVIATION.....GM..... | 190 | 70 | 150 | 151 | 154 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 62 | | 41 | 51 | 62 |
| 25TH.....GM..... | 83 | | 62 | 83 | 83 |
| 50TH.....GM..... | 165 | | 103 | 145 | 165 |
| 75TH.....GM..... | 269 | | 207 | 248 | 248 |
| 90TH.....GM..... | 413 | | 331 | 413 | 413 |
| 95TH.....GM..... | 559 | | 434 | 496 | 496 |
| 99TH.....GM..... | 993 | | 826 | 662 | 744 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 339 | 144 | 198 | 268 | 299 |
| STANDARD DEVIATION.....GM..... | 219 | 125 | 114 | 128 | 138 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 360 | 159 | 219 | 290 | 308 |
| STANDARD DEVIATION.....GM..... | 218 | 74 | 94 | 148 | 127 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 434 | 214 | 325 | 368 | 383 |
| STANDARD DEVIATION.....GM..... | 238 | 32 | 184 | 162 | 180 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 3,968 | 496 | 1,488 | 1,488 | 1,704 |
| MAXIMUM OVER 3 DAYS.....GM..... | 7,378 | 744 | 3,720 | 3,224 | 3,564 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 290 | 128 | 192 | 225 | 255 |
| STANDARD DEVIATION.....GM..... | 148 | 64 | 92 | 86 | 93 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 124 | | 93 | 124 | 124 |
| 25TH.....GM..... | 248 | | 124 | 186 | 248 |
| 50TH.....GM..... | 248 | | 186 | 248 | 248 |
| 75TH.....GM..... | 310 | | 248 | 248 | 248 |
| 90TH.....GM..... | 496 | | 248 | 310 | 372 |
| 95TH.....GM..... | 496 | | 372 | 372 | 496 |
| 99TH.....GM..... | 868 | | 496 | 496 | 496 |
| MAXIMUM.....GM..... | 3,968 | 248 | 992 | 744 | 930 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 42.9 | 0 | 3.4 | 10.6 | 4.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .5 | 0 | .5 | .8 | .4 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 20.8 | 19.9 | 13.5 | 14.6 | 9.6 | 10.5 | 4.1 | 4.6 | 2.0 | 2.3 | 2.1 | 1.0 |
| 10.7 | 9.7 | 7.2 | 7.3 | 5.0 | 5.9 | 2.3 | 2.8 | .7 | 1.3 | .7 | .6 |
| 6.3 | 6.1 | 3.9 | 4.3 | 2.5 | 2.8 | 1.4 | 1.2 | .4 | .4 | .6 | .2 |
| 3.9 | 4.1 | 2.4 | 3.0 | 2.1 | 1.9 | .5 | .7 | .9 | .6 | .7 | .2 |
| 2.2 | 2.2 | 2.3 | 2.1 | 2.2 | 2.1 | 1.8 | 1.8 | 3.3 | 2.1 | 2.9 | 1.7 |
| 1.6 | 1.5 | 1.6 | 1.4 | 1.7 | 1.5 | 1.3 | 1.2 | 2.3 | 1.6 | 2.0 | .8 |
| 10.0 | 9.0 | 9.0 | 8.0 | 12.0 | 9.0 | 7.0 | 8.0 | 7.0 | 7.0 | 6.0 | 3.0 |
| 238 | 212 | 285 | 226 | 280 | 215 | 222 | 173 | 396 | 231 | 261 | 172 |
| 197 | 161 | 234 | 170 | 256 | 189 | 250 | 132 | 345 | 226 | 200 | 86 |
| 83 | 62 | 83 | 83 | 83 | 62 | 83 | 55 | | | | |
| 88 | 83 | 103 | 83 | 124 | 83 | 83 | 83 | | | | |
| 165 | 165 | 227 | 165 | 207 | 165 | 165 | 124 | | | | |
| 310 | 248 | 372 | 289 | 331 | 248 | 248 | 217 | | | | |
| 475 | 413 | 496 | 413 | 578 | 413 | 413 | 331 | | | | |
| 620 | 579 | 827 | 496 | 703 | 496 | 537 | 413 | | | | |
| 993 | 745 | 1,117 | 827 | 1,158 | 993 | 1,591 | 662 | | | | |
| 383 | 342 | 458 | 363 | 463 | 341 | 350 | 305 | 325 | 287 | 248 | 392 |
| 205 | 191 | 305 | 202 | 402 | 174 | 178 | 144 | 93 | 94 | 0 | 244 |
| 432 | 334 | 520 | 373 | 455 | 393 | 426 | 313 | 316 | 593 | 248 | 403 |
| 289 | 149 | 287 | 135 | 182 | 183 | 429 | 108 | 52 | 474 | 0 | 0 |
| 461 | 429 | 585 | 449 | 551 | 463 | 558 | 396 | 713 | 437 | 516 | 201 |
| 220 | 176 | 266 | 204 | 358 | 267 | 340 | 159 | 293 | 177 | 92 | 41 |
| 2,790 | 1,984 | 2,480 | 1,488 | 3,968 | 1,984 | 3,968 | 1,240 | 1,736 | 1,984 | 930 | 806 |
| 3,906 | 2,976 | 3,906 | 3,224 | 6,448 | 3,968 | 7,378 | 2,232 | 2,976 | 2,728 | 1,860 | 806 |
| 320 | 286 | 371 | 316 | 385 | 310 | 365 | 283 | 356 | 333 | 273 | 298 |
| 145 | 126 | 181 | 126 | 217 | 127 | 250 | 96 | 190 | 146 | 30 | 122 |
| 186 | 124 | 218 | 186 | 186 | 186 | 248 | 155 | | | | |
| 248 | 248 | 248 | 248 | 248 | 248 | 248 | 248 | | | | |
| 248 | 248 | 310 | 248 | 310 | 248 | 279 | 248 | | | | |
| 372 | 310 | 496 | 372 | 496 | 372 | 466 | 310 | | | | |
| 496 | 496 | 496 | 496 | 496 | 496 | 496 | 496 | | | | |
| 496 | 496 | 621 | 496 | 869 | 496 | 621 | 496 | | | | |
| 745 | 744 | 993 | 745 | 993 | 745 | 993 | 496 | | | | |
| 1,488 | 1,984 | 1,488 | 992 | 1,984 | 992 | 3,968 | 744 | 992 | 992 | 310 | 558 |
| 8.1 | 6.6 | .8 | 3.2 | 1.4 | 2.4 | 0 | 1.8 | 0 | 0 | 0 | 0 |
| .8 | .7 | .2 | .7 | .2 | .2 | 0 | .3 | 0 | 0 | 0 | 0 |

TABLE 1.08.--FRUIT ADES AND DRINKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FCOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 13.7 | 3.5 | 22.2 | 22.4 | 23.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 8.2 | 2.1 | 10.7 | 11.9 | 13.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.2 | .8 | 6.2 | 6.2 | 5.7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.3 | .7 | 5.4 | 4.3 | 4.2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.9 | 1.8 | 2.3 | 2.2 | 2.0 |
| STANDARD DEVIATION.....DO..... | 1.3 | .8 | 1.9 | 1.5 | 1.4 |
| MAXIMUM.....DO..... | 23.0 | 3.0 | 18.0 | 10.0 | 8.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 170 | 91 | 133 | 146 | 153 |
| STANDARD DEVIATION.....GM..... | 149 | 77 | 110 | 109 | 113 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 42 | | 21 | 42 | 42 |
| 25TH.....GM..... | 83 | | 62 | 83 | 83 |
| 50TH.....GM..... | 125 | | 83 | 124 | 104 |
| 75TH.....GM..... | 208 | | 167 | 188 | 208 |
| 90TH.....GM..... | 333 | | 250 | 271 | 313 |
| 95TH.....GM..... | 417 | | 394 | 333 | 405 |
| 99TH.....GM..... | 707 | | 531 | 584 | 521 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 311 | 156 | 207 | 272 | 273 |
| STANDARD DEVIATION.....GM..... | 188 | 132 | 173 | 157 | 131 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 321 | 191 | 202 | 247 | 295 |
| STANDARD DEVIATION.....GM..... | 176 | 109 | 96 | 147 | 138 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 342 | 172 | 258 | 273 | 299 |
| STANDARD DEVIATION.....GM..... | 224 | 79 | 121 | 122 | 132 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 4,464 | 500 | 1,250 | 1,125 | 1,000 |
| MAXIMUM OVER 3 DAYS.....GM..... | 10,416 | 688 | 1,625 | 2,460 | 2,223 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 269 | 150 | 171 | 202 | 233 |
| STANDARD DEVIATION.....GM..... | 144 | 80 | 79 | 87 | 101 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 125 | | 63 | 95 | 125 |
| 25TH.....GM..... | 188 | | 125 | 125 | 186 |
| 50TH.....GM..... | 250 | | 155 | 188 | 249 |
| 75TH.....GM..... | 310 | | 249 | 250 | 250 |
| 90TH.....GM..... | 496 | | 250 | 250 | 372 |
| 95TH.....GM..... | 500 | | 250 | 375 | 496 |
| 99TH.....GM..... | 750 | | 500 | 500 | 500 |
| MAXIMUM.....GM..... | 3,000 | 250 | 750 | 750 | 871 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 41.1 | 0 | 3.2 | 1.1 | 1.7 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | .6 | .1 | .2 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 21.3 | 21.1 | 18.4 | 19.0 | 12.9 | 13.9 | 8.5 | 9.4 | 6.7 | 7.9 | 9.0 | 7.8 |
| 12.2 | 12.7 | 9.1 | 11.5 | 7.6 | 9.1 | 5.8 | 6.0 | 4.3 | 4.6 | 3.4 | 4.5 |
| 4.3 | 5.5 | 5.5 | 5.5 | 2.9 | 3.1 | 1.4 | 2.0 | 1.0 | 1.5 | 3.2 | 1.5 |
| 4.8 | 2.9 | 3.8 | 1.9 | 2.4 | 1.8 | 1.3 | 1.3 | 1.5 | 1.8 | 2.4 | 1.8 |
| 2.0 | 1.9 | 2.0 | 1.8 | 1.9 | 1.8 | 1.7 | 1.7 | 1.8 | 1.9 | 2.6 | 1.8 |
| 1.4 | 1.2 | 1.3 | 1.0 | 1.4 | 1.2 | 1.1 | 1.4 | 1.3 | 1.2 | 1.7 | 1.1 |
| 8.0 | 9.0 | 7.0 | 6.0 | 10.0 | 9.0 | 6.0 | 23.0 | 7.0 | 6.0 | 7.0 | 6.0 |
| 183 | 165 | 227 | 171 | 223 | 160 | 169 | 157 | 158 | 161 | 217 | 139 |
| 166 | 123 | 183 | 115 | 192 | 127 | 137 | 185 | 113 | 147 | 135 | 100 |
| 62 | 62 | 83 | 42 | 63 | 42 | 52 | 42 | 52 | 42 | | 42 |
| 83 | 83 | 103 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | | 83 |
| 125 | 125 | 167 | 125 | 165 | 125 | 125 | 103 | 103 | 104 | | 83 |
| 228 | 208 | 250 | 248 | 271 | 206 | 207 | 167 | 188 | 188 | | 167 |
| 372 | 333 | 457 | 333 | 496 | 313 | 331 | 292 | 333 | 372 | | 310 |
| 500 | 417 | 583 | 414 | 562 | 396 | 417 | 413 | 395 | 375 | | 333 |
| 745 | 625 | 998 | 583 | 1,001 | 706 | 833 | 661 | 496 | 1,000 | | 498 |
| 318 | 315 | 416 | 327 | 371 | 307 | 339 | 290 | 271 | 321 | 372 | 256 |
| 190 | 189 | 282 | 163 | 191 | 166 | 183 | 141 | 98 | 386 | 271 | 142 |
| 327 | 325 | 368 | 364 | 426 | 312 | 361 | 315 | 380 | 257 | 330 | 297 |
| 169 | 156 | 224 | 147 | 250 | 127 | 170 | 173 | 194 | 144 | 146 | 147 |
| 347 | 325 | 415 | 329 | 468 | 369 | 342 | 354 | 288 | 290 | 347 | 221 |
| 242 | 157 | 233 | 149 | 237 | 182 | 206 | 392 | 84 | 143 | 123 | 113 |
| 3,486 | 1,500 | 2,976 | 1,375 | 3,750 | 1,500 | 1,500 | 4,464 | 992 | 3,000 | 992 | 746 |
| 6,225 | 2,486 | 3,720 | 1,875 | 4,875 | 2,728 | 2,976 | 10,416 | 1,488 | 3,000 | 1,619 | 1,494 |
| 277 | 267 | 334 | 292 | 350 | 271 | 300 | 273 | 259 | 255 | 249 | 231 |
| 138 | 127 | 246 | 122 | 175 | 116 | 137 | 116 | 99 | 233 | 140 | 105 |
| 125 | 125 | 125 | 125 | 156 | 125 | 125 | 125 | 125 | 124 | | 63 |
| 188 | 247 | 248 | 248 | 249 | 190 | 248 | 247 | 188 | 188 | | 154 |
| 250 | 250 | 250 | 250 | 250 | 250 | 250 | 250 | 249 | 248 | | 250 |
| 310 | 250 | 375 | 372 | 375 | 313 | 374 | 310 | 250 | 250 | | 250 |
| 496 | 375 | 500 | 498 | 500 | 375 | 496 | 496 | 496 | 372 | | 313 |
| 500 | 500 | 625 | 500 | 745 | 500 | 500 | 500 | 496 | 496 | | 372 |
| 751 | 750 | 1,500 | 745 | 1,000 | 500 | 750 | 625 | 500 | 561 | | 563 |
| 1,500 | 1,500 | 2,976 | 1,000 | 1,500 | 1,488 | 1,000 | 750 | 500 | 3,000 | 992 | 620 |
| 7.9 | 4.6 | 2.7 | 2.2 | 3.4 | 2.2 | 6.0 | 3.1 | 0 | .9 | 0 | 2.2 |
| .9 | .5 | .5 | .5 | .3 | .2 | .9 | .3 | 0 | .4 | 0 | 1.6 |

TABLE 1.09.--BEER AND ALE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 8.4 | 0 | .3 | .1 | 0 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.3 | 0 | .2 | .1 | 0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.2 | 0 | 0 | 0 | 0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.9 | 0 | .1 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 2.4 | 0 | 1.6 | 1.0 | 0 |
| STANDARD DEVIATION.....DO..... | 2.3 | 0 | .9 | .0 | 0 |
| MAXIMUM.....DO..... | 27.0 | 0 | 3.0 | 1.0 | 0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 455 | 0 | 37 | 30 | 0 |
| STANDARD DEVIATION.....GM..... | 556 | 0 | 24 | 0 | 0 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 80 | | | | |
| 25TH.....GM..... | 120 | | | | |
| 50TH.....GM..... | 240 | | | | |
| 75TH.....GM..... | 600 | | | | |
| 90TH.....GM..... | 961 | | | | |
| 95TH.....GM..... | 1,321 | | | | |
| 99TH.....GM..... | 2,322 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 675 | 0 | 64 | 90 | 0 |
| STANDARD DEVIATION.....GM..... | 748 | 0 | 30 | 0 | 0 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 733 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 551 | 0 | 0 | 0 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 926 | 0 | 70 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 842 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 15,120 | 0 | 90 | 90 | 0 |
| MAXIMUM OVER 3 DAYS.....GM..... | 34,560 | 0 | 210 | 90 | 0 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 557 | 0 | 68 | 90 | 0 |
| STANDARD DEVIATION.....GM..... | 437 | 0 | 22 | 0 | 0 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 240 | | | | |
| 25TH.....GM..... | 360 | | | | |
| 50TH.....GM..... | 360 | | | | |
| 75TH.....GM..... | 720 | | | | |
| 90TH.....GM..... | 1,080 | | | | |
| 95TH.....GM..... | 1,441 | | | | |
| 99TH.....GM..... | 2,163 | | | | |
| MAXIMUM.....GM..... | 8,640 | 0 | 90 | 90 | 0 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 25.5 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|--------|--------|--------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 0 | (2) | 3.9 | 1.4 | 26.8 | 7.8 | 21.5 | 5.4 | 10.8 | 1.7 | 7.4 | 1.3 |
| 0 | (2) | 2.7 | 1.1 | 12.8 | 5.1 | 9.7 | 3.3 | 4.2 | .9 | 3.7 | .7 |
| 0 | 0 | .9 | .4 | 7.9 | 1.7 | 6.1 | 1.0 | 2.2 | .5 | .8 | .2 |
| 0 | 0 | .4 | 0 | 6.2 | .9 | 5.8 | 1.1 | 4.4 | .3 | 2.8 | .4 |
| 0 | 1.0 | 2.0 | 1.3 | 2.6 | 1.9 | 2.7 | 2.0 | 2.6 | 2.0 | 2.9 | 2.5 |
| 0 | .0 | 1.8 | .5 | 2.5 | 1.4 | 2.5 | 1.5 | 2.0 | 1.3 | 2.5 | 2.2 |
| 0 | 1.0 | 11.0 | 3.0 | 27.0 | 10.0 | 23.0 | 11.0 | 10.0 | 5.0 | 11.0 | 7.0 |
| 0 | 120 | 411 | 223 | 550 | 319 | 497 | 279 | 446 | 267 | 346 | 303 |
| 0 | 0 | 374 | 192 | 715 | 320 | 541 | 233 | 460 | 206 | 302 | 291 |
| | | 120 | | 120 | 80 | 100 | 70 | 110 | | | |
| | | 120 | | 160 | 120 | 160 | 120 | 120 | | | |
| | | 240 | | 360 | 220 | 360 | 240 | 300 | | | |
| | | 601 | | 720 | 420 | 601 | 360 | 661 | | | |
| | | 961 | | 1,081 | 721 | 1,081 | 600 | 921 | | | |
| | | 1,200 | | 1,441 | 960 | 1,482 | 721 | 1,201 | | | |
| | | 1,681 | | 2,761 | 1,441 | 2,524 | 1,082 | 2,641 | | | |
| 0 | 360 | 902 | 446 | 806 | 576 | 716 | 522 | 432 | 332 | 421 | 309 |
| 0 | 0 | 862 | 339 | 903 | 462 | 840 | 451 | 254 | 133 | 226 | 59 |
| 0 | 0 | 941 | 677 | 865 | 698 | 685 | 466 | 554 | 535 | 563 | 420 |
| 0 | 0 | 600 | 307 | 527 | 510 | 626 | 200 | 312 | 180 | 179 | 0 |
| 0 | 0 | 687 | 0 | 1,100 | 745 | 971 | 565 | 775 | 569 | 609 | 631 |
| 0 | 0 | 475 | 0 | 1,188 | 444 | 660 | 252 | 541 | 167 | 320 | 301 |
| 0 | 360 | 3,840 | 1,440 | 15,120 | 3,240 | 11,160 | 4,320 | 2,880 | 1,080 | 1,440 | 1,080 |
| 0 | 360 | 5,040 | 2,640 | 34,560 | 7,200 | 15,480 | 4,320 | 7,920 | 2,520 | 3,960 | 2,520 |
| 0 | 360 | 604 | 520 | 628 | 511 | 555 | 424 | 511 | 409 | 357 | 366 |
| 0 | 0 | 474 | 350 | 511 | 369 | 428 | 215 | 342 | 164 | 108 | 139 |
| | | 240 | | 240 | 209 | 240 | 210 | 210 | | | |
| | | 360 | | 360 | 360 | 360 | 360 | 360 | | | |
| | | 360 | | 360 | 360 | 360 | 360 | 360 | | | |
| | | 721 | | 721 | 720 | 720 | 360 | 720 | | | |
| | | 1,440 | | 1,082 | 961 | 1,080 | 721 | 1,080 | | | |
| | | 1,441 | | 1,621 | 1,441 | 1,262 | 721 | 1,440 | | | |
| | | 2,940 | | 2,164 | 1,922 | 2,163 | 1,082 | 1,800 | | | |
| 0 | 360 | 2,940 | 1,440 | 7,200 | 2,520 | 8,640 | 1,440 | 1,800 | 1,080 | 720 | 720 |
| 0 | 0 | 0 | 1.6 | 3.4 | 8.0 | 11.3 | 0 | 1.2 | 0 | 0 | 0 |
| 0 | 0 | 0 | 5.8 | .1 | 1.0 | .4 | 0 | .4 | 0 | 0 | 0 |

TABLE 1.10.--WINE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.8 | 0 | 0 | .1 | .2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.1 | 0 | 0 | .1 | .2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.0 | 0 | 0 | 0 | 0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .7 | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.7 | 0 | 0 | 1.0 | 1.0 |
| STANDARD DEVIATION.....DO..... | 1.1 | 0 | 0 | .0 | .0 |
| MAXIMUM.....DO..... | 9.0 | 0 | 0 | 1.0 | 1.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 113 | 0 | 0 | 56 | 45 |
| STANDARD DEVIATION.....GM..... | 118 | 0 | 0 | 33 | 27 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 19 | | | | |
| 25TH.....GM..... | 39 | | | | |
| 50TH.....GM..... | 77 | | | | |
| 75TH.....GM..... | 135 | | | | |
| 90TH.....GM..... | 232 | | | | |
| 95TH.....GM..... | 320 | | | | |
| 99TH.....GM..... | 618 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 221 | 0 | 0 | 167 | 136 |
| STANDARD DEVIATION.....GM..... | 211 | 0 | 0 | 99 | 80 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 221 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 181 | 0 | 0 | 0 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 237 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 173 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 3,213 | 0 | 0 | 232 | 232 |
| MAXIMUM OVER 3 DAYS.....GM..... | 4,634 | 0 | 0 | 232 | 232 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 200 | 0 | 0 | 167 | 136 |
| STANDARD DEVIATION.....GM..... | 163 | 0 | 0 | 99 | 80 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 58 | | | | |
| 25TH.....GM..... | 116 | | | | |
| 50TH.....GM..... | 174 | | | | |
| 75TH.....GM..... | 232 | | | | |
| 90TH.....GM..... | 348 | | | | |
| 95TH.....GM..... | 464 | | | | |
| 99TH.....GM..... | 721 | | | | |
| MAXIMUM.....GM..... | 2,227 | 0 | 0 | 232 | 232 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 5.1 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .2 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| .3 | .4 | 1.0 | 1.4 | 6.7 | 8.2 | 8.3 | 7.1 | 4.7 | 3.7 | 3.4 | 3.6 |
| .3 | .4 | .8 | 1.3 | 5.0 | 5.9 | 4.6 | 4.2 | 2.1 | 2.5 | .8 | 2.2 |
| 0 | 0 | 0 | .1 | 1.1 | 1.6 | 2.3 | 1.5 | 1.3 | .8 | 1.4 | .6 |
| 0 | 0 | .1 | 0 | .6 | .7 | 1.3 | 1.4 | 1.2 | .4 | 1.2 | .7 |
| 1.0 | 1.0 | 1.4 | 1.2 | 1.6 | 1.6 | 1.8 | 1.8 | 2.1 | 1.5 | 2.7 | 1.6 |
| .0 | .0 | 1.0 | .5 | 1.1 | 1.0 | 1.1 | 1.2 | 1.3 | .9 | 1.6 | .8 |
| 1.0 | 1.0 | 4.0 | 3.0 | 8.0 | 9.0 | 7.0 | 8.0 | 5.0 | 5.0 | 6.0 | 3.0 |
| 18 | 32 | 79 | 62 | 145 | 109 | 126 | 97 | 108 | 76 | 249 | 61 |
| 14 | 19 | 94 | 55 | 157 | 115 | 125 | 86 | 94 | 62 | 268 | 34 |
| | | | | 39 | 20 | 20 | 19 | 10 | 20 | | |
| | | | | 58 | 40 | 45 | 39 | 39 | 30 | | |
| | | | | 97 | 77 | 77 | 77 | 69 | 45 | | |
| | | | | 155 | 126 | 155 | 116 | 135 | 116 | | |
| | | | | 309 | 232 | 271 | 222 | 290 | 199 | | |
| | | | | 348 | 290 | 387 | 242 | 329 | 232 | | |
| | | | | 890 | 453 | 541 | 464 | 329 | 232 | | |
| 55 | 96 | 155 | 183 | 330 | 237 | 209 | 173 | 139 | 177 | 268 | 129 |
| 41 | 57 | 206 | 170 | 339 | 196 | 173 | 132 | 79 | 148 | 53 | 48 |
| 0 | 0 | 0 | 116 | 308 | 219 | 260 | 171 | 188 | 124 | 162 | 136 |
| 0 | 0 | 0 | 0 | 292 | 111 | 222 | 110 | 110 | 61 | 66 | 20 |
| 0 | 0 | 249 | 0 | 328 | 280 | 244 | 197 | 194 | 183 | 529 | 89 |
| 0 | 0 | 0 | 0 | 218 | 262 | 137 | 102 | 109 | 68 | 291 | 48 |
| 116 | 203 | 928 | 812 | 2,036 | 3,213 | 1,856 | 1,044 | 522 | 696 | 812 | 232 |
| 116 | 203 | 928 | 812 | 3,518 | 4,634 | 2,784 | 1,856 | 986 | 696 | 2,436 | 522 |
| 55 | 96 | 167 | 156 | 279 | 211 | 209 | 163 | 158 | 150 | 276 | 116 |
| 41 | 57 | 178 | 104 | 247 | 162 | 165 | 103 | 110 | 107 | 158 | 55 |
| | | | | 87 | 60 | 58 | 58 | 15 | 45 | | |
| | | | | 145 | 116 | 116 | 116 | 116 | 87 | | |
| | | | | 232 | 174 | 174 | 120 | 116 | 120 | | |
| | | | | 348 | 232 | 232 | 232 | 200 | 180 | | |
| | | | | 464 | 348 | 360 | 240 | 348 | 232 | | |
| | | | | 696 | 464 | 464 | 360 | 406 | 232 | | |
| | | | | 1,567 | 721 | 870 | 522 | 464 | 696 | | |
| 116 | 203 | 928 | 464 | 1,856 | 2,227 | 1,856 | 928 | 464 | 696 | 480 | 232 |
| 0 | 0 | 0 | 0 | 0 | 5.1 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | .8 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 1.11.--DISTILLED LIQUORS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.4 | 0 | 0 | 0 | 0 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.1 | 0 | 0 | 0 | 0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .7 | 0 | 0 | 0 | 0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .6 | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 2.0 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....DO..... | 1.7 | 0 | 0 | 0 | 0 |
| MAXIMUM.....DO..... | 17.0 | 0 | 0 | 0 | 0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 46 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 49 | 0 | 0 | 0 | 0 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 9 | | | | |
| 25TH.....GM..... | 14 | | | | |
| 50TH.....GM..... | 28 | | | | |
| 75TH.....GM..... | 56 | | | | |
| 90TH.....GM..... | 98 | | | | |
| 95TH.....GM..... | 140 | | | | |
| 99TH.....GM..... | 252 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 90 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 102 | 0 | 0 | 0 | 0 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 92 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 80 | 0 | 0 | 0 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 80 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 60 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,232 | 0 | 0 | 0 | 0 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,316 | 0 | 0 | 0 | 0 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 69 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 68 | 0 | 0 | 0 | 0 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 28 | | | | |
| 25TH.....GM..... | 28 | | | | |
| 50TH.....GM..... | 56 | | | | |
| 75TH.....GM..... | 84 | | | | |
| 90TH.....GM..... | 126 | | | | |
| 95TH.....GM..... | 168 | | | | |
| 99TH.....GM..... | 364 | | | | |
| MAXIMUM.....GM..... | 896 | 0 | 0 | 0 | 0 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 16.6 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| (2) | 0 | .3 | .4 | 5.4 | 3.7 | 7.4 | 5.1 | 5.8 | 3.3 | 3.1 | 1.5 |
| (2) | 0 | .3 | .4 | 3.8 | 2.5 | 4.3 | 3.0 | 1.8 | 2.1 | .3 | .6 |
| 0 | 0 | 0 | .1 | 1.1 | .8 | 1.8 | 1.1 | 1.0 | .6 | .9 | .6 |
| 0 | 0 | 0 | 0 | .5 | .4 | 1.2 | 1.0 | 3.0 | .6 | 1.9 | .3 |
| 1.0 | 0 | 1.0 | 1.1 | 1.9 | 1.8 | 2.0 | 2.0 | 2.8 | 1.8 | 3.2 | 1.8 |
| .0 | 0 | .0 | .3 | 2.2 | 1.4 | 1.8 | 1.5 | 1.6 | 1.1 | 2.2 | .7 |
| 1.0 | 0 | 1.0 | 2.0 | 17.0 | 9.0 | 15.0 | 11.0 | 6.0 | 5.0 | 9.0 | 3.0 |
| 149 | 0 | 9 | 9 | 44 | 40 | 55 | 41 | 60 | 26 | 48 | 32 |
| 0 | 0 | 0 | 4 | 46 | 48 | 56 | 44 | 47 | 15 | 47 | 28 |
| | | | | 9 | 9 | 9 | 9 | 9 | 5 | | |
| | | | | 14 | 14 | 19 | 14 | 25 | 14 | | |
| | | | | 28 | 23 | 37 | 28 | 47 | 25 | | |
| | | | | 56 | 56 | 70 | 47 | 93 | 37 | | |
| | | | | 103 | 75 | 112 | 89 | 149 | 42 | | |
| | | | | 121 | 135 | 168 | 112 | 149 | 42 | | |
| | | | | 224 | 299 | 299 | 280 | 159 | 75 | | |
| 448 | 0 | 28 | 24 | 89 | 81 | 116 | 81 | 55 | 58 | 19 | 32 |
| 0 | 0 | 0 | 6 | 107 | 111 | 121 | 74 | 27 | 34 | 0 | 20 |
| 0 | 0 | 0 | 28 | 127 | 83 | 107 | 76 | 67 | 55 | 41 | 43 |
| 0 | 0 | 0 | 0 | 82 | 60 | 87 | 84 | 36 | 16 | 10 | 18 |
| 0 | 0 | 0 | 0 | 73 | 94 | 92 | 71 | 90 | 40 | 63 | 84 |
| 0 | 0 | 0 | 0 | 34 | 75 | 77 | 52 | 45 | 14 | 54 | 0 |
| 448 | 0 | 28 | 28 | 896 | 896 | 1,232 | 448 | 224 | 126 | 168 | 84 |
| 448 | 0 | 28 | 56 | 896 | 980 | 1,316 | 896 | 476 | 224 | 504 | 252 |
| 448 | 0 | 28 | 25 | 71 | 67 | 83 | 62 | 64 | 44 | 44 | 54 |
| 0 | 0 | 0 | 6 | 68 | 73 | 81 | 59 | 33 | 23 | 22 | 27 |
| | | | | 28 | 28 | 28 | 28 | 28 | 14 | | |
| | | | | 28 | 42 | 42 | 28 | 42 | 28 | | |
| | | | | 56 | 56 | 56 | 42 | 56 | 42 | | |
| | | | | 84 | 56 | 112 | 56 | 84 | 56 | | |
| | | | | 168 | 112 | 140 | 112 | 112 | 56 | | |
| | | | | 168 | 168 | 224 | 168 | 112 | 112 | | |
| | | | | 252 | 448 | 448 | 448 | 168 | 112 | | |
| 448 | 0 | 28 | 28 | 896 | 896 | 896 | 448 | 224 | 112 | 112 | 84 |
| 0 | 0 | 0 | 0 | 2.3 | 3.3 | 10.0 | 0 | 0 | .9 | 0 | 0 |
| 0 | 0 | 0 | 0 | .6 | .9 | 1.4 | 0 | 0 | .9 | 0 | 0 |

TABLE 2.01.--EGGS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 54.3 | 17.7 | 61.3 | 55.2 | 48.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 28.6 | 8.9 | 31.2 | 32.4 | 32.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 15.5 | 5.7 | 19.7 | 15.7 | 11.8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 10.3 | 3.1 | 10.4 | 7.1 | 4.0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.7 | 1.7 | 1.7 | 1.6 | 1.4 |
| STANDARD DEVIATION.....DO..... | .9 | .8 | .8 | .8 | .7 |
| MAXIMUM.....DO..... | 9.0 | 4.0 | 4.0 | 5.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 47 | 28 | 34 | 35 | 34 |
| STANDARD DEVIATION.....GM..... | 33 | 20 | 22 | 24 | 23 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 15 | 5 | 11 | 13 | 15 |
| 25TH.....GM..... | 21 | 11 | 17 | 17 | 17 |
| 50TH.....GM..... | 40 | 23 | 29 | 29 | 29 |
| 75TH.....GM..... | 61 | 43 | 43 | 43 | 43 |
| 90TH.....GM..... | 92 | 50 | 64 | 64 | 64 |
| 95TH.....GM..... | 109 | 73 | 73 | 82 | 79 |
| 99TH.....GM..... | 165 | 81 | 116 | 128 | 110 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 85 | 47 | 62 | 67 | 72 |
| STANDARD DEVIATION.....GM..... | 50 | 35 | 33 | 34 | 42 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 86 | 55 | 59 | 69 | 73 |
| STANDARD DEVIATION.....GM..... | 41 | 26 | 22 | 32 | 28 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 83 | 46 | 61 | 69 | 68 |
| STANDARD DEVIATION.....GM..... | 38 | 19 | 25 | 28 | 32 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 728 | 220 | 248 | 440 | 440 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,144 | 284 | 448 | 526 | 526 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 82 | 49 | 59 | 66 | 70 |
| STANDARD DEVIATION.....GM..... | 44 | 30 | 27 | 34 | 37 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 40 | 14 | 25 | 35 | 40 |
| 25TH.....GM..... | 50 | 32 | 46 | 46 | 46 |
| 50TH.....GM..... | 64 | 50 | 52 | 64 | 64 |
| 75TH.....GM..... | 100 | 64 | 64 | 64 | 88 |
| 90TH.....GM..... | 128 | 64 | 92 | 128 | 128 |
| 95TH.....GM..... | 150 | 99 | 128 | 128 | 128 |
| 99TH.....GM..... | 237 | 220 | 141 | 192 | 220 |
| MAXIMUM.....GM..... | 715 | 220 | 220 | 440 | 440 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 88.1 | .9 | 1.6 | 4.4 | 2.3 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .2 | .6 | .1 | .3 | .2 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 49.1 | 44.3 | 52.3 | 44.4 | 54.8 | 51.1 | 62.0 | 56.7 | 66.7 | 57.4 | 71.7 | 57.4 |
| 31.2 | 28.2 | 27.0 | 27.5 | 27.3 | 29.1 | 27.1 | 29.5 | 25.6 | 28.9 | 21.3 | 29.5 |
| 12.6 | 11.1 | 15.6 | 13.0 | 16.6 | 15.4 | 18.0 | 16.0 | 19.2 | 16.3 | 21.6 | 14.2 |
| 5.4 | 5.0 | 9.6 | 3.9 | 10.9 | 6.6 | 16.9 | 11.3 | 21.9 | 12.2 | 28.8 | 13.7 |
| 1.5 | 1.5 | 1.7 | 1.5 | 1.8 | 1.6 | 1.9 | 1.8 | 2.0 | 1.8 | 2.2 | 1.8 |
| .7 | .8 | .8 | .7 | .9 | .8 | .9 | .9 | .9 | .9 | 1.0 | .9 |
| 5.0 | 4.0 | 6.0 | 5.0 | 5.0 | 6.0 | 8.0 | 6.0 | 5.0 | 6.0 | 9.0 | 5.0 |
| 43 | 38 | 58 | 40 | 62 | 45 | 59 | 43 | 54 | 38 | 54 | 37 |
| 30 | 27 | 39 | 27 | 40 | 31 | 37 | 30 | 34 | 25 | 36 | 22 |
| 15 | 13 | 15 | 13 | 15 | 15 | 15 | 15 | 15 | 15 | 15 | 13 |
| 21 | 17 | 31 | 19 | 31 | 21 | 31 | 21 | 31 | 17 | 31 | 17 |
| 33 | 31 | 46 | 33 | 52 | 38 | 50 | 33 | 46 | 33 | 46 | 33 |
| 56 | 46 | 76 | 49 | 85 | 59 | 83 | 56 | 73 | 49 | 69 | 48 |
| 80 | 73 | 100 | 75 | 115 | 85 | 104 | 85 | 100 | 70 | 97 | 64 |
| 100 | 92 | 128 | 87 | 135 | 100 | 128 | 100 | 123 | 85 | 123 | 85 |
| 160 | 132 | 197 | 128 | 192 | 144 | 180 | 141 | 140 | 128 | 166 | 109 |
| 89 | 77 | 108 | 82 | 112 | 88 | 100 | 77 | 88 | 68 | 69 | 69 |
| 49 | 45 | 60 | 49 | 61 | 51 | 56 | 41 | 41 | 37 | 29 | 31 |
| 86 | 76 | 102 | 81 | 109 | 85 | 100 | 80 | 87 | 65 | 84 | 67 |
| 42 | 36 | 41 | 35 | 47 | 36 | 46 | 40 | 39 | 26 | 44 | 25 |
| 86 | 79 | 104 | 81 | 107 | 85 | 93 | 75 | 80 | 66 | 76 | 60 |
| 38 | 35 | 45 | 35 | 43 | 39 | 37 | 31 | 35 | 26 | 35 | 22 |
| 572 | 406 | 566 | 440 | 534 | 715 | 728 | 660 | 400 | 287 | 457 | 256 |
| 704 | 605 | 768 | 584 | 998 | 826 | 1,144 | 880 | 800 | 534 | 828 | 384 |
| 85 | 75 | 101 | 79 | 105 | 83 | 93 | 74 | 81 | 64 | 73 | 63 |
| 47 | 40 | 49 | 41 | 52 | 45 | 46 | 39 | 39 | 30 | 36 | 28 |
| 40 | 40 | 46 | 40 | 46 | 40 | 44 | 40 | 40 | 40 | 40 | 40 |
| 50 | 46 | 64 | 46 | 80 | 50 | 50 | 46 | 50 | 46 | 46 | 46 |
| 80 | 64 | 92 | 64 | 92 | 64 | 92 | 64 | 80 | 50 | 64 | 50 |
| 100 | 92 | 128 | 100 | 128 | 104 | 128 | 92 | 100 | 64 | 92 | 64 |
| 128 | 128 | 150 | 128 | 165 | 128 | 128 | 128 | 128 | 110 | 128 | 100 |
| 165 | 128 | 192 | 128 | 192 | 138 | 160 | 128 | 136 | 128 | 128 | 128 |
| 240 | 220 | 256 | 220 | 316 | 234 | 256 | 220 | 200 | 134 | 192 | 128 |
| 480 | 360 | 486 | 440 | 512 | 715 | 486 | 480 | 330 | 256 | 365 | 200 |
| 4.9 | 4.7 | 1.5 | 5.0 | 2.0 | 16.5 | 15.3 | 16.6 | 5.1 | 4.5 | 1.5 | 1.3 |
| .3 | .3 | .1 | .5 | .1 | .4 | .3 | .2 | .3 | .3 | .2 | .1 |

TABLE 2.02.--BOILED AND POACHED EGGS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 12.4 | 3.7 | 11.8 | 8.9 | 7.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 8.9 | 2.1 | 8.8 | 7.2 | 6.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.2 | 1.1 | 1.9 | 1.3 | 1.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.3 | .5 | 1.1 | .4 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.4 | 1.6 | 1.4 | 1.3 | 1.2 |
| STANDARD DEVIATION.....DO..... | .7 | .7 | .6 | .5 | .4 |
| MAXIMUM.....OO..... | 7.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 31 | 21 | 22 | 23 | 22 |
| STANDARD DEVIATION.....GM..... | 21 | 11 | 12 | 13 | 11 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 15 | | 8 | 8 | 15 |
| 25TH.....GM..... | 17 | | 15 | 17 | 17 |
| 50TH.....GM..... | 29 | | 17 | 17 | 17 |
| 75TH.....GM..... | 33 | | 29 | 29 | 29 |
| 90TH.....GM..... | 50 | | 39 | 33 | 33 |
| 95TH.....GM..... | 67 | | 50 | 50 | 50 |
| 99TH.....GM..... | 100 | | 56 | 83 | 67 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 67 | 40 | 49 | 58 | 55 |
| STANDARD DEVIATION.....GM..... | 30 | 15 | 19 | 24 | 18 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 67 | 48 | 52 | 58 | 61 |
| STANDARD DEVIATION.....GM..... | 27 | 3 | 12 | 22 | 19 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 65 | 32 | 43 | 46 | 48 |
| STANDARD DEVIATION.....GM..... | 30 | 11 | 11 | 20 | 2 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 400 | 50 | 132 | 150 | 100 |
| MAXIMUM OVER 3 DAYS.....GM..... | 850 | 150 | 167 | 261 | 200 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 65 | 41 | 48 | 55 | 56 |
| STANDARD DEVIATION.....GM..... | 28 | 13 | 16 | 22 | 19 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 44 | | 25 | 25 | 44 |
| 25TH.....GM..... | 50 | | 44 | 44 | 50 |
| 50TH.....GM..... | 50 | | 50 | 50 | 50 |
| 75TH.....GM..... | 88 | | 50 | 50 | 50 |
| 90TH.....GM..... | 100 | | 58 | 100 | 100 |
| 95TH.....GM..... | 100 | | 88 | 100 | 100 |
| 99TH.....GM..... | 150 | | 100 | 116 | 100 |
| MAXIMUM.....GM..... | 250 | 50 | 100 | 150 | 100 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 29.6 | 0 | 0 | 1.0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | 0 | .5 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.9 | 7.5 | 4.0 | 7.4 | 7.8 | 12.5 | 13.1 | 17.7 | 19.3 | 22.6 | 22.2 | 24.2 |
| 5.0 | 6.2 | 3.4 | 5.8 | 6.1 | 9.2 | 9.4 | 12.6 | 10.7 | 15.2 | 12.9 | 13.8 |
| .7 | .8 | .5 | 1.2 | 1.2 | 2.4 | 2.3 | 3.2 | 3.9 | 4.9 | 4.3 | 5.7 |
| .1 | .6 | .1 | .4 | .5 | .9 | 1.5 | 1.9 | 4.7 | 2.5 | 5.0 | 4.7 |
| 1.2 | 1.3 | 1.2 | 1.3 | 1.3 | 1.4 | 1.5 | 1.4 | 1.8 | 1.5 | 1.7 | 1.6 |
| .4 | .6 | .6 | .6 | .6 | .7 | .8 | .7 | 1.0 | .7 | .9 | .9 |
| 3.0 | 3.0 | 4.0 | 3.0 | 4.0 | 6.0 | 7.0 | 6.0 | 5.0 | 4.0 | 4.0 | 5.0 |
| 26 | 26 | 34 | 29 | 37 | 31 | 35 | 29 | 42 | 27 | 36 | 32 |
| 15 | 17 | 24 | 18 | 25 | 20 | 27 | 17 | 34 | 17 | 18 | 19 |
| 11 | 15 | 8 | 13 | 8 | 15 | 15 | 15 | 15 | 13 | 15 | 15 |
| 17 | 17 | 17 | 17 | 17 | 17 | 17 | 17 | 17 | 17 | 17 | 17 |
| 17 | 17 | 33 | 25 | 33 | 29 | 33 | 25 | 33 | 17 | 33 | 29 |
| 33 | 33 | 33 | 33 | 45 | 33 | 39 | 33 | 50 | 33 | 50 | 44 |
| 42 | 50 | 59 | 50 | 67 | 51 | 67 | 50 | 83 | 50 | 59 | 50 |
| 59 | 58 | 67 | 67 | 96 | 67 | 88 | 67 | 100 | 59 | 67 | 72 |
| 67 | 100 | 133 | 100 | 117 | 100 | 117 | 100 | 133 | 92 | 100 | 100 |
| 68 | 63 | 86 | 67 | 82 | 69 | 75 | 64 | 70 | 57 | 73 | 58 |
| 30 | 29 | 54 | 26 | 42 | 31 | 35 | 25 | 30 | 24 | 28 | 23 |
| 67 | 56 | 84 | 79 | 92 | 68 | 76 | 62 | 72 | 61 | 78 | 59 |
| 20 | 15 | 46 | 34 | 37 | 25 | 31 | 23 | 29 | 22 | 22 | 20 |
| 77 | 66 | 85 | 58 | 86 | 71 | 78 | 59 | 80 | 54 | 53 | 58 |
| 25 | 21 | 17 | 17 | 22 | 26 | 45 | 17 | 44 | 14 | 12 | 18 |
| 200 | 174 | 400 | 200 | 232 | 250 | 350 | 232 | 400 | 200 | 150 | 116 |
| 300 | 300 | 400 | 350 | 400 | 500 | 850 | 348 | 800 | 300 | 300 | 300 |
| 68 | 62 | 82 | 67 | 84 | 68 | 73 | 61 | 72 | 56 | 64 | 57 |
| 28 | 27 | 39 | 27 | 39 | 29 | 33 | 23 | 32 | 23 | 23 | 21 |
| 38 | 38 | 33 | 38 | 25 | 44 | 44 | 44 | 44 | 44 | 44 | 44 |
| 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 |
| 50 | 50 | 100 | 50 | 88 | 50 | 50 | 50 | 50 | 50 | 50 | 50 |
| 100 | 88 | 100 | 100 | 100 | 100 | 100 | 76 | 100 | 50 | 100 | 50 |
| 100 | 100 | 150 | 100 | 132 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| 100 | 100 | 150 | 100 | 150 | 100 | 132 | 100 | 116 | 100 | 100 | 100 |
| 150 | 174 | 200 | 150 | 200 | 152 | 174 | 116 | 200 | 116 | 116 | 116 |
| 200 | 174 | 200 | 150 | 232 | 200 | 250 | 184 | 200 | 200 | 116 | 116 |
| .7 | 2.9 | .9 | 1.0 | 1.0 | 6.0 | 5.7 | 7.7 | 0 | 0 | 1.5 | 1.3 |
| .5 | 1.4 | 1.3 | .7 | .3 | .7 | .6 | .4 | 0 | 0 | .7 | .3 |

TABLE 2.03.--FRIED EGGS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 24.3 | 2.1 | 20.7 | 21.7 | 20.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 15.9 | 1.8 | 13.5 | 15.3 | 16.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 5.5 | 0 | 5.9 | 5.2 | 3.5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 3.0 | .3 | 1.3 | 1.2 | 1.1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.5 | 1.3 | 1.4 | 1.4 | 1.3 |
| STANDARD DEVIATION.....DO..... | .7 | .7 | .6 | .6 | .6 |
| MAXIMUM.....DO..... | 9.0 | 3.0 | 4.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 34 | 22 | 22 | 23 | 24 |
| STANDARD DEVIATION.....GM..... | 23 | 12 | 12 | 14 | 14 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 13 | | 12 | 13 | 13 |
| 25TH.....GM..... | 15 | | 15 | 15 | 15 |
| 50TH.....GM..... | 31 | | 15 | 15 | 15 |
| 75TH.....GM..... | 44 | | 31 | 31 | 31 |
| 90TH.....GM..... | 61 | | 33 | 40 | 43 |
| 95TH.....GM..... | 92 | | 46 | 46 | 61 |
| 99TH.....GM..... | 107 | | 61 | 77 | 80 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 69 | 54 | 47 | 52 | 55 |
| STANDARD DEVIATION.....GM..... | 29 | 25 | 15 | 19 | 20 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 71 | 0 | 45 | 51 | 59 |
| STANDARD DEVIATION.....GM..... | 27 | 0 | 11 | 16 | 20 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 71 | 46 | 50 | 53 | 57 |
| STANDARD DEVIATION.....GM..... | 28 | 0 | 16 | 23 | 18 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 368 | 92 | 104 | 184 | 138 |
| MAXIMUM OVER 3 DAYS.....GM..... | 828 | 138 | 312 | 368 | 276 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 69 | 51 | 46 | 51 | 56 |
| STANDARD DEVIATION.....GM..... | 28 | 21 | 15 | 18 | 21 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 40 | | 23 | 40 | 40 |
| 25TH.....GM..... | 46 | | 40 | 40 | 46 |
| 50TH.....GM..... | 52 | | 46 | 46 | 46 |
| 75TH.....GM..... | 92 | | 46 | 46 | 80 |
| 90TH.....GM..... | 92 | | 52 | 92 | 92 |
| 95TH.....GM..... | 104 | | 92 | 92 | 92 |
| 99TH.....GM..... | 138 | | 104 | 104 | 104 |
| MAXIMUM.....GM..... | 368 | 92 | 104 | 138 | 138 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 14.8 | 0 | 1.6 | 0 | 1.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .1 | 0 | .5 | 0 | .3 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 24.3 | 19.9 | 27.7 | 19.0 | 28.9 | 19.8 | 33.7 | 24.1 | 31.2 | 19.9 | 36.4 | 17.8 |
| 17.9 | 15.2 | 17.1 | 14.3 | 17.3 | 14.7 | 19.3 | 16.2 | 15.3 | 13.0 | 14.7 | 10.5 |
| 4.5 | 3.5 | 7.2 | 3.8 | 7.8 | 3.5 | 8.7 | 5.2 | 8.5 | 3.6 | 9.2 | 3.7 |
| 1.9 | 1.2 | 3.4 | .9 | 3.8 | 1.5 | 5.7 | 2.7 | 7.4 | 3.4 | 12.5 | 3.6 |
| 1.4 | 1.3 | 1.5 | 1.3 | 1.5 | 1.3 | 1.6 | 1.5 | 1.8 | 1.5 | 2.0 | 1.6 |
| .7 | .6 | .7 | .6 | .7 | .6 | .8 | .7 | .9 | .8 | 1.1 | .8 |
| 5.0 | 4.0 | 4.0 | 4.0 | 4.0 | 4.0 | 4.0 | 4.0 | 4.0 | 4.0 | 9.0 | 3.0 |
| 32 | 26 | 43 | 27 | 45 | 29 | 43 | 30 | 40 | 28 | 44 | 30 |
| 20 | 17 | 26 | 15 | 27 | 17 | 25 | 18 | 25 | 16 | 32 | 17 |
| 13 | 13 | 15 | 13 | 15 | 13 | 15 | 13 | 15 | 13 | 13 | 13 |
| 15 | 15 | 31 | 15 | 31 | 15 | 27 | 15 | 15 | 15 | 27 | 15 |
| 31 | 15 | 31 | 27 | 31 | 31 | 31 | 29 | 31 | 23 | 31 | 31 |
| 31 | 31 | 53 | 31 | 61 | 31 | 61 | 31 | 52 | 32 | 61 | 40 |
| 61 | 46 | 88 | 46 | 88 | 53 | 92 | 57 | 77 | 46 | 92 | 52 |
| 77 | 61 | 92 | 61 | 92 | 61 | 92 | 69 | 92 | 61 | 92 | 61 |
| 100 | 92 | 123 | 88 | 128 | 92 | 132 | 92 | 123 | 82 | 123 | 92 |
| 73 | 58 | 84 | 62 | 89 | 67 | 81 | 62 | 68 | 56 | 66 | 58 |
| 35 | 21 | 27 | 23 | 32 | 26 | 29 | 24 | 27 | 20 | 27 | 25 |
| 70 | 63 | 84 | 65 | 88 | 66 | 81 | 63 | 73 | 55 | 62 | 55 |
| 27 | 27 | 27 | 25 | 27 | 23 | 27 | 23 | 26 | 15 | 24 | 18 |
| 73 | 59 | 88 | 56 | 89 | 64 | 78 | 61 | 66 | 53 | 73 | 52 |
| 24 | 23 | 31 | 18 | 30 | 23 | 26 | 21 | 26 | 17 | 36 | 16 |
| 368 | 323 | 230 | 172 | 276 | 230 | 276 | 184 | 200 | 184 | 276 | 184 |
| 412 | 415 | 690 | 276 | 644 | 322 | 450 | 385 | 400 | 368 | 828 | 276 |
| 71 | 59 | 85 | 62 | 88 | 66 | 79 | 61 | 68 | 54 | 65 | 55 |
| 32 | 24 | 29 | 23 | 31 | 25 | 27 | 23 | 26 | 18 | 25 | 21 |
| 40 | 40 | 46 | 40 | 46 | 40 | 40 | 40 | 40 | 40 | 40 | 40 |
| 46 | 46 | 80 | 46 | 80 | 46 | 46 | 46 | 46 | 46 | 46 | 40 |
| 80 | 46 | 92 | 46 | 92 | 46 | 92 | 46 | 52 | 46 | 46 | 46 |
| 92 | 80 | 92 | 92 | 92 | 92 | 92 | 92 | 92 | 52 | 92 | 52 |
| 92 | 92 | 104 | 92 | 138 | 92 | 104 | 92 | 92 | 92 | 92 | 92 |
| 104 | 92 | 138 | 92 | 138 | 92 | 138 | 92 | 92 | 92 | 92 | 92 |
| 160 | 104 | 184 | 120 | 184 | 138 | 156 | 120 | 138 | 96 | 138 | 92 |
| 368 | 323 | 230 | 138 | 276 | 184 | 184 | 140 | 146 | 104 | 138 | 184 |
| 1.8 | 0 | 0 | 0 | 0 | 3.2 | 3.5 | 1.0 | 2.1 | 0 | 0 | 0 |
| .3 | 0 | 0 | 0 | 0 | .2 | .1 | (2) | .3 | 0 | 0 | 0 |

TABLE 2.04.--SCRAMBLED EGGS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 20.1 | 8.6 | 32.6 | 26.8 | 20.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 14.2 | 5.1 | 20.6 | 19.0 | 15.9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 4.1 | 3.3 | 8.7 | 5.1 | 3.6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.8 | .3 | 3.3 | 2.7 | 1.0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.4 | 1.5 | 1.5 | 1.4 | 1.3 |
| STANDARD DEVIATION.....DO..... | .7 | .6 | .7 | .7 | .6 |
| MAXIMUM.....DO..... | 4.0 | 3.0 | 4.0 | 4.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 49 | 31 | 36 | 37 | 38 |
| STANDARD DEVIATION.....GM..... | 32 | 19 | 22 | 24 | 24 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 21 | | 11 | 19 | 21 |
| 25TH.....GM..... | 21 | | 21 | 21 | 21 |
| 50TH.....GM..... | 43 | | 21 | 21 | 32 |
| 75TH.....GM..... | 64 | | 43 | 43 | 43 |
| 90TH.....GM..... | 85 | | 64 | 64 | 64 |
| 95TH.....GM..... | 128 | | 73 | 85 | 85 |
| 99TH.....GM..... | 157 | | 128 | 128 | 147 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 109 | 59 | 75 | 81 | 92 |
| STANDARD DEVIATION.....GM..... | 52 | 41 | 34 | 37 | 49 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 104 | 70 | 71 | 79 | 79 |
| STANDARD DEVIATION.....GM..... | 46 | 20 | 28 | 28 | 23 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 102 | 53 | 73 | 81 | 92 |
| STANDARD DEVIATION.....GM..... | 40 | 0 | 24 | 29 | 42 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 728 | 220 | 220 | 256 | 440 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,144 | 284 | 448 | 512 | 512 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 105 | 63 | 72 | 80 | 88 |
| STANDARD DEVIATION.....GM..... | 48 | 33 | 29 | 34 | 43 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 64 | | 32 | 55 | 64 |
| 25TH.....GM..... | 64 | | 64 | 64 | 64 |
| 50TH.....GM..... | 128 | | 64 | 64 | 64 |
| 75TH.....GM..... | 128 | | 64 | 96 | 128 |
| 90TH.....GM..... | 128 | | 128 | 128 | 128 |
| 95TH.....GM..... | 192 | | 128 | 128 | 165 |
| 99TH.....GM..... | 256 | | 165 | 220 | 220 |
| MAXIMUM.....GM..... | 715 | 220 | 220 | 220 | 440 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 7.2 | 0 | 0 | 1.9 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .1 | 0 | 0 | .3 | 0 |

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 19.1 | 16.7 | 21.5 | 17.6 | 20.7 | 20.6 | 20.5 | 18.8 | 19.9 | 18.5 | 18.3 | 17.7 |
| 14.3 | 12.4 | 15.3 | 13.7 | 14.5 | 15.0 | 13.5 | 13.5 | 12.2 | 12.8 | 9.3 | 13.5 |
| 3.5 | 3.3 | 4.0 | 3.1 | 4.3 | 4.3 | 4.6 | 3.7 | 4.1 | 3.4 | 4.8 | 2.4 |
| 1.3 | 1.0 | 2.2 | 1.0 | 1.9 | 1.3 | 2.3 | 1.6 | 3.6 | 2.2 | 4.3 | 1.8 |
| 1.3 | 1.3 | 1.4 | 1.3 | 1.4 | 1.3 | 1.5 | 1.4 | 1.6 | 1.4 | 1.7 | 1.3 |
| .6 | .6 | .7 | .6 | .7 | .6 | .7 | .7 | .8 | .7 | .8 | .7 |
| 4.0 | 3.0 | 3.0 | 3.0 | 3.0 | 4.0 | 4.0 | 4.0 | 3.0 | 4.0 | 3.0 | 3.0 |
| 48 | 42 | 60 | 45 | 63 | 48 | 60 | 45 | 58 | 42 | 59 | 39 |
| 29 | 26 | 37 | 28 | 37 | 30 | 35 | 28 | 33 | 25 | 41 | 22 |
| 21 | 21 | 21 | 21 | 21 | 21 | 21 | 21 | 21 | 21 | 21 | 21 |
| 21 | 21 | 43 | 21 | 43 | 21 | 43 | 21 | 43 | 21 | 21 | 21 |
| 43 | 43 | 43 | 43 | 43 | 43 | 43 | 43 | 43 | 43 | 43 | 37 |
| 64 | 43 | 73 | 43 | 85 | 61 | 85 | 43 | 79 | 43 | 85 | 43 |
| 85 | 73 | 107 | 85 | 128 | 85 | 107 | 85 | 128 | 85 | 128 | 64 |
| 107 | 85 | 147 | 110 | 147 | 90 | 128 | 97 | 128 | 85 | 128 | 85 |
| 165 | 128 | 192 | 147 | 192 | 171 | 192 | 128 | 149 | 128 | 192 | 128 |
| 115 | 96 | 133 | 105 | 141 | 110 | 132 | 103 | 116 | 90 | 91 | 89 |
| 56 | 45 | 58 | 52 | 58 | 50 | 53 | 42 | 39 | 38 | 35 | 34 |
| 101 | 94 | 135 | 103 | 136 | 103 | 123 | 98 | 118 | 86 | 96 | 87 |
| 41 | 34 | 51 | 45 | 54 | 36 | 52 | 43 | 36 | 27 | 37 | 26 |
| 101 | 95 | 111 | 103 | 125 | 112 | 114 | 93 | 101 | 85 | 113 | 81 |
| 44 | 37 | 49 | 28 | 40 | 46 | 36 | 36 | 37 | 27 | 38 | 24 |
| 568 | 348 | 440 | 440 | 512 | 715 | 728 | 660 | 256 | 256 | 198 | 256 |
| 704 | 605 | 768 | 512 | 714 | 715 | 1,144 | 880 | 576 | 384 | 576 | 384 |
| 108 | 95 | 129 | 104 | 135 | 107 | 124 | 99 | 111 | 87 | 102 | 86 |
| 51 | 39 | 55 | 47 | 54 | 48 | 47 | 41 | 38 | 34 | 39 | 31 |
| 64 | 64 | 64 | 64 | 64 | 64 | 64 | 64 | 64 | 64 | 64 | 64 |
| 64 | 64 | 64 | 64 | 128 | 64 | 128 | 64 | 64 | 64 | 64 | 64 |
| 128 | 64 | 128 | 128 | 128 | 128 | 128 | 85 | 128 | 64 | 128 | 64 |
| 128 | 128 | 128 | 128 | 128 | 128 | 128 | 128 | 128 | 128 | 128 | 128 |
| 165 | 128 | 192 | 128 | 192 | 128 | 192 | 128 | 136 | 128 | 128 | 128 |
| 220 | 165 | 220 | 192 | 256 | 192 | 192 | 158 | 192 | 128 | 192 | 128 |
| 256 | 220 | 320 | 256 | 320 | 256 | 256 | 220 | 192 | 192 | 198 | 158 |
| 440 | 256 | 440 | 440 | 512 | 715 | 468 | 440 | 220 | 256 | 198 | 158 |
| 1.1 | 0 | 0 | 1.1 | 0 | 2.0 | 1.0 | 0 | 0 | 0 | 0 | 0 |
| .2 | 0 | 0 | .3 | 0 | .1 | .1 | 0 | 0 | 0 | 0 | 0 |

TABLE 3.01.--BUTTER
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 31.4 | 5.2 | 29.2 | 28.7 | 31.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 16.7 | 3.8 | 17.4 | 15.9 | 17.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 8.4 | 1.0 | 7.2 | 7.7 | 8.6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 6.3 | .3 | 4.6 | 5.1 | 5.8 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 2.1 | 1.5 | 1.9 | 2.0 | 2.0 |
| STANDARD DEVIATION.....DO..... | 1.4 | .9 | 1.2 | 1.3 | 1.3 |
| MAXIMUM.....DO..... | 12.0 | 4.0 | 7.0 | 9.0 | 8.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 8 | 3 | 4 | 6 | 6 |
| STANDARD DEVIATION.....GM..... | 10 | 3 | 4 | 7 | 7 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 2 | | 1 | 2 | 2 |
| 25TH.....GM..... | 2 | | 2 | 2 | 2 |
| 50TH.....GM..... | 5 | | 3 | 4 | 5 |
| 75TH.....GM..... | 9 | | 5 | 7 | 7 |
| 90TH.....GM..... | 19 | | 9 | 13 | 13 |
| 95TH.....GM..... | 26 | | 14 | 19 | 19 |
| 99TH.....GM..... | 47 | | 19 | 29 | 38 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 13 | 5 | 8 | 10 | 11 |
| STANDARD DEVIATION.....GM..... | 13 | 4 | 8 | 8 | 12 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 14 | 6 | 9 | 11 | 11 |
| STANDARD DEVIATION.....GM..... | 14 | 1 | 7 | 9 | 12 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 17 | 13 | 8 | 13 | 12 |
| STANDARD DEVIATION.....GM..... | 15 | 4 | 4 | 12 | 10 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 341 | 28 | 57 | 128 | 227 |
| MAXIMUM OVER 3 DAYS.....GM..... | 454 | 50 | 57 | 298 | 236 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 12 | 6 | 7 | 9 | 10 |
| STANDARD DEVIATION.....GM..... | 13 | 4 | 6 | 10 | 11 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 2 | | 1 | 2 | 2 |
| 25TH.....GM..... | 5 | | 5 | 5 | 5 |
| 50TH.....GM..... | 7 | | 5 | 7 | 5 |
| 75TH.....GM..... | 14 | | 7 | 10 | 10 |
| 90TH.....GM..... | 28 | | 14 | 14 | 14 |
| 95TH.....GM..... | 28 | | 14 | 28 | 28 |
| 99TH.....GM..... | 57 | | 28 | 43 | 57 |
| MAXIMUM.....GM..... | 341 | 14 | 57 | 128 | 227 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 2890.4 | 4.2 | 89.4 | 157.6 | 186.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 11.6 | 10.7 | 15.2 | 16.0 | 16.1 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,472 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 32.4 | 30.9 | 32.4 | 32.0 | 34.1 | 30.8 | 35.6 | 31.4 | 30.0 | 27.2 | 29.9 | 29.4 |
| 17.2 | 18.3 | 18.3 | 20.2 | 18.1 | 16.7 | 18.1 | 16.8 | 13.5 | 12.3 | 9.1 | 12.4 |
| 9.3 | 8.3 | 8.1 | 8.4 | 9.9 | 9.1 | 8.8 | 8.3 | 7.9 | 7.4 | 7.0 | 6.5 |
| 5.9 | 4.2 | 6.0 | 3.5 | 6.2 | 5.1 | 8.8 | 6.3 | 8.6 | 7.5 | 13.9 | 10.5 |
| 2.0 | 1.8 | 1.9 | 1.7 | 2.0 | 2.0 | 2.2 | 2.1 | 2.5 | 2.3 | 3.2 | 2.8 |
| 1.3 | 1.2 | 1.3 | 1.1 | 1.4 | 1.2 | 1.6 | 1.4 | 1.7 | 1.5 | 2.2 | 2.0 |
| 9.0 | 8.0 | 7.0 | 8.0 | 9.0 | 8.0 | 9.0 | 9.0 | 9.0 | 8.0 | 12.0 | 9.0 |
| 8 | 6 | 9 | 7 | 10 | 8 | 10 | 7 | 9 | 7 | 13 | 10 |
| 10 | 6 | 11 | 11 | 11 | 10 | 13 | 8 | 11 | 8 | 14 | 11 |
| 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 2 | 2 | 3 | 2 | 3 | 2 | 3 | 2 | 3 | 3 | 3 | 2 |
| 5 | 4 | 5 | 5 | 6 | 5 | 6 | 5 | 6 | 5 | 8 | 6 |
| 9 | 8 | 10 | 9 | 12 | 9 | 13 | 9 | 11 | 9 | 17 | 13 |
| 19 | 13 | 19 | 14 | 20 | 19 | 24 | 16 | 20 | 14 | 28 | 21 |
| 24 | 19 | 26 | 19 | 29 | 27 | 32 | 24 | 30 | 20 | 40 | 35 |
| 52 | 28 | 52 | 80 | 44 | 47 | 59 | 38 | 52 | 36 | 69 | 66 |
| 13 | 11 | 14 | 12 | 15 | 13 | 15 | 12 | 13 | 13 | 11 | 12 |
| 14 | 10 | 15 | 11 | 14 | 15 | 15 | 13 | 12 | 12 | 14 | 15 |
| 15 | 11 | 16 | 15 | 18 | 15 | 17 | 13 | 15 | 11 | 15 | 14 |
| 17 | 7 | 13 | 15 | 16 | 14 | 16 | 11 | 12 | 7 | 14 | 9 |
| 16 | 13 | 19 | 19 | 19 | 18 | 21 | 15 | 17 | 12 | 20 | 17 |
| 15 | 9 | 16 | 23 | 14 | 16 | 19 | 11 | 16 | 11 | 15 | 14 |
| 298 | 123 | 200 | 170 | 227 | 341 | 227 | 184 | 142 | 199 | 85 | 113 |
| 313 | 170 | 370 | 288 | 397 | 364 | 454 | 213 | 312 | 220 | 208 | 203 |
| 12 | 10 | 14 | 13 | 14 | 13 | 14 | 11 | 11 | 9 | 12 | 10 |
| 15 | 9 | 12 | 14 | 13 | 15 | 14 | 10 | 11 | 11 | 9 | 10 |
| 4 | 2 | 5 | 5 | 5 | 2 | 5 | 2 | 2 | 2 | 2 | 2 |
| 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| 9 | 7 | 9 | 9 | 10 | 7 | 9 | 7 | 9 | 5 | 9 | 7 |
| 14 | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 10 | 14 | 14 |
| 28 | 15 | 28 | 28 | 28 | 28 | 28 | 27 | 28 | 14 | 28 | 15 |
| 28 | 28 | 42 | 43 | 43 | 28 | 43 | 28 | 28 | 28 | 28 | 28 |
| 71 | 43 | 57 | 57 | 57 | 57 | 71 | 57 | 57 | 43 | 43 | 57 |
| 298 | 114 | 114 | 114 | 170 | 341 | 199 | 160 | 114 | 199 | 43 | 113 |
| 147.5 | 171.5 | 127.9 | 127.9 | 297.1 | 379.3 | 473.9 | 466.8 | 65.1 | 105.9 | 28.7 | 61.5 |
| 10.8 | 14.0 | 14.5 | 15.9 | 11.0 | 11.7 | 12.0 | 10.0 | 7.8 | 9.6 | 5.6 | 7.6 |

TABLE 3.02.--MARGARINE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 43.1 | 8.5 | 43.8 | 46.1 | 42.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 18.7 | 4.9 | 21.0 | 21.9 | 18.9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 13.2 | 2.4 | 14.0 | 14.2 | 15.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 11.2 | 1.2 | 8.7 | 9.9 | 8.9 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 2.3 | 1.8 | 2.1 | 2.2 | 2.2 |
| STANDARD DEVIATION.....DO..... | 1.5 | 1.1 | 1.3 | 1.3 | 1.3 |
| MAXIMUM.....DO..... | 12.0 | 5.0 | 7.0 | 8.0 | 7.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 9 | 3 | 5 | 6 | 7 |
| STANDARD DEVIATION.....GM..... | 10 | 2 | 6 | 6 | 7 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 2 | | 1 | 1 | 2 |
| 25TH.....GM..... | 3 | | 2 | 2 | 2 |
| 50TH.....GM..... | 5 | | 3 | 5 | 5 |
| 75TH.....GM..... | 10 | | 6 | 7 | 9 |
| 90TH.....GM..... | 19 | | 9 | 12 | 15 |
| 95TH.....GM..... | 28 | | 13 | 16 | 20 |
| 99TH.....GM..... | 47 | | 19 | 31 | 30 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 13 | 6 | 7 | 9 | 11 |
| STANDARD DEVIATION.....GM..... | 14 | 6 | 6 | 8 | 13 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 13 | 5 | 8 | 10 | 11 |
| STANDARD DEVIATION.....GM..... | 11 | 3 | 7 | 8 | 8 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 16 | 5 | 9 | 10 | 12 |
| STANDARD DEVIATION.....GM..... | 14 | 2 | 10 | 7 | 9 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 227 | 27 | 113 | 114 | 113 |
| MAXIMUM OVER 3 DAYS.....GM..... | 383 | 28 | 269 | 119 | 184 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 11 | 5 | 6 | 8 | 9 |
| STANDARD DEVIATION.....GM..... | 11 | 4 | 6 | 8 | 8 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 2 | | 1 | 2 | 2 |
| 25TH.....GM..... | 5 | | 3 | 5 | 5 |
| 50TH.....GM..... | 7 | | 5 | 5 | 5 |
| 75TH.....GM..... | 14 | | 7 | 9 | 14 |
| 90TH.....GM..... | 28 | | 14 | 14 | 14 |
| 95TH.....GM..... | 28 | | 14 | 21 | 28 |
| 99TH.....GM..... | 57 | | 28 | 28 | 43 |
| MAXIMUM.....GM..... | 227 | 27 | 71 | 114 | 113 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 1598.5 | 5.5 | 50.7 | 73.3 | 90.2 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 4.2 | 7.3 | 5.2 | 4.3 | 5.2 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 44.8 | 40.7 | 41.4 | 38.6 | 39.8 | 39.8 | 44.0 | 44.1 | 54.0 | 53.2 | 48.1 | 51.7 |
| 20.8 | 19.6 | 19.8 | 18.3 | 20.2 | 19.2 | 17.3 | 18.0 | 16.9 | 19.1 | 14.6 | 16.2 |
| 13.9 | 13.5 | 12.8 | 12.8 | 11.7 | 12.5 | 11.9 | 13.9 | 12.8 | 17.0 | 12.9 | 15.2 |
| 10.0 | 7.6 | 8.8 | 7.5 | 7.9 | 8.1 | 14.8 | 12.2 | 24.3 | 17.1 | 20.6 | 20.3 |
| 2.2 | 2.1 | 2.2 | 2.1 | 2.1 | 2.1 | 2.5 | 2.4 | 3.0 | 2.6 | 2.9 | 2.9 |
| 1.4 | 1.3 | 1.5 | 1.3 | 1.4 | 1.3 | 1.7 | 1.5 | 1.7 | 1.6 | 1.8 | 1.8 |
| 8.0 | 7.0 | 9.0 | 7.0 | 10.0 | 8.0 | 11.0 | 12.0 | 9.0 | 9.0 | 8.0 | 9.0 |
| 9 | 8 | 12 | 7 | 10 | 8 | 11 | 8 | 11 | 8 | 11 | 9 |
| 10 | 8 | 14 | 8 | 12 | 9 | 12 | 8 | 10 | 8 | 13 | 9 |
| 2 | 1 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 3 | 3 | 3 | 3 | 3 | 2 | 3 | 2 | 5 | 3 | 5 | 3 |
| 5 | 5 | 8 | 5 | 6 | 5 | 7 | 5 | 8 | 5 | 7 | 5 |
| 11 | 9 | 14 | 9 | 14 | 9 | 14 | 9 | 16 | 9 | 13 | 11 |
| 21 | 17 | 27 | 15 | 22 | 19 | 27 | 16 | 24 | 16 | 20 | 22 |
| 31 | 25 | 38 | 20 | 28 | 24 | 36 | 22 | 28 | 24 | 38 | 28 |
| 52 | 43 | 74 | 47 | 61 | 47 | 58 | 42 | 52 | 42 | 71 | 40 |
| 13 | 13 | 17 | 12 | 16 | 13 | 15 | 11 | 14 | 10 | 17 | 10 |
| 13 | 13 | 16 | 11 | 14 | 16 | 18 | 10 | 12 | 8 | 19 | 10 |
| 15 | 14 | 19 | 12 | 17 | 13 | 15 | 11 | 14 | 12 | 12 | 12 |
| 14 | 13 | 17 | 10 | 14 | 11 | 11 | 9 | 12 | 11 | 9 | 10 |
| 18 | 14 | 25 | 15 | 21 | 16 | 19 | 13 | 16 | 12 | 16 | 14 |
| 13 | 10 | 20 | 12 | 18 | 14 | 15 | 12 | 11 | 10 | 16 | 11 |
| 142 | 170 | 142 | 184 | 173 | 170 | 227 | 175 | 170 | 120 | 142 | 227 |
| 255 | 175 | 341 | 255 | 383 | 227 | 284 | 350 | 213 | 227 | 270 | 236 |
| 12 | 11 | 16 | 11 | 14 | 11 | 13 | 10 | 11 | 9 | 11 | 9 |
| 12 | 12 | 14 | 9 | 13 | 11 | 13 | 9 | 10 | 9 | 12 | 11 |
| 2 | 2 | 5 | 2 | 5 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| 9 | 7 | 10 | 7 | 9 | 9 | 9 | 5 | 9 | 5 | 7 | 5 |
| 14 | 14 | 20 | 14 | 14 | 14 | 14 | 14 | 14 | 9 | 14 | 10 |
| 28 | 28 | 28 | 28 | 28 | 28 | 28 | 19 | 24 | 14 | 28 | 14 |
| 28 | 28 | 43 | 28 | 43 | 28 | 28 | 28 | 28 | 28 | 28 | 28 |
| 57 | 57 | 71 | 43 | 57 | 57 | 57 | 43 | 56 | 57 | 57 | 43 |
| 142 | 170 | 128 | 85 | 114 | 170 | 227 | 113 | 170 | 85 | 99 | 227 |
| 77.6 | 76.0 | 63.3 | 53.8 | 144.9 | 189.5 | 246.6 | 284.1 | 53.9 | 96.7 | 32.2 | 60.2 |
| 3.8 | 4.1 | 4.9 | 4.4 | 4.3 | 4.3 | 4.5 | 3.8 | 3.0 | 4.0 | 4.3 | 4.1 |

TABLE 3.03.--WHIPPED BUTTER AND MARGARINE
 CCNSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FCCD: | | | | | |
| AT LEAST CNCE IN 3 DAYS.....PCT..... | 1.9 | .5 | 1.8 | 2.0 | 1.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.1 | .2 | 1.0 | 1.0 | .9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .5 | 0 | .6 | .6 | .6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | .3 | .2 | .4 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$ /NUMBER..... | 1.9 | 2.7 | 2.2 | 2.3 | 2.0 |
| STANDARD DEVIATION.....CO..... | 1.3 | 1.5 | 1.2 | 1.4 | 1.2 |
| MAXIMUM.....CO..... | 10.0 | 4.0 | 5.0 | 5.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$ /GM..... | 5 | 3 | 4 | 5 | 5 |
| STANDARD DEVIATION.....GM..... | 6 | 2 | 3 | 4 | 5 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 1 | | | | |
| 25TH.....GM..... | 2 | | | | |
| 50TH.....GM..... | 3 | | | | |
| 75TH.....GM..... | 6 | | | | |
| 90TH.....GM..... | 13 | | | | |
| 95TH.....GM..... | 17 | | | | |
| 99TH.....GM..... | 31 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$ /GM..... | 11 | 2 | 6 | 7 | 10 |
| STANDARD DEVIATION.....GM..... | 12 | 0 | 4 | 5 | 9 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$ /GM..... | 10 | 0 | 8 | 8 | 7 |
| STANDARD DEVIATION.....GM..... | 9 | 0 | 6 | 8 | 4 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$ /GM..... | 10 | 4 | 5 | 9 | 14 |
| STANDARD DEVIATION.....GM..... | 10 | 0 | 2 | 4 | 7 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 104 | 5 | 22 | 44 | 38 |
| MAXIMUM OVER 3 DAYS.....GM..... | 179 | 13 | 40 | 63 | 76 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$ /GM..... | 8 | 3 | 5 | 6 | 8 |
| STANDARD DEVIATION.....GM..... | 9 | 1 | 4 | 5 | 5 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 2 | | | | |
| 25TH.....GM..... | 3 | | | | |
| 50TH.....GM..... | 6 | | | | |
| 75TH.....GM..... | 9 | | | | |
| 90TH.....GM..... | 19 | | | | |
| 95TH.....GM..... | 28 | | | | |
| 99TH.....GM..... | 38 | | | | |
| MAXIMUM.....GM..... | 76 | 5 | 22 | 28 | 19 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 18.2 | 0 | 1.7 | 1.2 | 2.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.3 | 0 | 4.4 | 1.6 | 4.1 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.3 | 1.4 | 1.9 | 2.3 | 1.9 | 2.1 | 2.3 | 1.7 | 2.7 | 2.3 | 2.3 | 1.5 |
| .6 | 1.1 | 1.2 | 1.5 | 1.2 | 1.4 | 1.2 | .9 | 1.4 | 1.6 | .6 | .7 |
| .3 | .3 | .4 | .6 | .4 | .6 | .8 | .4 | .7 | .5 | 1.3 | .5 |
| .4 | .1 | .3 | .2 | .3 | .1 | .3 | .3 | .6 | .2 | .4 | .3 |
| 2.3 | 1.4 | 1.8 | 1.5 | 1.9 | 1.7 | 2.1 | 1.9 | 2.3 | 1.6 | 2.6 | 2.2 |
| 1.4 | .7 | 1.1 | .7 | 1.3 | 1.1 | 1.3 | 1.4 | 1.4 | 1.0 | 1.0 | 1.3 |
| 5.0 | 4.0 | 6.0 | 3.0 | 6.0 | 6.0 | 6.0 | 10.0 | 5.0 | 4.0 | 4.0 | 4.0 |
| 7 | 4 | 5 | 5 | 6 | 5 | 7 | 4 | 8 | 6 | 9 | 5 |
| 10 | 4 | 5 | 5 | 5 | 5 | 6 | 4 | 13 | 9 | 11 | 4 |
| | | | | 1 | 1 | 1 | 1 | | | | |
| | | | | 2 | 2 | 3 | 1 | | | | |
| | | | | 4 | 3 | 5 | 3 | | | | |
| | | | | 6 | 6 | 9 | 5 | | | | |
| | | | | 13 | 9 | 13 | 9 | | | | |
| | | | | 19 | 14 | 19 | 13 | | | | |
| | | | | 20 | 31 | 31 | 23 | | | | |
| 9 | 9 | 7 | 13 | 14 | 11 | 14 | 10 | 11 | 11 | 8 | 5 |
| 7 | 6 | 5 | 15 | 13 | 12 | 16 | 9 | 11 | 16 | 4 | 4 |
| 8 | 13 | 11 | 10 | 9 | 10 | 14 | 6 | 11 | 23 | 7 | 13 |
| 3 | 10 | 7 | 8 | 7 | 10 | 9 | 5 | 7 | 17 | 6 | 4 |
| 15 | 5 | 12 | 6 | 9 | 8 | 10 | 7 | 18 | 4 | 31 | 6 |
| 14 | 2 | 3 | 2 | 6 | 3 | 7 | 6 | 23 | 1 | 0 | 1 |
| 57 | 38 | 28 | 57 | 57 | 76 | 85 | 38 | 104 | 57 | 38 | 19 |
| 142 | 66 | 47 | 57 | 60 | 94 | 94 | 69 | 179 | 113 | 94 | 38 |
| 9 | 9 | 8 | 10 | 9 | 8 | 10 | 7 | 10 | 12 | 10 | 7 |
| 10 | 7 | 5 | 12 | 9 | 9 | 8 | 7 | 12 | 14 | 10 | 5 |
| | | | | 3 | 2 | 2 | 2 | | | | |
| | | | | 3 | 3 | 3 | 3 | | | | |
| | | | | 6 | 6 | 8 | 3 | | | | |
| | | | | 9 | 9 | 16 | 9 | | | | |
| | | | | 19 | 19 | 19 | 19 | | | | |
| | | | | 28 | 19 | 19 | 28 | | | | |
| | | | | 57 | 38 | 47 | 28 | | | | |
| 57 | 28 | 19 | 57 | 57 | 76 | 47 | 38 | 38 | 57 | 38 | 19 |
| 0 | 0 | 0 | 0 | 0 | 4.2 | 2.0 | 4.2 | 0 | 0 | 2.3 | 0 |
| 0 | 0 | 0 | 0 | 0 | 2.2 | .8 | 1.9 | 0 | 0 | 7.3 | 0 |

TABLE 3.04.--SALAD DRESSINGS AND MAYONNAISE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 44.7 | 1.1 | 22.6 | 34.2 | 39.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 25.5 | 1.1 | 16.4 | 21.7 | 24.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 13.5 | 0 | 4.8 | 9.9 | 10.4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 5.7 | 0 | 1.3 | 2.5 | 4.0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.7 | 1.0 | 1.5 | 1.6 | 1.6 |
| STANDARD DEVIATION.....DO..... | 1.0 | .0 | .8 | .8 | .9 |
| MAXIMUM.....DO..... | 8.0 | 1.0 | 5.0 | 5.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 12 | 3 | 5 | 8 | 8 |
| STANDARD DEVIATION.....GM..... | 14 | 2 | 5 | 10 | 8 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 2 | | 1 | 2 | 2 |
| 25TH.....GM..... | 5 | | 2 | 2 | 3 |
| 50TH.....GM..... | 9 | | 3 | 5 | 5 |
| 75TH.....GM..... | 16 | | 7 | 9 | 10 |
| 90TH.....GM..... | 26 | | 11 | 16 | 18 |
| 95TH.....GM..... | 36 | | 15 | 22 | 21 |
| 99TH.....GM..... | 59 | | 24 | 43 | 40 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 23 | 8 | 11 | 16 | 17 |
| STANDARD DEVIATION.....GM..... | 21 | 5 | 12 | 18 | 15 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 23 | 0 | 10 | 14 | 17 |
| STANDARD DEVIATION.....GM..... | 19 | 0 | 8 | 12 | 14 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 25 | 0 | 11 | 20 | 15 |
| STANDARD DEVIATION.....GM..... | 23 | 0 | 5 | 24 | 11 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 768 | 15 | 83 | 165 | 125 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,536 | 15 | 83 | 289 | 189 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 21 | 8 | 10 | 15 | 15 |
| STANDARD DEVIATION.....GM..... | 22 | 5 | 10 | 18 | 14 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 5 | | 2 | 2 | 2 |
| 25TH.....GM..... | 9 | | 5 | 5 | 5 |
| 50TH.....GM..... | 15 | | 5 | 10 | 14 |
| 75TH.....GM..... | 29 | | 14 | 15 | 16 |
| 90TH.....GM..... | 44 | | 28 | 30 | 31 |
| 95TH.....GM..... | 60 | | 30 | 41 | 32 |
| 99TH.....GM..... | 94 | | 41 | 83 | 63 |
| MAXIMUM.....GM..... | 768 | 15 | 83 | 165 | 125 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 1601.6 | 0 | 20.5 | 46.9 | 67.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 5.4 | 0 | 5.8 | 5.1 | 5.8 |

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 37.6 | 41.9 | 40.3 | 43.5 | 49.5 | 51.7 | 51.7 | 50.3 | 42.3 | 41.5 | 28.0 | 34.9 |
| 24.1 | 26.0 | 23.9 | 24.4 | 26.9 | 28.3 | 27.0 | 28.3 | 26.3 | 23.7 | 15.4 | 22.0 |
| 10.3 | 11.9 | 11.0 | 14.1 | 14.6 | 17.7 | 15.5 | 15.7 | 10.9 | 12.5 | 8.6 | 8.2 |
| 3.2 | 4.0 | 5.4 | 5.0 | 8.1 | 5.8 | 9.1 | 6.3 | 5.1 | 5.3 | 4.0 | 4.7 |
| 1.6 | 1.6 | 1.7 | 1.7 | 1.9 | 1.7 | 1.9 | 1.8 | 1.6 | 1.7 | 1.7 | 1.7 |
| .9 | .9 | 1.0 | 1.0 | 1.1 | .9 | 1.1 | 1.0 | .9 | .9 | .9 | .9 |
| 6.0 | 6.0 | 6.0 | 6.0 | 6.0 | 6.0 | 8.0 | 6.0 | 5.0 | 6.0 | 4.0 | 6.0 |
| 10 | 10 | 13 | 13 | 15 | 13 | 15 | 12 | 12 | 9 | 12 | 10 |
| 10 | 10 | 12 | 16 | 15 | 12 | 19 | 12 | 12 | 9 | 14 | 9 |
| 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 3 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 3 | 5 | 4 |
| 7 | 6 | 10 | 9 | 10 | 10 | 10 | 10 | 9 | 6 | 9 | 6 |
| 12 | 12 | 18 | 16 | 20 | 18 | 20 | 15 | 14 | 11 | 16 | 12 |
| 21 | 20 | 29 | 27 | 31 | 26 | 31 | 25 | 26 | 20 | 24 | 21 |
| 29 | 27 | 40 | 37 | 41 | 36 | 41 | 36 | 39 | 26 | 29 | 26 |
| 46 | 46 | 51 | 61 | 73 | 60 | 63 | 61 | 56 | 44 | 56 | 41 |
| 22 | 19 | 27 | 23 | 28 | 24 | 26 | 23 | 20 | 19 | 21 | 18 |
| 26 | 17 | 26 | 24 | 25 | 21 | 22 | 21 | 16 | 17 | 16 | 14 |
| 20 | 22 | 26 | 25 | 27 | 25 | 27 | 22 | 24 | 17 | 24 | 20 |
| 13 | 20 | 19 | 33 | 20 | 18 | 19 | 17 | 21 | 14 | 27 | 15 |
| 22 | 18 | 24 | 26 | 28 | 25 | 28 | 26 | 28 | 18 | 25 | 19 |
| 14 | 12 | 16 | 17 | 22 | 17 | 37 | 18 | 16 | 12 | 14 | 11 |
| 279 | 270 | 151 | 470 | 278 | 245 | 768 | 245 | 169 | 201 | 250 | 78 |
| 279 | 297 | 340 | 940 | 495 | 360 | 1,536 | 311 | 250 | 256 | 365 | 184 |
| 19 | 18 | 23 | 22 | 24 | 22 | 24 | 21 | 21 | 16 | 22 | 17 |
| 19 | 18 | 20 | 27 | 22 | 20 | 29 | 19 | 18 | 16 | 17 | 14 |
| 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 3 | 5 | 5 |
| 7 | 5 | 11 | 9 | 11 | 9 | 10 | 9 | 10 | 5 | 14 | 5 |
| 14 | 14 | 15 | 15 | 16 | 15 | 16 | 15 | 15 | 14 | 15 | 15 |
| 28 | 28 | 29 | 30 | 31 | 30 | 31 | 29 | 30 | 21 | 29 | 28 |
| 32 | 32 | 46 | 44 | 55 | 46 | 47 | 44 | 32 | 31 | 46 | 32 |
| 59 | 49 | 63 | 60 | 63 | 61 | 63 | 60 | 59 | 41 | 59 | 47 |
| 83 | 83 | 118 | 94 | 118 | 94 | 118 | 83 | 113 | 64 | 83 | 63 |
| 250 | 256 | 139 | 470 | 250 | 245 | 768 | 245 | 125 | 188 | 125 | 78 |
| 75.0 | 76.2 | 58.4 | 78.5 | 226.0 | 261.5 | 282.7 | 286.9 | 43.0 | 42.2 | 7.4 | 29.2 |
| 6.1 | 5.2 | 6.0 | 7.1 | 6.3 | 5.4 | 5.9 | 4.6 | 5.6 | 3.4 | 2.9 | 5.1 |

TABLE 3.05.--MAYONNAISE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 21.4 | .4 | 13.3 | 19.4 | 20.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 14.4 | .4 | 10.0 | 13.7 | 13.9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 5.3 | 0 | 2.8 | 4.2 | 4.7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.7 | 0 | .6 | 1.5 | 1.7 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.5 | 1.0 | 1.4 | 1.4 | 1.5 |
| STANDARD DEVIATION.....DO..... | .8 | .0 | .7 | .7 | .8 |
| MAXIMUM.....DO..... | 8.0 | 1.0 | 5.0 | 4.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 7 | 3 | 4 | 6 | 5 |
| STANDARD DEVIATION.....GM..... | 7 | 2 | 3 | 10 | 5 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 2 | | 1 | 1 | 1 |
| 25TH.....GM..... | 2 | | 2 | 2 | 2 |
| 50TH.....GM..... | 5 | | 2 | 5 | 5 |
| 75TH.....GM..... | 9 | | 5 | 6 | 9 |
| 90TH.....GM..... | 14 | | 9 | 9 | 11 |
| 95TH.....GM..... | 18 | | 9 | 14 | 14 |
| 99TH.....GM..... | 31 | | 14 | 96 | 28 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 14 | 10 | 9 | 12 | 11 |
| STANDARD DEVIATION.....GM..... | 13 | 5 | 8 | 10 | 8 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 14 | 0 | 8 | 10 | 12 |
| STANDARD DEVIATION.....GM..... | 10 | 0 | 5 | 8 | 8 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 15 | 0 | 7 | 24 | 12 |
| STANDARD DEVIATION.....GM..... | 11 | 0 | 2 | 30 | 10 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 206 | 14 | 41 | 165 | 69 |
| MAXIMUM OVER 3 DAYS.....GM..... | 289 | 14 | 41 | 289 | 165 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 13 | 10 | 8 | 13 | 11 |
| STANDARD DEVIATION.....GM..... | 12 | 5 | 7 | 17 | 8 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 2 | | 2 | 2 | 2 |
| 25TH.....GM..... | 5 | | 5 | 5 | 5 |
| 50TH.....GM..... | 14 | | 5 | 9 | 9 |
| 75TH.....GM..... | 14 | | 14 | 14 | 14 |
| 90TH.....GM..... | 28 | | 14 | 28 | 28 |
| 95TH.....GM..... | 28 | | 28 | 28 | 28 |
| 99TH.....GM..... | 55 | | 28 | 83 | 41 |
| MAXIMUM.....GM..... | 193 | 14 | 41 | 165 | 55 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 774.6 | 0 | 9.5 | 28.7 | 42.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 6.4 | 0 | 5.0 | 6.0 | 7.7 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 17.7 | 20.6 | 21.0 | 21.9 | 25.6 | 25.4 | 24.1 | 22.2 | 15.9 | 18.4 | 12.6 | 13.3 |
| 12.7 | 13.7 | 13.4 | 14.7 | 16.1 | 17.6 | 15.1 | 15.4 | 11.7 | 11.8 | 7.8 | 10.0 |
| 3.9 | 5.4 | 5.1 | 5.7 | 6.8 | 6.4 | 6.2 | 5.4 | 2.8 | 5.4 | 3.2 | 2.4 |
| 1.0 | 1.5 | 2.5 | 1.4 | 2.7 | 1.4 | 2.9 | 1.3 | 1.4 | 1.3 | 1.5 | .8 |
| 1.4 | 1.5 | 1.6 | 1.5 | 1.6 | 1.4 | 1.6 | 1.4 | 1.4 | 1.5 | 1.5 | 1.4 |
| .8 | .7 | .9 | .8 | .9 | .7 | .9 | .7 | .7 | .8 | .7 | .7 |
| 6.0 | 4.0 | 6.0 | 6.0 | 6.0 | 5.0 | 8.0 | 5.0 | 4.0 | 5.0 | 3.0 | 4.0 |
| 6 | 6 | 9 | 6 | 9 | 6 | 7 | 5 | 7 | 6 | 8 | 5 |
| 7 | 6 | 9 | 6 | 8 | 5 | 7 | 5 | 8 | 5 | 5 | 4 |
| 1 | 2 | 2 | 2 | 2 | 2 | 2 | 1 | 2 | 1 | 2 | 1 |
| 2 | 2 | 5 | 2 | 5 | 2 | 3 | 2 | 2 | 2 | 5 | 2 |
| 5 | 5 | 6 | 5 | 6 | 5 | 5 | 5 | 5 | 5 | 9 | 5 |
| 9 | 7 | 10 | 9 | 12 | 9 | 9 | 8 | 9 | 9 | 9 | 6 |
| 14 | 14 | 18 | 14 | 18 | 14 | 14 | 11 | 18 | 12 | 12 | 14 |
| 18 | 18 | 28 | 18 | 23 | 18 | 23 | 14 | 21 | 18 | 14 | 14 |
| 38 | 28 | 46 | 28 | 37 | 25 | 37 | 28 | 53 | 25 | 28 | 18 |
| 14 | 13 | 18 | 13 | 19 | 14 | 16 | 12 | 14 | 12 | 16 | 11 |
| 14 | 12 | 18 | 11 | 17 | 12 | 13 | 11 | 12 | 11 | 9 | 10 |
| 15 | 13 | 16 | 13 | 19 | 14 | 15 | 12 | 16 | 11 | 14 | 12 |
| 12 | 10 | 9 | 8 | 13 | 9 | 11 | 10 | 7 | 9 | 2 | 7 |
| 16 | 12 | 21 | 15 | 16 | 14 | 14 | 12 | 22 | 14 | 15 | 12 |
| 11 | 9 | 12 | 12 | 8 | 7 | 9 | 7 | 15 | 7 | 8 | 5 |
| 113 | 110 | 151 | 110 | 110 | 124 | 110 | 206 | 73 | 62 | 41 | 55 |
| 138 | 138 | 154 | 142 | 220 | 138 | 124 | 234 | 159 | 83 | 83 | 55 |
| 14 | 12 | 17 | 12 | 16 | 13 | 14 | 11 | 15 | 11 | 15 | 11 |
| 12 | 11 | 13 | 10 | 14 | 10 | 11 | 10 | 13 | 10 | 8 | 8 |
| 2 | 2 | 5 | 2 | 5 | 2 | 2 | 2 | 5 | 2 | 5 | 2 |
| 5 | 5 | 9 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 14 | 5 |
| 14 | 14 | 14 | 14 | 14 | 14 | 14 | 9 | 14 | 9 | 14 | 14 |
| 14 | 14 | 28 | 14 | 28 | 14 | 14 | 14 | 14 | 14 | 14 | 14 |
| 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 |
| 28 | 28 | 41 | 28 | 41 | 28 | 28 | 28 | 41 | 28 | 28 | 28 |
| 55 | 55 | 69 | 41 | 69 | 55 | 55 | 41 | 73 | 55 | 41 | 41 |
| 113 | 110 | 110 | 110 | 110 | 84 | 110 | 193 | 73 | 62 | 41 | 41 |
| 30.1 | 27.8 | 38.1 | 41.7 | 132.6 | 132.6 | 120.5 | 121.4 | 18.9 | 19.2 | 1.5 | 8.9 |
| 5.7 | 4.3 | 8.2 | 8.6 | 8.2 | 6.7 | 6.4 | 5.4 | 7.4 | 4.0 | 1.4 | 5.0 |

TABLE 3.06.--SALAC DRESSINGS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST CNCE IN 3 DAYS.....PCT..... | 29.7 | .6 | 11.0 | 18.6 | 22.8 |
| CN ONLY 1 OF 3 DAYS.....PCT..... | 19.2 | .6 | 8.5 | 13.4 | 15.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 7.8 | 0 | 2.3 | 4.6 | 5.4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.7 | 0 | .3 | .6 | 1.6 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 1.6 | 1.0 | 1.4 | 1.4 | 1.5 |
| STANDARD DEVIATION.....DO..... | .8 | .0 | .7 | .6 | .8 |
| MAXIMUM.....DO..... | 6.0 | 1.0 | 5.0 | 4.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 14 | 2 | 6 | 8 | 9 |
| STANDARD DEVIATION.....GM..... | 14 | 2 | 6 | 8 | 8 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 2 | (2) | | 2 | 2 |
| 25TH.....GM..... | 5 | | 2 | 2 | 3 |
| 50TH.....GM..... | 10 | | 4 | 5 | 7 |
| 75TH.....GM..... | 20 | | 7 | 10 | 11 |
| 90TH.....GM..... | 29 | | 15 | 19 | 20 |
| 95TH.....GM..... | 39 | | 20 | 21 | 21 |
| 99TH.....GM..... | 62 | | 28 | 42 | 41 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 28 | 7 | 13 | 18 | 21 |
| STANDARD DEVIATION.....GM..... | 23 | 5 | 14 | 21 | 17 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 29 | 0 | 14 | 18 | 20 |
| STANDARD DEVIATION.....GM..... | 29 | 0 | 11 | 14 | 16 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 28 | 0 | 11 | 10 | 16 |
| STANDARD DEVIATION.....GM..... | 19 | 0 | 5 | 6 | 12 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 768 | 15 | 83 | 128 | 125 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,518 | 15 | 83 | 128 | 189 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 26 | 7 | 12 | 17 | 19 |
| STANDARD DEVIATION.....GM..... | 25 | 5 | 12 | 18 | 16 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 5 | | 2 | 3 | 5 |
| 25TH.....GM..... | 15 | | 5 | 5 | 5 |
| 50TH.....GM..... | 16 | | 7 | 15 | 15 |
| 75TH.....GM..... | 31 | | 16 | 16 | 29 |
| 90TH.....GM..... | 59 | | 30 | 31 | 32 |
| 95TH.....GM..... | 63 | | 32 | 59 | 59 |
| 99TH.....GM..... | 123 | | 59 | 123 | 64 |
| MAXIMUM.....GM..... | 768 | 15 | 83 | 128 | 125 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 829.6 | 0 | 11.0 | 18.1 | 24.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 4.7 | 0 | 6.8 | 4.1 | 4.0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 23.9 | 26.4 | 24.7 | 27.4 | 31.2 | 35.2 | 35.9 | 36.2 | 28.9 | 27.4 | 16.8 | 24.4 |
| 16.9 | 18.9 | 17.2 | 17.0 | 19.6 | 23.1 | 21.4 | 23.0 | 18.4 | 17.0 | 8.5 | 15.6 |
| 5.6 | 6.2 | 5.5 | 8.2 | 7.9 | 9.1 | 10.2 | 9.9 | 7.2 | 7.6 | 6.7 | 5.2 |
| 1.5 | 1.3 | 2.0 | 2.3 | 3.7 | 3.0 | 4.3 | 3.3 | 3.3 | 2.8 | 1.7 | 3.6 |
| 1.4 | 1.4 | 1.5 | 1.6 | 1.6 | 1.5 | 1.6 | 1.6 | 1.6 | 1.6 | 1.7 | 1.7 |
| .7 | .7 | .8 | .8 | .9 | .8 | .9 | .8 | .9 | .8 | .8 | .9 |
| 5.0 | 6.0 | 5.0 | 5.0 | 6.0 | 6.0 | 5.0 | 6.0 | 5.0 | 5.0 | 4.0 | 6.0 |
| 11 | 11 | 14 | 15 | 17 | 14 | 16 | 14 | 13 | 10 | 15 | 11 |
| 11 | 11 | 13 | 19 | 15 | 13 | 21 | 12 | 13 | 9 | 16 | 9 |
| 2 | 2 | 2 | 2 | 3 | 2 | 3 | 2 | 2 | 2 | 2 | 2 |
| 5 | 5 | 5 | 5 | 7 | 5 | 7 | 5 | 5 | 5 | 5 | 5 |
| 9 | 9 | 10 | 10 | 10 | 10 | 11 | 10 | 10 | 9 | 11 | 10 |
| 15 | 12 | 20 | 20 | 21 | 20 | 21 | 20 | 15 | 11 | 20 | 15 |
| 21 | 21 | 30 | 31 | 32 | 28 | 31 | 27 | 30 | 21 | 21 | 23 |
| 29 | 30 | 41 | 40 | 42 | 40 | 42 | 40 | 44 | 30 | 46 | 26 |
| 57 | 55 | 54 | 74 | 82 | 61 | 77 | 62 | 56 | 45 | 122 | 52 |
| 26 | 24 | 31 | 30 | 35 | 30 | 32 | 28 | 24 | 22 | 26 | 21 |
| 28 | 22 | 25 | 26 | 26 | 23 | 24 | 22 | 20 | 18 | 19 | 15 |
| 22 | 26 | 33 | 33 | 33 | 31 | 35 | 28 | 26 | 20 | 26 | 25 |
| 14 | 21 | 22 | 42 | 22 | 21 | 51 | 20 | 24 | 16 | 29 | 17 |
| 23 | 22 | 24 | 31 | 33 | 28 | 29 | 29 | 30 | 20 | 34 | 19 |
| 16 | 15 | 20 | 15 | 26 | 19 | 18 | 18 | 15 | 13 | 16 | 11 |
| 279 | 256 | 147 | 470 | 250 | 245 | 768 | 245 | 169 | 188 | 250 | 78 |
| 279 | 256 | 340 | 940 | 495 | 360 | 1,518 | 308 | 250 | 243 | 365 | 184 |
| 23 | 23 | 28 | 29 | 31 | 28 | 30 | 26 | 24 | 20 | 26 | 20 |
| 22 | 21 | 23 | 32 | 25 | 22 | 35 | 21 | 19 | 18 | 20 | 15 |
| 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| 10 | 10 | 15 | 15 | 15 | 15 | 15 | 15 | 15 | 8 | 15 | 8 |
| 16 | 16 | 21 | 29 | 29 | 29 | 29 | 16 | 16 | 15 | 16 | 16 |
| 30 | 30 | 31 | 31 | 32 | 31 | 31 | 31 | 31 | 29 | 31 | 30 |
| 47 | 47 | 61 | 59 | 61 | 59 | 61 | 59 | 44 | 32 | 46 | 32 |
| 63 | 63 | 64 | 63 | 66 | 63 | 63 | 63 | 59 | 47 | 59 | 59 |
| 90 | 120 | 125 | 125 | 125 | 120 | 123 | 120 | 125 | 104 | 125 | 64 |
| 250 | 256 | 128 | 470 | 250 | 245 | 768 | 245 | 125 | 188 | 125 | 78 |
| 44.9 | 48.3 | 20.2 | 36.8 | 93.4 | 131.4 | 162.1 | 165.4 | 24.1 | 23.0 | 5.9 | 20.3 |
| 6.3 | 6.0 | 4.0 | 5.9 | 4.7 | 4.5 | 5.6 | 4.1 | 4.6 | 3.0 | 3.9 | 5.0 |

TABLE 3.07.--MAYONNAISE-TYPE SALAD DRESSING
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.9 | .4 | 4.1 | 4.7 | 4.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.4 | .4 | 3.3 | 3.4 | 3.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.1 | 0 | .8 | 1.0 | .9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .4 | 0 | 0 | .3 | .6 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.5 | 1.0 | 1.3 | 1.4 | 1.5 |
| STANDARD DEVIATION.....DO..... | .8 | .0 | .5 | .7 | .8 |
| MAXIMUM.....DO..... | 6.0 | 1.0 | 3.0 | 3.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 7 | 3 | 4 | 5 | 6 |
| STANDARD DEVIATION.....GM..... | 7 | 2 | 4 | 4 | 5 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 2 | | | 1 | 2 |
| 25TH.....GM..... | 3 | | | 2 | 3 |
| 50TH.....GM..... | 5 | | | 3 | 5 |
| 75TH.....GM..... | 10 | | | 6 | 10 |
| 90TH.....GM..... | 15 | | | 10 | 13 |
| 95TH.....GM..... | 20 | | | 15 | 15 |
| 99TH.....GM..... | 39 | | | 15 | 21 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 17 | 9 | 10 | 11 | 14 |
| STANDARD DEVIATION.....GM..... | 17 | 5 | 9 | 9 | 10 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 16 | 0 | 11 | 10 | 10 |
| STANDARD DEVIATION.....GM..... | 12 | 0 | 11 | 6 | 6 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 14 | 0 | 0 | 8 | 11 |
| STANDARD DEVIATION.....GM..... | 10 | 0 | 0 | 3 | 7 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 147 | 15 | 44 | 44 | 44 |
| MAXIMUM OVER 3 DAYS.....GM..... | 162 | 15 | 73 | 44 | 87 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 15 | 9 | 9 | 9 | 12 |
| STANDARD DEVIATION.....GM..... | 13 | 5 | 8 | 7 | 9 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 2 | | | 2 | 2 |
| 25TH.....GM..... | 5 | | | 5 | 5 |
| 50TH.....GM..... | 15 | | | 5 | 10 |
| 75TH.....GM..... | 15 | | | 15 | 15 |
| 90TH.....GM..... | 29 | | | 15 | 29 |
| 95TH.....GM..... | 44 | | | 22 | 29 |
| 99TH.....GM..... | 59 | | | 29 | 44 |
| MAXIMUM.....GM..... | 120 | 15 | 30 | 29 | 44 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 102.3 | 0 | 3.0 | 3.5 | 1.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 3.7 | 0 | 5.5 | 3.0 | 1.5 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.2 | 4.6 | 3.5 | 4.1 | 5.9 | 5.1 | 5.6 | 4.7 | 4.5 | 4.9 | 3.3 | 5.9 |
| 3.4 | 3.4 | 2.7 | 3.3 | 3.3 | 3.4 | 3.6 | 3.4 | 3.8 | 3.3 | 2.7 | 4.9 |
| 1.2 | 1.0 | .5 | .8 | 1.8 | 1.3 | 1.1 | 1.2 | .7 | 1.2 | .6 | .5 |
| .5 | .3 | .3 | 0 | .9 | .3 | .9 | .2 | 0 | .4 | 0 | .5 |
| 1.5 | 1.5 | 1.5 | 1.3 | 1.7 | 1.5 | 1.6 | 1.4 | 1.2 | 1.6 | 1.2 | 1.5 |
| .7 | .8 | .8 | .6 | 1.0 | .7 | .8 | .6 | .4 | .9 | .4 | 1.2 |
| 4.0 | 5.0 | 4.0 | 4.0 | 5.0 | 3.0 | 5.0 | 3.0 | 2.0 | 5.0 | 2.0 | 6.0 |
| 7 | 7 | 11 | 8 | 10 | 7 | 8 | 7 | 7 | 8 | 8 | 6 |
| 6 | 7 | 12 | 6 | 8 | 6 | 8 | 6 | 6 | 10 | 7 | 5 |
| 2 | 2 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 | | 2 |
| 3 | 3 | | 3 | 5 | 2 | 3 | 2 | 3 | 3 | | 2 |
| 5 | 5 | | 5 | 9 | 5 | 5 | 5 | 5 | 5 | | 5 |
| 10 | 8 | | 10 | 15 | 10 | 10 | 10 | 10 | 10 | | 7 |
| 15 | 15 | | 20 | 20 | 15 | 16 | 15 | 15 | 20 | | 10 |
| 20 | 24 | | 20 | 29 | 20 | 20 | 20 | 20 | 39 | | 20 |
| 34 | 39 | | 24 | 34 | 29 | 39 | 29 | 29 | 44 | | 20 |
| 15 | 17 | 25 | 19 | 20 | 17 | 17 | 15 | 20 | 20 | 24 | 17 |
| 9 | 19 | 30 | 16 | 16 | 15 | 17 | 15 | 18 | 26 | 21 | 14 |
| 17 | 15 | 32 | 19 | 20 | 14 | 19 | 16 | 18 | 9 | 9 | 6 |
| 13 | 14 | 24 | 8 | 11 | 9 | 13 | 12 | 9 | 8 | 6 | 1 |
| 15 | 11 | 18 | 0 | 17 | 14 | 13 | 10 | 0 | 22 | 0 | 10 |
| 8 | 7 | 7 | 0 | 12 | 12 | 8 | 5 | 0 | 14 | 0 | (2) |
| 66 | 118 | 147 | 59 | 88 | 88 | 120 | 118 | 88 | 118 | 60 | 60 |
| 110 | 118 | 162 | 73 | 162 | 118 | 120 | 118 | 88 | 132 | 60 | 60 |
| 15 | 14 | 22 | 18 | 18 | 15 | 16 | 14 | 18 | 15 | 19 | 12 |
| 10 | 14 | 18 | 14 | 13 | 12 | 14 | 13 | 12 | 20 | 16 | 11 |
| 5 | 5 | | 5 | 5 | 2 | 5 | 2 | 5 | 2 | | 5 |
| 5 | 5 | | 10 | 10 | 5 | 5 | 5 | 10 | 5 | | 5 |
| 15 | 10 | | 15 | 15 | 14 | 15 | 15 | 15 | 10 | | 7 |
| 15 | 15 | | 28 | 28 | 15 | 15 | 15 | 29 | 15 | | 15 |
| 29 | 29 | | 44 | 30 | 29 | 29 | 29 | 29 | 29 | | 29 |
| 29 | 29 | | 44 | 44 | 30 | 44 | 39 | 44 | 44 | | 30 |
| 59 | 59 | | 59 | 59 | 59 | 59 | 59 | 59 | 118 | | 59 |
| 66 | 118 | 88 | 59 | 88 | 73 | 120 | 118 | 59 | 118 | 59 | 59 |
| 8.1 | 3.7 | 2.2 | 6.4 | 15.4 | 9.6 | 21.8 | 12.4 | 1.3 | 3.4 | 2.5 | 7.2 |
| 5.0 | 2.4 | 2.9 | 8.3 | 3.8 | 2.4 | 4.9 | 2.7 | 2.1 | 2.5 | 11.3 | 8.2 |

TABLE 3.08.--FRENCH DRESSING
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 6.6 | 0 | 1.5 | 3.9 | 5.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.2 | 0 | 1.3 | 3.5 | 5.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.1 | 0 | .2 | .4 | .5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | 0 | 0 | .1 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 0 | 1.1 | 1.1 | 1.2 |
| STANDARD DEVIATION.....DO..... | .6 | 0 | .3 | .4 | .5 |
| MAXIMUM.....DO..... | 4.0 | 0 | 2.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 13 | 0 | 8 | 8 | 9 |
| STANDARD DEVIATION.....GM..... | 13 | 0 | 8 | 8 | 7 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 2 | | | 2 | 2 |
| 25TH.....GM..... | 5 | | | 3 | 3 |
| 50TH.....GM..... | 10 | | | 5 | 5 |
| 75TH.....GM..... | 16 | | | 10 | 10 |
| 90TH.....GM..... | 21 | | | 21 | 21 |
| 95TH.....GM..... | 32 | | | 21 | 21 |
| 99TH.....GM..... | 48 | | | 42 | 21 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 31 | 0 | 22 | 21 | 25 |
| STANDARD DEVIATION.....GM..... | 34 | 0 | 22 | 21 | 20 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 29 | 0 | 26 | 18 | 18 |
| STANDARD DEVIATION.....GM..... | 20 | 0 | 4 | 20 | 8 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 27 | 0 | 0 | 16 | 11 |
| STANDARD DEVIATION.....GM..... | 17 | 0 | 0 | 0 | 6 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 750 | 0 | 83 | 125 | 125 |
| MAXIMUM OVER 3 DAYS.....GM..... | 750 | 0 | 83 | 125 | 125 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 30 | 0 | 23 | 20 | 22 |
| STANDARD DEVIATION.....GM..... | 30 | 0 | 20 | 20 | 19 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 5 | | | 5 | 5 |
| 25TH.....GM..... | 16 | | | 5 | 10 |
| 50TH.....GM..... | 30 | | | 16 | 16 |
| 75TH.....GM..... | 31 | | | 30 | 31 |
| 90TH.....GM..... | 63 | | | 63 | 47 |
| 95TH.....GM..... | 63 | | | 63 | 63 |
| 99TH.....GM..... | 125 | | | 125 | 63 |
| MAXIMUM.....GM..... | 750 | 0 | 83 | 125 | 125 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 103.2 | 0 | .9 | 3.1 | 2.4 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 3.3 | 0 | 5.1 | 4.1 | 1.9 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 6.3 | 7.2 | 6.0 | 7.6 | 7.2 | 7.0 | 8.4 | 7.0 | 6.7 | 5.9 | 3.8 | 5.0 |
| 5.3 | 5.8 | 4.7 | 6.1 | 5.9 | 5.6 | 6.6 | 5.5 | 4.4 | 4.1 | 2.5 | 3.9 |
| .9 | 1.3 | 1.2 | 1.2 | 1.2 | 1.1 | 1.4 | 1.3 | 1.9 | 1.0 | 1.2 | .9 |
| (2) | .2 | .1 | .3 | .1 | .3 | .4 | .2 | .3 | .7 | 0 | .2 |
| 1.2 | 1.3 | 1.2 | 1.2 | 1.2 | 1.3 | 1.3 | 1.3 | 1.4 | 1.4 | 1.5 | 1.3 |
| .5 | .5 | .5 | .5 | .5 | .6 | .6 | .6 | .7 | .7 | .8 | .6 |
| 3.0 | 4.0 | 3.0 | 4.0 | 4.0 | 4.0 | 3.0 | 4.0 | 3.0 | 4.0 | 3.0 | 4.0 |
| 11 | 10 | 13 | 13 | 15 | 13 | 16 | 11 | 14 | 9 | 12 | 10 |
| 13 | 9 | 11 | 12 | 13 | 10 | 20 | 9 | 11 | 7 | 17 | 7 |
| 2 | 2 | 2 | 2 | 3 | 2 | 3 | 3 | 5 | 2 | | 2 |
| 5 | 5 | 5 | 5 | 7 | 5 | 8 | 5 | 10 | 5 | | 5 |
| 8 | 5 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 5 | | 10 |
| 11 | 11 | 21 | 16 | 21 | 21 | 21 | 16 | 11 | 10 | | 11 |
| 21 | 21 | 31 | 21 | 31 | 21 | 26 | 21 | 38 | 21 | | 21 |
| 21 | 26 | 42 | 42 | 42 | 31 | 42 | 31 | 42 | 27 | | 26 |
| 83 | 42 | 42 | 83 | 63 | 47 | 63 | 42 | 47 | 42 | | 31 |
| 30 | 25 | 32 | 33 | 40 | 33 | 39 | 27 | 36 | 22 | 16 | 24 |
| 39 | 21 | 27 | 34 | 31 | 24 | 60 | 18 | 27 | 18 | 9 | 15 |
| 25 | 27 | 31 | 26 | 34 | 30 | 32 | 31 | 21 | 17 | 37 | 24 |
| 17 | 17 | 18 | 15 | 27 | 16 | 21 | 18 | 14 | 8 | 38 | 11 |
| 16 | 9 | 27 | 31 | 35 | 25 | 43 | 21 | 30 | 18 | 0 | 26 |
| 0 | 5 | 8 | 15 | 8 | 20 | 18 | 7 | 18 | 9 | 0 | 0 |
| 250 | 125 | 125 | 250 | 250 | 125 | 750 | 125 | 125 | 125 | 125 | 63 |
| 250 | 125 | 126 | 250 | 281 | 188 | 750 | 141 | 141 | 125 | 240 | 94 |
| 28 | 24 | 31 | 30 | 37 | 31 | 37 | 27 | 28 | 19 | 24 | 24 |
| 34 | 20 | 24 | 30 | 31 | 22 | 49 | 18 | 23 | 15 | 25 | 14 |
| 5 | 5 | 5 | 5 | 8 | 5 | 10 | 5 | 5 | 5 | | 5 |
| 10 | 16 | 16 | 16 | 16 | 16 | 16 | 16 | 16 | 16 | | 16 |
| 16 | 16 | 31 | 30 | 31 | 31 | 31 | 16 | 16 | 16 | | 16 |
| 31 | 31 | 32 | 31 | 39 | 32 | 47 | 31 | 31 | 28 | | 31 |
| 63 | 63 | 63 | 63 | 63 | 63 | 63 | 48 | 47 | 31 | | 47 |
| 63 | 63 | 94 | 63 | 95 | 63 | 63 | 63 | 63 | 32 | | 47 |
| 250 | 83 | 125 | 125 | 125 | 125 | 125 | 94 | 125 | 63 | | 63 |
| 250 | 125 | 125 | 250 | 250 | 125 | 750 | 125 | 125 | 125 | 125 | 63 |
| 5.7 | 6.1 | 1.8 | 4.7 | 9.2 | 11.0 | 25.6 | 21.0 | 4.0 | 2.9 | 1.1 | 3.6 |
| 3.7 | 3.1 | 1.7 | 3.4 | 2.7 | 2.4 | 4.8 | 3.3 | 3.7 | 2.0 | 3.7 | 5.6 |

TABLE 3.09.--ITALIAN DRESSING
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 8.1 | 0 | 1.3 | 3.9 | 5.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 6.1 | 0 | 1.3 | 3.0 | 4.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.7 | 0 | 0 | .7 | 1.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .4 | 0 | 0 | .1 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.3 | 0 | 1.0 | 1.3 | 1.2 |
| STANDARD DEVIATION.....DO..... | .6 | 0 | .0 | .6 | .5 |
| MAXIMUM.....DO..... | 5.0 | 0 | 1.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 12 | 0 | 4 | 9 | 9 |
| STANDARD DEVIATION.....GM..... | 11 | 0 | 5 | 9 | 9 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 2 | | | 1 | 2 |
| 25TH.....GM..... | 5 | | | 2 | 5 |
| 50TH.....GM..... | 10 | | | 5 | 7 |
| 75TH.....GM..... | 15 | | | 10 | 10 |
| 90TH.....GM..... | 20 | | | 20 | 20 |
| 95TH.....GM..... | 34 | | | 29 | 20 |
| 99TH.....GM..... | 52 | | | 39 | 49 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 29 | 0 | 12 | 19 | 23 |
| STANDARD DEVIATION.....GM..... | 22 | 0 | 15 | 17 | 21 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 28 | 0 | 0 | 28 | 20 |
| STANDARD DEVIATION.....GM..... | 23 | 0 | 0 | 17 | 13 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 29 | 0 | 0 | 7 | 49 |
| STANDARD DEVIATION.....GM..... | 17 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 470 | 0 | 59 | 59 | 118 |
| MAXIMUM OVER 3 DAYS.....GM..... | 940 | 0 | 59 | 118 | 146 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 28 | 0 | 12 | 21 | 22 |
| STANDARD DEVIATION.....GM..... | 21 | 0 | 15 | 18 | 19 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 5 | | | 2 | 5 |
| 25TH.....GM..... | 15 | | | 5 | 10 |
| 50TH.....GM..... | 29 | | | 15 | 15 |
| 75TH.....GM..... | 30 | | | 29 | 29 |
| 90TH.....GM..... | 59 | | | 59 | 59 |
| 95TH.....GM..... | 59 | | | 59 | 59 |
| 99TH.....GM..... | 118 | | | 59 | 118 |
| MAXIMUM.....GM..... | 470 | 0 | 59 | 59 | 118 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 275.2 | 0 | 0 | 8.2 | 7.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 6.7 | 0 | 0 | 9.8 | 6.3 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.8 | 6.9 | 6.5 | 6.7 | 6.6 | 11.0 | 10.2 | 11.3 | 7.6 | 7.3 | 4.5 | 5.7 |
| 4.5 | 5.9 | 5.3 | 5.1 | 4.9 | 8.5 | 7.2 | 8.2 | 5.4 | 5.5 | 3.5 | 3.4 |
| 1.1 | .8 | .9 | 1.2 | 1.6 | 2.3 | 2.2 | 2.3 | 1.8 | 1.4 | .9 | 1.7 |
| .2 | .2 | .3 | .4 | .1 | .2 | .8 | .8 | .5 | .4 | .2 | .5 |
| 1.3 | 1.2 | 1.3 | 1.3 | 1.3 | 1.3 | 1.4 | 1.4 | 1.4 | 1.3 | 1.3 | 1.6 |
| .5 | .4 | .6 | .7 | .5 | .5 | .7 | .7 | .9 | .6 | .6 | .7 |
| 4.0 | 3.0 | 4.0 | 4.0 | 3.0 | 4.0 | 5.0 | 5.0 | 5.0 | 4.0 | 3.0 | 3.0 |
| 12 | 10 | 12 | 17 | 13 | 13 | 15 | 12 | 11 | 9 | 11 | 12 |
| 11 | 9 | 10 | 30 | 10 | 10 | 12 | 9 | 9 | 8 | 10 | 11 |
| 2 | 2 | 2 | 2 | 3 | 3 | 3 | 2 | 2 | 2 | | 2 |
| 5 | 5 | 5 | 10 | 5 | 5 | 7 | 5 | 5 | 5 | | 5 |
| 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 7 | | 10 |
| 15 | 10 | 15 | 20 | 20 | 15 | 20 | 15 | 15 | 10 | | 15 |
| 20 | 20 | 26 | 29 | 20 | 20 | 30 | 20 | 20 | 20 | | 29 |
| 39 | 20 | 39 | 44 | 39 | 29 | 39 | 29 | 29 | 20 | | 39 |
| 65 | 39 | 59 | 78 | 49 | 49 | 59 | 39 | 49 | 49 | | 52 |
| 25 | 25 | 31 | 31 | 33 | 32 | 33 | 28 | 22 | 20 | 24 | 22 |
| 23 | 18 | 23 | 19 | 26 | 26 | 24 | 18 | 11 | 14 | 15 | 14 |
| 29 | 31 | 34 | 58 | 29 | 28 | 30 | 25 | 29 | 24 | 32 | 30 |
| 11 | 26 | 25 | 95 | 17 | 13 | 18 | 13 | 18 | 20 | 21 | 22 |
| 48 | 25 | 22 | 45 | 35 | 23 | 35 | 27 | 24 | 14 | 29 | 16 |
| 19 | 5 | 8 | 21 | 9 | 17 | 17 | 17 | 13 | 7 | 0 | 6 |
| 118 | 118 | 118 | 470 | 206 | 235 | 147 | 118 | 103 | 118 | 78 | 78 |
| 195 | 235 | 176 | 940 | 206 | 235 | 235 | 294 | 147 | 146 | 122 | 155 |
| 28 | 26 | 30 | 39 | 31 | 30 | 31 | 26 | 23 | 20 | 25 | 23 |
| 21 | 20 | 23 | 53 | 22 | 20 | 21 | 17 | 15 | 17 | 18 | 18 |
| 5 | 5 | 5 | 10 | 5 | 5 | 7 | 5 | 5 | 5 | | 5 |
| 15 | 15 | 15 | 15 | 15 | 15 | 15 | 15 | 10 | 10 | | 15 |
| 29 | 28 | 29 | 29 | 29 | 29 | 29 | 28 | 21 | 15 | | 15 |
| 29 | 29 | 29 | 44 | 30 | 30 | 44 | 29 | 29 | 29 | | 29 |
| 59 | 59 | 59 | 59 | 59 | 59 | 59 | 59 | 44 | 30 | | 59 |
| 59 | 59 | 78 | 78 | 59 | 59 | 73 | 59 | 59 | 59 | | 59 |
| 118 | 118 | 118 | 470 | 118 | 118 | 118 | 78 | 59 | 118 | | 78 |
| 118 | 118 | 118 | 470 | 118 | 146 | 118 | 118 | 59 | 118 | 78 | 78 |
| 14.8 | 16.4 | 2.9 | 6.4 | 21.4 | 46.8 | 67.0 | 59.1 | 9.3 | 9.1 | 0 | 6.1 |
| 9.6 | 9.4 | 2.6 | 4.8 | 6.3 | 6.1 | 9.4 | 5.3 | 7.6 | 5.3 | 0 | 6.9 |

TABLE 4.01.--FRESH GRAPEFRUIT
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.7 | 0 | 1.1 | 1.0 | 1.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.6 | 0 | 1.1 | .7 | 1.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.2 | 0 | 0 | .2 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .9 | 0 | 0 | .1 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.7 | 0 | 1.1 | 1.3 | 1.2 |
| STANDARD DEVIATION.....DO..... | .9 | 0 | .3 | .6 | .5 |
| MAXIMUM.....DO..... | 7.0 | 0 | 2.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 90 | 0 | 55 | 66 | 64 |
| STANDARD DEVIATION.....GM..... | 58 | 0 | 24 | 31 | 39 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 38 | | | | |
| 25TH.....GM..... | 45 | | | | |
| 50TH.....GM..... | 89 | | | | |
| 75TH.....GM..... | 110 | | | | |
| 90TH.....GM..... | 165 | | | | |
| 95TH.....GM..... | 205 | | | | |
| 99TH.....GM..... | 268 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 163 | 0 | 164 | 159 | 155 |
| STANDARD DEVIATION.....GM..... | 67 | 0 | 73 | 71 | 55 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 162 | 0 | 0 | 139 | 195 |
| STANDARD DEVIATION.....GM..... | 59 | 0 | 0 | 15 | 75 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 167 | 0 | 0 | 134 | 0 |
| STANDARD DEVIATION.....GM..... | 65 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 804 | 0 | 268 | 330 | 402 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,876 | 0 | 268 | 402 | 670 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 159 | 0 | 145 | 149 | 158 |
| STANDARD DEVIATION.....GM..... | 58 | 0 | 57 | 56 | 64 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 106 | | | | |
| 25TH.....GM..... | 134 | | | | |
| 50TH.....GM..... | 134 | | | | |
| 75TH.....GM..... | 165 | | | | |
| 90TH.....GM..... | 268 | | | | |
| 95TH.....GM..... | 268 | | | | |
| 99TH.....GM..... | 330 | | | | |
| MAXIMUM.....GM..... | 660 | 0 | 268 | 330 | 402 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 4.9 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .2 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,354 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.6 | 2.4 | 2.2 | 2.3 | 2.7 | 3.7 | 6.0 | 7.4 | 8.9 | 12.5 | 7.9 | 10.4 |
| 1.3 | 1.7 | 1.7 | 1.5 | 2.0 | 2.1 | 3.2 | 3.8 | 3.5 | 5.7 | 2.9 | 5.6 |
| .2 | .5 | .4 | .4 | .4 | 1.1 | 1.8 | 2.2 | 2.8 | 3.0 | 2.4 | 2.1 |
| .1 | .2 | .1 | .4 | .3 | .5 | 1.0 | 1.4 | 2.6 | 3.7 | 2.6 | 2.8 |
| 1.3 | 1.4 | 1.3 | 1.7 | 1.4 | 1.6 | 1.7 | 1.8 | 2.0 | 1.8 | 2.0 | 1.8 |
| .9 | .6 | .5 | 1.0 | .9 | .7 | .8 | 1.0 | 1.0 | .9 | .9 | 1.0 |
| 5.0 | 3.0 | 3.0 | 4.0 | 5.0 | 4.0 | 5.0 | 7.0 | 7.0 | 3.0 | 4.0 | 5.0 |
| 72 | 70 | 63 | 93 | 85 | 90 | 90 | 94 | 110 | 93 | 93 | 83 |
| 44 | 33 | 33 | 45 | 55 | 52 | 53 | 68 | 76 | 51 | 53 | 52 |
| | 35 | | | 45 | 35 | 45 | 37 | 45 | 38 | | 22 |
| | 45 | | | 45 | 45 | 45 | 45 | 45 | 45 | | 45 |
| | 55 | | | 55 | 89 | 89 | 89 | 89 | 89 | | 55 |
| | 89 | | | 89 | 110 | 110 | 110 | 134 | 134 | | 134 |
| | 110 | | | 155 | 165 | 165 | 160 | 268 | 165 | | 165 |
| | 134 | | | 223 | 199 | 179 | 223 | 291 | 179 | | 165 |
| | 179 | | | 268 | 268 | 268 | 330 | 330 | 231 | | 244 |
| 169 | 165 | 151 | 200 | 183 | 182 | 160 | 162 | 148 | 149 | 142 | 133 |
| 63 | 63 | 55 | 75 | 69 | 103 | 61 | 60 | 35 | 50 | 23 | 43 |
| 162 | 150 | 154 | 196 | 197 | 176 | 170 | 158 | 166 | 152 | 133 | 132 |
| 58 | 38 | 56 | 74 | 78 | 64 | 61 | 58 | 77 | 40 | 8 | 13 |
| 186 | 134 | 134 | 142 | 187 | 149 | 160 | 183 | 194 | 153 | 149 | 157 |
| 44 | 0 | 0 | 13 | 62 | 46 | 49 | 91 | 69 | 34 | 56 | 33 |
| 330 | 330 | 536 | 469 | 495 | 660 | 460 | 804 | 660 | 330 | 536 | 330 |
| 670 | 536 | 620 | 603 | 804 | 804 | 804 | 1,876 | 990 | 804 | 1,072 | 732 |
| 160 | 153 | 150 | 159 | 178 | 169 | 160 | 160 | 167 | 152 | 141 | 136 |
| 56 | 50 | 68 | 57 | 67 | 77 | 61 | 58 | 59 | 42 | 30 | 28 |
| | 106 | | | 134 | 106 | 106 | 106 | 134 | 115 | | 83 |
| | 134 | | | 134 | 134 | 134 | 134 | 134 | 134 | | 134 |
| | 134 | | | 134 | 134 | 134 | 134 | 134 | 134 | | 134 |
| | 134 | | | 230 | 165 | 165 | 165 | 165 | 165 | | 134 |
| | 268 | | | 268 | 268 | 268 | 268 | 268 | 212 | | 165 |
| | 268 | | | 330 | 330 | 268 | 268 | 330 | 268 | | 165 |
| | 330 | | | 330 | 454 | 330 | 330 | 330 | 268 | | 268 |
| 330 | 330 | 536 | 330 | 495 | 660 | 460 | 495 | 330 | 330 | 268 | 268 |
| 0 | 0 | 0 | 0 | 0 | 0 | 2.2 | 0 | 0 | 1.2 | 0 | 1.5 |
| 0 | 0 | 0 | 0 | 0 | 0 | .4 | 0 | 0 | .3 | 0 | .8 |

TABLE 4.02.--GRAPEFRUIT JUICE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS 1/ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.6 | .6 | 1.0 | 1.2 | 1.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.9 | .3 | .6 | .6 | 1.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.0 | 0 | .3 | .5 | .3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .8 | .3 | .1 | .1 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE 1/NUMBER..... | 1.9 | 2.8 | 1.5 | 1.6 | 1.6 |
| STANDARD DEVIATION.....DO..... | 1.4 | 2.0 | .8 | .6 | 1.1 |
| MAXIMUM.....DO..... | 14.0 | 5.0 | 3.0 | 3.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY 1/GM..... | 131 | 133 | 79 | 95 | 100 |
| STANDARD DEVIATION.....GM..... | 115 | 103 | 66 | 44 | 70 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: 1/ | | | | | |
| 5TH.....GM..... | 41 | | | | |
| 25TH.....GM..... | 63 | | | | |
| 50TH.....GM..... | 83 | | | | |
| 75TH.....GM..... | 165 | | | | |
| 90TH.....GM..... | 250 | | | | |
| 95TH.....GM..... | 329 | | | | |
| 99TH.....GM..... | 658 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE 1/GM..... | 219 | 123 | 110 | 205 | 185 |
| STANDARD DEVIATION.....GM..... | 121 | 61 | 60 | 119 | 59 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE 1/GM..... | 223 | 0 | 212 | 183 | 239 |
| STANDARD DEVIATION.....GM..... | 114 | 0 | 29 | 38 | 86 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE 1/GM..... | 245 | 247 | 226 | 146 | 214 |
| STANDARD DEVIATION.....GM..... | 167 | 0 | 0 | 0 | 77 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,976 | 371 | 250 | 438 | 556 |
| MAXIMUM OVER 3 DAYS.....GM..... | 3,976 | 741 | 679 | 494 | 926 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE 1/GM..... | 202 | 143 | 156 | 174 | 184 |
| STANDARD DEVIATION.....GM..... | 99 | 44 | 66 | 47 | 52 |
| PERCENTILES: 1/ | | | | | |
| 5TH.....GM..... | 95 | | | | |
| 25TH.....GM..... | 125 | | | | |
| 50TH.....GM..... | 186 | | | | |
| 75TH.....GM..... | 247 | | | | |
| 90TH.....GM..... | 250 | | | | |
| 95TH.....GM..... | 375 | | | | |
| 99TH.....GM..... | 500 | | | | |
| MAXIMUM.....GM..... | 1,500 | 217 | 250 | 250 | 309 |
| QUANTITIES NOT SPECIFIED BY USERS: 1/ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 4.9 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .2 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.3 | 1.5 | 1.7 | 2.2 | 2.5 | 3.8 | 5.0 | 6.4 | 5.0 | 5.4 | 4.2 | 3.7 |
| .8 | 1.0 | 1.0 | 1.3 | 1.7 | 2.2 | 2.2 | 2.9 | 2.7 | 2.4 | 2.8 | 2.0 |
| .3 | .3 | .6 | .6 | .7 | 1.0 | 1.5 | 1.6 | 1.4 | 1.7 | .7 | .8 |
| .2 | .2 | .2 | .3 | .1 | .6 | 1.3 | 1.9 | .9 | 1.3 | .7 | .9 |
| 1.6 | 1.8 | 1.6 | 1.9 | 1.5 | 1.9 | 2.0 | 2.2 | 1.8 | 2.0 | 1.5 | 1.8 |
| .9 | 1.4 | .8 | 1.3 | .6 | 1.4 | 1.3 | 1.6 | .8 | 1.2 | .8 | 1.1 |
| 3.0 | 8.0 | 3.0 | 5.0 | 3.0 | 9.0 | 10.0 | 14.0 | 3.0 | 7.0 | 3.0 | 6.0 |
| 106 | 103 | 136 | 133 | 116 | 145 | 137 | 141 | 110 | 118 | 90 | 109 |
| 68 | 80 | 100 | 95 | 66 | 158 | 110 | 121 | 61 | 103 | 49 | 90 |
| | | | | 31 | 42 | 41 | 41 | 41 | 41 | | |
| | | | | 72 | 82 | 63 | 63 | 62 | 62 | | |
| | | | | 83 | 83 | 83 | 102 | 83 | 82 | | |
| | | | | 165 | 165 | 167 | 185 | 165 | 125 | | |
| | | | | 188 | 268 | 319 | 250 | 188 | 207 | | |
| | | | | 250 | 362 | 371 | 330 | 227 | 323 | | |
| | | | | 329 | 792 | 515 | 708 | 269 | 658 | | |
| 194 | 230 | 294 | 238 | 262 | 256 | 208 | 208 | 235 | 173 | 175 | 185 |
| 76 | 162 | 307 | 91 | 139 | 125 | 99 | 108 | 154 | 58 | 48 | 59 |
| 249 | 204 | 263 | 224 | 250 | 230 | 253 | 207 | 203 | 183 | 237 | 210 |
| 100 | 148 | 100 | 82 | 96 | 112 | 158 | 103 | 69 | 56 | 25 | 107 |
| 182 | 191 | 223 | 353 | 200 | 338 | 219 | 255 | 162 | 229 | 148 | 183 |
| 55 | 82 | 72 | 30 | 41 | 296 | 129 | 152 | 54 | 152 | 31 | 128 |
| 500 | 813 | 1,500 | 556 | 750 | 1,976 | 988 | 1,235 | 682 | 741 | 313 | 741 |
| 750 | 993 | 1,500 | 1,235 | 988 | 3,976 | 2,161 | 2,717 | 806 | 1,976 | 563 | 1,616 |
| 194 | 173 | 248 | 210 | 234 | 229 | 202 | 196 | 187 | 179 | 181 | 178 |
| 73 | 72 | 202 | 66 | 104 | 107 | 108 | 93 | 82 | 92 | 53 | 72 |
| | | | | 124 | 124 | 94 | 98 | 98 | 94 | | |
| | | | | 185 | 185 | 125 | 125 | 125 | 124 | | |
| | | | | 216 | 188 | 185 | 185 | 185 | 185 | | |
| | | | | 250 | 250 | 247 | 247 | 247 | 247 | | |
| | | | | 438 | 375 | 313 | 250 | 250 | 247 | | |
| | | | | 494 | 500 | 375 | 313 | 250 | 250 | | |
| | | | | 500 | 588 | 500 | 625 | 494 | 741 | | |
| 371 | 438 | 1,500 | 500 | 500 | 750 | 988 | 988 | 494 | 741 | 313 | 494 |
| 0 | 0 | 0 | 0 | 4.9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 3.4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 4.03.--FRESH ORANGES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 9.0 | .9 | 8.1 | 10.0 | 12.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.7 | .4 | 5.2 | 7.1 | 9.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.2 | 0 | 2.6 | 1.8 | 2.5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.1 | .5 | .3 | 1.1 | .4 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.6 | 2.2 | 1.5 | 1.5 | 1.4 |
| STANDARD DEVIATION.....DO..... | .9 | 1.0 | .7 | .9 | .9 |
| MAXIMUM.....DO..... | 8.0 | 3.0 | 6.0 | 6.0 | 7.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 78 | 63 | 57 | 68 | 63 |
| STANDARD DEVIATION.....GM..... | 54 | 25 | 30 | 46 | 48 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 24 | | 24 | 21 | 22 |
| 25TH.....GM..... | 48 | | 33 | 48 | 48 |
| 50TH.....GM..... | 48 | | 48 | 48 | 48 |
| 75TH.....GM..... | 97 | | 67 | 82 | 73 |
| 90TH.....GM..... | 145 | | 97 | 145 | 115 |
| 95TH.....GM..... | 180 | | 120 | 169 | 145 |
| 99TH.....GM..... | 290 | | 145 | 265 | 338 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 147 | 110 | 125 | 140 | 135 |
| STANDARD DEVIATION.....GM..... | 66 | 36 | 40 | 47 | 57 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 158 | 0 | 123 | 143 | 159 |
| STANDARD DEVIATION.....GM..... | 63 | 0 | 42 | 56 | 62 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 167 | 82 | 120 | 161 | 212 |
| STANDARD DEVIATION.....GM..... | 61 | 8 | 34 | 50 | 97 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,160 | 145 | 290 | 470 | 580 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,595 | 268 | 435 | 870 | 1,015 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 146 | 87 | 117 | 134 | 134 |
| STANDARD DEVIATION.....GM..... | 57 | 34 | 45 | 44 | 46 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 73 | | 50 | 63 | 65 |
| 25TH.....GM..... | 145 | | 73 | 100 | 100 |
| 50TH.....GM..... | 145 | | 145 | 145 | 145 |
| 75TH.....GM..... | 145 | | 145 | 145 | 145 |
| 90TH.....GM..... | 180 | | 145 | 145 | 180 |
| 95TH.....GM..... | 228 | | 180 | 195 | 180 |
| 99TH.....GM..... | 360 | | 228 | 290 | 290 |
| MAXIMUM.....GM..... | 1,160 | 145 | 290 | 290 | 435 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 1.9 | 0 | 1.0 | 1.0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | (2) | 0 | .8 | .4 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 10.7 | 11.2 | 8.9 | 9.4 | 7.3 | 8.0 | 9.1 | 8.9 | 7.1 | 9.4 | 10.7 | 10.7 |
| 7.1 | 7.7 | 5.5 | 6.8 | 4.4 | 5.3 | 4.6 | 5.7 | 4.8 | 4.8 | 4.8 | 6.0 |
| 3.1 | 2.5 | 2.4 | 1.9 | 1.9 | 1.9 | 2.9 | 1.9 | 1.1 | 2.9 | 1.8 | 2.5 |
| .5 | 1.0 | 1.0 | .8 | 1.1 | .8 | 1.6 | 1.3 | 1.2 | 1.7 | 4.1 | 2.2 |
| 1.5 | 1.5 | 1.6 | 1.4 | 1.6 | 1.5 | 1.7 | 1.6 | 1.5 | 1.8 | 2.2 | 1.7 |
| .7 | .9 | 1.0 | .8 | .8 | .8 | .9 | .9 | .9 | .9 | 1.4 | 1.1 |
| 5.0 | 6.0 | 6.0 | 6.0 | 5.0 | 5.0 | 5.0 | 5.0 | 5.0 | 4.0 | 6.0 | 8.0 |
| 75 | 69 | 87 | 68 | 85 | 79 | 88 | 78 | 79 | 88 | 103 | 78 |
| 47 | 50 | 76 | 43 | 56 | 53 | 52 | 52 | 73 | 59 | 90 | 49 |
| 30 | 17 | 24 | 24 | 33 | 30 | 30 | 24 | 24 | 30 | 8 | 10 |
| 48 | 48 | 48 | 48 | 48 | 48 | 48 | 48 | 48 | 48 | 48 | 48 |
| 48 | 48 | 60 | 48 | 60 | 56 | 67 | 48 | 48 | 67 | 73 | 48 |
| 97 | 97 | 97 | 97 | 97 | 97 | 120 | 97 | 97 | 108 | 102 | 98 |
| 145 | 145 | 145 | 120 | 153 | 145 | 157 | 145 | 145 | 145 | 290 | 145 |
| 168 | 152 | 196 | 167 | 180 | 150 | 180 | 193 | 277 | 193 | 337 | 178 |
| 205 | 242 | 420 | 205 | 290 | 242 | 242 | 242 | 360 | 338 | 362 | 180 |
| 155 | 134 | 149 | 145 | 158 | 159 | 160 | 146 | 133 | 141 | 137 | 133 |
| 68 | 66 | 53 | 63 | 54 | 77 | 95 | 60 | 42 | 44 | 58 | 52 |
| 164 | 156 | 182 | 143 | 171 | 172 | 150 | 159 | 137 | 165 | 121 | 143 |
| 53 | 60 | 119 | 36 | 78 | 89 | 40 | 50 | 22 | 57 | 29 | 31 |
| 184 | 167 | 209 | 169 | 167 | 161 | 160 | 166 | 207 | 166 | 178 | 151 |
| 73 | 56 | 98 | 43 | 51 | 37 | 46 | 53 | 95 | 66 | 103 | 41 |
| 580 | 580 | 1,160 | 580 | 540 | 1,080 | 900 | 456 | 540 | 435 | 505 | 300 |
| 1,160 | 1,085 | 1,595 | 870 | 900 | 1,440 | 1,015 | 905 | 1,080 | 1,080 | 1,085 | 800 |
| 150 | 137 | 158 | 142 | 157 | 156 | 151 | 145 | 155 | 150 | 141 | 136 |
| 51 | 49 | 84 | 51 | 54 | 72 | 57 | 52 | 74 | 51 | 60 | 41 |
| 73 | 50 | 73 | 73 | 84 | 84 | 72 | 73 | 74 | 90 | 73 | 73 |
| 145 | 145 | 145 | 145 | 145 | 145 | 145 | 145 | 145 | 145 | 90 | 145 |
| 145 | 145 | 145 | 145 | 145 | 145 | 145 | 145 | 145 | 145 | 145 | 145 |
| 145 | 145 | 145 | 145 | 145 | 145 | 180 | 145 | 145 | 145 | 180 | 145 |
| 180 | 180 | 180 | 180 | 218 | 180 | 180 | 180 | 180 | 180 | 180 | 180 |
| 228 | 180 | 290 | 180 | 290 | 290 | 200 | 228 | 360 | 290 | 290 | 180 |
| 293 | 290 | 435 | 290 | 360 | 360 | 360 | 290 | 435 | 360 | 360 | 228 |
| 580 | 540 | 1,160 | 580 | 390 | 1,080 | 900 | 456 | 435 | 435 | 360 | 290 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 4.04.--ORANGE JUICE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 35.5 | 20.9 | 40.9 | 41.7 | 43.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 13.9 | 9.8 | 14.3 | 17.1 | 17.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 9.3 | 5.6 | 10.5 | 11.5 | 11.8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 12.4 | 5.5 | 16.2 | 13.1 | 14.6 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER.... | 2.1 | 2.0 | 2.4 | 2.1 | 2.1 |
| STANDARD DEVIATION.....DO..... | 1.1 | 1.3 | 1.4 | 1.2 | 1.1 |
| MAXIMUM.....DO..... | 12.0 | 8.0 | 11.0 | 10.0 | 8.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 133 | 81 | 121 | 119 | 124 |
| STANDARD DEVIATION.....GM..... | 92 | 61 | 94 | 82 | 87 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 42 | 21 | 28 | 42 | 42 |
| 25TH.....GM..... | 73 | 42 | 42 | 62 | 62 |
| 50TH.....GM..... | 125 | 73 | 104 | 93 | 104 |
| 75TH.....GM..... | 166 | 104 | 166 | 166 | 166 |
| 90TH.....GM..... | 249 | 145 | 228 | 208 | 228 |
| 95TH.....GM..... | 290 | 167 | 291 | 249 | 249 |
| 99TH.....GM..... | 496 | 332 | 415 | 415 | 415 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 207 | 116 | 141 | 187 | 188 |
| STANDARD DEVIATION.....GM..... | 107 | 49 | 72 | 82 | 77 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 209 | 144 | 174 | 185 | 187 |
| STANDARD DEVIATION.....GM..... | 98 | 49 | 84 | 90 | 73 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 201 | 142 | 189 | 190 | 197 |
| STANDARD DEVIATION.....GM..... | 102 | 74 | 101 | 89 | 100 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,952 | 498 | 996 | 1,058 | 1,245 |
| MAXIMUM OVER 3 DAYS.....GM..... | 3,237 | 1,370 | 2,739 | 1,994 | 3,237 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 190 | 122 | 153 | 167 | 178 |
| STANDARD DEVIATION.....GM..... | 84 | 51 | 70 | 73 | 68 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 95 | 62 | 62 | 93 | 95 |
| 25TH.....GM..... | 125 | 93 | 125 | 125 | 125 |
| 50TH.....GM..... | 187 | 125 | 125 | 125 | 187 |
| 75TH.....GM..... | 249 | 131 | 187 | 187 | 248 |
| 90TH.....GM..... | 249 | 187 | 249 | 249 | 249 |
| 95TH.....GM..... | 311 | 249 | 249 | 249 | 249 |
| 99TH.....GM..... | 498 | 250 | 374 | 498 | 374 |
| MAXIMUM.....GM..... | 1,240 | 250 | 500 | 747 | 747 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER.... | 126.0 | 0 | 2.9 | 8.3 | 4.8 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | .3 | .5 | .3 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 39.4 | 41.0 | 37.3 | 36.6 | 32.9 | 31.8 | 32.0 | 35.1 | 32.7 | 40.4 | 28.6 | 36.7 |
| 16.7 | 16.4 | 15.2 | 18.3 | 14.8 | 13.8 | 12.1 | 12.7 | 9.5 | 11.0 | 8.0 | 10.2 |
| 9.7 | 11.8 | 10.0 | 9.4 | 8.7 | 9.0 | 8.0 | 9.4 | 7.9 | 8.5 | 5.2 | 9.2 |
| 13.0 | 12.8 | 12.0 | 8.9 | 9.4 | 9.1 | 11.9 | 13.0 | 15.3 | 20.9 | 15.4 | 17.2 |
| 2.1 | 2.1 | 2.0 | 1.9 | 2.0 | 2.0 | 2.1 | 2.1 | 2.3 | 2.4 | 2.3 | 2.3 |
| 1.1 | 1.1 | 1.0 | 1.0 | 1.0 | 1.2 | 1.0 | 1.1 | 1.1 | 1.1 | .9 | 1.0 |
| 8.0 | 10.0 | 7.0 | 7.0 | 7.0 | 12.0 | 12.0 | 12.0 | 9.0 | 9.0 | 5.0 | 8.0 |
| 134 | 129 | 155 | 129 | 148 | 139 | 133 | 128 | 134 | 143 | 141 | 129 |
| 87 | 84 | 108 | 82 | 112 | 106 | 86 | 84 | 85 | 81 | 72 | 87 |
| 42 | 42 | 42 | 42 | 42 | 42 | 42 | 42 | 42 | 42 | 42 | 42 |
| 83 | 63 | 83 | 62 | 83 | 83 | 83 | 72 | 83 | 83 | 83 | 83 |
| 124 | 124 | 125 | 104 | 125 | 125 | 125 | 124 | 125 | 125 | 125 | 125 |
| 167 | 166 | 187 | 166 | 187 | 166 | 166 | 166 | 166 | 187 | 187 | 166 |
| 249 | 249 | 249 | 249 | 249 | 249 | 249 | 228 | 248 | 249 | 249 | 208 |
| 280 | 270 | 332 | 280 | 332 | 331 | 270 | 249 | 249 | 291 | 249 | 249 |
| 436 | 415 | 540 | 372 | 498 | 498 | 415 | 435 | 498 | 415 | 331 | 457 |
| 222 | 202 | 256 | 218 | 238 | 218 | 213 | 197 | 185 | 185 | 192 | 166 |
| 117 | 97 | 150 | 101 | 127 | 103 | 134 | 85 | 72 | 59 | 94 | 63 |
| 209 | 206 | 246 | 232 | 257 | 226 | 217 | 186 | 183 | 195 | 248 | 182 |
| 89 | 87 | 147 | 92 | 128 | 101 | 95 | 68 | 71 | 88 | 112 | 101 |
| 206 | 202 | 236 | 216 | 234 | 229 | 189 | 192 | 185 | 190 | 173 | 178 |
| 93 | 87 | 111 | 81 | 138 | 136 | 89 | 92 | 90 | 73 | 55 | 89 |
| 1,245 | 996 | 1,992 | 992 | 1,494 | 1,183 | 1,984 | 1,183 | 996 | 996 | 498 | 996 |
| 2,241 | 2,054 | 2,988 | 1,991 | 3,113 | 3,175 | 2,988 | 2,771 | 2,241 | 1,494 | 996 | 2,241 |
| 195 | 188 | 228 | 208 | 227 | 207 | 191 | 181 | 173 | 179 | 180 | 168 |
| 80 | 77 | 116 | 81 | 122 | 86 | 80 | 67 | 63 | 63 | 71 | 65 |
| 124 | 124 | 124 | 124 | 124 | 124 | 95 | 124 | 93 | 124 | 93 | 95 |
| 125 | 125 | 155 | 125 | 154 | 125 | 125 | 125 | 125 | 125 | 125 | 125 |
| 187 | 187 | 248 | 187 | 187 | 187 | 187 | 187 | 156 | 186 | 186 | 125 |
| 249 | 249 | 249 | 249 | 249 | 249 | 249 | 248 | 248 | 248 | 248 | 187 |
| 249 | 249 | 311 | 249 | 374 | 249 | 249 | 249 | 249 | 249 | 249 | 249 |
| 311 | 249 | 498 | 374 | 498 | 374 | 311 | 249 | 249 | 249 | 249 | 249 |
| 498 | 498 | 747 | 498 | 750 | 498 | 498 | 496 | 374 | 374 | 496 | 436 |
| 747 | 747 | 1,240 | 747 | 1,000 | 996 | 996 | 623 | 498 | 498 | 498 | 498 |
| 8.1 | 4.7 | 12.0 | 5.5 | 14.7 | 15.1 | 20.2 | 13.4 | 6.8 | 4.5 | 1.6 | 3.5 |
| .5 | .3 | 1.1 | .6 | .6 | .4 | .6 | .3 | .8 | .3 | .5 | .4 |

TABLE 4.05.--RAW APPLES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 18.2 | 1.7 | 23.6 | 23.8 | 25.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 11.3 | 1.3 | 14.9 | 16.8 | 16.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 4.8 | .3 | 6.0 | 5.8 | 6.8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.2 | 0 | 2.6 | 1.2 | 3.0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.6 | 1.2 | 1.6 | 1.4 | 1.6 |
| STANDARD DEVIATION.....DO..... | .9 | .4 | .9 | .7 | .9 |
| MAXIMUM.....DO..... | 8.0 | 2.0 | 6.0 | 6.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 75 | 37 | 56 | 59 | 71 |
| STANDARD DEVIATION.....GM..... | 53 | 31 | 42 | 37 | 45 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 23 | | 12 | 23 | 23 |
| 25TH.....GM..... | 46 | | 23 | 35 | 46 |
| 50TH.....GM..... | 46 | | 46 | 46 | 46 |
| 75TH.....GM..... | 92 | | 78 | 71 | 92 |
| 90TH.....GM..... | 138 | | 92 | 92 | 138 |
| 95TH.....GM..... | 184 | | 138 | 138 | 141 |
| 99TH.....GM..... | 276 | | 184 | 184 | 230 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 142 | 73 | 107 | 129 | 134 |
| STANDARD DEVIATION.....GM..... | 58 | 50 | 58 | 51 | 58 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 149 | 138 | 115 | 133 | 147 |
| STANDARD DEVIATION.....GM..... | 52 | 0 | 44 | 48 | 48 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 166 | 0 | 123 | 142 | 150 |
| STANDARD DEVIATION.....GM..... | 71 | 0 | 64 | 56 | 39 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,272 | 138 | 848 | 552 | 700 |
| MAXIMUM OVER 3 DAYS.....GM..... | 3,816 | 276 | 1,272 | 966 | 976 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 141 | 94 | 105 | 124 | 132 |
| STANDARD DEVIATION.....GM..... | 49 | 51 | 44 | 39 | 41 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 69 | | 35 | 69 | 69 |
| 25TH.....GM..... | 138 | | 69 | 106 | 106 |
| 50TH.....GM..... | 138 | | 106 | 138 | 138 |
| 75TH.....GM..... | 138 | | 138 | 138 | 138 |
| 90TH.....GM..... | 212 | | 138 | 138 | 138 |
| 95TH.....GM..... | 212 | | 138 | 212 | 212 |
| 99TH.....GM..... | 276 | | 212 | 212 | 276 |
| MAXIMUM.....GM..... | 636 | 138 | 276 | 424 | 530 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 4.0 | 0 | 0 | 0 | 1.7 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | (2) | 0 | 0 | 0 | .2 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 22.0 | 24.5 | 16.7 | 19.1 | 15.0 | 17.0 | 19.2 | 16.8 | 13.0 | 16.3 | 14.0 | 14.0 |
| 13.7 | 16.4 | 10.4 | 12.7 | 8.5 | 11.2 | 10.6 | 9.9 | 6.7 | 11.3 | 7.7 | 7.0 |
| 5.8 | 5.4 | 4.3 | 4.8 | 4.3 | 4.5 | 5.3 | 4.9 | 3.1 | 3.1 | 3.1 | 4.0 |
| 2.6 | 2.7 | 2.0 | 1.6 | 2.1 | 1.3 | 3.2 | 1.9 | 3.1 | 1.9 | 3.1 | 3.0 |
| 1.6 | 1.5 | 1.6 | 1.5 | 1.7 | 1.5 | 1.7 | 1.6 | 1.8 | 1.5 | 1.8 | 1.9 |
| .9 | .8 | 1.0 | .8 | .9 | .8 | 1.0 | .9 | 1.0 | .9 | 1.0 | .9 |
| 8.0 | 6.0 | 8.0 | 5.0 | 7.0 | 7.0 | 8.0 | 7.0 | 6.0 | 6.0 | 6.0 | 6.0 |
| 78 | 71 | 83 | 71 | 85 | 71 | 86 | 75 | 90 | 70 | 97 | 81 |
| 58 | 45 | 55 | 45 | 54 | 45 | 61 | 44 | 62 | 48 | 160 | 53 |
| 23 | 35 | 26 | 26 | 35 | 23 | 35 | 26 | 18 | 23 | 10 | 21 |
| 46 | 46 | 46 | 46 | 46 | 46 | 46 | 46 | 46 | 46 | 46 | 46 |
| 46 | 46 | 71 | 46 | 71 | 46 | 58 | 71 | 46 | 46 | 46 | 69 |
| 92 | 92 | 92 | 92 | 92 | 92 | 92 | 92 | 138 | 92 | 106 | 92 |
| 138 | 138 | 141 | 138 | 141 | 138 | 141 | 138 | 163 | 138 | 141 | 141 |
| 184 | 141 | 212 | 141 | 212 | 141 | 209 | 163 | 212 | 187 | 212 | 212 |
| 353 | 230 | 283 | 230 | 276 | 258 | 325 | 212 | 279 | 230 | 1,273 | 230 |
| 145 | 144 | 155 | 144 | 157 | 146 | 149 | 144 | 137 | 135 | 126 | 133 |
| 62 | 52 | 68 | 47 | 68 | 66 | 50 | 55 | 60 | 47 | 49 | 53 |
| 152 | 150 | 168 | 155 | 157 | 148 | 156 | 148 | 156 | 154 | 137 | 136 |
| 45 | 42 | 58 | 59 | 55 | 48 | 70 | 42 | 46 | 42 | 42 | 41 |
| 186 | 157 | 182 | 158 | 173 | 166 | 178 | 157 | 171 | 163 | 237 | 154 |
| 81 | 53 | 55 | 48 | 55 | 54 | 64 | 39 | 55 | 57 | 292 | 54 |
| 690 | 636 | 690 | 828 | 742 | 848 | 1,272 | 636 | 424 | 424 | 1,272 | 276 |
| 1,380 | 966 | 1,104 | 966 | 966 | 1,008 | 1,484 | 966 | 976 | 828 | 3,816 | 828 |
| 146 | 140 | 151 | 142 | 153 | 140 | 150 | 140 | 147 | 137 | 164 | 131 |
| 55 | 41 | 48 | 46 | 52 | 45 | 50 | 44 | 48 | 47 | 121 | 50 |
| 69 | 78 | 106 | 69 | 106 | 69 | 106 | 69 | 69 | 63 | 69 | 63 |
| 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 | 106 |
| 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 |
| 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 | 138 |
| 212 | 207 | 212 | 212 | 212 | 212 | 212 | 212 | 212 | 212 | 212 | 212 |
| 212 | 212 | 212 | 212 | 276 | 212 | 212 | 212 | 212 | 212 | 636 | 212 |
| 414 | 276 | 276 | 276 | 276 | 276 | 312 | 276 | 276 | 276 | 636 | 276 |
| 636 | 552 | 424 | 552 | 530 | 424 | 636 | 636 | 276 | 424 | 636 | 276 |
| 0 | 0 | 0 | 0 | 2.3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | .2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 4.06.--APPLESAUCE AND COOKED APPLES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 9.8 | 35.6 | 13.6 | 10.4 | 14.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 7.6 | 18.5 | 8.3 | 8.1 | 11.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.7 | 11.6 | 3.9 | 1.8 | 2.5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .5 | 5.5 | 1.4 | .5 | .3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.4 | 2.2 | 1.7 | 1.3 | 1.2 |
| STANDARD DEVIATION.....CO..... | .7 | 1.5 | 1.1 | .7 | .5 |
| MAXIMUM.....DO..... | 9.0 | 9.0 | 8.0 | 5.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 61 | 52 | 59 | 56 | 55 |
| STANDARD DEVIATION.....GM..... | 47 | 39 | 49 | 35 | 37 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 11 | 9 | 9 | 11 | 16 |
| 25TH.....GM..... | 38 | 23 | 21 | 41 | 32 |
| 50TH.....GM..... | 43 | 43 | 43 | 43 | 43 |
| 75TH.....GM..... | 85 | 66 | 85 | 85 | 83 |
| 90TH.....GM..... | 127 | 99 | 128 | 85 | 85 |
| 95TH.....GM..... | 160 | 135 | 170 | 128 | 128 |
| 99TH.....GM..... | 253 | 209 | 255 | 213 | 170 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 146 | 102 | 109 | 135 | 136 |
| STANDARD DEVIATION.....GM..... | 95 | 68 | 79 | 75 | 80 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 139 | 95 | 114 | 125 | 130 |
| STANDARD DEVIATION.....GM..... | 78 | 51 | 63 | 41 | 68 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 137 | 89 | 141 | 139 | 140 |
| STANDARD DEVIATION.....GM..... | 88 | 52 | 62 | 45 | 30 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,016 | 510 | 510 | 383 | 510 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,034 | 638 | 797 | 638 | 638 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 134 | 71 | 104 | 126 | 132 |
| STANDARD DEVIATION.....GM..... | 86 | 49 | 65 | 61 | 76 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 28 | 14 | 16 | 32 | 32 |
| 25TH.....GM..... | 64 | 42 | 57 | 65 | 64 |
| 50TH.....GM..... | 128 | 57 | 97 | 128 | 128 |
| 75TH.....GM..... | 130 | 92 | 128 | 128 | 128 |
| 90TH.....GM..... | 255 | 128 | 213 | 255 | 255 |
| 95TH.....GM..... | 255 | 135 | 255 | 255 | 255 |
| 99TH.....GM..... | 488 | 255 | 255 | 255 | 480 |
| MAXIMUM.....GM..... | 765 | 510 | 255 | 383 | 510 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 32.6 | .6 | 2.9 | 0 | 6.2 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | .2 | 1.2 | 0 | 1.9 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 13.6 | 11.1 | 10.2 | 7.7 | 5.1 | 5.8 | 8.1 | 9.5 | 10.7 | 11.5 | 15.1 | 17.4 |
| 11.8 | 9.2 | 8.5 | 6.7 | 4.3 | 5.1 | 6.8 | 7.5 | 7.5 | 7.9 | 10.5 | 9.2 |
| 1.5 | 1.7 | 1.5 | .9 | .8 | .6 | 1.2 | 1.7 | 2.5 | 2.9 | 2.2 | 5.3 |
| .2 | .3 | .2 | .1 | (2) | .1 | .2 | .4 | .7 | .7 | 2.4 | 2.9 |
| 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.3 | 1.4 | 1.5 | 1.7 | 1.8 |
| .4 | .5 | .5 | .4 | .4 | .5 | .5 | .6 | .6 | .7 | .9 | 1.0 |
| 3.0 | 4.0 | 4.0 | 4.0 | 3.0 | 4.0 | 3.0 | 5.0 | 4.0 | 4.0 | 4.0 | 6.0 |
| 58 | 55 | 69 | 57 | 68 | 51 | 65 | 59 | 56 | 62 | 89 | 85 |
| 53 | 42 | 73 | 33 | 46 | 40 | 45 | 47 | 30 | 41 | 67 | 57 |
| 10 | 11 | 15 | 16 | 21 | 10 | 11 | 10 | 11 | 11 | 21 | 16 |
| 31 | 28 | 32 | 41 | 43 | 21 | 41 | 32 | 40 | 41 | 40 | 43 |
| 43 | 43 | 43 | 43 | 53 | 43 | 43 | 43 | 43 | 43 | 85 | 84 |
| 83 | 80 | 85 | 85 | 85 | 71 | 85 | 85 | 85 | 85 | 127 | 112 |
| 128 | 85 | 149 | 85 | 128 | 85 | 128 | 127 | 85 | 127 | 202 | 170 |
| 128 | 138 | 170 | 128 | 169 | 128 | 166 | 168 | 95 | 129 | 213 | 198 |
| 340 | 213 | 255 | 170 | 339 | 170 | 255 | 254 | 128 | 170 | 340 | 255 |
| 142 | 137 | 157 | 158 | 184 | 129 | 165 | 142 | 147 | 139 | 207 | 170 |
| 92 | 86 | 110 | 86 | 130 | 82 | 99 | 97 | 74 | 84 | 130 | 90 |
| 187 | 130 | 184 | 129 | 154 | 156 | 179 | 135 | 108 | 132 | 136 | 157 |
| 129 | 78 | 90 | 54 | 67 | 56 | 94 | 73 | 56 | 52 | 54 | 88 |
| 222 | 158 | 376 | 150 | 127 | 146 | 72 | 153 | 77 | 131 | 174 | 141 |
| 106 | 80 | 235 | 0 | 0 | 121 | 47 | 92 | 23 | 56 | 98 | 63 |
| 765 | 510 | 1,016 | 383 | 1,016 | 510 | 638 | 638 | 255 | 454 | 636 | 510 |
| 1,275 | 765 | 2,034 | 510 | 1,016 | 893 | 893 | 1,020 | 510 | 760 | 1,020 | 765 |
| 151 | 134 | 171 | 146 | 171 | 130 | 158 | 134 | 121 | 128 | 161 | 141 |
| 107 | 82 | 125 | 73 | 98 | 79 | 92 | 84 | 71 | 67 | 89 | 80 |
| 32 | 26 | 48 | 48 | 64 | 16 | 31 | 23 | 21 | 32 | 30 | 26 |
| 84 | 64 | 96 | 122 | 128 | 64 | 120 | 64 | 64 | 65 | 113 | 90 |
| 128 | 128 | 128 | 128 | 128 | 128 | 128 | 128 | 128 | 128 | 128 | 128 |
| 191 | 128 | 244 | 191 | 199 | 128 | 255 | 130 | 128 | 128 | 255 | 191 |
| 255 | 255 | 255 | 255 | 255 | 255 | 255 | 255 | 255 | 255 | 255 | 255 |
| 255 | 255 | 510 | 255 | 383 | 255 | 255 | 255 | 255 | 255 | 255 | 255 |
| 512 | 383 | 510 | 383 | 508 | 284 | 510 | 383 | 255 | 255 | 383 | 383 |
| 765 | 510 | 762 | 383 | 510 | 510 | 510 | 610 | 255 | 386 | 383 | 383 |
| 2.6 | 1.5 | 1.0 | 0 | 3.7 | 3.0 | 3.1 | 2.4 | .9 | 0 | 0 | 4.6 |
| .8 | .5 | .6 | 0 | 1.5 | .8 | .6 | .3 | .6 | 0 | 0 | 1.5 |

TABLE 4.07.--APPLEJUICE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.8 | 19.2 | 13.1 | 8.5 | 5.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.2 | 9.3 | 5.1 | 3.8 | 3.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .9 | 4.5 | 4.6 | 2.8 | 1.6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .6 | 5.4 | 3.5 | 2.0 | .9 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER.... | 1.8 | 2.2 | 2.6 | 2.3 | 1.8 |
| STANDARD DEVIATION.....DO..... | 1.4 | 1.5 | 2.4 | 1.8 | 1.1 |
| MAXIMUM.....DO..... | 15.0 | 8.0 | 15.0 | 11.0 | 7.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 116 | 91 | 128 | 132 | 115 |
| STANDARD DEVIATION.....GM..... | 98 | 71 | 117 | 141 | 88 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 41 | 21 | 31 | 41 | 41 |
| 25TH.....GM..... | 62 | 42 | 62 | 62 | 62 |
| 50TH.....GM..... | 83 | 78 | 103 | 83 | 83 |
| 75TH.....GM..... | 145 | 124 | 165 | 155 | 145 |
| 90TH.....GM..... | 227 | 176 | 227 | 238 | 207 |
| 95TH.....GM..... | 289 | 248 | 372 | 331 | 289 |
| 99TH.....GM..... | 496 | 331 | 651 | 889 | 516 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 219 | 128 | 165 | 175 | 201 |
| STANDARD DEVIATION.....GM..... | 130 | 50 | 81 | 81 | 91 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 223 | 164 | 200 | 193 | 209 |
| STANDARD DEVIATION.....GM..... | 113 | 90 | 104 | 71 | 83 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 223 | 157 | 230 | 279 | 239 |
| STANDARD DEVIATION.....GM..... | 152 | 76 | 162 | 223 | 128 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,302 | 496 | 868 | 1,302 | 1,178 |
| MAXIMUM OVER 3 DAYS.....GM..... | 3,224 | 1,240 | 2,356 | 3,224 | 1,550 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 191 | 125 | 148 | 170 | 193 |
| STANDARD DEVIATION.....GM..... | 101 | 56 | 64 | 65 | 87 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 63 | 47 | 62 | 93 | 124 |
| 25TH.....GM..... | 124 | 93 | 124 | 124 | 124 |
| 50TH.....GM..... | 186 | 124 | 124 | 155 | 186 |
| 75TH.....GM..... | 248 | 131 | 186 | 186 | 248 |
| 90TH.....GM..... | 248 | 217 | 248 | 248 | 248 |
| 95TH.....GM..... | 372 | 248 | 248 | 248 | 248 |
| 99TH.....GM..... | 496 | 250 | 250 | 372 | 496 |
| MAXIMUM.....GM..... | 992 | 250 | 496 | 496 | 992 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER.... | 14.4 | 0 | 2.9 | 3.7 | 1.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | 0 | .8 | 1.1 | .9 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.0 | 4.0 | 2.7 | 3.1 | 2.4 | 3.7 | 2.3 | 2.7 | 1.6 | 2.8 | 3.8 | 3.1 |
| 2.0 | 2.8 | 1.9 | 2.4 | 1.7 | 2.6 | 1.4 | 1.7 | .9 | 1.3 | 2.0 | 1.5 |
| .9 | .8 | .5 | .7 | .5 | .7 | .6 | .6 | .4 | .6 | 1.0 | .6 |
| .1 | .3 | .4 | .1 | .2 | .3 | .4 | .4 | .3 | .9 | .8 | 1.1 |
| 1.6 | 1.5 | 1.6 | 1.4 | 1.4 | 1.5 | 1.7 | 1.7 | 1.8 | 2.0 | 1.7 | 1.9 |
| .8 | .7 | .8 | .9 | .8 | .9 | .9 | 1.2 | 1.1 | 1.2 | .8 | 1.0 |
| 5.0 | 5.0 | 3.0 | 5.0 | 4.0 | 7.0 | 4.0 | 10.0 | 4.0 | 5.0 | 3.0 | 4.0 |
| 98 | 99 | 138 | 112 | 126 | 118 | 120 | 118 | 131 | 102 | 110 | 97 |
| 63 | 67 | 124 | 88 | 128 | 86 | 76 | 89 | 124 | 60 | 68 | 46 |
| 41 | 41 | 41 | 41 | 41 | 41 | 52 | 41 | 31 | 41 | 41 | 41 |
| 62 | 62 | 62 | 62 | 62 | 62 | 83 | 62 | 41 | 62 | 41 | 41 |
| 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | 62 | 103 | 93 | 83 |
| 114 | 124 | 165 | 103 | 145 | 155 | 165 | 165 | 186 | 124 | 145 | 124 |
| 165 | 186 | 227 | 248 | 289 | 248 | 186 | 207 | 413 | 186 | 227 | 165 |
| 181 | 248 | 434 | 331 | 331 | 310 | 248 | 248 | 413 | 248 | 248 | 165 |
| 393 | 372 | 662 | 413 | 662 | 413 | 413 | 569 | 413 | 310 | 248 | 186 |
| 219 | 211 | 267 | 264 | 237 | 249 | 260 | 244 | 125 | 197 | 189 | 167 |
| 90 | 98 | 124 | 183 | 159 | 152 | 121 | 148 | 25 | 87 | 113 | 55 |
| 199 | 229 | 371 | 289 | 291 | 274 | 198 | 228 | 393 | 178 | 250 | 194 |
| 87 | 94 | 198 | 172 | 164 | 119 | 58 | 89 | 199 | 39 | 75 | 58 |
| 237 | 243 | 241 | 118 | 419 | 263 | 213 | 223 | 226 | 146 | 154 | 136 |
| 129 | 65 | 207 | 92 | 228 | 92 | 113 | 136 | 30 | 76 | 59 | 22 |
| 496 | 558 | 992 | 992 | 992 | 992 | 744 | 992 | 744 | 496 | 496 | 248 |
| 1,178 | 1,116 | 1,984 | 1,240 | 1,984 | 1,488 | 1,240 | 1,736 | 1,240 | 930 | 744 | 558 |
| 190 | 204 | 259 | 236 | 262 | 233 | 216 | 211 | 215 | 153 | 196 | 152 |
| 69 | 74 | 180 | 139 | 152 | 114 | 94 | 102 | 91 | 81 | 100 | 49 |
| 124 | 124 | 124 | 41 | 124 | 124 | 94 | 124 | 93 | 30 | 93 | 124 |
| 124 | 155 | 186 | 155 | 186 | 186 | 124 | 124 | 124 | 124 | 124 | 124 |
| 186 | 186 | 186 | 248 | 248 | 248 | 186 | 186 | 248 | 186 | 186 | 124 |
| 248 | 248 | 310 | 248 | 248 | 248 | 248 | 248 | 248 | 186 | 248 | 186 |
| 248 | 248 | 496 | 372 | 496 | 372 | 372 | 248 | 248 | 248 | 248 | 248 |
| 248 | 372 | 496 | 496 | 496 | 496 | 372 | 310 | 496 | 248 | 434 | 248 |
| 496 | 496 | 992 | 992 | 992 | 744 | 496 | 558 | 496 | 496 | 496 | 248 |
| 496 | 496 | 992 | 992 | 992 | 992 | 620 | 992 | 496 | 496 | 496 | 248 |
| 1.1 | .9 | 0 | 0 | 0 | 0 | 2.2 | 0 | 0 | 2.0 | 0 | 0 |
| 1.1 | .7 | 0 | 0 | 0 | 0 | 1.2 | 0 | 0 | 2.0 | 0 | 0 |

TABLE 4.08.--BANANAS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1: | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 16.1 | 9.0 | 23.7 | 18.1 | 17.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 10.4 | 4.6 | 13.1 | 11.8 | 12.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.9 | 2.0 | 7.9 | 4.8 | 4.7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.9 | 2.4 | 2.8 | 1.4 | .6 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.6 | 2.0 | 1.8 | 1.5 | 1.4 |
| STANDARD DEVIATION.....DO..... | .9 | 1.3 | 1.0 | .8 | .6 |
| MAXIMUM.....DO..... | 7.0 | 7.0 | 6.0 | 5.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 55 | 40 | 51 | 51 | 50 |
| STANDARD DEVIATION.....GM..... | 34 | 30 | 35 | 30 | 26 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 20 | | 15 | 20 | 20 |
| 25TH.....GM..... | 40 | | 25 | 32 | 40 |
| 50TH.....GM..... | 40 | | 40 | 40 | 40 |
| 75TH.....GM..... | 79 | | 65 | 71 | 63 |
| 90TH.....GM..... | 108 | | 96 | 91 | 79 |
| 95TH.....GM..... | 119 | | 119 | 119 | 99 |
| 99TH.....GM..... | 170 | | 170 | 136 | 119 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 113 | 63 | 90 | 107 | 110 |
| STANDARD DEVIATION.....GM..... | 44 | 34 | 38 | 46 | 32 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 110 | 76 | 102 | 105 | 111 |
| STANDARD DEVIATION.....GM..... | 37 | 35 | 49 | 35 | 28 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 115 | 67 | 99 | 109 | 127 |
| STANDARD DEVIATION.....GM..... | 40 | 35 | 38 | 28 | 24 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 544 | 179 | 357 | 276 | 272 |
| MAXIMUM OVER 3 DAYS.....GM..... | 816 | 476 | 714 | 476 | 595 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 106 | 61 | 86 | 99 | 109 |
| STANDARD DEVIATION.....GM..... | 37 | 31 | 34 | 34 | 30 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 50 | | 30 | 48 | 56 |
| 25TH.....GM..... | 95 | | 60 | 60 | 95 |
| 50TH.....GM..... | 119 | | 95 | 119 | 119 |
| 75TH.....GM..... | 119 | | 119 | 119 | 119 |
| 90TH.....GM..... | 136 | | 119 | 119 | 136 |
| 95TH.....GM..... | 136 | | 119 | 136 | 136 |
| 99TH.....GM..... | 238 | | 136 | 238 | 224 |
| MAXIMUM.....GM..... | 476 | 136 | 272 | 276 | 238 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 32.4 | .9 | 0 | .9 | 5.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 1.1 | 0 | .2 | 1.3 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 14.6 | 14.6 | 12.8 | 12.4 | 11.7 | 11.1 | 17.9 | 15.8 | 24.4 | 26.4 | 28.7 | 26.1 |
| 10.5 | 10.4 | 9.1 | 9.3 | 7.7 | 8.0 | 10.8 | 10.3 | 13.5 | 15.7 | 15.0 | 12.6 |
| 3.4 | 3.2 | 2.8 | 2.5 | 2.6 | 2.1 | 4.4 | 3.8 | 5.6 | 5.9 | 9.1 | 6.7 |
| .6 | 1.0 | .9 | .6 | 1.4 | .9 | 2.7 | 1.7 | 5.2 | 4.8 | 4.6 | 6.8 |
| 1.4 | 1.4 | 1.4 | 1.3 | 1.6 | 1.4 | 1.6 | 1.6 | 1.7 | 1.7 | 1.7 | 1.9 |
| .7 | .6 | .7 | .6 | .9 | .7 | .9 | .9 | .9 | .9 | .8 | 1.0 |
| 5.0 | 4.0 | 5.0 | 4.0 | 6.0 | 5.0 | 6.0 | 7.0 | 6.0 | 5.0 | 4.0 | 6.0 |
| 53 | 52 | 57 | 48 | 65 | 50 | 62 | 53 | 61 | 57 | 57 | 62 |
| 30 | 27 | 35 | 26 | 43 | 31 | 36 | 34 | 36 | 35 | 36 | 38 |
| 20 | 20 | 20 | 13 | 32 | 20 | 20 | 20 | 20 | 20 | 20 | 20 |
| 40 | 40 | 40 | 40 | 40 | 40 | 40 | 40 | 40 | 40 | 40 | 40 |
| 40 | 40 | 40 | 40 | 40 | 40 | 45 | 40 | 45 | 40 | 40 | 40 |
| 79 | 71 | 79 | 48 | 79 | 60 | 79 | 65 | 79 | 79 | 79 | 79 |
| 79 | 79 | 119 | 79 | 119 | 85 | 119 | 95 | 119 | 103 | 95 | 119 |
| 119 | 119 | 136 | 95 | 136 | 119 | 125 | 119 | 119 | 119 | 119 | 127 |
| 159 | 125 | 181 | 159 | 238 | 159 | 181 | 182 | 198 | 198 | 136 | 159 |
| 119 | 121 | 124 | 110 | 131 | 110 | 121 | 109 | 113 | 108 | 103 | 105 |
| 44 | 46 | 62 | 36 | 56 | 37 | 51 | 39 | 37 | 36 | 38 | 34 |
| 118 | 110 | 130 | 108 | 124 | 103 | 116 | 110 | 105 | 102 | 104 | 98 |
| 39 | 34 | 38 | 33 | 35 | 36 | 37 | 38 | 25 | 28 | 30 | 37 |
| 128 | 103 | 129 | 113 | 149 | 122 | 120 | 112 | 111 | 111 | 108 | 108 |
| 34 | 29 | 26 | 32 | 51 | 32 | 30 | 45 | 36 | 39 | 45 | 33 |
| 397 | 408 | 456 | 272 | 476 | 357 | 544 | 357 | 271 | 272 | 272 | 285 |
| 595 | 544 | 595 | 476 | 812 | 598 | 714 | 714 | 714 | 680 | 816 | 714 |
| 114 | 110 | 121 | 109 | 125 | 106 | 113 | 102 | 106 | 103 | 102 | 98 |
| 38 | 32 | 50 | 35 | 49 | 35 | 37 | 32 | 30 | 33 | 37 | 31 |
| 60 | 60 | 59 | 48 | 60 | 39 | 60 | 48 | 57 | 48 | 57 | 60 |
| 119 | 95 | 119 | 95 | 119 | 95 | 95 | 68 | 95 | 68 | 68 | 60 |
| 119 | 119 | 119 | 119 | 119 | 119 | 119 | 119 | 119 | 119 | 119 | 119 |
| 119 | 119 | 119 | 119 | 119 | 119 | 119 | 119 | 119 | 119 | 119 | 119 |
| 136 | 136 | 136 | 136 | 136 | 136 | 136 | 119 | 119 | 136 | 119 | 136 |
| 136 | 136 | 140 | 136 | 238 | 136 | 136 | 136 | 136 | 136 | 136 | 136 |
| 238 | 238 | 456 | 238 | 357 | 238 | 238 | 152 | 190 | 238 | 272 | 136 |
| 357 | 357 | 456 | 272 | 476 | 357 | 476 | 285 | 266 | 238 | 272 | 238 |
| 5.2 | 0 | 0 | 0 | 0 | 1.7 | 5.1 | 3.6 | 1.6 | 3.6 | 1.6 | 2.2 |
| 1.2 | 0 | 0 | 0 | 0 | .2 | .4 | .2 | .3 | .5 | .6 | .5 |

TABLE 4.09.--CANTALOUPE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.3 | .2 | 1.1 | 1.5 | 2.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.4 | .2 | .9 | 1.0 | 2.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .7 | 0 | .2 | .5 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.4 | 1.0 | 1.1 | 1.4 | 1.1 |
| STANDARD DEVIATION.....CO..... | .8 | .0 | .4 | .5 | .3 |
| MAXIMUM.....DO..... | 6.0 | 1.0 | 2.0 | 2.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 81 | 45 | 26 | 57 | 49 |
| STANDARD DEVIATION.....GM..... | 61 | 0 | 11 | 29 | 31 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 21 | | | | |
| 25TH.....GM..... | 45 | | | | |
| 50TH.....GM..... | 68 | | | | |
| 75TH.....GM..... | 91 | | | | |
| 90TH.....GM..... | 181 | | | | |
| 95TH.....GM..... | 181 | | | | |
| 99TH.....GM..... | 272 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 180 | 136 | 79 | 164 | 133 |
| STANDARD DEVIATION.....GM..... | 107 | 0 | 31 | 91 | 75 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 181 | 0 | 35 | 92 | 148 |
| STANDARD DEVIATION.....GM..... | 87 | 0 | 19 | 37 | 64 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 192 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 96 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,362 | 136 | 136 | 363 | 272 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,632 | 136 | 136 | 363 | 408 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 171 | 136 | 68 | 125 | 135 |
| STANDARD DEVIATION.....GM..... | 91 | 0 | 35 | 73 | 76 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 61 | | | | |
| 25TH.....GM..... | 136 | | | | |
| 50TH.....GM..... | 136 | | | | |
| 75TH.....GM..... | 272 | | | | |
| 90TH.....GM..... | 272 | | | | |
| 95TH.....GM..... | 272 | | | | |
| 99TH.....GM..... | 529 | | | | |
| MAXIMUM.....GM..... | 896 | 136 | 136 | 363 | 272 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 52.7 | 0 | 1.0 | 0 | 2.0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 3.0 | 0 | 7.3 | 0 | 4.5 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 2.2 | 2.5 | 2.0 | 2.5 | 2.5 | 2.2 | 3.9 | 5.4 | 5.3 | 6.2 | 3.9 | 3.3 |
| 1.9 | 1.9 | 1.8 | 1.8 | 2.2 | 1.7 | 2.8 | 3.8 | 3.8 | 3.6 | 2.5 | 2.4 |
| .3 | .6 | .2 | .7 | .3 | .4 | .9 | 1.1 | .9 | 1.8 | 1.0 | .3 |
| 0 | .1 | 0 | 0 | (2) | .1 | .2 | .5 | .6 | .8 | .3 | .5 |
| 1.2 | 1.3 | 1.1 | 1.4 | 1.1 | 1.3 | 1.4 | 1.5 | 1.5 | 1.7 | 1.5 | 1.5 |
| .4 | .7 | .3 | .6 | .4 | .6 | .7 | .9 | .8 | 1.0 | .6 | .7 |
| 2.0 | 4.0 | 2.0 | 3.0 | 3.0 | 3.0 | 4.0 | 6.0 | 4.0 | 5.0 | 3.0 | 3.0 |
| 63 | 68 | 77 | 89 | 74 | 71 | 87 | 84 | 97 | 98 | 123 | 77 |
| 46 | 44 | 59 | 68 | 38 | 50 | 72 | 65 | 59 | 60 | 87 | 42 |
| | 6 | | | 27 | 11 | 23 | 21 | 45 | 45 | | |
| | 45 | | | 45 | 45 | 45 | 45 | 45 | 45 | | |
| | 57 | | | 85 | 68 | 91 | 68 | 91 | 91 | | |
| | 91 | | | 91 | 91 | 91 | 91 | 91 | 112 | | |
| | 91 | | | 107 | 107 | 160 | 181 | 181 | 181 | | |
| | 107 | | | 136 | 181 | 181 | 227 | 213 | 181 | | |
| | 272 | | | 208 | 272 | 453 | 272 | 272 | 227 | | |
| 146 | 166 | 182 | 188 | 201 | 176 | 192 | 171 | 207 | 212 | 242 | 185 |
| 83 | 87 | 79 | 104 | 83 | 96 | 154 | 95 | 73 | 85 | 224 | 87 |
| 218 | 150 | 332 | 237 | 175 | 177 | 179 | 179 | 212 | 171 | 272 | 217 |
| 67 | 85 | 111 | 129 | 88 | 113 | 70 | 81 | 64 | 63 | 0 | 67 |
| 0 | 181 | 0 | 0 | 160 | 112 | 215 | 199 | 220 | 185 | 272 | 113 |
| 0 | 0 | 0 | 0 | 0 | 48 | 139 | 89 | 46 | 91 | 0 | 33 |
| 272 | 544 | 544 | 544 | 544 | 544 | 1,362 | 544 | 408 | 1,088 | 896 | 272 |
| 544 | 816 | 816 | 1,088 | 624 | 816 | 1,362 | 1,360 | 816 | 1,632 | 896 | 544 |
| 165 | 152 | 209 | 189 | 193 | 164 | 180 | 165 | 201 | 173 | 252 | 155 |
| 85 | 77 | 111 | 113 | 94 | 96 | 101 | 84 | 66 | 74 | 143 | 73 |
| | 32 | | | 68 | 32 | 68 | 40 | 136 | 80 | | |
| | 136 | | | 136 | 80 | 136 | 136 | 136 | 136 | | |
| | 136 | | | 160 | 136 | 136 | 136 | 160 | 136 | | |
| | 204 | | | 272 | 272 | 272 | 272 | 272 | 272 | | |
| | 272 | | | 272 | 272 | 272 | 272 | 272 | 272 | | |
| | 272 | | | 320 | 272 | 272 | 272 | 272 | 272 | | |
| | 272 | | | 544 | 544 | 681 | 320 | 272 | 408 | | |
| 272 | 272 | 544 | 544 | 544 | 544 | 681 | 544 | 272 | 544 | 896 | 272 |
| 3.1 | 1.8 | 2.2 | 0 | 0 | 10.1 | 2.9 | 17.9 | 1.0 | 8.1 | .7 | 1.8 |
| 5.9 | 2.5 | 7.2 | 0 | 0 | 6.6 | 1.1 | 3.1 | 1.2 | 4.4 | 2.4 | 3.6 |

TABLE 4.10.--FRUIT COCKTAIL
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.9 | 1.4 | 4.6 | 4.5 | 8.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.4 | 1.4 | 3.4 | 3.9 | 7.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | 0 | 1.0 | .5 | .6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | .1 | .1 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.2 | 1.0 | 1.4 | 1.2 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .0 | .7 | .5 | .4 |
| MAXIMUM.....DO..... | 4.0 | 1.0 | 4.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 62 | 32 | 58 | 53 | 57 |
| STANDARD DEVIATION.....GM..... | 41 | 14 | 41 | 30 | 33 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 11 | | | 14 | 21 |
| 25TH.....GM..... | 43 | | | 28 | 42 |
| 50TH.....GM..... | 43 | | | 43 | 43 |
| 75TH.....GM..... | 85 | | | 85 | 85 |
| 90TH.....GM..... | 85 | | | 85 | 85 |
| 95TH.....GM..... | 149 | | | 99 | 126 |
| 99TH.....GM..... | 184 | | | 149 | 170 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 163 | 96 | 130 | 139 | 157 |
| STANDARD DEVIATION.....GM..... | 93 | 43 | 69 | 74 | 84 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 158 | 0 | 149 | 142 | 155 |
| STANDARD DEVIATION.....GM..... | 75 | 0 | 71 | 34 | 52 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 180 | 0 | 170 | 127 | 150 |
| STANDARD DEVIATION.....GM..... | 82 | 0 | 0 | 0 | 21 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 701 | 128 | 510 | 319 | 510 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,403 | 128 | 606 | 446 | 510 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 159 | 96 | 125 | 135 | 152 |
| STANDARD DEVIATION.....GM..... | 88 | 43 | 61 | 66 | 76 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 32 | | | 63 | 63 |
| 25TH.....GM..... | 125 | | | 84 | 125 |
| 50TH.....GM..... | 128 | | | 128 | 128 |
| 75TH.....GM..... | 255 | | | 155 | 250 |
| 90TH.....GM..... | 255 | | | 255 | 255 |
| 95TH.....GM..... | 255 | | | 255 | 255 |
| 99TH.....GM..... | 510 | | | 279 | 319 |
| MAXIMUM.....GM..... | 510 | 128 | 255 | 279 | 510 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 10.8 | 0 | .9 | 1.6 | 3.4 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | 0 | 1.4 | 1.7 | 2.0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 7.2 | 7.5 | 5.2 | 3.8 | 1.9 | 2.6 | 3.3 | 3.1 | 5.5 | 3.2 | 2.6 | 3.9 |
| 6.5 | 6.7 | 4.9 | 3.5 | 1.7 | 2.4 | 2.8 | 2.8 | 3.7 | 2.5 | 1.7 | 3.4 |
| .6 | .7 | .3 | .3 | .2 | .2 | .4 | .3 | 1.4 | .6 | .6 | .3 |
| (2) | .1 | 0 | 0 | 0 | 0 | .1 | (2) | .4 | .2 | .3 | .2 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.1 | 1.4 | 1.3 | 1.4 | 1.2 |
| .3 | .4 | .2 | .3 | .4 | .3 | .5 | .4 | .7 | .6 | .7 | .5 |
| 3.0 | 3.0 | 2.0 | 2.0 | 3.0 | 2.0 | 4.0 | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| 56 | 60 | 61 | 59 | 76 | 58 | 73 | 55 | 81 | 71 | 86 | 61 |
| 30 | 35 | 35 | 38 | 57 | 31 | 44 | 33 | 56 | 72 | 83 | 41 |
| 21 | 21 | 21 | 10 | 21 | 11 | 21 | 10 | 16 | 5 | | |
| 43 | 43 | 43 | 42 | 43 | 42 | 43 | 42 | 43 | 30 | | |
| 43 | 43 | 43 | 43 | 64 | 43 | 74 | 43 | 64 | 57 | | |
| 85 | 85 | 85 | 85 | 85 | 85 | 85 | 85 | 85 | 85 | | |
| 85 | 85 | 85 | 85 | 170 | 85 | 128 | 85 | 170 | 135 | | |
| 128 | 128 | 128 | 170 | 170 | 93 | 170 | 85 | 191 | 170 | | |
| 170 | 170 | 170 | 170 | 276 | 170 | 255 | 170 | 255 | 468 | | |
| 160 | 161 | 179 | 164 | 203 | 168 | 184 | 153 | 180 | 144 | 91 | 161 |
| 86 | 79 | 101 | 104 | 131 | 91 | 88 | 90 | 91 | 101 | 56 | 115 |
| 121 | 169 | 132 | 162 | 233 | 136 | 192 | 120 | 148 | 161 | 255 | 169 |
| 45 | 79 | 71 | 67 | 142 | 49 | 70 | 46 | 87 | 62 | 0 | 31 |
| 127 | 133 | 0 | 0 | 0 | 0 | 216 | 141 | 199 | 272 | 255 | 103 |
| 0 | 5 | 0 | 0 | 0 | 0 | 42 | 65 | 50 | 147 | 0 | 0 |
| 510 | 510 | 510 | 510 | 701 | 510 | 510 | 510 | 510 | 510 | 255 | 510 |
| 510 | 893 | 510 | 510 | 829 | 510 | 765 | 551 | 765 | 1,403 | 765 | 510 |
| 152 | 160 | 175 | 163 | 202 | 156 | 182 | 145 | 169 | 162 | 179 | 153 |
| 76 | 77 | 100 | 101 | 134 | 80 | 87 | 85 | 86 | 108 | 90 | 103 |
| 64 | 64 | 64 | 31 | 45 | 32 | 64 | 31 | 32 | 16 | | |
| 125 | 128 | 125 | 123 | 128 | 123 | 128 | 90 | 97 | 90 | | |
| 128 | 128 | 128 | 128 | 128 | 128 | 180 | 128 | 171 | 128 | | |
| 250 | 255 | 255 | 255 | 255 | 255 | 255 | 191 | 255 | 250 | | |
| 255 | 255 | 255 | 255 | 510 | 255 | 255 | 255 | 255 | 255 | | |
| 255 | 255 | 383 | 255 | 510 | 255 | 255 | 255 | 255 | 383 | | |
| 383 | 510 | 510 | 510 | 510 | 383 | 510 | 510 | 510 | 510 | | |
| 510 | 510 | 510 | 510 | 510 | 383 | 510 | 510 | 510 | 510 | 255 | 510 |
| .7 | 2.4 | 1.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | .8 | 0 |
| .4 | 1.3 | 1.3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3.8 | 0 |

TABLE 4.11.--GRAPEJUICE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.8 | 2.6 | 5.2 | 6.8 | 5.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.9 | 1.8 | 3.3 | 4.8 | 3.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .7 | .7 | 1.5 | 1.2 | 1.4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | .1 | .4 | .8 | .7 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.6 | 1.6 | 1.7 | 1.5 | 1.7 |
| STANDARD DEVIATION.....DO..... | .9 | .9 | .9 | .9 | 1.2 |
| MAXIMUM.....DO..... | 7.0 | 4.0 | 4.0 | 7.0 | 7.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 114 | 66 | 93 | 92 | 106 |
| STANDARD DEVIATION.....GM..... | 95 | 57 | 66 | 68 | 86 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 32 | | 21 | 32 | 32 |
| 25TH.....GM..... | 63 | | 42 | 42 | 63 |
| 50TH.....GM..... | 84 | | 84 | 63 | 84 |
| 75TH.....GM..... | 148 | | 127 | 116 | 127 |
| 90TH.....GM..... | 211 | | 169 | 179 | 190 |
| 95TH.....GM..... | 253 | | 211 | 190 | 253 |
| 99TH.....GM..... | 464 | | 337 | 316 | 569 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 233 | 181 | 198 | 193 | 221 |
| STANDARD DEVIATION.....GM..... | 149 | 151 | 125 | 97 | 103 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 232 | 79 | 211 | 186 | 164 |
| STANDARD DEVIATION.....GM..... | 126 | 54 | 117 | 81 | 57 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 246 | 211 | 142 | 201 | 264 |
| STANDARD DEVIATION.....GM..... | 157 | 0 | 56 | 102 | 141 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,265 | 506 | 633 | 759 | 759 |
| MAXIMUM OVER 3 DAYS.....GM..... | 3,036 | 633 | 1,012 | 1,771 | 1,708 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 212 | 121 | 166 | 180 | 183 |
| STANDARD DEVIATION.....GM..... | 110 | 69 | 85 | 75 | 78 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 83 | | 63 | 63 | 63 |
| 25TH.....GM..... | 127 | | 127 | 127 | 127 |
| 50TH.....GM..... | 190 | | 127 | 190 | 190 |
| 75TH.....GM..... | 253 | | 190 | 190 | 253 |
| 90TH.....GM..... | 316 | | 253 | 253 | 253 |
| 95TH.....GM..... | 380 | | 253 | 253 | 253 |
| 99TH.....GM..... | 569 | | 506 | 380 | 506 |
| MAXIMUM.....GM..... | 1,012 | 253 | 506 | 569 | 506 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 4.5 | 0 | 0 | 1.4 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 0 | 0 | .8 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 2.9 | 4.8 | 2.2 | 2.0 | 2.1 | 3.5 | 1.3 | 1.9 | 1.9 | 2.2 | .8 | 1.6 |
| 2.1 | 3.4 | 1.8 | 1.5 | 1.3 | 2.3 | .8 | 1.2 | 1.1 | 1.5 | .2 | .5 |
| .6 | 1.0 | .2 | .3 | .3 | 1.0 | .3 | .5 | .5 | .5 | .3 | .9 |
| .2 | .4 | .2 | .2 | .5 | .2 | .2 | .2 | .3 | .2 | .3 | .2 |
| 1.5 | 1.6 | 1.4 | 1.4 | 1.8 | 1.6 | 1.6 | 1.6 | 1.7 | 1.5 | 3.6 | 2.0 |
| .8 | .8 | .9 | .7 | 1.0 | .9 | 1.0 | .8 | .8 | 1.0 | 2.6 | .8 |
| 4.0 | 6.0 | 5.0 | 3.0 | 5.0 | 6.0 | 6.0 | 4.0 | 3.0 | 5.0 | 7.0 | 3.0 |
| 100 | 106 | 141 | 126 | 170 | 122 | 114 | 100 | 122 | 105 | 322 | 150 |
| 70 | 79 | 117 | 93 | 104 | 120 | 88 | 66 | 104 | 97 | 304 | 85 |
| 32 | 42 | | | 42 | 42 | 21 | 21 | | | | |
| 63 | 63 | | | 84 | 63 | 63 | 42 | | | | |
| 84 | 84 | | | 127 | 84 | 84 | 84 | | | | |
| 127 | 127 | | | 253 | 169 | 158 | 169 | | | | |
| 200 | 210 | | | 337 | 211 | 211 | 190 | | | | |
| 211 | 274 | | | 337 | 253 | 295 | 211 | | | | |
| 464 | 506 | | | 464 | 1,012 | 422 | 253 | | | | |
| 218 | 238 | 324 | 267 | 370 | 238 | 248 | 188 | 193 | 190 | 190 | 183 |
| 119 | 130 | 250 | 111 | 241 | 111 | 240 | 99 | 72 | 61 | 0 | 52 |
| 254 | 205 | 525 | 358 | 335 | 252 | 217 | 241 | 272 | 181 | 158 | 298 |
| 142 | 118 | 160 | 203 | 161 | 105 | 129 | 101 | 221 | 35 | 0 | 107 |
| 156 | 243 | 220 | 244 | 266 | 395 | 200 | 175 | 245 | 354 | 717 | 155 |
| 38 | 118 | 121 | 85 | 77 | 338 | 37 | 17 | 10 | 134 | 0 | 62 |
| 1,012 | 822 | 1,012 | 759 | 1,012 | 1,012 | 1,265 | 633 | 759 | 633 | 759 | 506 |
| 1,392 | 1,518 | 1,392 | 1,265 | 1,392 | 3,036 | 1,265 | 949 | 1,012 | 1,392 | 2,151 | 759 |
| 202 | 205 | 301 | 274 | 285 | 234 | 209 | 193 | 214 | 205 | 268 | 229 |
| 105 | 96 | 207 | 136 | 135 | 101 | 142 | 86 | 104 | 60 | 102 | 95 |
| 95 | 95 | | | 127 | 127 | 83 | 63 | | | | |
| 127 | 127 | | | 253 | 190 | 127 | 127 | | | | |
| 190 | 190 | | | 253 | 253 | 190 | 190 | | | | |
| 253 | 253 | | | 316 | 253 | 253 | 253 | | | | |
| 316 | 253 | | | 506 | 316 | 253 | 253 | | | | |
| 380 | 316 | | | 506 | 506 | 380 | 380 | | | | |
| 506 | 506 | | | 759 | 506 | 1,012 | 380 | | | | |
| 759 | 822 | 1,012 | 759 | 759 | 506 | 1,012 | 633 | 506 | 316 | 506 | 380 |
| 0 | 0 | 0 | 0 | 1.3 | 1.8 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | .9 | .6 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 4.12.--RAW PEACHES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.5 | 1.2 | 3.5 | 3.8 | 4.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.0 | 1.0 | 2.7 | 2.6 | 3.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.2 | .2 | .6 | 1.0 | 1.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .4 | 0 | .1 | .2 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER.... | 1.6 | 1.2 | 1.4 | 1.6 | 1.5 |
| STANDARD DEVIATION.....DO..... | 1.0 | .4 | 1.0 | 1.1 | .9 |
| MAXIMUM.....DO..... | 7.0 | 2.0 | 7.0 | 7.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 84 | 46 | 60 | 67 | 74 |
| STANDARD DEVIATION.....GM..... | 66 | 28 | 58 | 50 | 52 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 25 | | | 25 | 30 |
| 25TH.....GM..... | 51 | | | 33 | 51 |
| 50TH.....GM..... | 51 | | | 51 | 51 |
| 75TH.....GM..... | 101 | | | 96 | 101 |
| 90TH.....GM..... | 152 | | | 101 | 152 |
| 95TH.....GM..... | 203 | | | 135 | 152 |
| 99TH.....GM..... | 355 | | | 355 | 304 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 164 | 104 | 121 | 135 | 160 |
| STANDARD DEVIATION.....GM..... | 86 | 38 | 55 | 58 | 82 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 176 | 152 | 152 | 155 | 179 |
| STANDARD DEVIATION.....GM..... | 74 | 0 | 15 | 33 | 95 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 224 | 0 | 355 | 197 | 152 |
| STANDARD DEVIATION.....GM..... | 100 | 0 | 0 | 115 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 760 | 152 | 456 | 456 | 500 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,520 | 304 | 1,064 | 1,064 | 912 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 160 | 118 | 129 | 128 | 145 |
| STANDARD DEVIATION.....GM..... | 75 | 39 | 48 | 36 | 68 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 76 | | | 76 | 76 |
| 25TH.....GM..... | 152 | | | 100 | 100 |
| 50TH.....GM..... | 152 | | | 152 | 152 |
| 75TH.....GM..... | 152 | | | 152 | 152 |
| 90TH.....GM..... | 304 | | | 152 | 152 |
| 95TH.....GM..... | 304 | | | 152 | 250 |
| 99TH.....GM..... | 456 | | | 200 | 456 |
| MAXIMUM.....GM..... | 760 | 152 | 300 | 200 | 500 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER.... | 7.8 | 0 | 0 | 0 | 1.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 0 | 0 | 0 | 1.2 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.5 | 4.9 | 4.0 | 3.3 | 2.8 | 4.0 | 4.2 | 5.7 | 5.8 | 8.9 | 4.1 | 6.9 |
| 2.9 | 3.3 | 3.1 | 2.1 | 1.9 | 2.7 | 2.8 | 3.6 | 3.8 | 5.3 | 2.1 | 4.3 |
| .5 | 1.3 | .8 | 1.0 | .6 | 1.0 | 1.1 | 1.5 | 1.6 | 2.8 | 2.0 | 2.3 |
| .2 | .4 | .1 | .2 | .3 | .3 | .3 | .7 | .4 | .8 | 0 | .3 |
| 1.5 | 1.6 | 1.3 | 1.5 | 1.6 | 1.6 | 1.6 | 1.7 | 1.7 | 1.6 | 1.5 | 1.6 |
| 1.1 | 1.1 | .5 | .7 | .9 | 1.0 | 1.0 | 1.0 | 1.1 | .8 | .5 | .8 |
| 6.0 | 6.0 | 3.0 | 4.0 | 6.0 | 7.0 | 7.0 | 6.0 | 5.0 | 4.0 | 2.0 | 4.0 |
| 83 | 84 | 87 | 71 | 86 | 83 | 87 | 92 | 96 | 86 | 69 | 81 |
| 69 | 65 | 65 | 42 | 64 | 64 | 63 | 77 | 82 | 62 | 25 | 59 |
| 25 | 25 | 30 | | 25 | 25 | 33 | 30 | 30 | 30 | | 17 |
| 51 | 51 | 51 | | 51 | 51 | 51 | 51 | 51 | 51 | | 51 |
| 51 | 51 | 51 | | 59 | 51 | 51 | 59 | 51 | 59 | | 59 |
| 101 | 101 | 101 | | 101 | 101 | 101 | 101 | 109 | 101 | | 101 |
| 203 | 152 | 203 | | 193 | 152 | 153 | 152 | 253 | 152 | | 203 |
| 253 | 203 | 203 | | 253 | 178 | 203 | 262 | 279 | 203 | | 211 |
| 304 | 355 | 304 | | 304 | 402 | 355 | 405 | 304 | 456 | | 226 |
| 167 | 157 | 197 | 156 | 161 | 170 | 183 | 169 | 150 | 159 | 143 | 141 |
| 92 | 90 | 134 | 88 | 59 | 104 | 88 | 83 | 43 | 58 | 40 | 82 |
| 279 | 192 | 228 | 132 | 185 | 166 | 166 | 161 | 250 | 181 | 137 | 196 |
| 97 | 101 | 100 | 28 | 85 | 60 | 52 | 45 | 134 | 61 | 17 | 78 |
| 235 | 207 | 253 | 181 | 211 | 206 | 255 | 242 | 247 | 182 | 0 | 192 |
| 64 | 48 | 0 | 21 | 60 | 101 | 85 | 116 | 39 | 117 | 0 | 50 |
| 608 | 760 | 608 | 608 | 608 | 760 | 608 | 760 | 608 | 608 | 200 | 456 |
| 912 | 1,064 | 912 | 608 | 912 | 1,216 | 1,260 | 1,520 | 912 | 1,368 | 304 | 678 |
| 170 | 153 | 205 | 142 | 165 | 156 | 166 | 165 | 173 | 160 | 139 | 152 |
| 77 | 68 | 111 | 66 | 58 | 71 | 76 | 84 | 75 | 67 | 33 | 70 |
| 89 | 76 | 89 | | 89 | 76 | 76 | 76 | 76 | 76 | | 50 |
| 152 | 152 | 152 | | 152 | 152 | 152 | 152 | 152 | 150 | | 89 |
| 152 | 152 | 152 | | 152 | 152 | 152 | 152 | 152 | 152 | | 152 |
| 152 | 152 | 304 | | 152 | 152 | 152 | 152 | 178 | 152 | | 178 |
| 304 | 178 | 304 | | 304 | 253 | 304 | 304 | 304 | 304 | | 304 |
| 304 | 304 | 456 | | 304 | 304 | 304 | 304 | 304 | 304 | | 304 |
| 456 | 456 | 608 | | 304 | 356 | 504 | 456 | 304 | 304 | | 356 |
| 608 | 608 | 608 | 608 | 304 | 760 | 608 | 760 | 304 | 456 | 200 | 356 |
| 0 | 1.5 | 0 | 1.2 | 0 | 0 | 0 | 1.7 | 1.8 | 0 | 0 | 0 |
| 0 | .9 | 0 | 1.7 | 0 | 0 | 0 | .3 | 1.6 | 0 | 0 | 0 |

TABLE 4.13.--COOKED, CANNED, AND FROZEN PEACHES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 8.1 | 28.3 | 10.5 | 6.9 | 12.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 6.4 | 16.4 | 7.2 | 6.1 | 11.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.3 | 8.7 | 3.0 | .7 | 1.3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 3.2 | .3 | .2 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 1.9 | 1.4 | 1.2 | 1.1 |
| STANDARD DEVIATION.....DO..... | .6 | 1.1 | .6 | .4 | .4 |
| MAXIMUM.....DC..... | 7.0 | 7.0 | 4.0 | 3.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 68 | 51 | 57 | 58 | 54 |
| STANDARD DEVIATION.....GM..... | 52 | 35 | 39 | 35 | 35 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 20 | 9 | 10 | 16 | 15 |
| 25TH.....GM..... | 42 | 28 | 28 | 32 | 32 |
| 50TH.....GM..... | 49 | 42 | 43 | 43 | 43 |
| 75TH.....GM..... | 85 | 70 | 82 | 83 | 71 |
| 90TH.....GM..... | 128 | 96 | 96 | 85 | 91 |
| 95TH.....GM..... | 171 | 113 | 128 | 128 | 123 |
| 99TH.....GM..... | 256 | 162 | 192 | 171 | 171 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 166 | 107 | 135 | 158 | 143 |
| STANDARD DEVIATION.....GM..... | 104 | 59 | 84 | 93 | 81 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 153 | 104 | 119 | 132 | 132 |
| STANDARD DEVIATION.....GM..... | 76 | 54 | 68 | 61 | 38 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 178 | 84 | 123 | 106 | 205 |
| STANDARD DEVIATION.....GM..... | 120 | 45 | 42 | 30 | 86 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,280 | 340 | 425 | 512 | 512 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,816 | 616 | 638 | 512 | 1,024 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 156 | 82 | 123 | 149 | 141 |
| STANDARD DEVIATION.....GM..... | 96 | 49 | 74 | 87 | 72 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 48 | 14 | 31 | 57 | 48 |
| 25TH.....GM..... | 96 | 56 | 64 | 96 | 96 |
| 50TH.....GM..... | 128 | 68 | 112 | 128 | 128 |
| 75TH.....GM..... | 192 | 113 | 128 | 192 | 192 |
| 90TH.....GM..... | 256 | 135 | 220 | 256 | 256 |
| 95TH.....GM..... | 288 | 170 | 288 | 256 | 256 |
| 99TH.....GM..... | 512 | 224 | 384 | 512 | 384 |
| MAXIMUM.....GM..... | 1,280 | 288 | 384 | 512 | 384 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 32.3 | 0 | 0 | 1.1 | 1.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .8 | 0 | 0 | .8 | .6 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 10.7 | 10.6 | 8.4 | 6.9 | 4.2 | 4.0 | 8.1 | 6.7 | 13.1 | 11.3 | 12.0 | 13.2 |
| 9.8 | 9.7 | 7.9 | 6.3 | 3.5 | 3.4 | 6.3 | 5.2 | 8.8 | 7.3 | 9.1 | 8.7 |
| .9 | .9 | .5 | .6 | .6 | .6 | 1.4 | 1.2 | 3.1 | 2.8 | 2.0 | 3.7 |
| 0 | .1 | 0 | 0 | .1 | (2) | .4 | .2 | 1.3 | 1.1 | .8 | .9 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.2 | 1.3 | 1.3 | 1.5 | 1.5 | 1.4 | 1.6 |
| .3 | .3 | .2 | .3 | .4 | .5 | .6 | .6 | .8 | .8 | .7 | .9 |
| 2.0 | 3.0 | 2.0 | 2.0 | 3.0 | 4.0 | 5.0 | 4.0 | 6.0 | 5.0 | 4.0 | 4.0 |
| 51 | 54 | 58 | 53 | 83 | 67 | 79 | 74 | 93 | 76 | 76 | 77 |
| 31 | 33 | 35 | 30 | 46 | 41 | 57 | 65 | 88 | 55 | 44 | 59 |
| 11 | 21 | 16 | 16 | 21 | 21 | 21 | 21 | 21 | 21 | 25 | 21 |
| 32 | 32 | 42 | 32 | 43 | 42 | 43 | 42 | 43 | 42 | 43 | 42 |
| 43 | 43 | 43 | 43 | 83 | 64 | 64 | 53 | 83 | 64 | 64 | 57 |
| 64 | 64 | 85 | 83 | 96 | 85 | 85 | 85 | 113 | 85 | 91 | 85 |
| 85 | 85 | 96 | 85 | 160 | 125 | 161 | 128 | 171 | 152 | 148 | 169 |
| 107 | 115 | 128 | 96 | 171 | 171 | 203 | 182 | 229 | 189 | 171 | 192 |
| 167 | 171 | 192 | 147 | 213 | 213 | 256 | 293 | 341 | 213 | 197 | 296 |
| 148 | 149 | 164 | 145 | 217 | 173 | 194 | 173 | 215 | 151 | 193 | 147 |
| 89 | 82 | 93 | 73 | 115 | 94 | 132 | 113 | 142 | 74 | 112 | 84 |
| 114 | 142 | 183 | 151 | 200 | 167 | 172 | 166 | 155 | 154 | 157 | 160 |
| 45 | 79 | 63 | 58 | 56 | 58 | 88 | 87 | 81 | 77 | 43 | 69 |
| 0 | 111 | 0 | 0 | 213 | 213 | 181 | 238 | 214 | 176 | 138 | 227 |
| 0 | 0 | 0 | 0 | 0 | 0 | 63 | 177 | 199 | 63 | 54 | 53 |
| 512 | 512 | 576 | 288 | 640 | 512 | 1,280 | 1,024 | 1,024 | 672 | 512 | 531 |
| 512 | 768 | 576 | 512 | 640 | 768 | 1,280 | 2,048 | 2,816 | 1,152 | 592 | 888 |
| 142 | 145 | 166 | 146 | 210 | 166 | 180 | 168 | 185 | 148 | 169 | 145 |
| 85 | 81 | 91 | 71 | 106 | 86 | 115 | 109 | 120 | 71 | 81 | 63 |
| 32 | 63 | 48 | 48 | 64 | 63 | 60 | 63 | 48 | 61 | 74 | 63 |
| 96 | 96 | 125 | 96 | 128 | 125 | 125 | 96 | 125 | 97 | 128 | 96 |
| 128 | 128 | 128 | 128 | 192 | 128 | 128 | 128 | 148 | 128 | 148 | 128 |
| 192 | 192 | 250 | 192 | 256 | 250 | 256 | 222 | 250 | 192 | 218 | 192 |
| 256 | 256 | 256 | 256 | 375 | 256 | 256 | 256 | 256 | 256 | 256 | 250 |
| 256 | 256 | 288 | 256 | 384 | 288 | 313 | 375 | 384 | 256 | 273 | 256 |
| 500 | 384 | 576 | 288 | 513 | 512 | 577 | 513 | 768 | 320 | 500 | 288 |
| 512 | 512 | 576 | 288 | 640 | 512 | 1,280 | 1,024 | 768 | 640 | 500 | 339 |
| 2.9 | 2.0 | 2.1 | 0 | 1.1 | 6.3 | 4.3 | 6.3 | .9 | 0 | 1.3 | 2.2 |
| 1.2 | .8 | 1.7 | 0 | .6 | 2.4 | .8 | 1.0 | .4 | 0 | 1.5 | 1.1 |

TABLE 4.14.--RAW PEARS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.1 | 1.2 | 2.3 | 2.9 | 4.0 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.4 | .3 | 1.7 | 2.3 | 3.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .6 | .7 | .4 | .4 | .6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | .2 | .2 | .1 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.4 | 2.0 | 1.4 | 1.2 | 1.3 |
| STANDARD DEVIATION.....DO..... | .8 | .6 | .7 | .5 | .7 |
| MAXIMUM.....DO..... | 6.0 | 3.0 | 3.0 | 3.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 75 | 37 | 61 | 62 | 71 |
| STANDARD DEVIATION.....GM..... | 50 | 35 | 43 | 34 | 40 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 27 | | | 18 | 28 |
| 25TH.....GM..... | 55 | | | 55 | 55 |
| 50TH.....GM..... | 55 | | | 55 | 55 |
| 75TH.....GM..... | 109 | | | 55 | 82 |
| 90TH.....GM..... | 114 | | | 110 | 109 |
| 95TH.....GM..... | 164 | | | 149 | 164 |
| 99TH.....GM..... | 273 | | | 164 | 219 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 168 | 20 | 117 | 147 | 173 |
| STANDARD DEVIATION.....GM..... | 85 | 0 | 69 | 51 | 58 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 173 | 38 | 164 | 184 | 167 |
| STANDARD DEVIATION.....GM..... | 58 | 5 | 0 | 37 | 57 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 184 | 110 | 148 | 103 | 193 |
| STANDARD DEVIATION.....GM..... | 63 | 0 | 14 | 40 | 27 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 2,132 | 164 | 328 | 328 | 492 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,132 | 329 | 492 | 492 | 656 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 163 | 56 | 131 | 150 | 163 |
| STANDARD DEVIATION.....GM..... | 69 | 40 | 43 | 57 | 42 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 82 | | | 55 | 82 |
| 25TH.....GM..... | 164 | | | 141 | 164 |
| 50TH.....GM..... | 164 | | | 164 | 164 |
| 75TH.....GM..... | 164 | | | 164 | 164 |
| 90TH.....GM..... | 164 | | | 164 | 164 |
| 95TH.....GM..... | 328 | | | 200 | 164 |
| 99TH.....GM..... | 328 | | | 328 | 328 |
| MAXIMUM.....GM..... | 2,132 | 164 | 164 | 328 | 328 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | .6 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | (2) | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 2.7 | 3.3 | 3.2 | 1.4 | 1.8 | 2.1 | 4.0 | 3.9 | 4.9 | 4.3 | 3.5 | 4.2 |
| 2.4 | 2.6 | 2.8 | 1.1 | 1.1 | 1.6 | 2.9 | 3.0 | 3.0 | 3.4 | 1.7 | 3.2 |
| .2 | .7 | .3 | .2 | .5 | .4 | .8 | .6 | 1.6 | .7 | .6 | 1.0 |
| (2) | .1 | .1 | 0 | .2 | .2 | .3 | .3 | .3 | .2 | 1.2 | 0 |
| 1.2 | 1.3 | 1.3 | 1.2 | 1.6 | 1.4 | 1.4 | 1.4 | 1.5 | 1.3 | 2.2 | 1.3 |
| .5 | .7 | .7 | .5 | .9 | .7 | .7 | .9 | .6 | .6 | 1.1 | .4 |
| 4.0 | 5.0 | 5.0 | 3.0 | 5.0 | 4.0 | 5.0 | 6.0 | 3.0 | 3.0 | 4.0 | 2.0 |
| 63 | 71 | 83 | 69 | 92 | 70 | 80 | 77 | 74 | 69 | 111 | 65 |
| 34 | 40 | 86 | 27 | 59 | 44 | 49 | 57 | 28 | 38 | 42 | 40 |
| 27 | 27 | | | 27 | 14 | 27 | 28 | 27 | 14 | | |
| 55 | 55 | | | 55 | 55 | 55 | 55 | 55 | 55 | | |
| 55 | 55 | | | 55 | 55 | 55 | 55 | 55 | 55 | | |
| 55 | 55 | | | 109 | 91 | 109 | 72 | 109 | 82 | | |
| 109 | 109 | | | 164 | 109 | 164 | 164 | 109 | 109 | | |
| 164 | 164 | | | 273 | 164 | 164 | 219 | 109 | 164 | | |
| 219 | 273 | | | 273 | 219 | 273 | 273 | 110 | 164 | | |
| 165 | 163 | 230 | 179 | 168 | 155 | 177 | 166 | 172 | 157 | 252 | 157 |
| 65 | 51 | 268 | 62 | 51 | 71 | 89 | 44 | 57 | 48 | 126 | 36 |
| 182 | 177 | 164 | 165 | 181 | 159 | 180 | 192 | 153 | 182 | 164 | 160 |
| 34 | 42 | 0 | 1 | 40 | 13 | 49 | 87 | 27 | 42 | 0 | 96 |
| 219 | 195 | 168 | 0 | 210 | 178 | 187 | 218 | 82 | 164 | 150 | 0 |
| 0 | 82 | 5 | 0 | 57 | 48 | 46 | 76 | 0 | 0 | 14 | 0 |
| 492 | 492 | 2,132 | 328 | 328 | 492 | 656 | 656 | 330 | 328 | 492 | 328 |
| 656 | 820 | 2,132 | 332 | 820 | 656 | 820 | 1,230 | 330 | 492 | 492 | 656 |
| 163 | 161 | 195 | 167 | 171 | 156 | 171 | 166 | 149 | 162 | 151 | 153 |
| 46 | 42 | 219 | 57 | 47 | 56 | 57 | 63 | 39 | 46 | 44 | 64 |
| 82 | 82 | | | 82 | 82 | 82 | 82 | 82 | 82 | | |
| 164 | 164 | | | 164 | 164 | 164 | 164 | 152 | 164 | | |
| 164 | 164 | | | 164 | 164 | 164 | 164 | 164 | 164 | | |
| 164 | 164 | | | 164 | 164 | 164 | 164 | 164 | 164 | | |
| 164 | 164 | | | 168 | 164 | 165 | 164 | 164 | 164 | | |
| 200 | 165 | | | 328 | 164 | 328 | 328 | 165 | 246 | | |
| 328 | 328 | | | 328 | 492 | 328 | 410 | 246 | 328 | | |
| 328 | 492 | 2,132 | 328 | 328 | 492 | 328 | 656 | 246 | 328 | 246 | 328 |
| 0 | .6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | .7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 4.15.--COOKED AND CANNED PEARS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.9 | 23.1 | 5.4 | 3.2 | 4.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.1 | 13.8 | 4.1 | 2.7 | 4.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .6 | 6.0 | 1.2 | .5 | .1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 3.2 | .1 | (2) | (2) |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.3 | 2.1 | 1.3 | 1.2 | 1.1 |
| STANDARD DEVIATION.....DO..... | .7 | 1.2 | .5 | .5 | .3 |
| MAXIMUM.....DO..... | 8.0 | 8.0 | 4.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 65 | 52 | 47 | 63 | 49 |
| STANDARD DEVIATION.....GM..... | 46 | 32 | 29 | 50 | 27 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 21 | 9 | 16 | 19 | 11 |
| 25TH.....GM..... | 31 | 28 | 28 | 31 | 31 |
| 50TH.....GM..... | 53 | 43 | 43 | 45 | 43 |
| 75TH.....GM..... | 84 | 72 | 63 | 63 | 63 |
| 90TH.....GM..... | 125 | 99 | 75 | 101 | 94 |
| 95TH.....GM..... | 157 | 128 | 104 | 188 | 94 |
| 99TH.....GM..... | 251 | 135 | 207 | 302 | 128 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 156 | 111 | 111 | 149 | 141 |
| STANDARD DEVIATION.....GM..... | 86 | 65 | 47 | 81 | 78 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 166 | 101 | 112 | 189 | 144 |
| STANDARD DEVIATION.....GM..... | 90 | 46 | 31 | 119 | 36 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 148 | 89 | 207 | 188 | 94 |
| STANDARD DEVIATION.....GM..... | 73 | 31 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 752 | 424 | 250 | 564 | 383 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,504 | 424 | 620 | 907 | 383 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 149 | 75 | 112 | 154 | 138 |
| STANDARD DEVIATION.....GM..... | 85 | 44 | 47 | 89 | 74 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 42 | 14 | 47 | 47 | 32 |
| 25TH.....GM..... | 94 | 42 | 77 | 94 | 94 |
| 50TH.....GM..... | 128 | 64 | 97 | 128 | 128 |
| 75TH.....GM..... | 188 | 112 | 133 | 188 | 188 |
| 90TH.....GM..... | 255 | 128 | 188 | 255 | 255 |
| 95TH.....GM..... | 282 | 135 | 188 | 282 | 282 |
| 99TH.....GM..... | 383 | 224 | 250 | 510 | 383 |
| MAXIMUM.....GM..... | 752 | 224 | 250 | 510 | 383 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 6.5 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 4.9 | 5.5 | 3.1 | 2.7 | 2.0 | 1.8 | 3.9 | 3.7 | 5.2 | 6.2 | 5.8 | 5.6 |
| 4.3 | 5.1 | 2.7 | 2.3 | 1.6 | 1.5 | 2.8 | 3.1 | 4.4 | 3.9 | 4.2 | 4.0 |
| .4 | .4 | .3 | .3 | .4 | .3 | 1.0 | .4 | .6 | 1.6 | 1.6 | 1.3 |
| .1 | .1 | .1 | .1 | (2) | 0 | .1 | .1 | .2 | .6 | 0 | .3 |
| 1.2 | 1.1 | 1.1 | 1.2 | 1.2 | 1.2 | 1.3 | 1.2 | 1.2 | 1.5 | 1.3 | 1.4 |
| .4 | .4 | .4 | .4 | .5 | .4 | .5 | .6 | .5 | .8 | .4 | .6 |
| 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 6.0 | 3.0 | 4.0 | 2.0 | 3.0 |
| 52 | 52 | 73 | 63 | 73 | 64 | 79 | 64 | 78 | 86 | 71 | 72 |
| 39 | 30 | 51 | 37 | 41 | 36 | 51 | 47 | 74 | 63 | 47 | 42 |
| 21 | 13 | | | 21 | 21 | 21 | 19 | 31 | 19 | | 25 |
| 31 | 31 | | | 43 | 31 | 43 | 31 | 43 | 43 | | 42 |
| 43 | 43 | | | 63 | 63 | 63 | 56 | 63 | 64 | | 63 |
| 63 | 63 | | | 101 | 83 | 105 | 77 | 85 | 113 | | 85 |
| 85 | 85 | | | 128 | 125 | 157 | 125 | 163 | 148 | | 149 |
| 128 | 105 | | | 146 | 137 | 170 | 157 | 179 | 188 | | 149 |
| 251 | 128 | | | 188 | 170 | 251 | 245 | 501 | 340 | | 181 |
| 129 | 148 | 179 | 166 | 190 | 171 | 179 | 159 | 171 | 160 | 152 | 164 |
| 67 | 87 | 84 | 92 | 97 | 88 | 102 | 86 | 61 | 79 | 74 | 88 |
| 162 | 127 | 224 | 152 | 165 | 157 | 191 | 155 | 294 | 198 | 185 | 161 |
| 82 | 30 | 99 | 69 | 66 | 61 | 79 | 69 | 227 | 92 | 76 | 47 |
| 208 | 77 | 251 | 126 | 188 | 0 | 167 | 194 | 163 | 173 | 0 | 146 |
| 41 | 0 | 0 | 0 | 0 | 0 | 15 | 86 | 0 | 75 | 0 | 48 |
| 510 | 658 | 397 | 383 | 454 | 510 | 510 | 638 | 752 | 510 | 383 | 446 |
| 792 | 658 | 773 | 564 | 680 | 510 | 752 | 1,024 | 1,504 | 1,020 | 609 | 542 |
| 135 | 142 | 191 | 160 | 179 | 160 | 179 | 156 | 189 | 169 | 166 | 151 |
| 68 | 82 | 90 | 85 | 90 | 69 | 84 | 87 | 125 | 82 | 81 | 64 |
| 63 | 38 | | | 64 | 64 | 64 | 63 | 77 | 58 | | 64 |
| 94 | 94 | | | 125 | 94 | 122 | 94 | 128 | 128 | | 94 |
| 128 | 128 | | | 188 | 152 | 188 | 128 | 188 | 154 | | 128 |
| 188 | 188 | | | 191 | 213 | 250 | 191 | 255 | 188 | | 188 |
| 191 | 255 | | | 304 | 255 | 282 | 256 | 255 | 255 | | 255 |
| 255 | 255 | | | 376 | 255 | 376 | 304 | 282 | 282 | | 255 |
| 383 | 383 | | | 383 | 282 | 383 | 383 | 752 | 510 | | 282 |
| 510 | 658 | 397 | 383 | 454 | 282 | 448 | 638 | 752 | 510 | 383 | 282 |
| 0 | 0 | 0 | 0 | 0 | 1.0 | 0 | 1.8 | 0 | 0 | 1.6 | 2.2 |
| 0 | 0 | 0 | 0 | 0 | .9 | 0 | .6 | 0 | 0 | 4.0 | 2.7 |

TABLE 4.16.--COOKED AND CANNED PINEAPPLE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1: | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.6 | .5 | 2.2 | 1.6 | 2.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.2 | .5 | 2.0 | 1.3 | 2.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .3 | 0 | .2 | .4 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 1.2 | 1.0 | 1.1 | 1.3 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .0 | .3 | .5 | .4 |
| MAXIMUM.....DO..... | 4.0 | 1.0 | 2.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 41 | 18 | 24 | 40 | 35 |
| STANDARD DEVIATION.....GM..... | 33 | 20 | 15 | 26 | 19 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 7 | | | | 8 |
| 25TH.....GM..... | 21 | | | | 21 |
| 50TH.....GM..... | 39 | | | | 39 |
| 75TH.....GM..... | 42 | | | | 42 |
| 90TH.....GM..... | 83 | | | | 63 |
| 95TH.....GM..... | 85 | | | | 82 |
| 99TH.....GM..... | 170 | | | | 83 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 102 | 53 | 68 | 87 | 99 |
| STANDARD DEVIATION.....GM..... | 62 | 60 | 34 | 45 | 51 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 121 | 0 | 59 | 113 | 97 |
| STANDARD DEVIATION.....GM..... | 80 | 0 | 43 | 35 | 30 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 134 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 52 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 515 | 125 | 125 | 206 | 250 |
| MAXIMUM OVER 3 DAYS.....GM..... | 824 | 125 | 204 | 312 | 250 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 105 | 53 | 66 | 94 | 96 |
| STANDARD DEVIATION.....GM..... | 64 | 60 | 37 | 44 | 50 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 23 | | | | 23 |
| 25TH.....GM..... | 58 | | | | 58 |
| 50TH.....GM..... | 106 | | | | 103 |
| 75TH.....GM..... | 125 | | | | 125 |
| 90TH.....GM..... | 206 | | | | 128 |
| 95TH.....GM..... | 250 | | | | 188 |
| 99TH.....GM..... | 308 | | | | 250 |
| MAXIMUM.....GM..... | 375 | 125 | 125 | 206 | 250 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 35.0 | 0 | 2.8 | 2.2 | 1.2 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 3.0 | 0 | 11.2 | 6.2 | 2.1 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 2.6 | 3.2 | 2.0 | 1.9 | 1.7 | 2.6 | 2.2 | 3.4 | 2.8 | 3.7 | 2.0 | 4.6 |
| 2.5 | 2.9 | 1.8 | 1.9 | 1.4 | 2.4 | 1.8 | 2.7 | 2.4 | 3.3 | 1.7 | 3.5 |
| .1 | .3 | .2 | (2) | .2 | .1 | .4 | .7 | .2 | .3 | .3 | 1.0 |
| 0 | 0 | 0 | 0 | (2) | .1 | 0 | (2) | .2 | .1 | 0 | .1 |
| 1.1 | 1.1 | 1.1 | 1.0 | 1.2 | 1.1 | 1.2 | 1.2 | 1.3 | 1.2 | 1.3 | 1.3 |
| .2 | .4 | .3 | .2 | .4 | .4 | .4 | .5 | .6 | .5 | .7 | .6 |
| 2.0 | 3.0 | 2.0 | 2.0 | 3.0 | 3.0 | 2.0 | 3.0 | 3.0 | 3.0 | 3.0 | 4.0 |
| 36 | 51 | 40 | 35 | 47 | 38 | 43 | 42 | 47 | 36 | 46 | 49 |
| 24 | 51 | 28 | 20 | 32 | 30 | 32 | 38 | 29 | 23 | 21 | 39 |
| 10 | 12 | | | 15 | 8 | 6 | 5 | | 10 | | |
| 19 | 34 | | | 28 | 19 | 21 | 19 | | 20 | | |
| 39 | 42 | | | 41 | 34 | 39 | 37 | | 34 | | |
| 42 | 42 | | | 63 | 42 | 55 | 43 | | 42 | | |
| 61 | 82 | | | 83 | 83 | 83 | 82 | | 83 | | |
| 83 | 123 | | | 109 | 85 | 83 | 104 | | 85 | | |
| 167 | 275 | | | 168 | 170 | 167 | 250 | | 96 | | |
| 99 | 118 | 104 | 102 | 126 | 96 | 108 | 95 | 120 | 94 | 135 | 105 |
| 52 | 51 | 59 | 55 | 70 | 65 | 73 | 64 | 63 | 53 | 67 | 62 |
| 150 | 261 | 130 | 125 | 119 | 152 | 115 | 113 | 70 | 93 | 76 | 118 |
| 77 | 156 | 69 | 0 | 84 | 61 | 60 | 76 | 0 | 47 | 0 | 60 |
| 0 | 0 | 0 | 0 | 78 | 125 | 0 | 255 | 126 | 85 | 0 | 180 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 250 | 515 | 250 | 250 | 328 | 375 | 412 | 500 | 250 | 254 | 250 | 255 |
| 500 | 824 | 412 | 250 | 505 | 510 | 500 | 765 | 378 | 287 | 250 | 540 |
| 102 | 135 | 109 | 104 | 122 | 103 | 110 | 101 | 111 | 91 | 109 | 112 |
| 57 | 72 | 61 | 54 | 73 | 65 | 68 | 65 | 49 | 52 | 68 | 67 |
| 20 | 37 | | | 23 | 23 | 20 | 24 | | 23 | | |
| 58 | 103 | | | 63 | 58 | 58 | 58 | | 58 | | |
| 116 | 123 | | | 116 | 103 | 103 | 95 | | 103 | | |
| 125 | 125 | | | 142 | 125 | 125 | 125 | | 125 | | |
| 183 | 250 | | | 246 | 206 | 246 | 206 | | 128 | | |
| 250 | 309 | | | 255 | 250 | 250 | 250 | | 159 | | |
| 250 | 309 | | | 328 | 255 | 255 | 255 | | 254 | | |
| 250 | 309 | 250 | 250 | 328 | 375 | 309 | 313 | 250 | 254 | 250 | 255 |
| 1.0 | 0 | 1.1 | 0 | 6.3 | 11.5 | 0 | 3.9 | 0 | 1.0 | 4.2 | 0 |
| 1.8 | 0 | 3.6 | 0 | 8.3 | 7.5 | 0 | 1.3 | 0 | 1.3 | 31.0 | 0 |

TABLE 4.17.--RAISINS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.2 | 0 | 7.0 | 4.8 | 3.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.6 | 0 | 4.2 | 3.5 | 2.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .5 | 0 | 1.9 | .8 | .9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | 0 | .9 | .5 | .3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.4 | 0 | 1.6 | 1.4 | 1.5 |
| STANDARD DEVIATION.....DO..... | .7 | 0 | .8 | .7 | .7 |
| MAXIMUM.....DO..... | 5.0 | 0 | 4.0 | 3.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 16 | 0 | 15 | 14 | 15 |
| STANDARD DEVIATION.....GM..... | 14 | 0 | 12 | 11 | 10 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 2 | | 1 | 1 | 1 |
| 25TH.....GM..... | 6 | | 6 | 6 | 6 |
| 50TH.....GM..... | 12 | | 12 | 12 | 12 |
| 75TH.....GM..... | 24 | | 21 | 22 | 21 |
| 90TH.....GM..... | 30 | | 35 | 28 | 28 |
| 95TH.....GM..... | 48 | | 36 | 29 | 29 |
| 99TH.....GM..... | 73 | | 48 | 37 | 57 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 37 | 0 | 29 | 32 | 31 |
| STANDARD DEVIATION.....GM..... | 33 | 0 | 23 | 20 | 20 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 35 | 0 | 28 | 31 | 36 |
| STANDARD DEVIATION.....GM..... | 23 | 0 | 18 | 12 | 16 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 28 | 0 | 29 | 27 | 22 |
| STANDARD DEVIATION.....GM..... | 21 | 0 | 17 | 20 | 10 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 290 | 0 | 86 | 109 | 129 |
| MAXIMUM OVER 3 DAYS.....GM..... | 290 | 0 | 218 | 254 | 172 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 33 | 0 | 28 | 30 | 30 |
| STANDARD DEVIATION.....GM..... | 28 | 0 | 20 | 19 | 16 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 3 | | 3 | 3 | 3 |
| 25TH.....GM..... | 14 | | 14 | 17 | 17 |
| 50TH.....GM..... | 28 | | 18 | 28 | 28 |
| 75TH.....GM..... | 43 | | 36 | 43 | 43 |
| 90TH.....GM..... | 73 | | 73 | 55 | 43 |
| 95TH.....GM..... | 73 | | 73 | 73 | 55 |
| 99TH.....GM..... | 145 | | 86 | 73 | 73 |
| MAXIMUM.....GM..... | 290 | 0 | 86 | 109 | 73 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 35.8 | 0 | 2.3 | 2.6 | 1.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 3.0 | 0 | 2.0 | 2.2 | 1.6 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.8 | 1.5 | 1.7 | 1.2 | 1.7 | 2.8 | 1.9 | 1.6 | 2.1 | 2.0 | 2.3 | 1.8 |
| 1.2 | 1.2 | 1.2 | 1.0 | 1.2 | 1.8 | 1.4 | 1.2 | 1.8 | 1.5 | 1.7 | 1.3 |
| .5 | .2 | .5 | .2 | .2 | .7 | .3 | .3 | .2 | .3 | 0 | .6 |
| .1 | .1 | 0 | 0 | .2 | .3 | .1 | .1 | .1 | .2 | .6 | 0 |
| 1.5 | 1.2 | 1.3 | 1.2 | 1.4 | 1.5 | 1.3 | 1.5 | 1.2 | 1.4 | 1.5 | 1.3 |
| .7 | .5 | .6 | .4 | .8 | .8 | .7 | .7 | .5 | .7 | .9 | .5 |
| 4.0 | 3.0 | 3.0 | 2.0 | 4.0 | 5.0 | 4.0 | 4.0 | 3.0 | 3.0 | 3.0 | 2.0 |
| 20 | 19 | 18 | 19 | 18 | 18 | 15 | 17 | 9 | 11 | 8 | 9 |
| 18 | 17 | 12 | 17 | 16 | 18 | 13 | 13 | 6 | 8 | 7 | 12 |
| | | | | 1 | 2 | 2 | 2 | | | | |
| | | | | 6 | 6 | 6 | 6 | | | | |
| | | | | 14 | 12 | 12 | 15 | | | | |
| | | | | 19 | 24 | 18 | 24 | | | | |
| | | | | 48 | 30 | 36 | 29 | | | | |
| | | | | 48 | 64 | 48 | 36 | | | | |
| | | | | 73 | 97 | 48 | 63 | | | | |
| 43 | 52 | 47 | 40 | 43 | 40 | 37 | 43 | 23 | 27 | 22 | 19 |
| 47 | 43 | 37 | 23 | 36 | 43 | 34 | 33 | 13 | 12 | 16 | 15 |
| 48 | 49 | 35 | 73 | 28 | 34 | 35 | 37 | 35 | 19 | 0 | 22 |
| 26 | 37 | 11 | 28 | 28 | 23 | 22 | 18 | 8 | 14 | 0 | 29 |
| 31 | 12 | 0 | 0 | 35 | 37 | 17 | 25 | 9 | 26 | 12 | 0 |
| 8 | 0 | 0 | 0 | 20 | 26 | 14 | 24 | 0 | 13 | 9 | 0 |
| 181 | 145 | 145 | 145 | 145 | 290 | 145 | 188 | 43 | 43 | 43 | 73 |
| 181 | 231 | 145 | 218 | 218 | 290 | 181 | 243 | 85 | 109 | 71 | 145 |
| 41 | 46 | 40 | 49 | 37 | 35 | 33 | 35 | 23 | 25 | 16 | 21 |
| 41 | 41 | 29 | 31 | 33 | 34 | 30 | 25 | 14 | 13 | 14 | 23 |
| | | | | 4 | 5 | 2 | 3 | | | | |
| | | | | 17 | 17 | 14 | 14 | | | | |
| | | | | 28 | 28 | 28 | 36 | | | | |
| | | | | 43 | 36 | 43 | 48 | | | | |
| | | | | 73 | 73 | 73 | 73 | | | | |
| | | | | 109 | 96 | 73 | 73 | | | | |
| | | | | 145 | 145 | 145 | 145 | | | | |
| 181 | 145 | 145 | 145 | 145 | 290 | 145 | 145 | 43 | 43 | 43 | 73 |
| 0 | .9 | .9 | .9 | 2.4 | 0 | 5.2 | 7.5 | 3.4 | 2.0 | 3.2 | 3.0 |
| 0 | 2.2 | 2.9 | 4.3 | 2.5 | 0 | 4.2 | 4.6 | 12.1 | 4.1 | 16.3 | 12.8 |

TABLE 4.18.--RAW STRAWBERRIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.1 | .2 | 1.5 | 1.2 | 1.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.6 | 0 | .9 | 1.0 | 1.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | .2 | .6 | .1 | .1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER.... | 1.4 | 2.0 | 1.5 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .8 | .0 | .8 | .6 | .4 |
| MAXIMUM.....DO..... | 6.0 | 2.0 | 4.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 46 | 80 | 45 | 28 | 34 |
| STANDARD DEVIATION.....GM..... | 44 | 0 | 45 | 17 | 27 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 12 | | | | |
| 25TH.....GM..... | 25 | | | | |
| 50TH.....GM..... | 37 | | | | |
| 75TH.....GM..... | 50 | | | | |
| 90TH.....GM..... | 99 | | | | |
| 95TH.....GM..... | 120 | | | | |
| 99TH.....GM..... | 225 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 101 | 0 | 75 | 70 | 88 |
| STANDARD DEVIATION.....GM..... | 60 | 0 | 34 | 38 | 42 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 115 | 120 | 110 | 87 | 134 |
| STANDARD DEVIATION.....GM..... | 76 | 0 | 88 | 18 | 96 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 139 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 93 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 600 | 150 | 298 | 149 | 373 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,273 | 240 | 596 | 225 | 522 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 100 | 120 | 87 | 69 | 87 |
| STANDARD DEVIATION.....GM..... | 58 | 30 | 41 | 34 | 44 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 37 | | | | |
| 25TH.....GM..... | 75 | | | | |
| 50TH.....GM..... | 75 | | | | |
| 75TH.....GM..... | 149 | | | | |
| 90TH.....GM..... | 149 | | | | |
| 95TH.....GM..... | 180 | | | | |
| 99TH.....GM..... | 298 | | | | |
| MAXIMUM.....GM..... | 447 | 150 | 149 | 149 | 224 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER.... | 16.4 | 0 | 0 | 1.3 | 1.3 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.5 | 0 | 0 | 5.2 | 3.7 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.2 | 2.2 | 1.6 | 1.9 | 1.6 | 2.2 | 2.2 | 2.2 | 4.9 | 3.1 | 2.8 | 2.7 |
| 1.0 | 2.0 | 1.2 | 1.6 | 1.5 | 1.8 | 1.8 | 1.7 | 2.8 | 1.9 | 2.1 | 1.6 |
| .1 | .3 | .4 | .2 | .1 | .4 | .3 | .4 | 1.5 | .8 | .5 | .8 |
| .1 | 0 | 0 | .1 | (2) | (2) | .1 | .1 | .6 | .4 | .2 | .3 |
| 1.5 | 1.2 | 1.4 | 1.2 | 1.2 | 1.3 | 1.4 | 1.4 | 1.8 | 1.6 | 1.4 | 1.6 |
| 1.2 | .4 | .7 | .6 | .7 | .8 | .8 | .7 | 1.1 | .8 | .9 | .7 |
| 5.0 | 3.0 | 3.0 | 3.0 | 6.0 | 6.0 | 6.0 | 4.0 | 6.0 | 4.0 | 4.0 | 3.0 |
| 49 | 36 | 57 | 34 | 45 | 43 | 48 | 47 | 62 | 53 | 51 | 53 |
| 47 | 24 | 39 | 28 | 49 | 49 | 49 | 41 | 44 | 44 | 25 | 55 |
| | | | | 11 | 9 | 12 | 15 | 6 | 9 | | |
| | | | | 25 | 19 | 25 | 25 | 25 | 25 | | |
| | | | | 40 | 30 | 37 | 37 | 50 | 38 | | |
| | | | | 50 | 50 | 50 | 50 | 99 | 63 | | |
| | | | | 50 | 90 | 87 | 75 | 137 | 113 | | |
| | | | | 100 | 99 | 99 | 114 | 149 | 149 | | |
| | | | | 348 | 225 | 302 | 273 | 150 | 186 | | |
| 94 | 96 | 130 | 74 | 105 | 98 | 106 | 103 | 117 | 107 | 131 | 91 |
| 60 | 55 | 69 | 43 | 54 | 61 | 61 | 48 | 101 | 66 | 63 | 40 |
| 137 | 94 | 141 | 127 | 190 | 97 | 141 | 139 | 114 | 79 | 111 | 69 |
| 31 | 55 | 73 | 50 | 134 | 56 | 63 | 106 | 40 | 50 | 40 | 31 |
| 139 | 0 | 0 | 74 | 221 | 320 | 142 | 74 | 137 | 136 | 62 | 194 |
| 79 | 0 | 0 | 0 | 111 | 99 | 132 | 39 | 13 | 47 | 0 | 54 |
| 298 | 315 | 298 | 199 | 447 | 600 | 596 | 447 | 447 | 338 | 298 | 298 |
| 671 | 447 | 522 | 348 | 1,043 | 1,273 | 1,192 | 820 | 450 | 559 | 298 | 745 |
| 95 | 91 | 121 | 82 | 115 | 97 | 106 | 101 | 103 | 98 | 105 | 102 |
| 53 | 50 | 63 | 45 | 66 | 60 | 65 | 59 | 46 | 55 | 60 | 69 |
| | | | | 37 | 15 | 37 | 37 | 28 | 28 | | |
| | | | | 75 | 57 | 75 | 75 | 75 | 75 | | |
| | | | | 120 | 75 | 75 | 75 | 90 | 75 | | |
| | | | | 149 | 149 | 149 | 149 | 149 | 149 | | |
| | | | | 149 | 150 | 149 | 149 | 149 | 149 | | |
| | | | | 150 | 180 | 298 | 224 | 162 | 224 | | |
| | | | | 447 | 300 | 298 | 298 | 224 | 338 | | |
| 298 | 315 | 298 | 149 | 447 | 300 | 298 | 298 | 224 | 338 | 298 | 298 |
| 0 | 0 | 0 | 0 | 2.5 | 6.2 | 2.2 | 3.0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 3.2 | 4.0 | 1.5 | 1.4 | 0 | 0 | 0 | 0 |

TABLE 5.01.--YEAST BREADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 93.7 | 17.6 | 88.0 | 95.1 | 97.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 11.2 | 6.5 | 14.7 | 10.1 | 9.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 24.5 | 5.7 | 29.1 | 28.3 | 27.7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 58.0 | 5.5 | 44.1 | 56.7 | 60.1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 4.1 | 2.5 | 3.5 | 4.0 | 4.1 |
| STANDARD DEVIATION.....DO..... | 2.0 | 1.6 | 1.8 | 1.8 | 1.8 |
| MAXIMUM.....DO..... | 14.0 | 8.0 | 9.0 | 10.0 | 11.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 62 | 17 | 33 | 47 | 55 |
| STANDARD DEVIATION.....GM..... | 40 | 12 | 22 | 26 | 29 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 15 | 2 | 7 | 13 | 15 |
| 25TH.....GM..... | 33 | 7 | 17 | 27 | 33 |
| 50TH.....GM..... | 55 | 13 | 29 | 43 | 50 |
| 75TH.....GM..... | 82 | 25 | 46 | 62 | 72 |
| 90TH.....GM..... | 113 | 32 | 63 | 81 | 92 |
| 95TH.....GM..... | 137 | 45 | 75 | 92 | 106 |
| 99TH.....GM..... | 194 | 53 | 97 | 123 | 144 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 57 | 21 | 35 | 45 | 53 |
| STANDARD DEVIATION.....GM..... | 38 | 16 | 23 | 23 | 34 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 64 | 24 | 39 | 52 | 59 |
| STANDARD DEVIATION.....GM..... | 34 | 14 | 22 | 25 | 25 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 79 | 29 | 46 | 59 | 67 |
| STANDARD DEVIATION.....GM..... | 39 | 11 | 21 | 24 | 26 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 838 | 100 | 262 | 277 | 352 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,588 | 160 | 427 | 520 | 578 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 46 | 20 | 28 | 36 | 40 |
| STANDARD DEVIATION.....GM..... | 26 | 11 | 16 | 17 | 19 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 21 | 7 | 11 | 13 | 21 |
| 25TH.....GM..... | 25 | 11 | 21 | 25 | 25 |
| 50TH.....GM..... | 44 | 20 | 25 | 28 | 40 |
| 75TH.....GM..... | 50 | 25 | 40 | 50 | 50 |
| 90TH.....GM..... | 75 | 28 | 50 | 50 | 56 |
| 95TH.....GM..... | 100 | 50 | 50 | 57 | 75 |
| 99TH.....GM..... | 140 | 50 | 80 | 100 | 100 |
| MAXIMUM.....GM..... | 700 | 75 | 193 | 242 | 228 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 638.4 | 0 | 9.5 | 29.2 | 36.0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | .3 | .5 | .5 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 96.9 | 96.4 | 96.2 | 93.7 | 96.1 | 92.9 | 95.5 | 92.9 | 97.2 | 95.3 | 96.7 | 97.4 |
| 9.9 | 13.4 | 9.2 | 14.5 | 11.1 | 15.8 | 8.9 | 11.8 | 7.0 | 7.4 | 7.3 | 8.2 |
| 27.3 | 28.4 | 23.6 | 28.8 | 24.8 | 29.5 | 19.5 | 23.5 | 18.4 | 22.9 | 13.9 | 16.5 |
| 59.6 | 54.5 | 63.4 | 50.3 | 60.2 | 47.6 | 67.2 | 57.7 | 71.7 | 64.9 | 75.5 | 72.7 |
| 4.2 | 3.8 | 4.4 | 3.8 | 4.1 | 3.5 | 4.5 | 3.9 | 4.7 | 4.3 | 5.0 | 4.5 |
| 1.9 | 1.8 | 2.0 | 1.9 | 1.9 | 1.8 | 2.1 | 1.9 | 2.1 | 2.0 | 2.2 | 2.1 |
| 11.0 | 10.0 | 14.0 | 10.0 | 12.0 | 10.0 | 14.0 | 14.0 | 12.0 | 10.0 | 12.0 | 12.0 |
| 69 | 57 | 87 | 56 | 83 | 52 | 80 | 54 | 71 | 54 | 68 | 54 |
| 41 | 32 | 52 | 32 | 50 | 32 | 47 | 33 | 40 | 31 | 36 | 29 |
| 17 | 13 | 17 | 13 | 17 | 12 | 17 | 13 | 17 | 15 | 17 | 14 |
| 40 | 33 | 50 | 31 | 47 | 29 | 47 | 31 | 43 | 32 | 42 | 32 |
| 63 | 52 | 79 | 50 | 75 | 47 | 73 | 49 | 65 | 50 | 65 | 50 |
| 90 | 75 | 115 | 75 | 111 | 70 | 105 | 72 | 92 | 71 | 92 | 72 |
| 121 | 98 | 155 | 98 | 150 | 93 | 138 | 95 | 121 | 94 | 119 | 91 |
| 143 | 113 | 178 | 114 | 177 | 111 | 162 | 111 | 147 | 110 | 136 | 107 |
| 193 | 152 | 250 | 148 | 237 | 150 | 232 | 149 | 188 | 161 | 168 | 144 |
| 64 | 60 | 80 | 58 | 78 | 55 | 66 | 50 | 69 | 43 | 57 | 48 |
| 39 | 42 | 59 | 34 | 47 | 33 | 40 | 29 | 35 | 22 | 31 | 35 |
| 70 | 64 | 87 | 65 | 86 | 59 | 75 | 58 | 63 | 53 | 60 | 52 |
| 34 | 29 | 47 | 30 | 44 | 26 | 37 | 30 | 33 | 26 | 29 | 26 |
| 87 | 73 | 106 | 73 | 105 | 71 | 96 | 68 | 83 | 66 | 78 | 63 |
| 40 | 30 | 49 | 29 | 48 | 30 | 44 | 31 | 38 | 30 | 33 | 27 |
| 767 | 535 | 838 | 390 | 740 | 526 | 549 | 554 | 340 | 400 | 380 | 300 |
| 1,588 | 755 | 1,363 | 644 | 1,269 | 757 | 1,012 | 1,099 | 776 | 658 | 575 | 650 |
| 49 | 44 | 59 | 44 | 61 | 44 | 53 | 41 | 45 | 38 | 41 | 36 |
| 28 | 23 | 35 | 21 | 35 | 22 | 29 | 21 | 23 | 19 | 22 | 17 |
| 22 | 22 | 23 | 22 | 25 | 22 | 22 | 21 | 22 | 20 | 22 | 18 |
| 28 | 25 | 40 | 28 | 44 | 28 | 40 | 25 | 25 | 25 | 25 | 23 |
| 50 | 44 | 50 | 44 | 50 | 44 | 50 | 42 | 44 | 34 | 44 | 28 |
| 50 | 50 | 70 | 50 | 75 | 50 | 56 | 50 | 50 | 50 | 50 | 50 |
| 80 | 57 | 100 | 57 | 100 | 57 | 100 | 56 | 70 | 56 | 56 | 50 |
| 100 | 84 | 114 | 80 | 114 | 80 | 100 | 75 | 88 | 60 | 75 | 57 |
| 142 | 125 | 170 | 113 | 170 | 114 | 150 | 113 | 124 | 105 | 113 | 100 |
| 617 | 510 | 638 | 340 | 700 | 476 | 454 | 454 | 234 | 340 | 280 | 236 |
| 30.5 | 36.1 | 22.2 | 20.7 | 91.2 | 97.9 | 90.9 | 100.5 | 22.3 | 22.5 | 11.0 | 17.9 |
| .4 | .5 | .4 | .4 | .6 | .6 | .4 | .4 | .4 | .3 | .4 | .4 |

TABLE 5.02.--WHITE BREADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 75.7 | 12.8 | 76.1 | 83.7 | 86.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 20.3 | 4.4 | 20.5 | 16.9 | 20.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 23.4 | 4.9 | 25.6 | 29.8 | 30.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 32.0 | 3.4 | 30.0 | 37.0 | 36.2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 3.1 | 2.4 | 3.0 | 3.2 | 3.1 |
| STANDARD DEVIATION.....DO..... | 1.8 | 1.3 | 1.7 | 1.7 | 1.7 |
| MAXIMUM.....DO..... | 14.0 | 7.0 | 9.0 | 9.0 | 10.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 45 | 17 | 27 | 37 | 40 |
| STANDARD DEVIATION.....GM..... | 33 | 12 | 19 | 23 | 25 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 8 | 4 | 4 | 8 | 8 |
| 25TH.....GM..... | 19 | 7 | 13 | 19 | 20 |
| 50TH.....GM..... | 37 | 16 | 24 | 33 | 33 |
| 75TH.....GM..... | 61 | 25 | 37 | 50 | 53 |
| 90TH.....GM..... | 87 | 32 | 53 | 67 | 75 |
| 95TH.....GM..... | 108 | 39 | 65 | 81 | 87 |
| 99TH.....GM..... | 158 | 53 | 93 | 112 | 113 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 51 | 22 | 31 | 40 | 46 |
| STANDARD DEVIATION.....GM..... | 31 | 17 | 20 | 20 | 24 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 56 | 28 | 36 | 48 | 51 |
| STANDARD DEVIATION.....GM..... | 29 | 14 | 20 | 23 | 21 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 68 | 28 | 41 | 52 | 59 |
| STANDARD DEVIATION.....GM..... | 34 | 10 | 19 | 21 | 23 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 838 | 100 | 227 | 242 | 249 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,363 | 160 | 366 | 421 | 485 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 43 | 21 | 27 | 34 | 39 |
| STANDARD DEVIATION.....GM..... | 23 | 12 | 15 | 16 | 17 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 22 | 11 | 11 | 13 | 20 |
| 25TH.....GM..... | 25 | 13 | 22 | 22 | 25 |
| 50TH.....GM..... | 44 | 22 | 25 | 25 | 44 |
| 75TH.....GM..... | 50 | 25 | 25 | 50 | 50 |
| 90TH.....GM..... | 57 | 27 | 50 | 50 | 50 |
| 95TH.....GM..... | 100 | 50 | 50 | 50 | 57 |
| 99TH.....GM..... | 113 | 75 | 57 | 100 | 100 |
| MAXIMUM.....GM..... | 638 | 75 | 160 | 242 | 227 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 265.5 | 0 | 5.5 | 12.6 | 27.0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 0 | .2 | .3 | .5 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 85.6 | 84.4 | 84.8 | 80.8 | 76.5 | 70.7 | 76.8 | 71.5 | 74.1 | 69.9 | 74.1 | 72.3 |
| 21.8 | 23.7 | 19.0 | 24.5 | 21.9 | 23.3 | 20.0 | 20.1 | 15.7 | 17.5 | 14.0 | 13.4 |
| 28.4 | 28.4 | 27.6 | 28.0 | 23.0 | 23.8 | 19.7 | 21.7 | 19.2 | 21.4 | 13.5 | 15.9 |
| 35.5 | 32.2 | 38.1 | 28.3 | 31.6 | 23.6 | 37.2 | 29.7 | 39.2 | 31.1 | 46.5 | 43.1 |
| 3.1 | 3.0 | 3.3 | 2.9 | 3.1 | 2.7 | 3.4 | 3.0 | 3.7 | 3.2 | 4.2 | 3.8 |
| 1.7 | 1.7 | 1.8 | 1.7 | 1.8 | 1.6 | 2.0 | 1.8 | 2.1 | 1.9 | 2.3 | 2.1 |
| 9.0 | 9.0 | 10.0 | 9.0 | 10.0 | 9.0 | 14.0 | 14.0 | 12.0 | 9.0 | 9.0 | 12.0 |
| 48 | 41 | 61 | 40 | 59 | 38 | 56 | 39 | 51 | 38 | 52 | 43 |
| 33 | 27 | 43 | 27 | 43 | 26 | 40 | 27 | 35 | 25 | 33 | 27 |
| 8 | 8 | 15 | 8 | 15 | 8 | 11 | 7 | 8 | 7 | 9 | 8 |
| 24 | 17 | 30 | 17 | 29 | 17 | 25 | 17 | 24 | 17 | 25 | 22 |
| 42 | 33 | 50 | 33 | 49 | 33 | 48 | 33 | 46 | 33 | 50 | 37 |
| 65 | 56 | 81 | 54 | 81 | 50 | 75 | 50 | 73 | 50 | 69 | 58 |
| 91 | 75 | 120 | 77 | 115 | 73 | 110 | 73 | 96 | 73 | 94 | 77 |
| 109 | 92 | 144 | 94 | 142 | 86 | 132 | 89 | 115 | 83 | 110 | 94 |
| 161 | 127 | 193 | 121 | 206 | 123 | 190 | 129 | 163 | 121 | 151 | 129 |
| 55 | 50 | 65 | 49 | 66 | 49 | 58 | 45 | 51 | 42 | 46 | 41 |
| 32 | 28 | 39 | 24 | 39 | 27 | 37 | 23 | 31 | 23 | 25 | 20 |
| 62 | 55 | 71 | 55 | 77 | 52 | 66 | 49 | 55 | 47 | 54 | 48 |
| 30 | 23 | 35 | 25 | 38 | 23 | 32 | 24 | 27 | 22 | 26 | 24 |
| 72 | 63 | 90 | 64 | 90 | 62 | 81 | 59 | 72 | 56 | 68 | 56 |
| 33 | 27 | 43 | 27 | 44 | 25 | 40 | 28 | 33 | 25 | 30 | 26 |
| 667 | 350 | 838 | 340 | 498 | 300 | 454 | 554 | 300 | 245 | 350 | 300 |
| 789 | 725 | 1,363 | 594 | 1,140 | 680 | 1,012 | 908 | 625 | 483 | 575 | 650 |
| 47 | 42 | 56 | 42 | 58 | 42 | 50 | 39 | 42 | 35 | 38 | 34 |
| 24 | 18 | 30 | 18 | 31 | 17 | 25 | 18 | 19 | 15 | 16 | 14 |
| 22 | 22 | 22 | 22 | 22 | 22 | 22 | 20 | 21 | 20 | 22 | 17 |
| 25 | 25 | 44 | 25 | 44 | 25 | 40 | 25 | 25 | 22 | 25 | 22 |
| 50 | 44 | 50 | 44 | 50 | 44 | 50 | 44 | 44 | 25 | 40 | 25 |
| 50 | 50 | 57 | 50 | 66 | 50 | 50 | 50 | 50 | 50 | 50 | 46 |
| 75 | 50 | 100 | 50 | 100 | 50 | 88 | 50 | 50 | 50 | 50 | 50 |
| 100 | 66 | 100 | 60 | 100 | 60 | 100 | 50 | 75 | 50 | 60 | 50 |
| 113 | 100 | 150 | 100 | 150 | 100 | 150 | 100 | 100 | 85 | 100 | 66 |
| 567 | 227 | 638 | 340 | 454 | 227 | 454 | 454 | 227 | 170 | 150 | 113 |
| 16.3 | 15.8 | 9.3 | 9.6 | 29.5 | 23.9 | 43.3 | 34.3 | 12.2 | 11.5 | 6.8 | 8.0 |
| .3 | .3 | .2 | .3 | .3 | .2 | .3 | .2 | .4 | .3 | .4 | .3 |

TABLE 5.03.--WHITE ROLLS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 36.8 | 1.9 | 24.5 | 35.4 | 46.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 25.4 | 1.8 | 19.7 | 26.8 | 30.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 8.8 | .2 | 4.1 | 7.4 | 12.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.6 | 0 | .7 | 1.3 | 3.5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.5 | 1.1 | 1.3 | 1.4 | 1.5 |
| STANDARD DEVIATION.....DO..... | .8 | .3 | .6 | .6 | .8 |
| MAXIMUM.....DO..... | 8.0 | 2.0 | 4.0 | 4.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 25 | 7 | 14 | 18 | 21 |
| STANDARD DEVIATION.....GM..... | 18 | 4 | 9 | 10 | 13 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 9 | | 5 | 7 | 9 |
| 25TH.....GM..... | 13 | | 9 | 13 | 13 |
| 50TH.....GM..... | 19 | | 13 | 13 | 15 |
| 75TH.....GM..... | 28 | | 17 | 23 | 27 |
| 90TH.....GM..... | 47 | | 27 | 31 | 40 |
| 95TH.....GM..... | 60 | | 28 | 38 | 48 |
| 99TH.....GM..... | 93 | | 40 | 50 | 69 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 52 | 19 | 35 | 41 | 46 |
| STANDARD DEVIATION.....GM..... | 30 | 11 | 20 | 18 | 23 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 54 | 16 | 35 | 42 | 46 |
| STANDARD DEVIATION.....GM..... | 26 | 0 | 10 | 11 | 17 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 58 | 0 | 33 | 40 | 43 |
| STANDARD DEVIATION.....GM..... | 26 | 0 | 6 | 12 | 14 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 454 | 40 | 190 | 170 | 227 |
| MAXIMUM OVER 3 DAYS.....GM..... | 628 | 40 | 190 | 185 | 254 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 49 | 19 | 33 | 39 | 42 |
| STANDARD DEVIATION.....GM..... | 26 | 10 | 15 | 15 | 18 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 28 | | 14 | 19 | 28 |
| 25TH.....GM..... | 35 | | 20 | 28 | 28 |
| 50TH.....GM..... | 40 | | 30 | 40 | 40 |
| 75TH.....GM..... | 56 | | 40 | 40 | 40 |
| 90TH.....GM..... | 80 | | 40 | 56 | 64 |
| 95TH.....GM..... | 100 | | 56 | 70 | 80 |
| 99TH.....GM..... | 140 | | 80 | 85 | 100 |
| MAXIMUM.....GM..... | 354 | 40 | 160 | 170 | 227 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 37.6 | 0 | 0 | 2.2 | 2.3 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .2 | 0 | 0 | .3 | .2 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 52.9 | 46.8 | 49.7 | 42.7 | 43.0 | 36.7 | 38.7 | 31.5 | 26.1 | 23.8 | 16.8 | 18.3 |
| 34.1 | 31.6 | 30.6 | 27.9 | 28.0 | 27.1 | 26.6 | 23.1 | 17.9 | 16.1 | 12.0 | 11.8 |
| 14.2 | 11.7 | 13.7 | 12.1 | 11.8 | 8.0 | 8.4 | 6.5 | 6.2 | 6.0 | 3.6 | 5.0 |
| 4.6 | 3.5 | 5.3 | 2.7 | 3.3 | 1.5 | 3.7 | 1.8 | 1.9 | 1.7 | 1.3 | 1.5 |
| 1.6 | 1.5 | 1.7 | 1.5 | 1.6 | 1.4 | 1.6 | 1.4 | 1.6 | 1.5 | 1.5 | 1.6 |
| .8 | .8 | 1.0 | .8 | .9 | .7 | .9 | .8 | .9 | .9 | .8 | .9 |
| 5.0 | 7.0 | 8.0 | 6.0 | 8.0 | 5.0 | 6.0 | 6.0 | 6.0 | 6.0 | 5.0 | 6.0 |
| 26 | 22 | 33 | 24 | 33 | 21 | 28 | 21 | 25 | 21 | 24 | 21 |
| 18 | 15 | 24 | 17 | 24 | 14 | 20 | 16 | 19 | 15 | 20 | 16 |
| 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 7 |
| 13 | 13 | 13 | 13 | 13 | 13 | 13 | 13 | 12 | 13 | 13 | 9 |
| 19 | 15 | 27 | 19 | 27 | 13 | 24 | 13 | 21 | 13 | 19 | 13 |
| 32 | 27 | 40 | 28 | 43 | 27 | 37 | 27 | 31 | 27 | 31 | 27 |
| 53 | 42 | 64 | 47 | 67 | 38 | 53 | 40 | 45 | 40 | 40 | 40 |
| 59 | 53 | 80 | 56 | 80 | 47 | 69 | 49 | 56 | 49 | 53 | 55 |
| 81 | 72 | 120 | 80 | 107 | 75 | 98 | 76 | 105 | 80 | 113 | 82 |
| 52 | 47 | 64 | 50 | 68 | 48 | 60 | 46 | 54 | 41 | 52 | 41 |
| 28 | 22 | 40 | 27 | 42 | 23 | 32 | 23 | 28 | 15 | 28 | 29 |
| 52 | 48 | 68 | 55 | 69 | 50 | 62 | 47 | 55 | 45 | 46 | 45 |
| 23 | 21 | 37 | 29 | 31 | 19 | 29 | 19 | 25 | 20 | 16 | 16 |
| 58 | 50 | 66 | 49 | 76 | 55 | 60 | 56 | 57 | 54 | 72 | 51 |
| 20 | 23 | 28 | 14 | 27 | 28 | 26 | 29 | 31 | 24 | 35 | 19 |
| 300 | 315 | 420 | 284 | 361 | 454 | 340 | 284 | 235 | 205 | 160 | 236 |
| 447 | 542 | 628 | 425 | 480 | 556 | 528 | 567 | 430 | 324 | 340 | 245 |
| 49 | 45 | 59 | 47 | 63 | 46 | 55 | 44 | 48 | 41 | 49 | 40 |
| 24 | 20 | 32 | 23 | 34 | 22 | 29 | 21 | 24 | 17 | 27 | 22 |
| 28 | 28 | 28 | 28 | 28 | 28 | 28 | 25 | 25 | 25 | 21 | 14 |
| 37 | 35 | 40 | 37 | 40 | 37 | 37 | 28 | 28 | 28 | 28 | 28 |
| 40 | 40 | 40 | 40 | 56 | 40 | 40 | 40 | 40 | 40 | 40 | 35 |
| 56 | 40 | 80 | 56 | 80 | 56 | 74 | 50 | 57 | 40 | 64 | 40 |
| 80 | 80 | 100 | 80 | 112 | 80 | 84 | 74 | 80 | 57 | 85 | 57 |
| 84 | 80 | 120 | 100 | 120 | 84 | 113 | 84 | 90 | 80 | 112 | 80 |
| 120 | 113 | 168 | 125 | 170 | 120 | 160 | 113 | 135 | 113 | 120 | 105 |
| 300 | 280 | 280 | 227 | 354 | 236 | 255 | 224 | 170 | 135 | 120 | 236 |
| 7.3 | 3.1 | 0 | 1.5 | 8.8 | 2.0 | 3.9 | 4.6 | 1.0 | .8 | 0 | 0 |
| .4 | .2 | 0 | .2 | .3 | .1 | .1 | .1 | .2 | .1 | 0 | 0 |

TABLE 5.04.--FRENCH, VIENNA, AND ITALIAN BREADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 6.2 | .2 | 2.8 | 4.2 | 6.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.7 | .2 | 2.1 | 3.5 | 5.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.0 | 0 | .6 | .4 | 1.1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .5 | 0 | .1 | .2 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER.... | 1.5 | 1.0 | 1.3 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | 1.0 | .0 | .5 | .5 | .5 |
| MAXIMUM.....DO..... | 12.0 | 1.0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 26 | 9 | 15 | 15 | 16 |
| STANDARD DEVIATION.....GM..... | 31 | 0 | 18 | 17 | 15 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 7 | | | 5 | 6 |
| 25TH.....GM..... | 8 | | | 7 | 8 |
| 50TH.....GM..... | 17 | | | 8 | 10 |
| 75TH.....GM..... | 30 | | | 17 | 19 |
| 90TH.....GM..... | 53 | | | 33 | 33 |
| 95TH.....GM..... | 80 | | | 38 | 38 |
| 99TH.....GM..... | 151 | | | 113 | 95 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 51 | 28 | 30 | 34 | 36 |
| STANDARD DEVIATION.....GM..... | 42 | 0 | 26 | 32 | 24 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 66 | 0 | 36 | 48 | 52 |
| STANDARD DEVIATION.....GM..... | 56 | 0 | 22 | 28 | 38 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 80 | 0 | 94 | 47 | 25 |
| STANDARD DEVIATION.....GM..... | 54 | 0 | 0 | 34 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 567 | 28 | 113 | 227 | 227 |
| MAXIMUM OVER 3 DAYS.....GM..... | 908 | 28 | 284 | 340 | 340 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 54 | 28 | 37 | 38 | 39 |
| STANDARD DEVIATION.....GM..... | 45 | 0 | 30 | 32 | 30 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 18 | | | 15 | 18 |
| 25TH.....GM..... | 25 | | | 20 | 25 |
| 50TH.....GM..... | 40 | | | 25 | 25 |
| 75TH.....GM..... | 60 | | | 40 | 50 |
| 90TH.....GM..... | 113 | | | 61 | 80 |
| 95TH.....GM..... | 113 | | | 113 | 113 |
| 99TH.....GM..... | 227 | | | 227 | 198 |
| MAXIMUM.....GM..... | 567 | 28 | 113 | 227 | 227 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER.... | 19.5 | 0 | 0 | 1.1 | 1.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | 0 | 0 | 1.2 | .7 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 7.2 | 7.6 | 8.1 | 6.8 | 6.9 | 6.0 | 8.1 | 6.3 | 5.8 | 2.9 | 3.9 | 2.6 |
| 6.0 | 6.5 | 6.5 | 5.4 | 5.1 | 4.6 | 6.0 | 4.5 | 3.1 | 2.3 | 2.4 | 1.3 |
| 1.1 | 1.1 | 1.1 | 1.0 | 1.2 | 1.0 | 1.4 | 1.0 | 1.7 | .3 | .2 | .9 |
| .2 | (2) | .5 | .4 | .6 | .4 | .7 | .7 | 1.1 | .3 | 1.3 | .4 |
| 1.3 | 1.2 | 1.4 | 1.3 | 1.5 | 1.4 | 1.6 | 1.6 | 2.2 | 1.6 | 2.8 | 2.3 |
| .6 | .5 | .7 | .7 | 1.0 | .7 | 1.1 | 1.1 | 1.8 | 1.4 | 2.5 | 1.8 |
| 5.0 | 3.0 | 4.0 | 5.0 | 7.0 | 4.0 | 7.0 | 8.0 | 12.0 | 7.0 | 8.0 | 7.0 |
| 21 | 17 | 27 | 22 | 40 | 23 | 32 | 26 | 28 | 23 | 42 | 28 |
| 22 | 14 | 26 | 25 | 48 | 26 | 32 | 33 | 25 | 27 | 45 | 23 |
| 7 | 6 | 7 | 7 | 8 | 7 | 8 | 5 | 7 | 4 | | |
| 8 | 8 | 13 | 8 | 13 | 9 | 13 | 10 | 9 | 8 | | |
| 15 | 13 | 17 | 13 | 23 | 17 | 20 | 17 | 20 | 13 | | |
| 25 | 17 | 28 | 25 | 42 | 30 | 38 | 27 | 33 | 24 | | |
| 40 | 38 | 66 | 57 | 95 | 44 | 76 | 43 | 63 | 57 | | |
| 64 | 45 | 79 | 85 | 151 | 58 | 85 | 108 | 76 | 104 | | |
| 95 | 76 | 140 | 113 | 265 | 151 | 173 | 151 | 148 | 111 | | |
| 52 | 42 | 56 | 41 | 64 | 46 | 66 | 48 | 37 | 43 | 40 | 27 |
| 56 | 31 | 39 | 28 | 38 | 28 | 53 | 49 | 16 | 36 | 15 | 10 |
| 55 | 44 | 76 | 69 | 111 | 51 | 80 | 54 | 65 | 32 | 113 | 65 |
| 39 | 26 | 52 | 47 | 86 | 23 | 52 | 67 | 29 | 9 | 0 | 26 |
| 59 | 85 | 80 | 79 | 129 | 88 | 73 | 72 | 53 | 80 | 89 | 56 |
| 21 | 0 | 34 | 35 | 69 | 57 | 56 | 44 | 34 | 37 | 47 | 18 |
| 567 | 227 | 250 | 340 | 454 | 227 | 454 | 454 | 148 | 170 | 170 | 113 |
| 567 | 255 | 425 | 390 | 908 | 680 | 709 | 908 | 444 | 332 | 454 | 240 |
| 51 | 42 | 60 | 51 | 80 | 50 | 61 | 50 | 38 | 43 | 45 | 37 |
| 49 | 31 | 37 | 41 | 59 | 33 | 46 | 49 | 22 | 30 | 22 | 14 |
| 20 | 15 | 20 | 15 | 20 | 20 | 20 | 15 | 20 | 15 | | |
| 25 | 20 | 36 | 25 | 40 | 25 | 30 | 25 | 25 | 23 | | |
| 40 | 30 | 50 | 40 | 60 | 46 | 50 | 40 | 36 | 36 | | |
| 57 | 50 | 75 | 57 | 113 | 57 | 75 | 57 | 40 | 50 | | |
| 113 | 85 | 113 | 113 | 142 | 113 | 113 | 113 | 60 | 69 | | |
| 120 | 113 | 142 | 113 | 184 | 113 | 142 | 113 | 80 | 113 | | |
| 227 | 151 | 170 | 227 | 227 | 227 | 227 | 227 | 120 | 170 | | |
| 567 | 227 | 227 | 340 | 454 | 227 | 454 | 454 | 120 | 170 | 113 | 70 |
| 0 | 1.2 | 0 | 0 | 4.1 | 4.5 | 1.7 | 3.8 | 2.0 | 0 | 0 | 0 |
| 0 | .6 | 0 | 0 | 1.0 | 1.0 | .3 | .5 | 1.4 | 0 | 0 | 0 |

TABLE 5.05.--WHOLE AND CRACKED WHEAT BREADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 17.8 | 2.4 | 13.7 | 13.5 | 12.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 7.0 | 1.5 | 5.3 | 5.3 | 5.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 5.1 | .6 | 4.9 | 4.2 | 3.9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 5.7 | .4 | 3.4 | 4.0 | 3.1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 2.6 | 1.6 | 2.3 | 2.6 | 2.2 |
| STANDARD DEVIATION.....DO..... | 1.7 | .9 | 1.3 | 1.6 | 1.3 |
| MAXIMUM.....DO..... | 12.0 | 4.0 | 6.0 | 8.0 | 7.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 38 | 12 | 21 | 29 | 29 |
| STANDARD DEVIATION.....GM..... | 30 | 8 | 14 | 20 | 21 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 8 | | 4 | 5 | 8 |
| 25TH.....GM..... | 18 | | 9 | 17 | 15 |
| 50TH.....GM..... | 29 | | 19 | 25 | 19 |
| 75TH.....GM..... | 50 | | 28 | 36 | 44 |
| 90TH.....GM..... | 75 | | 37 | 61 | 56 |
| 95TH.....GM..... | 94 | | 52 | 74 | 65 |
| 99TH.....GM..... | 149 | | 63 | 87 | 109 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 51 | 25 | 27 | 40 | 41 |
| STANDARD DEVIATION.....GM..... | 30 | 14 | 13 | 22 | 17 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 55 | 23 | 37 | 44 | 48 |
| STANDARD DEVIATION.....GM..... | 26 | 7 | 14 | 17 | 19 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 65 | 25 | 33 | 50 | 55 |
| STANDARD DEVIATION.....GM..... | 34 | 10 | 15 | 21 | 20 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 362 | 53 | 128 | 140 | 150 |
| MAXIMUM OVER 3 DAYS.....GM..... | 972 | 103 | 196 | 268 | 373 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 44 | 23 | 27 | 34 | 40 |
| STANDARD DEVIATION.....GM..... | 22 | 10 | 14 | 16 | 16 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 21 | | 11 | 13 | 21 |
| 25TH.....GM..... | 25 | | 14 | 25 | 25 |
| 50TH.....GM..... | 50 | | 25 | 28 | 35 |
| 75TH.....GM..... | 56 | | 28 | 50 | 56 |
| 90TH.....GM..... | 56 | | 56 | 56 | 56 |
| 95TH.....GM..... | 84 | | 56 | 56 | 56 |
| 99TH.....GM..... | 112 | | 56 | 57 | 84 |
| MAXIMUM.....GM..... | 227 | 50 | 100 | 112 | 100 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 13.2 | 0 | .9 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .1 | 0 | .3 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 10.7 | 10.4 | 12.5 | 11.0 | 19.0 | 18.8 | 19.9 | 20.5 | 24.5 | 28.4 | 25.7 | 26.3 |
| 4.7 | 5.9 | 5.5 | 6.0 | 8.5 | 7.9 | 7.0 | 8.4 | 7.3 | 7.6 | 5.3 | 6.8 |
| 3.4 | 2.4 | 3.7 | 3.0 | 5.4 | 6.4 | 6.0 | 5.7 | 5.2 | 7.7 | 5.8 | 5.0 |
| 2.5 | 2.1 | 3.3 | 2.0 | 5.2 | 4.4 | 6.9 | 6.4 | 12.0 | 13.1 | 14.6 | 14.4 |
| 2.4 | 2.2 | 2.4 | 2.1 | 2.4 | 2.3 | 2.7 | 2.5 | 3.3 | 3.2 | 3.7 | 3.4 |
| 1.6 | 1.5 | 1.7 | 1.4 | 1.6 | 1.3 | 1.8 | 1.6 | 1.9 | 1.9 | 2.1 | 2.2 |
| 8.0 | 9.0 | 9.0 | 8.0 | 10.0 | 8.0 | 9.0 | 10.0 | 9.0 | 9.0 | 9.0 | 12.0 |
| 40 | 31 | 50 | 29 | 49 | 32 | 46 | 31 | 49 | 38 | 54 | 38 |
| 32 | 23 | 47 | 20 | 39 | 21 | 36 | 22 | 31 | 26 | 36 | 26 |
| 8 | 8 | 9 | 8 | 12 | 8 | 9 | 8 | 8 | 8 | 9 | 8 |
| 17 | 9 | 19 | 17 | 19 | 17 | 19 | 17 | 25 | 19 | 25 | 19 |
| 28 | 19 | 35 | 19 | 37 | 28 | 37 | 25 | 44 | 33 | 50 | 33 |
| 56 | 44 | 69 | 37 | 67 | 45 | 58 | 44 | 65 | 52 | 73 | 48 |
| 83 | 64 | 112 | 57 | 108 | 60 | 93 | 59 | 92 | 70 | 105 | 75 |
| 96 | 81 | 140 | 66 | 120 | 75 | 117 | 74 | 113 | 86 | 113 | 85 |
| 141 | 112 | 231 | 99 | 200 | 105 | 170 | 101 | 131 | 125 | 168 | 140 |
| 55 | 50 | 61 | 49 | 71 | 48 | 55 | 46 | 66 | 48 | 42 | 44 |
| 42 | 28 | 30 | 27 | 43 | 25 | 26 | 21 | 50 | 25 | 13 | 27 |
| 67 | 53 | 70 | 54 | 75 | 53 | 61 | 46 | 63 | 46 | 54 | 45 |
| 32 | 22 | 37 | 18 | 36 | 18 | 30 | 19 | 26 | 17 | 16 | 16 |
| 74 | 65 | 101 | 58 | 91 | 57 | 80 | 53 | 69 | 55 | 75 | 52 |
| 36 | 21 | 57 | 17 | 42 | 23 | 38 | 23 | 29 | 27 | 32 | 26 |
| 346 | 165 | 324 | 170 | 362 | 224 | 282 | 196 | 255 | 217 | 224 | 154 |
| 636 | 336 | 972 | 302 | 724 | 411 | 728 | 492 | 501 | 584 | 504 | 420 |
| 49 | 43 | 62 | 43 | 62 | 42 | 52 | 38 | 45 | 36 | 44 | 34 |
| 23 | 18 | 33 | 16 | 31 | 16 | 24 | 15 | 21 | 14 | 18 | 15 |
| 25 | 23 | 25 | 23 | 25 | 23 | 24 | 21 | 24 | 21 | 25 | 20 |
| 28 | 28 | 50 | 28 | 50 | 26 | 28 | 25 | 28 | 25 | 28 | 25 |
| 50 | 50 | 56 | 50 | 56 | 50 | 50 | 28 | 50 | 28 | 50 | 28 |
| 56 | 56 | 57 | 56 | 56 | 56 | 56 | 50 | 56 | 50 | 56 | 50 |
| 73 | 56 | 112 | 56 | 112 | 56 | 84 | 56 | 69 | 56 | 56 | 56 |
| 100 | 57 | 112 | 56 | 112 | 56 | 112 | 56 | 84 | 56 | 75 | 56 |
| 112 | 112 | 168 | 75 | 168 | 100 | 112 | 75 | 112 | 63 | 112 | 56 |
| 224 | 138 | 224 | 170 | 227 | 138 | 225 | 140 | 170 | 100 | 125 | 125 |
| 0 | 1.5 | 0 | 0 | 2.3 | 2.6 | 2.3 | 1.0 | 0 | 0 | 0 | 2.6 |
| 0 | .3 | 0 | 0 | .1 | .1 | .1 | (2) | 0 | 0 | 0 | .3 |

TABLE 5.06.--RYE AND PUMPERNICKEL BREADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 7.0 | .4 | 1.9 | 1.7 | 2.0 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.3 | .4 | 1.4 | 1.2 | 1.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.6 | 0 | .5 | .3 | .6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.2 | 0 | .1 | .1 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.9 | 1.0 | 1.6 | 1.5 | 1.7 |
| STANDARD DEVIATION.....DO..... | 1.3 | .0 | 1.1 | 1.0 | 1.0 |
| MAXIMUM.....DO..... | 9.0 | 1.0 | 5.0 | 5.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 28 | 3 | 17 | 17 | 21 |
| STANDARD DEVIATION.....GM..... | 24 | 1 | 17 | 14 | 15 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 8 | | | | |
| 25TH.....GM..... | 17 | | | | |
| 50TH.....GM..... | 17 | | | | |
| 75TH.....GM..... | 33 | | | | |
| 90TH.....GM..... | 55 | | | | |
| 95TH.....GM..... | 73 | | | | |
| 99TH.....GM..... | 112 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 48 | 9 | 25 | 29 | 38 |
| STANDARD DEVIATION.....GM..... | 24 | 2 | 13 | 13 | 22 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 54 | 0 | 54 | 55 | 47 |
| STANDARD DEVIATION.....GM..... | 26 | 0 | 34 | 26 | 24 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 61 | 0 | 37 | 38 | 43 |
| STANDARD DEVIATION.....GM..... | 34 | 0 | 0 | 4 | 7 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 567 | 11 | 123 | 123 | 122 |
| MAXIMUM OVER 3 DAYS.....GM..... | 794 | 11 | 223 | 198 | 200 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 45 | 9 | 31 | 33 | 37 |
| STANDARD DEVIATION.....GM..... | 22 | 2 | 17 | 14 | 16 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 22 | | | | |
| 25TH.....GM..... | 25 | | | | |
| 50TH.....GM..... | 50 | | | | |
| 75TH.....GM..... | 50 | | | | |
| 90TH.....GM..... | 57 | | | | |
| 95TH.....GM..... | 100 | | | | |
| 99TH.....GM..... | 113 | | | | |
| MAXIMUM.....GM..... | 340 | 11 | 73 | 57 | 112 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 1.1 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | (2) | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 2.9 | 2.7 | 4.4 | 3.8 | 7.9 | 5.9 | 12.6 | 9.6 | 11.9 | 9.3 | 8.6 | 7.6 |
| 1.9 | 1.9 | 2.9 | 3.1 | 5.2 | 4.4 | 7.4 | 5.6 | 5.8 | 4.9 | 3.8 | 2.7 |
| .7 | .7 | .8 | .2 | 1.4 | 1.1 | 2.5 | 2.5 | 3.4 | 2.1 | 2.2 | 2.6 |
| .2 | .1 | .6 | .4 | 1.3 | .4 | 2.6 | 1.5 | 2.8 | 2.3 | 2.6 | 2.4 |
| 1.5 | 1.4 | 1.8 | 1.4 | 1.7 | 1.5 | 2.0 | 1.9 | 2.1 | 2.1 | 2.5 | 2.4 |
| .9 | .7 | 1.3 | 1.1 | 1.1 | 1.0 | 1.5 | 1.2 | 1.4 | 1.4 | 1.8 | 1.7 |
| 6.0 | 5.0 | 6.0 | 6.0 | 7.0 | 7.0 | 8.0 | 7.0 | 7.0 | 7.0 | 6.0 | 9.0 |
| 26 | 21 | 34 | 20 | 31 | 21 | 36 | 24 | 32 | 26 | 35 | 29 |
| 33 | 14 | 28 | 15 | 25 | 15 | 32 | 18 | 25 | 20 | 25 | 22 |
| 7 | 7 | 8 | 7 | 8 | 7 | 8 | 8 | 8 | 8 | | 8 |
| 15 | 8 | 17 | 9 | 17 | 15 | 17 | 15 | 17 | 8 | | 17 |
| 17 | 17 | 17 | 17 | 17 | 17 | 23 | 17 | 25 | 17 | | 25 |
| 33 | 31 | 50 | 17 | 36 | 25 | 48 | 33 | 38 | 33 | | 33 |
| 41 | 33 | 79 | 48 | 67 | 36 | 67 | 46 | 50 | 53 | | 67 |
| 57 | 50 | 93 | 50 | 83 | 50 | 100 | 55 | 81 | 67 | | 83 |
| 265 | 92 | 108 | 83 | 112 | 95 | 183 | 88 | 144 | 77 | | 100 |
| 43 | 45 | 56 | 44 | 53 | 45 | 56 | 45 | 47 | 41 | 46 | 39 |
| 20 | 22 | 38 | 16 | 25 | 19 | 26 | 21 | 20 | 31 | 22 | 17 |
| 58 | 46 | 83 | 43 | 69 | 54 | 61 | 46 | 54 | 41 | 70 | 43 |
| 39 | 13 | 39 | 20 | 32 | 28 | 25 | 18 | 19 | 16 | 46 | 11 |
| 79 | 71 | 74 | 55 | 69 | 46 | 78 | 49 | 61 | 49 | 56 | 48 |
| 77 | 21 | 19 | 14 | 25 | 15 | 43 | 24 | 33 | 20 | 12 | 29 |
| 567 | 100 | 227 | 100 | 225 | 170 | 300 | 340 | 255 | 200 | 163 | 150 |
| 794 | 275 | 325 | 250 | 468 | 296 | 700 | 590 | 432 | 294 | 300 | 300 |
| 51 | 44 | 58 | 42 | 54 | 41 | 52 | 39 | 47 | 36 | 42 | 36 |
| 41 | 17 | 30 | 16 | 21 | 16 | 24 | 18 | 21 | 14 | 23 | 16 |
| 22 | 22 | 25 | 22 | 22 | 22 | 24 | 22 | 22 | 22 | | 22 |
| 25 | 25 | 46 | 25 | 50 | 25 | 44 | 25 | 25 | 25 | | 25 |
| 50 | 50 | 50 | 50 | 50 | 50 | 50 | 44 | 50 | 28 | | 25 |
| 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | | 50 |
| 57 | 56 | 100 | 50 | 100 | 50 | 100 | 50 | 57 | 50 | | 57 |
| 113 | 57 | 100 | 64 | 100 | 56 | 100 | 56 | 100 | 50 | | 75 |
| 227 | 100 | 150 | 100 | 113 | 113 | 128 | 88 | 113 | 50 | | 75 |
| 340 | 100 | 227 | 100 | 142 | 113 | 200 | 227 | 113 | 113 | 150 | 75 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1.1 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | .1 | 0 | 0 | 0 | 0 |

TABLE 5.07.--HIGH FIBER, WHEAT GERM, BRAN, OATMEAL, AND MULTIGRAIN BREADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 1.6 | .4 | .9 | .8 | 1.0 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | .7 | .4 | .4 | .5 | .5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .5 | 0 | .4 | .1 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .4 | 0 | .1 | .1 | .3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 2.4 | 1.0 | 1.8 | 1.7 | 1.8 |
| STANDARD DEVIATION.....DD..... | 1.6 | .0 | .6 | 1.1 | .9 |
| MAXIMUM.....DD..... | 9.0 | 1.0 | 3.0 | 5.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 38 | 7 | 15 | 20 | 25 |
| STANDARD DEVIATION.....GM..... | 31 | 4 | 8 | 20 | 13 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 8 | | | | |
| 25TH.....GM..... | 17 | | | | |
| 50TH.....GM..... | 31 | | | | |
| 75TH.....GM..... | 49 | | | | |
| 90TH.....GM..... | 79 | | | | |
| 95TH.....GM..... | 102 | | | | |
| 99TH.....GM..... | 139 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 53 | 22 | 22 | 29 | 47 |
| STANDARD DEVIATION.....GM..... | 31 | 11 | 5 | 18 | 16 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 61 | 0 | 34 | 40 | 44 |
| STANDARD DEVIATION.....GM..... | 29 | 0 | 6 | 7 | 18 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 72 | 0 | 15 | 62 | 41 |
| STANDARD DEVIATION.....GM..... | 36 | 0 | 0 | 11 | 7 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 248 | 34 | 56 | 84 | 81 |
| MAXIMUM OVER 3 DAYS.....GM..... | 627 | 34 | 81 | 221 | 156 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 48 | 22 | 25 | 36 | 41 |
| STANDARD DEVIATION.....GM..... | 24 | 11 | 13 | 14 | 15 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 23 | | | | |
| 25TH.....GM..... | 28 | | | | |
| 50TH.....GM..... | 50 | | | | |
| 75TH.....GM..... | 57 | | | | |
| 90TH.....GM..... | 68 | | | | |
| 95TH.....GM..... | 100 | | | | |
| 99TH.....GM..... | 136 | | | | |
| MAXIMUM.....GM..... | 200 | 34 | 56 | 56 | 75 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 0 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 0 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| .7 | 1.0 | .8 | .7 | 1.4 | 1.4 | 1.9 | 2.4 | 2.6 | 3.4 | 2.1 | 2.4 |
| .4 | .6 | .4 | .3 | .5 | .6 | .7 | 1.2 | 1.2 | 1.3 | 1.2 | 1.0 |
| .3 | .2 | .1 | .2 | .5 | .6 | .7 | .7 | .4 | .7 | .2 | 1.1 |
| (2) | .2 | .3 | .2 | .4 | .2 | .6 | .5 | 1.0 | 1.4 | .8 | .3 |
| 1.7 | 1.6 | 3.0 | 2.4 | 2.6 | 2.2 | 2.8 | 2.2 | 2.8 | 2.6 | 3.0 | 2.6 |
| 1.0 | .8 | 2.5 | 1.4 | 1.7 | 1.4 | 1.9 | 1.4 | 2.0 | 1.7 | 2.3 | 1.5 |
| 5.0 | 3.0 | 8.0 | 5.0 | 9.0 | 7.0 | 9.0 | 6.0 | 8.0 | 8.0 | 7.0 | 7.0 |
| 29 | 23 | 55 | 40 | 59 | 31 | 49 | 31 | 59 | 41 | 41 | 32 |
| 19 | 12 | 38 | 26 | 42 | 21 | 30 | 22 | 51 | 34 | 43 | 17 |
| | | | | 17 | 8 | 9 | 8 | | 8 | | |
| | | | | 25 | 15 | 25 | 15 | | 15 | | |
| | | | | 52 | 24 | 45 | 23 | | 37 | | |
| | | | | 89 | 45 | 65 | 39 | | 57 | | |
| | | | | 102 | 65 | 88 | 59 | | 66 | | |
| | | | | 139 | 79 | 121 | 79 | | 99 | | |
| | | | | 209 | 83 | 136 | 95 | | 170 | | |
| 49 | 54 | 72 | 51 | 77 | 39 | 79 | 49 | 47 | 42 | 48 | 60 |
| 30 | 25 | 47 | 12 | 33 | 15 | 46 | 23 | 16 | 25 | 33 | 27 |
| 60 | 36 | 68 | 54 | 79 | 61 | 73 | 54 | 68 | 60 | 72 | 52 |
| 24 | 11 | 0 | 8 | 34 | 29 | 37 | 20 | 41 | 18 | 0 | 16 |
| 64 | 42 | 99 | 80 | 105 | 54 | 75 | 59 | 112 | 65 | 76 | 55 |
| 0 | 9 | 19 | 2 | 41 | 13 | 26 | 24 | 33 | 38 | 53 | 23 |
| 130 | 112 | 146 | 112 | 227 | 173 | 204 | 136 | 248 | 204 | 168 | 126 |
| 200 | 169 | 357 | 248 | 627 | 250 | 408 | 316 | 496 | 510 | 392 | 270 |
| 50 | 44 | 54 | 51 | 69 | 42 | 52 | 42 | 62 | 47 | 40 | 36 |
| 23 | 16 | 21 | 16 | 32 | 22 | 26 | 17 | 28 | 17 | 15 | 13 |
| | | | | 25 | 23 | 23 | 23 | | 24 | | |
| | | | | 50 | 25 | 31 | 25 | | 31 | | |
| | | | | 68 | 34 | 50 | 46 | | 46 | | |
| | | | | 92 | 50 | 68 | 56 | | 66 | | |
| | | | | 108 | 68 | 75 | 62 | | 68 | | |
| | | | | 113 | 68 | 100 | 68 | | 68 | | |
| | | | | 200 | 136 | 136 | 102 | | 68 | | |
| 112 | 69 | 100 | 68 | 200 | 136 | 170 | 136 | 124 | 84 | 56 | 62 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 5.08.--ENGLISH MUFFINS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.0 | 0 | 2.7 | 4.2 | 4.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.2 | 0 | 2.4 | 2.9 | 3.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.3 | 0 | .3 | 1.0 | 1.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .5 | 0 | .1 | .4 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1</u> /NUMBER..... | 1.5 | 0 | 1.2 | 1.4 | 1.3 |
| STANDARD DEVIATION.....CO..... | .7 | 0 | .5 | .6 | .5 |
| MAXIMUM.....DO..... | 5.0 | 0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1</u> /GM..... | 28 | 0 | 15 | 20 | 25 |
| STANDARD DEVIATION.....GM..... | 18 | 0 | 11 | 13 | 17 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> / | | | | | |
| 5TH.....GM..... | 9 | | | 8 | 9 |
| 25TH.....GM..... | 19 | | | 9 | 17 |
| 50TH.....GM..... | 19 | | | 19 | 19 |
| 75TH.....GM..... | 38 | | | 19 | 33 |
| 90TH.....GM..... | 57 | | | 38 | 38 |
| 95TH.....GM..... | 57 | | | 50 | 57 |
| 99TH.....GM..... | 95 | | | 76 | 114 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 60 | 0 | 38 | 42 | 62 |
| STANDARD DEVIATION.....GM..... | 27 | 0 | 22 | 15 | 34 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 60 | 0 | 38 | 43 | 60 |
| STANDARD DEVIATION.....GM..... | 24 | 0 | 12 | 17 | 35 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 57 | 0 | 52 | 43 | 30 |
| STANDARD DEVIATION.....GM..... | 23 | 0 | 0 | 19 | 14 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 228 | 0 | 113 | 113 | 228 |
| MAXIMUM OVER 3 DAYS.....GM..... | 528 | 0 | 157 | 227 | 341 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1</u> /GM..... | 57 | 0 | 39 | 43 | 58 |
| STANDARD DEVIATION.....GM..... | 25 | 0 | 20 | 18 | 36 |
| PERCENTILES: <u>1</u> / | | | | | |
| 5TH.....GM..... | 28 | | | 14 | 25 |
| 25TH.....GM..... | 50 | | | 28 | 28 |
| 50TH.....GM..... | 57 | | | 50 | 57 |
| 75TH.....GM..... | 57 | | | 57 | 57 |
| 90TH.....GM..... | 100 | | | 57 | 114 |
| 95TH.....GM..... | 114 | | | 57 | 114 |
| 99TH.....GM..... | 142 | | | 113 | 227 |
| MAXIMUM.....GM..... | 228 | 0 | 113 | 113 | 228 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> / | | | | | |
| EATING OCCASIONS.....NUMBER..... | 0 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 0 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 4.7 | 4.6 | 4.9 | 4.4 | 5.9 | 5.5 | 5.2 | 5.9 | 4.7 | 4.8 | 1.3 | 3.3 |
| 3.2 | 3.4 | 2.9 | 2.7 | 3.8 | 3.9 | 3.1 | 3.7 | 2.8 | 2.9 | .9 | 1.9 |
| 1.3 | .9 | 1.4 | 1.6 | 1.4 | 1.4 | 1.4 | 1.4 | 1.5 | 1.4 | 0 | .7 |
| .3 | .3 | .5 | .1 | .6 | .2 | .7 | .8 | .4 | .5 | .4 | .7 |
| 1.4 | 1.4 | 1.5 | 1.4 | 1.5 | 1.4 | 1.6 | 1.6 | 1.5 | 1.5 | 1.6 | 1.7 |
| .7 | .6 | .7 | .6 | .7 | .6 | .8 | .8 | .6 | .7 | .9 | .8 |
| 4.0 | 3.0 | 4.0 | 3.0 | 4.0 | 4.0 | 5.0 | 5.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| 31 | 27 | 39 | 29 | 32 | 26 | 31 | 28 | 30 | 27 | 24 | 29 |
| 24 | 15 | 26 | 19 | 20 | 14 | 16 | 18 | 15 | 19 | 11 | 16 |
| 9 | 9 | 17 | 9 | 9 | 10 | 17 | 9 | 17 | 9 | | |
| 19 | 19 | 19 | 19 | 19 | 19 | 19 | 17 | 19 | 17 | | |
| 19 | 19 | 33 | 19 | 19 | 19 | 25 | 19 | 19 | 19 | | |
| 38 | 38 | 55 | 38 | 38 | 36 | 38 | 38 | 36 | 36 | | |
| 57 | 50 | 71 | 38 | 57 | 48 | 57 | 55 | 57 | 50 | | |
| 76 | 57 | 105 | 67 | 76 | 57 | 57 | 57 | 57 | 57 | | |
| 85 | 76 | 114 | 133 | 114 | 75 | 76 | 95 | 67 | 113 | | |
| 60 | 64 | 77 | 60 | 69 | 59 | 65 | 55 | 58 | 49 | 52 | 57 |
| 25 | 28 | 43 | 22 | 34 | 21 | 25 | 23 | 16 | 18 | 3 | 20 |
| 70 | 61 | 76 | 68 | 61 | 60 | 59 | 56 | 63 | 51 | 0 | 59 |
| 23 | 30 | 30 | 32 | 24 | 19 | 24 | 22 | 18 | 14 | 0 | 18 |
| 88 | 49 | 84 | 36 | 67 | 57 | 54 | 53 | 54 | 67 | 42 | 47 |
| 38 | 9 | 27 | 0 | 25 | 7 | 10 | 22 | 3 | 26 | 0 | 13 |
| 228 | 171 | 228 | 228 | 228 | 171 | 171 | 171 | 114 | 114 | 57 | 100 |
| 528 | 228 | 397 | 399 | 342 | 227 | 342 | 342 | 200 | 340 | 125 | 171 |
| 66 | 59 | 78 | 61 | 64 | 58 | 58 | 52 | 58 | 53 | 46 | 52 |
| 30 | 28 | 38 | 23 | 29 | 19 | 23 | 21 | 15 | 20 | 10 | 19 |
| 28 | 28 | 40 | 28 | 28 | 28 | 28 | 25 | 50 | 28 | | |
| 51 | 50 | 57 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | | |
| 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 50 | | |
| 57 | 57 | 112 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | | |
| 114 | 114 | 114 | 100 | 114 | 57 | 100 | 57 | 57 | 57 | | |
| 114 | 114 | 114 | 113 | 114 | 112 | 114 | 100 | 100 | 113 | | |
| 150 | 171 | 228 | 171 | 171 | 114 | 114 | 142 | 114 | 114 | | |
| 228 | 171 | 228 | 171 | 228 | 171 | 171 | 171 | 114 | 114 | 57 | 100 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 5.09.--BISCUITS OTHER THAN REFRIGERATOR TYPE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 7.5 | .8 | 6.6 | 7.3 | 9.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.1 | .6 | 4.9 | 5.6 | 7.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.5 | .1 | 1.3 | 1.3 | 1.7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .9 | 0 | .5 | .4 | .5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.6 | 1.4 | 1.5 | 1.4 | 1.4 |
| STANDARD DEVIATION.....DO..... | 1.0 | .8 | 1.0 | .9 | .8 |
| MAXIMUM.....DO..... | 8.0 | 3.0 | 6.0 | 6.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 35 | 8 | 17 | 21 | 25 |
| STANDARD DEVIATION.....GM..... | 31 | 7 | 14 | 14 | 17 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 9 | | 5 | 5 | 9 |
| 25TH.....GM..... | 19 | | 9 | 9 | 13 |
| 50TH.....GM..... | 25 | | 13 | 19 | 19 |
| 75TH.....GM..... | 42 | | 25 | 25 | 28 |
| 90TH.....GM..... | 75 | | 38 | 38 | 49 |
| 95TH.....GM..... | 93 | | 47 | 51 | 63 |
| 99TH.....GM..... | 168 | | 75 | 78 | 89 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 71 | 16 | 42 | 50 | 56 |
| STANDARD DEVIATION.....GM..... | 53 | 7 | 29 | 27 | 27 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 73 | 35 | 29 | 42 | 66 |
| STANDARD DEVIATION.....GM..... | 42 | 0 | 18 | 20 | 29 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 81 | 0 | 47 | 55 | 48 |
| STANDARD DEVIATION.....GM..... | 46 | 0 | 14 | 20 | 19 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 748 | 56 | 156 | 168 | 192 |
| MAXIMUM OVER 3 DAYS.....GM..... | 888 | 70 | 224 | 280 | 268 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 67 | 19 | 35 | 44 | 53 |
| STANDARD DEVIATION.....GM..... | 43 | 8 | 24 | 25 | 28 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 28 | | 11 | 14 | 28 |
| 25TH.....GM..... | 38 | | 21 | 28 | 38 |
| 50TH.....GM..... | 56 | | 28 | 38 | 56 |
| 75TH.....GM..... | 84 | | 38 | 56 | 60 |
| 90TH.....GM..... | 114 | | 57 | 84 | 84 |
| 95TH.....GM..... | 140 | | 76 | 84 | 112 |
| 99TH.....GM..... | 228 | | 113 | 114 | 170 |
| MAXIMUM.....GM..... | 448 | 28 | 156 | 152 | 192 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 18.2 | 0 | 0 | .8 | 1.7 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | 0 | .5 | .7 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 9.3 | 7.1 | 8.3 | 6.9 | 6.8 | 6.9 | 7.9 | 7.7 | 10.0 | 6.8 | 8.5 | 5.3 |
| 6.3 | 5.3 | 5.6 | 5.1 | 4.7 | 5.1 | 4.8 | 5.0 | 6.3 | 4.5 | 4.3 | 2.7 |
| 2.3 | 1.3 | 1.4 | 1.0 | 1.4 | 1.3 | 1.9 | 1.7 | 1.6 | 1.2 | 2.7 | 1.2 |
| .7 | .5 | 1.3 | .7 | .7 | .6 | 1.2 | 1.0 | 2.1 | 1.1 | 1.5 | 1.4 |
| 1.5 | 1.4 | 1.6 | 1.5 | 1.5 | 1.4 | 1.8 | 1.6 | 1.9 | 1.8 | 2.2 | 2.0 |
| .7 | .7 | 1.0 | .8 | .9 | .8 | 1.1 | 1.0 | 1.4 | 1.4 | 1.3 | 1.5 |
| 6.0 | 4.0 | 5.0 | 5.0 | 6.0 | 5.0 | 6.0 | 7.0 | 7.0 | 8.0 | 7.0 | 8.0 |
| 33 | 28 | 47 | 29 | 47 | 31 | 49 | 32 | 48 | 32 | 55 | 34 |
| 23 | 17 | 43 | 24 | 38 | 23 | 42 | 26 | 43 | 28 | 40 | 28 |
| 9 | 9 | 9 | 9 | 13 | 9 | 9 | 9 | 9 | 9 | | 9 |
| 19 | 19 | 19 | 13 | 25 | 14 | 24 | 14 | 19 | 13 | | 14 |
| 28 | 25 | 37 | 19 | 37 | 25 | 37 | 24 | 28 | 25 | | 25 |
| 42 | 37 | 63 | 40 | 56 | 38 | 63 | 38 | 67 | 35 | | 45 |
| 56 | 56 | 89 | 57 | 93 | 56 | 93 | 65 | 103 | 80 | | 56 |
| 75 | 65 | 131 | 84 | 112 | 79 | 126 | 76 | 114 | 103 | | 122 |
| 129 | 89 | 240 | 114 | 189 | 127 | 196 | 139 | 224 | 149 | | 126 |
| 74 | 67 | 85 | 55 | 104 | 72 | 87 | 62 | 83 | 58 | 84 | 53 |
| 39 | 35 | 58 | 34 | 95 | 55 | 51 | 40 | 52 | 27 | 32 | 27 |
| 68 | 59 | 108 | 77 | 96 | 63 | 92 | 60 | 92 | 57 | 122 | 52 |
| 21 | 23 | 57 | 32 | 43 | 33 | 47 | 33 | 48 | 27 | 65 | 17 |
| 75 | 61 | 98 | 72 | 94 | 63 | 106 | 72 | 98 | 75 | 81 | 67 |
| 47 | 13 | 62 | 24 | 40 | 25 | 59 | 36 | 57 | 41 | 32 | 35 |
| 240 | 224 | 520 | 210 | 748 | 380 | 468 | 380 | 336 | 168 | 308 | 169 |
| 660 | 266 | 820 | 354 | 748 | 420 | 888 | 504 | 672 | 448 | 532 | 378 |
| 68 | 61 | 87 | 60 | 93 | 64 | 83 | 58 | 74 | 53 | 75 | 51 |
| 37 | 29 | 54 | 36 | 58 | 43 | 46 | 34 | 43 | 26 | 38 | 25 |
| 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | | 28 |
| 42 | 38 | 56 | 38 | 56 | 38 | 56 | 28 | 56 | 28 | | 28 |
| 56 | 56 | 75 | 56 | 84 | 56 | 76 | 56 | 60 | 56 | | 56 |
| 84 | 76 | 112 | 76 | 112 | 76 | 112 | 76 | 84 | 76 | | 60 |
| 112 | 84 | 168 | 112 | 152 | 113 | 128 | 84 | 120 | 84 | | 76 |
| 140 | 114 | 224 | 120 | 170 | 114 | 168 | 120 | 152 | 114 | | 84 |
| 228 | 168 | 280 | 196 | 336 | 240 | 240 | 168 | 227 | 120 | | 168 |
| 240 | 190 | 284 | 210 | 448 | 380 | 360 | 380 | 240 | 120 | 224 | 168 |
| 2.5 | .7 | 0 | 0 | 3.8 | 0 | 1.5 | 4.9 | 0 | 2.2 | 0 | 0 |
| .9 | .3 | 0 | 0 | .9 | 0 | .2 | .6 | 0 | 1.0 | 0 | 0 |

TABLE 5.10.--REFRIGERATOR-TYPE BISCUITS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.5 | .1 | 3.1 | 4.6 | 5.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.6 | .1 | 2.5 | 3.5 | 4.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .7 | 0 | .6 | .9 | .9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | 0 | .1 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.4 | 1.0 | 1.2 | 1.3 | 1.3 |
| STANDARD DEVIATION.....DO..... | .8 | .0 | .5 | .7 | .7 |
| MAXIMUM.....DO..... | 7.0 | 1.0 | 3.0 | 5.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 24 | 7 | 10 | 14 | 16 |
| STANDARD DEVIATION.....GM..... | 18 | 0 | 5 | 9 | 11 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 7 | | | 7 | 7 |
| 25TH.....GM..... | 13 | | | 7 | 7 |
| 50TH.....GM..... | 20 | | | 13 | 13 |
| 75TH.....GM..... | 28 | | | 19 | 20 |
| 90TH.....GM..... | 47 | | | 27 | 27 |
| 95TH.....GM..... | 60 | | | 27 | 40 |
| 99TH.....GM..... | 82 | | | 40 | 60 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 52 | 20 | 28 | 33 | 38 |
| STANDARD DEVIATION.....GM..... | 30 | 0 | 14 | 15 | 20 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 53 | 0 | 20 | 35 | 36 |
| STANDARD DEVIATION.....GM..... | 26 | 0 | 7 | 13 | 12 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 59 | 0 | 0 | 35 | 56 |
| STANDARD DEVIATION.....GM..... | 29 | 0 | 0 | 12 | 6 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 340 | 20 | 93 | 80 | 120 |
| MAXIMUM OVER 3 DAYS.....GM..... | 553 | 20 | 93 | 160 | 180 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 50 | 20 | 25 | 32 | 36 |
| STANDARD DEVIATION.....GM..... | 28 | 0 | 13 | 14 | 18 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 20 | | | 20 | 20 |
| 25TH.....GM..... | 40 | | | 20 | 20 |
| 50TH.....GM..... | 40 | | | 30 | 40 |
| 75TH.....GM..... | 60 | | | 40 | 40 |
| 90TH.....GM..... | 80 | | | 50 | 60 |
| 95TH.....GM..... | 100 | | | 60 | 80 |
| 99TH.....GM..... | 125 | | | 80 | 100 |
| MAXIMUM.....GM..... | 340 | 20 | 93 | 80 | 100 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 4.3 | 0 | 0 | 0 | 1.2 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .2 | 0 | 0 | 0 | 1.0 |

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 4.1 | 3.8 | 3.9 | 3.1 | 4.3 | 3.6 | 3.4 | 2.9 | 3.7 | 3.0 | 3.3 | 2.9 |
| 3.1 | 2.9 | 3.1 | 2.2 | 2.9 | 3.0 | 2.5 | 2.0 | 2.2 | 1.4 | 1.0 | .9 |
| .9 | .7 | .6 | .8 | 1.1 | .4 | .6 | .6 | .9 | 1.3 | .7 | 1.4 |
| .1 | .1 | .3 | .1 | .2 | .1 | .3 | .3 | .6 | .3 | 1.5 | .5 |
| 1.3 | 1.3 | 1.4 | 1.3 | 1.4 | 1.3 | 1.4 | 1.5 | 1.7 | 1.8 | 3.2 | 2.1 |
| .7 | .7 | .9 | .6 | .7 | .6 | .7 | .9 | 1.1 | .8 | 2.0 | 1.0 |
| 4.0 | 5.0 | 6.0 | 4.0 | 4.0 | 4.0 | 4.0 | 6.0 | 7.0 | 3.0 | 7.0 | 5.0 |
| 24 | 20 | 29 | 20 | 34 | 19 | 29 | 21 | 42 | 26 | 54 | 27 |
| 18 | 13 | 22 | 13 | 18 | 11 | 17 | 15 | 38 | 19 | 34 | 14 |
| 7 | 7 | 7 | | 13 | 7 | 13 | 7 | | 7 | | |
| 13 | 13 | 13 | | 20 | 13 | 13 | 13 | | 13 | | |
| 19 | 19 | 20 | | 33 | 13 | 27 | 13 | | 19 | | |
| 27 | 27 | 33 | | 47 | 20 | 40 | 27 | | 40 | | |
| 40 | 40 | 67 | | 60 | 36 | 57 | 40 | | 40 | | |
| 53 | 42 | 76 | | 67 | 47 | 67 | 53 | | 53 | | |
| 80 | 73 | 107 | | 80 | 58 | 80 | 80 | | 112 | | |
| 53 | 47 | 63 | 45 | 79 | 47 | 67 | 42 | 65 | 43 | 48 | 65 |
| 32 | 26 | 38 | 20 | 36 | 23 | 32 | 19 | 27 | 13 | 18 | 32 |
| 61 | 51 | 74 | 47 | 74 | 46 | 60 | 46 | 75 | 45 | 70 | 39 |
| 36 | 23 | 38 | 20 | 22 | 19 | 24 | 19 | 31 | 16 | 7 | 17 |
| 49 | 43 | 75 | 71 | 63 | 48 | 59 | 44 | 111 | 59 | 83 | 40 |
| 9 | 3 | 18 | 0 | 21 | 8 | 16 | 24 | 44 | 28 | 24 | 17 |
| 340 | 120 | 227 | 113 | 200 | 140 | 170 | 180 | 220 | 112 | 142 | 128 |
| 440 | 220 | 320 | 213 | 325 | 173 | 300 | 380 | 553 | 336 | 369 | 200 |
| 53 | 46 | 63 | 46 | 72 | 45 | 62 | 41 | 74 | 45 | 50 | 40 |
| 37 | 24 | 37 | 20 | 30 | 22 | 28 | 18 | 33 | 21 | 15 | 25 |
| 20 | 20 | 20 | | 40 | 20 | 20 | 20 | | 20 | | |
| 40 | 20 | 40 | | 50 | 20 | 40 | 25 | | 40 | | |
| 40 | 40 | 60 | | 60 | 40 | 60 | 40 | | 40 | | |
| 60 | 60 | 80 | | 80 | 60 | 80 | 50 | | 50 | | |
| 80 | 80 | 100 | | 120 | 80 | 100 | 60 | | 60 | | |
| 100 | 100 | 120 | | 120 | 80 | 120 | 80 | | 100 | | |
| 240 | 120 | 227 | | 160 | 113 | 170 | 85 | | 112 | | |
| 340 | 120 | 227 | 113 | 200 | 120 | 170 | 113 | 180 | 112 | 85 | 128 |
| 0 | 0 | 0 | 0 | 0 | 0 | 1.5 | 1.5 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | .7 | .5 | 0 | 0 | 0 | 0 |

TABLE 5.11.--CORNBREAD
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 10.1 | 2.0 | 7.9 | 9.9 | 10.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 6.9 | 1.5 | 4.9 | 6.6 | 7.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.4 | .2 | 2.1 | 2.5 | 2.3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .8 | .3 | .9 | .7 | .4 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.5 | 1.7 | 1.6 | 1.5 | 1.4 |
| STANDARD DEVIATION.....DO..... | .9 | 1.4 | .8 | .8 | .7 |
| MAXIMUM.....DO..... | 9.0 | 5.0 | 4.0 | 4.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 49 | 15 | 28 | 33 | 33 |
| STANDARD DEVIATION.....GM..... | 51 | 13 | 25 | 29 | 28 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 9 | | 3 | 6 | 6 |
| 25TH.....GM..... | 22 | | 11 | 13 | 15 |
| 50TH.....GM..... | 33 | | 22 | 26 | 26 |
| 75TH.....GM..... | 57 | | 37 | 47 | 40 |
| 90TH.....GM..... | 100 | | 63 | 66 | 69 |
| 95TH.....GM..... | 138 | | 83 | 91 | 84 |
| 99TH.....GM..... | 266 | | 102 | 133 | 121 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 103 | 26 | 56 | 71 | 77 |
| STANDARD DEVIATION.....GM..... | 87 | 13 | 47 | 63 | 61 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 106 | 61 | 54 | 72 | 67 |
| STANDARD DEVIATION.....GM..... | 80 | 0 | 25 | 45 | 32 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 116 | 30 | 63 | 72 | 106 |
| STANDARD DEVIATION.....GM..... | 94 | 16 | 39 | 33 | 47 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,062 | 78 | 266 | 398 | 531 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,188 | 144 | 448 | 515 | 681 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 98 | 25 | 53 | 67 | 73 |
| STANDARD DEVIATION.....GM..... | 78 | 16 | 39 | 45 | 53 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 20 | | 8 | 9 | 17 |
| 25TH.....GM..... | 46 | | 31 | 33 | 40 |
| 50TH.....GM..... | 78 | | 40 | 60 | 75 |
| 75TH.....GM..... | 133 | | 78 | 78 | 78 |
| 90TH.....GM..... | 180 | | 100 | 128 | 133 |
| 95TH.....GM..... | 266 | | 120 | 149 | 156 |
| 99TH.....GM..... | 398 | | 199 | 199 | 224 |
| MAXIMUM.....GM..... | 797 | 78 | 266 | 266 | 531 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 142.0 | 0 | 3.1 | 6.6 | 5.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.4 | 0 | 2.3 | 2.6 | 2.3 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 10.5 | 10.4 | 12.4 | 9.1 | 7.2 | 8.7 | 10.6 | 11.1 | 14.8 | 12.9 | 14.1 | 11.2 |
| 7.0 | 7.4 | 8.8 | 6.9 | 5.6 | 6.0 | 7.3 | 7.6 | 8.8 | 7.9 | 8.7 | 6.3 |
| 2.9 | 2.1 | 2.7 | 1.7 | 1.3 | 2.4 | 2.4 | 2.6 | 4.0 | 3.0 | 3.4 | 3.0 |
| .6 | .9 | .8 | .4 | .3 | .3 | .9 | .9 | 2.0 | 2.0 | 2.0 | 2.0 |
| 1.5 | 1.5 | 1.4 | 1.3 | 1.3 | 1.4 | 1.5 | 1.5 | 1.8 | 1.8 | 1.8 | 2.0 |
| .8 | .8 | .7 | .6 | .6 | .7 | .9 | .9 | 1.1 | 1.0 | 1.3 | 1.3 |
| 6.0 | 4.0 | 4.0 | 4.0 | 4.0 | 4.0 | 6.0 | 6.0 | 6.0 | 6.0 | 9.0 | 6.0 |
| 49 | 43 | 57 | 39 | 56 | 44 | 61 | 48 | 53 | 52 | 80 | 67 |
| 47 | 39 | 55 | 31 | 45 | 38 | 63 | 53 | 50 | 47 | 109 | 64 |
| 11 | 7 | 11 | 6 | 13 | 11 | 11 | 11 | 11 | 9 | 11 | 10 |
| 22 | 19 | 25 | 17 | 26 | 22 | 26 | 19 | 24 | 22 | 27 | 25 |
| 36 | 27 | 39 | 27 | 44 | 30 | 44 | 28 | 39 | 41 | 52 | 50 |
| 61 | 52 | 66 | 50 | 77 | 52 | 71 | 52 | 61 | 69 | 103 | 80 |
| 104 | 89 | 105 | 78 | 104 | 92 | 138 | 94 | 100 | 103 | 160 | 177 |
| 127 | 104 | 177 | 104 | 133 | 117 | 177 | 133 | 156 | 151 | 177 | 199 |
| 211 | 216 | 299 | 174 | 183 | 185 | 302 | 266 | 252 | 266 | 730 | 291 |
| 101 | 89 | 121 | 88 | 133 | 93 | 125 | 93 | 95 | 108 | 174 | 133 |
| 73 | 68 | 99 | 69 | 82 | 69 | 111 | 79 | 61 | 90 | 162 | 125 |
| 108 | 106 | 123 | 96 | 140 | 106 | 131 | 101 | 109 | 103 | 114 | 110 |
| 83 | 76 | 82 | 53 | 106 | 72 | 95 | 80 | 78 | 75 | 37 | 101 |
| 117 | 93 | 146 | 94 | 112 | 97 | 142 | 128 | 106 | 89 | 183 | 126 |
| 79 | 45 | 99 | 15 | 42 | 29 | 112 | 100 | 75 | 62 | 240 | 73 |
| 598 | 531 | 664 | 531 | 598 | 598 | 797 | 1,062 | 623 | 531 | 844 | 623 |
| 1,195 | 789 | 930 | 531 | 1,195 | 797 | 1,793 | 1,793 | 936 | 890 | 2,188 | 872 |
| 97 | 90 | 118 | 89 | 127 | 93 | 120 | 93 | 89 | 87 | 137 | 101 |
| 74 | 66 | 89 | 61 | 86 | 64 | 100 | 74 | 60 | 71 | 112 | 86 |
| 29 | 21 | 33 | 28 | 33 | 28 | 23 | 25 | 17 | 17 | 39 | 17 |
| 45 | 45 | 57 | 52 | 78 | 50 | 66 | 45 | 50 | 45 | 72 | 40 |
| 78 | 78 | 85 | 78 | 117 | 78 | 85 | 75 | 75 | 75 | 90 | 78 |
| 133 | 112 | 149 | 113 | 156 | 120 | 156 | 120 | 133 | 100 | 160 | 117 |
| 156 | 149 | 266 | 149 | 234 | 156 | 227 | 156 | 156 | 156 | 299 | 224 |
| 240 | 224 | 284 | 199 | 266 | 224 | 266 | 266 | 234 | 249 | 398 | 266 |
| 448 | 312 | 415 | 299 | 448 | 299 | 532 | 300 | 266 | 332 | 531 | 531 |
| 598 | 531 | 664 | 531 | 598 | 531 | 797 | 797 | 398 | 531 | 531 | 531 |
| 15.7 | 6.8 | 5.0 | 5.4 | 13.6 | 12.2 | 26.4 | 26.3 | 3.8 | 1.4 | 2.6 | 7.3 |
| 4.7 | 2.1 | 2.0 | 3.0 | 3.6 | 1.9 | 3.3 | 2.2 | 1.3 | .4 | 1.9 | 3.3 |

TAELE 5.12.--PANCAKES
 CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|------------|------------|------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE | MALE | MALE | MALE |
| | | AND FEMALE | AND FEMALE | AND FEMALE | AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 8.3 | 3.0 | 12.2 | 12.7 | 11.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 7.3 | 2.7 | 11.0 | 11.4 | 10.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .8 | .3 | 1.1 | 1.2 | 1.1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | .1 | .1 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .3 | .3 | .3 | .4 |
| MAXIMUM.....DO..... | 3.0 | 2.0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 43 | 14 | 22 | 28 | 36 |
| STANDARD DEVIATION.....GM..... | 36 | 12 | 21 | 21 | 27 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 9 | | 5 | 9 | 9 |
| 25TH.....GM..... | 18 | | 9 | 18 | 18 |
| 50TH.....GM..... | 33 | | 18 | 24 | 27 |
| 75TH.....GM..... | 50 | | 24 | 36 | 49 |
| 90TH.....GM..... | 74 | | 49 | 54 | 73 |
| 95TH.....GM..... | 108 | | 54 | 73 | 74 |
| 99TH.....GM..... | 195 | | 110 | 98 | 146 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 113 | 36 | 57 | 77 | 96 |
| STANDARD DEVIATION.....GM..... | 88 | 29 | 47 | 53 | 60 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 114 | 54 | 67 | 74 | 99 |
| STANDARD DEVIATION.....GM..... | 76 | 0 | 60 | 44 | 62 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 119 | 0 | 66 | 74 | 109 |
| STANDARD DEVIATION.....GM..... | 66 | 0 | 43 | 37 | 25 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 940 | 85 | 329 | 294 | 365 |
| MAXIMUM OVER 3 DAYS.....GM..... | 940 | 108 | 504 | 329 | 504 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 113 | 39 | 59 | 76 | 96 |
| STANDARD DEVIATION.....GM..... | 85 | 27 | 50 | 52 | 59 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 27 | | 14 | 27 | 27 |
| 25TH.....GM..... | 54 | | 27 | 41 | 54 |
| 50TH.....GM..... | 81 | | 50 | 54 | 81 |
| 75TH.....GM..... | 146 | | 73 | 95 | 113 |
| 90TH.....GM..... | 219 | | 146 | 146 | 189 |
| 95TH.....GM..... | 282 | | 146 | 198 | 219 |
| 99TH.....GM..... | 438 | | 284 | 229 | 252 |
| MAXIMUM.....GM..... | 940 | 85 | 329 | 294 | 365 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 9.2 | 0 | .7 | 1.4 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 0 | .5 | .6 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 13.5 | 10.7 | 9.8 | 9.8 | 8.5 | 7.6 | 8.0 | 5.8 | 5.6 | 4.9 | 6.0 | 4.3 |
| 12.0 | 9.6 | 8.5 | 8.4 | 7.6 | 6.9 | 6.4 | 5.2 | 4.6 | 4.7 | 5.4 | 4.2 |
| 1.2 | .9 | 1.0 | 1.3 | .9 | .7 | 1.3 | .5 | .8 | 0 | .5 | .1 |
| .3 | .2 | .3 | 0 | (2) | (2) | .3 | .1 | .2 | .2 | .1 | 0 |
| 1.2 | 1.1 | 1.2 | 1.1 | 1.1 | 1.1 | 1.2 | 1.1 | 1.2 | 1.1 | 1.2 | 1.0 |
| .4 | .4 | .5 | .3 | .3 | .3 | .5 | .4 | .5 | .4 | .4 | .1 |
| 3.0 | 3.0 | 3.0 | 2.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 2.0 |
| 45 | 38 | 63 | 46 | 59 | 36 | 55 | 38 | 52 | 33 | 47 | 31 |
| 31 | 35 | 48 | 40 | 41 | 24 | 45 | 36 | 42 | 20 | 46 | 14 |
| 18 | 9 | 18 | 18 | 18 | 9 | 18 | 9 | 9 | 12 | | |
| 25 | 18 | 33 | 18 | 36 | 18 | 27 | 18 | 24 | 18 | | |
| 36 | 27 | 50 | 33 | 49 | 27 | 37 | 27 | 36 | 27 | | |
| 54 | 45 | 73 | 49 | 73 | 49 | 72 | 45 | 66 | 42 | | |
| 91 | 72 | 122 | 97 | 97 | 72 | 108 | 73 | 122 | 57 | | |
| 108 | 97 | 173 | 146 | 146 | 73 | 153 | 97 | 157 | 73 | | |
| 146 | 188 | 243 | 210 | 195 | 146 | 219 | 197 | 189 | 97 | | |
| 123 | 103 | 165 | 114 | 158 | 101 | 139 | 100 | 132 | 98 | 115 | 94 |
| 80 | 94 | 120 | 96 | 95 | 62 | 105 | 89 | 103 | 60 | 54 | 41 |
| 103 | 99 | 134 | 144 | 163 | 92 | 125 | 104 | 119 | 0 | 227 | 40 |
| 38 | 76 | 53 | 78 | 101 | 53 | 77 | 64 | 77 | 0 | 155 | 0 |
| 127 | 85 | 192 | 0 | 189 | 73 | 139 | 132 | 116 | 39 | 27 | 0 |
| 46 | 15 | 66 | 0 | 0 | 0 | 65 | 77 | 54 | 3 | 0 | 0 |
| 438 | 940 | 730 | 752 | 584 | 564 | 584 | 940 | 470 | 292 | 378 | 185 |
| 616 | 940 | 886 | 752 | 882 | 564 | 752 | 940 | 567 | 292 | 756 | 185 |
| 118 | 101 | 161 | 121 | 159 | 98 | 134 | 102 | 125 | 92 | 121 | 92 |
| 72 | 89 | 110 | 93 | 97 | 61 | 96 | 87 | 83 | 60 | 89 | 41 |
| 54 | 27 | 54 | 49 | 54 | 27 | 37 | 27 | 27 | 27 | | |
| 73 | 54 | 85 | 54 | 81 | 54 | 81 | 54 | 54 | 54 | | |
| 100 | 81 | 146 | 81 | 144 | 81 | 108 | 81 | 98 | 81 | | |
| 146 | 108 | 170 | 146 | 219 | 128 | 162 | 113 | 162 | 108 | | |
| 200 | 188 | 292 | 219 | 292 | 185 | 219 | 219 | 219 | 162 | | |
| 282 | 252 | 370 | 292 | 365 | 219 | 296 | 219 | 252 | 219 | | |
| 438 | 514 | 588 | 564 | 441 | 292 | 565 | 441 | 438 | 292 | | |
| 438 | 940 | 730 | 752 | 584 | 564 | 584 | 940 | 438 | 292 | 378 | 185 |
| 0 | .9 | 0 | 2.5 | 0 | 0 | 2.2 | 0 | 0 | 1.5 | 0 | 0 |
| 0 | .3 | 0 | 1.5 | 0 | 0 | .5 | 0 | 0 | 1.6 | 0 | 0 |

TABLE 5.13.--WAFFLES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.9 | .6 | 3.4 | 5.7 | 5.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.4 | .6 | 2.4 | 4.4 | 4.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | 0 | .9 | 1.3 | .9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | .1 | .1 | .3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 1.0 | 1.3 | 1.3 | 1.3 |
| STANDARD DEVIATION.....DO..... | .5 | .0 | .5 | .5 | .5 |
| MAXIMUM.....DO..... | 4.0 | 1.0 | 3.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 35 | 10 | 24 | 30 | 29 |
| STANDARD DEVIATION.....GM..... | 30 | 4 | 23 | 26 | 24 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 7 | | | 7 | 7 |
| 25TH.....GM..... | 13 | | | 13 | 13 |
| 50TH.....GM..... | 26 | | | 23 | 26 |
| 75TH.....GM..... | 47 | | | 33 | 33 |
| 90TH.....GM..... | 67 | | | 67 | 53 |
| 95TH.....GM..... | 95 | | | 82 | 83 |
| 99TH.....GM..... | 158 | | | 153 | 121 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 92 | 30 | 53 | 62 | 68 |
| STANDARD DEVIATION.....GM..... | 82 | 13 | 44 | 36 | 42 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 80 | 0 | 59 | 80 | 71 |
| STANDARD DEVIATION.....GM..... | 48 | 0 | 45 | 45 | 46 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 71 | 0 | 66 | 153 | 80 |
| STANDARD DEVIATION.....GM..... | 36 | 0 | 0 | 0 | 33 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 711 | 43 | 200 | 258 | 283 |
| MAXIMUM OVER 3 DAYS.....GM..... | 711 | 43 | 350 | 458 | 400 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 87 | 30 | 56 | 69 | 69 |
| STANDARD DEVIATION.....GM..... | 74 | 13 | 45 | 41 | 45 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 20 | | | 20 | 20 |
| 25TH.....GM..... | 40 | | | 39 | 40 |
| 50TH.....GM..... | 78 | | | 50 | 50 |
| 75TH.....GM..... | 100 | | | 100 | 85 |
| 90TH.....GM..... | 158 | | | 113 | 100 |
| 95TH.....GM..... | 200 | | | 158 | 158 |
| 99TH.....GM..... | 400 | | | 200 | 250 |
| MAXIMUM.....GM..... | 711 | 43 | 200 | 200 | 283 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 31.5 | 0 | 0 | 4.2 | 3.7 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.4 | 0 | 0 | 3.3 | 2.7 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ . MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.2 | 4.1 | 3.5 | 2.4 | 2.6 | 2.5 | 2.2 | 1.8 | 2.3 | 1.6 | 2.6 | 1.3 |
| 4.2 | 3.4 | 2.9 | 2.3 | 2.1 | 2.2 | 2.0 | 1.6 | 2.1 | 1.0 | 1.5 | 1.0 |
| .8 | .7 | .4 | .1 | .4 | .2 | .2 | .1 | .1 | .5 | .3 | .3 |
| .2 | (2) | .1 | 0 | .2 | (2) | .1 | .1 | .1 | .1 | .8 | 0 |
| 1.2 | 1.2 | 1.2 | 1.0 | 1.3 | 1.1 | 1.2 | 1.1 | 1.2 | 1.4 | 1.7 | 1.3 |
| .5 | .4 | .5 | .2 | .6 | .4 | .4 | .4 | .5 | .6 | .9 | .4 |
| 3.0 | 3.0 | 3.0 | 2.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 2.0 |
| 36 | 32 | 50 | 27 | 48 | 35 | 43 | 29 | 39 | 28 | 41 | 42 |
| 28 | 29 | 29 | 18 | 38 | 36 | 33 | 22 | 30 | 16 | 21 | 37 |
| 10 | 7 | | | 13 | 7 | 13 | 7 | | | | |
| 17 | 13 | | | 17 | 17 | 20 | 13 | | | | |
| 27 | 23 | | | 33 | 26 | 33 | 26 | | | | |
| 50 | 33 | | | 67 | 33 | 52 | 33 | | | | |
| 75 | 75 | | | 100 | 67 | 80 | 57 | | | | |
| 94 | 95 | | | 117 | 94 | 105 | 76 | | | | |
| 133 | 133 | | | 237 | 189 | 189 | 100 | | | | |
| 88 | 75 | 134 | 83 | 129 | 105 | 122 | 81 | 99 | 59 | 98 | 108 |
| 68 | 74 | 77 | 56 | 121 | 111 | 99 | 56 | 56 | 22 | 49 | 55 |
| 81 | 92 | 115 | 39 | 98 | 49 | 83 | 80 | 200 | 67 | 35 | 91 |
| 36 | 44 | 48 | 0 | 36 | 32 | 49 | 48 | 0 | 23 | 0 | 97 |
| 100 | 59 | 84 | 0 | 65 | 79 | 59 | 52 | 20 | 40 | 64 | 0 |
| 44 | 0 | 18 | 0 | 21 | 0 | 32 | 42 | 0 | 0 | 14 | 0 |
| 400 | 494 | 340 | 284 | 711 | 566 | 566 | 267 | 234 | 100 | 158 | 316 |
| 524 | 494 | 400 | 284 | 711 | 566 | 566 | 316 | 400 | 200 | 234 | 474 |
| 87 | 80 | 125 | 79 | 112 | 95 | 110 | 78 | 101 | 61 | 71 | 101 |
| 62 | 68 | 70 | 55 | 100 | 102 | 91 | 55 | 65 | 23 | 37 | 80 |
| 20 | 20 | | | 40 | 20 | 39 | 20 | | | | |
| 50 | 40 | | | 50 | 40 | 50 | 40 | | | | |
| 78 | 57 | | | 80 | 78 | 85 | 78 | | | | |
| 100 | 100 | | | 150 | 100 | 150 | 100 | | | | |
| 150 | 158 | | | 200 | 200 | 200 | 158 | | | | |
| 200 | 200 | | | 300 | 235 | 300 | 170 | | | | |
| 400 | 400 | | | 711 | 566 | 566 | 267 | | | | |
| 400 | 494 | 340 | 284 | 711 | 566 | 566 | 267 | 234 | 100 | 158 | 316 |
| 4.3 | 2.1 | 2.0 | 0 | 11.6 | 0 | 3.6 | 0 | 0 | 0 | 0 | 0 |
| 3.2 | 2.0 | 3.5 | 0 | 8.8 | 0 | 2.8 | 0 | 0 | 0 | 0 | 0 |

TABLE 5.14.--TORTILLAS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.9 | .8 | 3.9 | 5.1 | 4.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.6 | .8 | 2.2 | 2.4 | 2.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .5 | 0 | .3 | .8 | .8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .8 | 0 | 1.4 | 1.9 | 1.2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 2.4 | 1.0 | 2.5 | 2.9 | 2.3 |
| STANDARD DEVIATION.....DO..... | 2.0 | .0 | 1.8 | 2.3 | 2.0 |
| MAXIMUM.....DO..... | 9.0 | 1.0 | 7.0 | 9.0 | 9.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 55 | 5 | 21 | 35 | 43 |
| STANDARD DEVIATION.....GM..... | 62 | 2 | 17 | 31 | 54 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 10 | | | 6 | 10 |
| 25TH.....GM..... | 20 | | | 10 | 10 |
| 50TH.....GM..... | 30 | | | 25 | 23 |
| 75TH.....GM..... | 63 | | | 53 | 47 |
| 90TH.....GM..... | 130 | | | 76 | 100 |
| 95TH.....GM..... | 180 | | | 95 | 110 |
| 99TH.....GM..... | 284 | | | 180 | 280 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 68 | 16 | 30 | 42 | 50 |
| STANDARD DEVIATION.....GM..... | 46 | 7 | 13 | 21 | 26 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 76 | 0 | 41 | 50 | 50 |
| STANDARD DEVIATION.....GM..... | 40 | 0 | 17 | 22 | 21 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 121 | 0 | 37 | 62 | 108 |
| STANDARD DEVIATION.....GM..... | 79 | 0 | 16 | 33 | 72 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 630 | 28 | 70 | 180 | 350 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,680 | 28 | 175 | 540 | 840 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 69 | 16 | 26 | 36 | 55 |
| STANDARD DEVIATION.....GM..... | 39 | 7 | 11 | 16 | 29 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 28 | | | 15 | 28 |
| 25TH.....GM..... | 30 | | | 30 | 30 |
| 50TH.....GM..... | .60 | | | 30 | 57 |
| 75TH.....GM..... | 90 | | | 35 | 70 |
| 90TH.....GM..... | 120 | | | 60 | 90 |
| 95TH.....GM..... | 140 | | | 60 | 105 |
| 99TH.....GM..... | 210 | | | 90 | 140 |
| MAXIMUM.....GM..... | 227 | 28 | 60 | 90 | 140 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 10.8 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 4.0 | 4.3 | 3.4 | 4.0 | 3.2 | 3.4 | 2.2 | 2.3 | .4 | .7 | 1.7 | .3 |
| 2.2 | 2.2 | 2.1 | 2.3 | 1.6 | 1.8 | 1.2 | 1.4 | .3 | .2 | .7 | .1 |
| .7 | .8 | .6 | .9 | .7 | .6 | .3 | .3 | 0 | .4 | .3 | .2 |
| 1.1 | 1.3 | .6 | .9 | .9 | 1.0 | .7 | .6 | .2 | .1 | .8 | 0 |
| 2.2 | 2.3 | 1.9 | 2.1 | 2.5 | 2.4 | 2.3 | 2.5 | 2.9 | 2.3 | 2.5 | 2.0 |
| 1.6 | 1.9 | 1.6 | 1.5 | 2.1 | 1.9 | 1.8 | 2.2 | 2.4 | 1.5 | 1.6 | .0 |
| 6.0 | 9.0 | 9.0 | 9.0 | 9.0 | 9.0 | 8.0 | 9.0 | 6.0 | 6.0 | 5.0 | 2.0 |
| 54 | 50 | 63 | 48 | 81 | 54 | 76 | 52 | 53 | 56 | 53 | 47 |
| 42 | 52 | 60 | 42 | 82 | 55 | 92 | 58 | 29 | 53 | 32 | 4 |
| 10 | 10 | | 10 | 10 | 10 | 10 | 10 | | | | |
| 20 | 20 | | 20 | 23 | 20 | 20 | 19 | | | | |
| 40 | 30 | | 38 | 47 | 30 | 47 | 30 | | | | |
| 80 | 60 | | 60 | 133 | 60 | 93 | 67 | | | | |
| 117 | 110 | | 82 | 240 | 120 | 150 | 132 | | | | |
| 140 | 170 | | 120 | 260 | 180 | 268 | 150 | | | | |
| 190 | 292 | | 240 | 330 | 280 | 560 | 315 | | | | |
| 82 | 67 | 91 | 82 | 83 | 67 | 82 | 67 | 90 | 41 | 60 | 120 |
| 46 | 44 | 43 | 63 | 50 | 38 | 57 | 43 | 0 | 14 | 0 | 0 |
| 87 | 67 | 123 | 77 | 96 | 64 | 88 | 72 | 0 | 77 | 85 | 75 |
| 39 | 29 | 61 | 28 | 44 | 31 | 38 | 43 | 0 | 24 | 0 | 0 |
| 107 | 103 | 158 | 99 | 193 | 119 | 165 | 128 | 90 | 195 | 82 | 0 |
| 38 | 65 | 67 | 56 | 75 | 63 | 118 | 66 | 0 | 0 | 20 | 0 |
| 280 | 315 | 420 | 245 | 420 | 360 | 630 | 385 | 150 | 195 | 120 | 120 |
| 570 | 875 | 840 | 720 | 990 | 840 | 1,680 | 945 | 270 | 585 | 300 | 150 |
| 74 | 66 | 100 | 69 | 97 | 67 | 97 | 64 | 54 | 72 | 63 | 71 |
| 31 | 33 | 48 | 33 | 42 | 31 | 51 | 31 | 35 | 23 | 8 | 14 |
| 30 | 30 | | 30 | 30 | 30 | 30 | 30 | | | | |
| 60 | 35 | | 35 | 60 | 35 | 60 | 35 | | | | |
| 60 | 60 | | 60 | 105 | 60 | 90 | 60 | | | | |
| 90 | 75 | | 90 | 120 | 90 | 120 | 90 | | | | |
| 120 | 105 | | 120 | 142 | 105 | 180 | 105 | | | | |
| 128 | 140 | | 120 | 180 | 120 | 210 | 113 | | | | |
| 140 | 210 | | 140 | 180 | 180 | 210 | 175 | | | | |
| 180 | 227 | 210 | 150 | 227 | 227 | 210 | 210 | 120 | 105 | 85 | 90 |
| 0 | .6 | 2.2 | .7 | 2.4 | .9 | 1.7 | 2.3 | 0 | 0 | 0 | 0 |
| 0 | .3 | 2.5 | .5 | .8 | .2 | .7 | .6 | 0 | 0 | 0 | 0 |

TABLE 5.15.--CAKES AND CUPCAKES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 25.5 | 1.6 | 17.4 | 25.3 | 34.4 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 17.3 | 1.1 | 14.7 | 19.1 | 22.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 6.2 | .5 | 2.0 | 4.8 | 9.6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.9 | 0 | .6 | 1.5 | 2.5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.5 | 1.3 | 1.3 | 1.5 | 1.6 |
| STANDARD DEVIATION.....DO..... | .9 | .5 | .6 | .9 | .9 |
| MAXIMUM.....DO..... | 8.0 | 2.0 | 4.0 | 6.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 41 | 23 | 22 | 30 | 35 |
| STANDARD DEVIATION.....GM..... | 38 | 16 | 20 | 26 | 26 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 9 | | 3 | 7 | 9 |
| 25TH.....GM..... | 16 | | 9 | 13 | 16 |
| 50TH.....GM..... | 31 | | 15 | 21 | 28 |
| 75TH.....GM..... | 52 | | 30 | 38 | 42 |
| 90TH.....GM..... | 83 | | 48 | 60 | 65 |
| 95TH.....GM..... | 112 | | 57 | 77 | 88 |
| 99TH.....GM..... | 192 | | 114 | 137 | 128 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 84 | 55 | 55 | 66 | 73 |
| STANDARD DEVIATION.....GM..... | 64 | 44 | 42 | 51 | 50 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 90 | 51 | 57 | 74 | 74 |
| STANDARD DEVIATION.....GM..... | 63 | 22 | 38 | 53 | 38 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 95 | 0 | 65 | 63 | 69 |
| STANDARD DEVIATION.....GM..... | 57 | 0 | 38 | 33 | 40 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,382 | 142 | 259 | 411 | 460 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,460 | 142 | 405 | 570 | 610 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 79 | 53 | 51 | 61 | 66 |
| STANDARD DEVIATION.....GM..... | 59 | 37 | 38 | 45 | 42 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 23 | | 9 | 19 | 25 |
| 25TH.....GM..... | 41 | | 28 | 35 | 38 |
| 50TH.....GM..... | 63 | | 41 | 46 | 52 |
| 75TH.....GM..... | 99 | | 65 | 78 | 88 |
| 90TH.....GM..... | 144 | | 100 | 113 | 113 |
| 95TH.....GM..... | 184 | | 138 | 147 | 139 |
| 99TH.....GM..... | 284 | | 179 | 228 | 227 |
| MAXIMUM.....GM..... | 1,382 | 142 | 259 | 358 | 358 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 2357.2 | 2.6 | 28.0 | 104.8 | 121.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 15.8 | 24.7 | 11.9 | 16.6 | 12.2 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 36.4 | 35.2 | 31.0 | 26.5 | 24.5 | 20.2 | 27.2 | 22.8 | 24.8 | 22.7 | 22.4 | 25.8 |
| 23.5 | 24.5 | 20.7 | 19.5 | 15.5 | 15.0 | 17.2 | 16.2 | 16.3 | 14.5 | 14.5 | 16.1 |
| 9.9 | 8.8 | 8.7 | 5.7 | 6.4 | 4.2 | 6.9 | 5.2 | 5.8 | 6.0 | 5.5 | 8.1 |
| 3.1 | 1.9 | 1.7 | 1.3 | 2.5 | 1.0 | 3.1 | 1.4 | 2.7 | 2.2 | 2.4 | 1.7 |
| 1.6 | 1.5 | 1.6 | 1.5 | 1.7 | 1.4 | 1.7 | 1.5 | 1.7 | 1.6 | 1.8 | 1.5 |
| .9 | .8 | .8 | .8 | 1.0 | .7 | 1.0 | .8 | 1.0 | .9 | 1.1 | .8 |
| 6.0 | 7.0 | 6.0 | 8.0 | 6.0 | 5.0 | 6.0 | 7.0 | 6.0 | 7.0 | 6.0 | 5.0 |
| 43 | 39 | 48 | 39 | 53 | 37 | 49 | 35 | 43 | 39 | 53 | 35 |
| 36 | 33 | 45 | 37 | 48 | 30 | 44 | 31 | 39 | 31 | 55 | 47 |
| 9 | 9 | 11 | 9 | 12 | 8 | 10 | 7 | 8 | 7 | 11 | 6 |
| 19 | 16 | 19 | 18 | 23 | 15 | 19 | 14 | 19 | 16 | 19 | 15 |
| 32 | 30 | 33 | 30 | 38 | 30 | 36 | 27 | 34 | 31 | 35 | 27 |
| 55 | 50 | 62 | 49 | 65 | 48 | 62 | 44 | 53 | 55 | 76 | 42 |
| 86 | 76 | 95 | 76 | 108 | 76 | 108 | 71 | 91 | 79 | 118 | 61 |
| 116 | 104 | 118 | 96 | 143 | 93 | 140 | 90 | 113 | 95 | 147 | 77 |
| 189 | 157 | 222 | 155 | 244 | 139 | 217 | 170 | 203 | 160 | 339 | 131 |
| 87 | 81 | 103 | 88 | 101 | 83 | 94 | 75 | 86 | 81 | 94 | 67 |
| 62 | 58 | 93 | 61 | 73 | 62 | 76 | 54 | 53 | 68 | 69 | 45 |
| 90 | 89 | 105 | 94 | 121 | 83 | 101 | 77 | 80 | 77 | 96 | 84 |
| 58 | 48 | 61 | 84 | 87 | 44 | 63 | 50 | 45 | 42 | 43 | 111 |
| 93 | 95 | 103 | 81 | 101 | 93 | 110 | 91 | 107 | 82 | 164 | 61 |
| 53 | 61 | 97 | 55 | 56 | 43 | 59 | 46 | 69 | 37 | 89 | 28 |
| 546 | 438 | 875 | 677 | 743 | 372 | 796 | 560 | 737 | 496 | 498 | 1,382 |
| 823 | 944 | 1,460 | 1,340 | 1,080 | 710 | 1,040 | 960 | 907 | 709 | 1,017 | 1,427 |
| 80 | 77 | 93 | 80 | 96 | 78 | 89 | 71 | 77 | 73 | 92 | 68 |
| 56 | 55 | 71 | 59 | 67 | 52 | 62 | 50 | 53 | 52 | 57 | 105 |
| 25 | 26 | 28 | 24 | 29 | 22 | 28 | 20 | 22 | 22 | 29 | 19 |
| 41 | 39 | 43 | 42 | 48 | 41 | 44 | 37 | 40 | 38 | 45 | 38 |
| 66 | 57 | 72 | 64 | 85 | 66 | 80 | 57 | 58 | 57 | 84 | 51 |
| 96 | 96 | 113 | 101 | 113 | 100 | 113 | 92 | 105 | 95 | 114 | 79 |
| 158 | 140 | 179 | 143 | 174 | 139 | 170 | 137 | 147 | 125 | 172 | 106 |
| 186 | 179 | 230 | 185 | 227 | 179 | 200 | 173 | 179 | 173 | 180 | 132 |
| 284 | 302 | 346 | 302 | 340 | 269 | 306 | 257 | 240 | 227 | 319 | 232 |
| 410 | 410 | 614 | 648 | 626 | 355 | 614 | 480 | 369 | 496 | 319 | 1,382 |
| 157.4 | 155.1 | 77.0 | 92.8 | 231.2 | 301.6 | 392.1 | 409.8 | 83.1 | 123.1 | 24.0 | 52.7 |
| 12.8 | 13.5 | 11.5 | 16.2 | 14.5 | 19.8 | 17.6 | 17.3 | 18.0 | 19.5 | 11.4 | 13.3 |

TABLE 5.16.--CHOCOLATE CAKES AND CUPCAKES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS 1/ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 8.8 | .3 | 5.8 | 10.5 | 16.0 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 6.7 | 0 | 5.2 | 8.8 | 12.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.7 | .3 | .5 | 1.5 | 3.4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | .2 | .2 | .5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE 1/NUMBER..... | 1.4 | 2.0 | 1.3 | 1.3 | 1.4 |
| STANDARD DEVIATION.....DO..... | .7 | .0 | .6 | .8 | .7 |
| MAXIMUM.....DO..... | 6.0 | 2.0 | 3.0 | 5.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY 1/GM..... | 36 | 46 | 22 | 26 | 29 |
| STANDARD DEVIATION.....GM..... | 31 | 0 | 16 | 22 | 20 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: 1/ | | | | | |
| 5TH.....GM..... | 9 | | 8 | 8 | 9 |
| 25TH.....GM..... | 15 | | 12 | 13 | 15 |
| 50TH.....GM..... | 30 | | 15 | 19 | 24 |
| 75TH.....GM..... | 46 | | 29 | 31 | 35 |
| 90TH.....GM..... | 73 | | 46 | 53 | 57 |
| 95TH.....GM..... | 92 | | 57 | 59 | 62 |
| 99TH.....GM..... | 170 | | 85 | 92 | 94 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE 1/GM..... | 84 | 0 | 58 | 62 | 67 |
| STANDARD DEVIATION.....GM..... | 61 | 0 | 40 | 37 | 39 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE 1/GM..... | 89 | 69 | 46 | 82 | 70 |
| STANDARD DEVIATION.....GM..... | 59 | 0 | 17 | 60 | 38 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE 1/GM..... | 85 | 0 | 67 | 34 | 55 |
| STANDARD DEVIATION.....GM..... | 52 | 0 | 17 | 17 | 29 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 737 | 92 | 184 | 284 | 460 |
| MAXIMUM OVER 3 DAYS.....GM..... | 884 | 138 | 255 | 567 | 495 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE 1/GM..... | 79 | 69 | 52 | 59 | 62 |
| STANDARD DEVIATION.....GM..... | 53 | 23 | 31 | 41 | 35 |
| PERCENTILES: 1/ | | | | | |
| 5TH.....GM..... | 28 | | 23 | 19 | 28 |
| 25TH.....GM..... | 41 | | 35 | 35 | 38 |
| 50TH.....GM..... | 70 | | 41 | 46 | 46 |
| 75TH.....GM..... | 92 | | 57 | 70 | 85 |
| 90TH.....GM..... | 142 | | 92 | 104 | 92 |
| 95TH.....GM..... | 177 | | 118 | 113 | 133 |
| 99TH.....GM..... | 278 | | 171 | 284 | 184 |
| MAXIMUM.....GM..... | 626 | 92 | 171 | 284 | 244 |
| QUANTITIES NOT SPECIFIED BY USERS: 1/ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 543.8 | 1.4 | 5.1 | 37.2 | 41.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 11.8 | 50.0 | 6.6 | 15.6 | 10.2 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 14.6 | 13.9 | 12.4 | 9.6 | 9.2 | 7.3 | 9.2 | 6.5 | 5.1 | 4.1 | 6.0 | 4.9 |
| 11.0 | 11.1 | 9.3 | 7.8 | 6.4 | 5.8 | 6.5 | 5.2 | 3.8 | 2.6 | 5.2 | 4.0 |
| 2.9 | 2.3 | 2.9 | 1.6 | 2.2 | 1.4 | 2.0 | 1.1 | 1.2 | 1.4 | .6 | .9 |
| .7 | .5 | .2 | .2 | .6 | .1 | .7 | .2 | .1 | .1 | .2 | .1 |
| 1.4 | 1.4 | 1.4 | 1.3 | 1.5 | 1.3 | 1.5 | 1.3 | 1.4 | 1.5 | 1.4 | 1.3 |
| .8 | .7 | .7 | .7 | .8 | .6 | .8 | .7 | .7 | .7 | .7 | .5 |
| 6.0 | 5.0 | 4.0 | 4.0 | 5.0 | 4.0 | 4.0 | 6.0 | 3.0 | 4.0 | 3.0 | 3.0 |
| 36 | 32 | 46 | 37 | 51 | 32 | 44 | 31 | 49 | 42 | 41 | 27 |
| 26 | 25 | 38 | 33 | 46 | 26 | 35 | 22 | 39 | 30 | 33 | 19 |
| 11 | 9 | 12 | 11 | 13 | 9 | 12 | 7 | 19 | 6 | | |
| 16 | 15 | 23 | 19 | 24 | 15 | 19 | 15 | 31 | 17 | | |
| 30 | 27 | 31 | 28 | 36 | 28 | 31 | 27 | 38 | 35 | | |
| 47 | 44 | 61 | 46 | 61 | 38 | 58 | 38 | 52 | 61 | | |
| 67 | 61 | 90 | 62 | 93 | 61 | 87 | 61 | 92 | 78 | | |
| 88 | 76 | 113 | 89 | 139 | 92 | 96 | 77 | 123 | 94 | | |
| 150 | 124 | 220 | 141 | 244 | 139 | 207 | 104 | 246 | 189 | | |
| 90 | 77 | 108 | 86 | 106 | 77 | 96 | 76 | 134 | 88 | 106 | 68 |
| 59 | 51 | 97 | 51 | 75 | 48 | 64 | 50 | 120 | 55 | 93 | 51 |
| 72 | 82 | 112 | 109 | 121 | 86 | 95 | 73 | 96 | 87 | 109 | 60 |
| 38 | 45 | 61 | 87 | 87 | 60 | 51 | 38 | 52 | 34 | 31 | 18 |
| 76 | 70 | 70 | 67 | 110 | 72 | 98 | 76 | 46 | 141 | 81 | 81 |
| 49 | 48 | 16 | 28 | 60 | 29 | 56 | 28 | 0 | 47 | 0 | 0 |
| 371 | 371 | 726 | 663 | 696 | 355 | 454 | 348 | 737 | 227 | 354 | 232 |
| 547 | 650 | 726 | 884 | 812 | 710 | 650 | 487 | 737 | 567 | 354 | 244 |
| 75 | 71 | 95 | 83 | 103 | 76 | 90 | 70 | 103 | 85 | 87 | 64 |
| 47 | 47 | 64 | 49 | 68 | 50 | 54 | 45 | 66 | 47 | 64 | 41 |
| 28 | 25 | 28 | 30 | 35 | 23 | 35 | 20 | 41 | 23 | | |
| 39 | 38 | 46 | 46 | 57 | 41 | 46 | 37 | 57 | 46 | | |
| 62 | 57 | 84 | 70 | 91 | 64 | 85 | 57 | 92 | 91 | | |
| 92 | 92 | 114 | 92 | 133 | 92 | 112 | 92 | 122 | 108 | | |
| 138 | 133 | 184 | 162 | 184 | 138 | 162 | 128 | 162 | 157 | | |
| 168 | 162 | 227 | 186 | 227 | 157 | 184 | 157 | 183 | 177 | | |
| 266 | 278 | 340 | 221 | 278 | 278 | 278 | 227 | 369 | 186 | | |
| 371 | 371 | 369 | 313 | 626 | 355 | 454 | 348 | 369 | 227 | 313 | 232 |
| 37.7 | 35.0 | 29.3 | 26.4 | 67.6 | 98.9 | 77.9 | 45.0 | 7.2 | 20.9 | 2.0 | 10.7 |
| 8.6 | 8.6 | 11.8 | 14.0 | 12.6 | 19.8 | 11.8 | 7.4 | 8.8 | 19.7 | 4.5 | 17.3 |

TABLE 5.17.--COOKIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 30.8 | 11.9 | 46.3 | 48.1 | 53.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 17.4 | 4.7 | 22.5 | 23.7 | 27.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 9.0 | 5.3 | 13.9 | 16.0 | 17.5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 4.4 | 1.9 | 9.9 | 8.4 | 8.6 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.9 | 2.0 | 2.1 | 2.0 | 2.0 |
| STANDARD DEVIATION.....DO..... | 1.2 | 1.1 | 1.3 | 1.2 | 1.3 |
| MAXIMUM.....DO..... | 12.0 | 6.0 | 8.0 | 8.0 | 9.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 20 | 10 | 14 | 17 | 19 |
| STANDARD DEVIATION.....GM..... | 21 | 9 | 13 | 16 | 18 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 3 | 2 | 2 | 3 | 3 |
| 25TH.....GM..... | 7 | 4 | 7 | 7 | 7 |
| 50TH.....GM..... | 14 | 7 | 11 | 12 | 13 |
| 75TH.....GM..... | 25 | 13 | 18 | 23 | 25 |
| 90TH.....GM..... | 42 | 26 | 28 | 33 | 38 |
| 95TH.....GM..... | 57 | 30 | 38 | 47 | 53 |
| 99TH.....GM..... | 103 | 36 | 66 | 76 | 82 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 36 | 15 | 23 | 32 | 31 |
| STANDARD DEVIATION.....GM..... | 37 | 13 | 17 | 33 | 26 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 38 | 21 | 26 | 32 | 33 |
| STANDARD DEVIATION.....GM..... | 31 | 15 | 15 | 24 | 21 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 40 | 12 | 26 | 29 | 39 |
| STANDARD DEVIATION.....GM..... | 29 | 7 | 18 | 17 | 26 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 648 | 71 | 165 | 432 | 369 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,358 | 108 | 312 | 474 | 759 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 32 | 15 | 21 | 25 | 28 |
| STANDARD DEVIATION.....GM..... | 30 | 13 | 15 | 22 | 21 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 7 | 7 | 5 | 7 | 7 |
| 25TH.....GM..... | 14 | 7 | 10 | 14 | 14 |
| 50TH.....GM..... | 26 | 10 | 16 | 20 | 24 |
| 75TH.....GM..... | 40 | 14 | 26 | 30 | 36 |
| 90TH.....GM..... | 60 | 35 | 39 | 48 | 56 |
| 95TH.....GM..... | 84 | 42 | 57 | 57 | 60 |
| 99TH.....GM..... | 144 | 57 | 64 | 106 | 112 |
| MAXIMUM.....GM..... | 648 | 71 | 165 | 432 | 216 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 79.1 | 1.5 | 3.6 | 10.0 | 5.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 1.2 | .4 | .6 | .3 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 44.4 | 43.1 | 37.9 | 34.9 | 24.5 | 24.4 | 27.3 | 23.8 | 27.6 | 22.6 | 29.6 | 27.4 |
| 23.7 | 23.8 | 20.9 | 21.4 | 14.5 | 15.5 | 14.8 | 15.4 | 13.3 | 14.0 | 15.8 | 12.3 |
| 14.5 | 13.2 | 11.8 | 9.2 | 6.8 | 6.3 | 7.7 | 6.4 | 8.1 | 5.5 | 7.6 | 8.4 |
| 6.1 | 6.1 | 5.2 | 4.3 | 3.2 | 2.5 | 4.7 | 2.0 | 6.2 | 3.1 | 6.2 | 6.7 |
| 1.9 | 1.9 | 1.9 | 1.8 | 1.8 | 1.7 | 1.9 | 1.6 | 2.1 | 1.7 | 2.0 | 2.1 |
| 1.2 | 1.2 | 1.2 | 1.1 | 1.1 | 1.1 | 1.2 | 1.0 | 1.3 | 1.1 | 1.5 | 1.3 |
| 8.0 | 8.0 | 7.0 | 6.0 | 9.0 | 8.0 | 11.0 | 8.0 | 8.0 | 7.0 | 12.0 | 6.0 |
| 23 | 20 | 29 | 19 | 27 | 18 | 24 | 15 | 21 | 16 | 21 | 15 |
| 27 | 19 | 29 | 18 | 26 | 21 | 22 | 14 | 18 | 19 | 26 | 13 |
| 3 | 3 | 4 | 3 | 3 | 2 | 3 | 2 | 3 | 3 | 3 | 3 |
| 8 | 8 | 9 | 7 | 9 | 7 | 9 | 6 | 9 | 7 | 7 | 6 |
| 15 | 15 | 20 | 14 | 19 | 12 | 17 | 11 | 16 | 11 | 14 | 11 |
| 30 | 27 | 37 | 24 | 36 | 22 | 32 | 19 | 27 | 19 | 27 | 19 |
| 47 | 43 | 59 | 39 | 57 | 38 | 52 | 30 | 45 | 32 | 37 | 30 |
| 63 | 57 | 81 | 54 | 82 | 56 | 67 | 40 | 58 | 45 | 64 | 47 |
| 113 | 88 | 156 | 80 | 113 | 104 | 115 | 69 | 86 | 73 | 196 | 59 |
| 37 | 36 | 58 | 37 | 53 | 36 | 41 | 30 | 34 | 32 | 35 | 23 |
| 36 | 32 | 74 | 32 | 52 | 33 | 41 | 25 | 27 | 28 | 30 | 17 |
| 44 | 40 | 49 | 35 | 53 | 40 | 45 | 31 | 36 | 25 | 39 | 25 |
| 34 | 33 | 36 | 28 | 46 | 43 | 28 | 19 | 25 | 11 | 27 | 18 |
| 49 | 39 | 57 | 40 | 53 | 37 | 47 | 33 | 37 | 41 | 41 | 27 |
| 48 | 21 | 36 | 24 | 26 | 26 | 28 | 23 | 21 | 36 | 44 | 14 |
| 624 | 336 | 648 | 317 | 540 | 621 | 395 | 255 | 228 | 364 | 227 | 113 |
| 1,358 | 672 | 693 | 439 | 756 | 684 | 427 | 454 | 318 | 510 | 588 | 227 |
| 36 | 32 | 45 | 31 | 46 | 33 | 38 | 27 | 30 | 28 | 32 | 21 |
| 36 | 29 | 50 | 26 | 41 | 33 | 29 | 21 | 20 | 28 | 27 | 14 |
| 7 | 7 | 7 | 7 | 9 | 7 | 8 | 7 | 9 | 7 | 7 | 7 |
| 16 | 14 | 21 | 14 | 21 | 14 | 18 | 14 | 16 | 14 | 16 | 13 |
| 27 | 26 | 32 | 25 | 36 | 25 | 30 | 21 | 27 | 22 | 26 | 16 |
| 42 | 40 | 56 | 41 | 56 | 40 | 48 | 32 | 39 | 32 | 36 | 28 |
| 64 | 60 | 84 | 60 | 85 | 64 | 72 | 53 | 50 | 48 | 54 | 36 |
| 85 | 81 | 113 | 81 | 113 | 85 | 94 | 63 | 57 | 57 | 81 | 48 |
| 168 | 165 | 227 | 113 | 227 | 170 | 142 | 113 | 136 | 144 | 170 | 78 |
| 624 | 336 | 648 | 240 | 384 | 432 | 284 | 192 | 170 | 338 | 192 | 113 |
| 5.9 | 9.8 | 3.0 | 4.7 | 11.5 | 0 | 11.8 | 6.8 | 0 | 3.5 | 0 | 2.1 |
| .3 | .6 | .3 | .5 | .7 | 0 | .5 | .2 | 0 | .5 | 0 | .4 |

TABLE 5.18.--CHOCOLATE CHIP COOKIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 8.2 | 3.1 | 16.1 | 17.6 | 16.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.7 | 2.3 | 10.8 | 11.1 | 11.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.8 | .5 | 3.7 | 5.3 | 3.7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .6 | .3 | 1.5 | 1.2 | 1.4 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER.... | 1.6 | 1.5 | 1.7 | 1.7 | 1.6 |
| STANDARD DEVIATION.....DO..... | 1.0 | 1.1 | 1.2 | 1.0 | .9 |
| MAXIMUM.....DO..... | 12.0 | 5.0 | 8.0 | 8.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 12 | 11 | 9 | 12 | 12 |
| STANDARD DEVIATION.....GM..... | 13 | 6 | 8 | 11 | 11 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 2 | | 2 | 2 | 2 |
| 25TH.....GM..... | 5 | | 5 | 5 | 5 |
| 50TH.....GM..... | 9 | | 7 | 9 | 9 |
| 75TH.....GM..... | 14 | | 12 | 14 | 14 |
| 90TH.....GM..... | 28 | | 19 | 23 | 26 |
| 95TH.....GM..... | 37 | | 24 | 28 | 33 |
| 99TH.....GM..... | 61 | | 45 | 57 | 57 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 26 | 30 | 17 | 23 | 24 |
| STANDARD DEVIATION.....GM..... | 25 | 17 | 11 | 19 | 22 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 29 | 15 | 22 | 26 | 26 |
| STANDARD DEVIATION.....GM..... | 24 | 1 | 11 | 20 | 19 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 26 | 17 | 23 | 23 | 25 |
| STANDARD DEVIATION.....GM..... | 18 | 9 | 13 | 13 | 10 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 340 | 57 | 113 | 180 | 170 |
| MAXIMUM OVER 3 DAYS.....GM..... | 567 | 77 | 198 | 294 | 227 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 24 | 21 | 17 | 21 | 22 |
| STANDARD DEVIATION.....GM..... | 20 | 15 | 11 | 14 | 16 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 7 | | 7 | 7 | 7 |
| 25TH.....GM..... | 14 | | 10 | 14 | 14 |
| 50TH.....GM..... | 21 | | 14 | 14 | 15 |
| 75TH.....GM..... | 28 | | 21 | 28 | 28 |
| 90TH.....GM..... | 42 | | 28 | 35 | 35 |
| 95TH.....GM..... | 57 | | 30 | 44 | 48 |
| 99TH.....GM..... | 113 | | 57 | 85 | 105 |
| MAXIMUM.....GM..... | 227 | 57 | 85 | 113 | 170 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER.... | 10.4 | .7 | 0 | 1.2 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .2 | 2.7 | 0 | .2 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 14.0 | 12.3 | 12.7 | 11.6 | 6.4 | 6.4 | 6.2 | 5.0 | 3.6 | 3.0 | 3.5 | 3.7 |
| 9.5 | 8.5 | 8.8 | 8.3 | 4.1 | 4.7 | 4.4 | 3.9 | 2.1 | 2.1 | 2.4 | 1.9 |
| 3.6 | 2.7 | 3.0 | 2.6 | 1.6 | 1.2 | 1.1 | 1.0 | .6 | .9 | .5 | .9 |
| .9 | 1.2 | .9 | .6 | .7 | .4 | .7 | .1 | .9 | 0 | .6 | .9 |
| 1.6 | 1.6 | 1.6 | 1.4 | 1.7 | 1.5 | 1.5 | 1.3 | 1.7 | 1.4 | 2.4 | 1.9 |
| .9 | 1.0 | 1.0 | .7 | .9 | 1.0 | .9 | .6 | 1.0 | .7 | 2.8 | 1.0 |
| 7.0 | 7.0 | 6.0 | 5.0 | 6.0 | 7.0 | 6.0 | 4.0 | 4.0 | 4.0 | 12.0 | 4.0 |
| 15 | 13 | 16 | 11 | 18 | 13 | 13 | 9 | 13 | 7 | 16 | 8 |
| 16 | 12 | 16 | 11 | 15 | 16 | 11 | 9 | 12 | 4 | 25 | 4 |
| 2 | 2 | 2 | 2 | 3 | 2 | 3 | 2 | | 2 | | |
| 5 | 5 | 7 | 5 | 7 | 5 | 5 | 5 | | 5 | | |
| 9 | 9 | 12 | 9 | 14 | 8 | 9 | 7 | | 7 | | |
| 19 | 14 | 19 | 12 | 28 | 14 | 15 | 9 | | 9 | | |
| 28 | 28 | 35 | 19 | 37 | 28 | 28 | 16 | | 12 | | |
| 47 | 38 | 43 | 23 | 44 | 38 | 38 | 22 | | 14 | | |
| 76 | 70 | 57 | 57 | 66 | 76 | 57 | 38 | | 16 | | |
| 30 | 26 | 38 | 23 | 35 | 29 | 26 | 21 | 22 | 17 | 26 | 14 |
| 36 | 23 | 35 | 14 | 27 | 31 | 20 | 17 | 9 | 8 | 24 | 10 |
| 35 | 30 | 30 | 27 | 40 | 32 | 34 | 21 | 25 | 16 | 25 | 12 |
| 24 | 24 | 15 | 28 | 19 | 41 | 18 | 22 | 8 | 5 | 10 | 2 |
| 28 | 25 | 38 | 18 | 37 | 24 | 25 | 22 | 24 | 0 | 48 | 14 |
| 24 | 15 | 34 | 8 | 17 | 13 | 14 | 12 | 19 | 0 | 47 | (2) |
| 340 | 170 | 255 | 198 | 227 | 340 | 170 | 255 | 98 | 28 | 126 | 49 |
| 397 | 227 | 468 | 340 | 290 | 567 | 175 | 354 | 182 | 49 | 308 | 49 |
| 27 | 24 | 30 | 22 | 33 | 26 | 26 | 20 | 23 | 15 | 21 | 12 |
| 24 | 19 | 25 | 21 | 20 | 28 | 16 | 16 | 11 | 6 | 12 | 4 |
| 7 | 7 | 7 | 7 | 10 | 7 | 7 | 7 | | 7 | | |
| 14 | 14 | 14 | 14 | 21 | 14 | 14 | 14 | | 14 | | |
| 21 | 21 | 22 | 21 | 28 | 21 | 21 | 14 | | 14 | | |
| 28 | 28 | 35 | 28 | 42 | 28 | 33 | 21 | | 14 | | |
| 42 | 42 | 57 | 35 | 57 | 44 | 42 | 35 | | 28 | | |
| 71 | 56 | 84 | 42 | 84 | 66 | 56 | 44 | | 28 | | |
| 125 | 113 | 168 | 170 | 105 | 170 | 85 | 85 | | 28 | | |
| 170 | 170 | 170 | 198 | 113 | 227 | 126 | 170 | 49 | 28 | 42 | 28 |
| 0 | 4.0 | 1.9 | 1.0 | 0 | 0 | 1.6 | 0 | 0 | 0 | 0 | 0 |
| 0 | .9 | .7 | .4 | 0 | 0 | .3 | 0 | 0 | 0 | 0 | 0 |

TABLE 5.19.--OATMEAL COOKIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.5 | .9 | 4.6 | 4.3 | 6.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.5 | .6 | 3.4 | 3.5 | 4.9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .7 | .3 | 1.0 | .5 | 1.3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | .2 | .3 | .4 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.6 | 1.7 | 1.5 | 1.4 | 1.5 |
| STANDARD DEVIATION.....DO..... | 1.0 | .9 | 1.1 | .9 | .9 |
| MAXIMUM.....DO..... | 7.0 | 3.0 | 7.0 | 5.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 18 | 9 | 14 | 13 | 18 |
| STANDARD DEVIATION.....GM..... | 18 | 4 | 14 | 8 | 16 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 4 | | | 4 | 4 |
| 25TH.....GM..... | 9 | | | 9 | 9 |
| 50TH.....GM..... | 13 | | | 9 | 13 |
| 75TH.....GM..... | 22 | | | 17 | 22 |
| 90TH.....GM..... | 39 | | | 26 | 39 |
| 95TH.....GM..... | 52 | | | 30 | 52 |
| 99TH.....GM..... | 91 | | | 43 | 82 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 38 | 19 | 26 | 31 | 36 |
| STANDARD DEVIATION.....GM..... | 29 | 6 | 12 | 15 | 24 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 44 | 19 | 40 | 35 | 44 |
| STANDARD DEVIATION.....GM..... | 31 | 0 | 25 | 11 | 26 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 44 | 0 | 54 | 32 | 44 |
| STANDARD DEVIATION.....GM..... | 28 | 0 | 15 | 7 | 25 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 364 | 26 | 165 | 91 | 170 |
| MAXIMUM OVER 3 DAYS.....GM..... | 429 | 39 | 196 | 130 | 281 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 36 | 15 | 28 | 27 | 34 |
| STANDARD DEVIATION.....GM..... | 26 | 5 | 18 | 12 | 20 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 13 | | | 13 | 13 |
| 25TH.....GM..... | 23 | | | 13 | 26 |
| 50TH.....GM..... | 26 | | | 26 | 26 |
| 75TH.....GM..... | 39 | | | 39 | 39 |
| 90TH.....GM..... | 65 | | | 39 | 57 |
| 95TH.....GM..... | 78 | | | 52 | 65 |
| 99TH.....GM..... | 130 | | | 57 | 113 |
| MAXIMUM.....GM..... | 338 | 26 | 165 | 65 | 170 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 7.2 | 0 | .9 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 0 | 1.2 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.0 | 4.1 | 4.3 | 3.8 | 3.0 | 2.7 | 3.4 | 2.5 | 4.4 | 2.5 | 5.1 | 3.9 |
| 3.8 | 3.2 | 3.6 | 3.4 | 2.3 | 1.9 | 2.1 | 1.8 | 2.7 | 1.9 | 3.0 | 2.2 |
| 1.1 | .6 | .7 | .3 | .6 | .7 | .9 | .7 | 1.3 | .3 | 1.6 | .9 |
| .2 | .3 | .1 | .2 | .1 | .2 | .5 | .1 | .5 | .2 | .5 | .7 |
| 1.5 | 1.4 | 1.4 | 1.2 | 1.5 | 1.6 | 1.8 | 1.5 | 1.7 | 1.5 | 1.9 | 2.0 |
| .9 | .8 | .8 | .6 | 1.1 | 1.2 | 1.0 | .7 | .9 | 1.0 | 1.3 | 1.4 |
| 5.0 | 4.0 | 4.0 | 3.0 | 7.0 | 7.0 | 5.0 | 5.0 | 4.0 | 5.0 | 5.0 | 6.0 |
| 19 | 16 | 22 | 14 | 23 | 20 | 25 | 14 | 16 | 17 | 18 | 19 |
| 19 | 11 | 20 | 11 | 18 | 21 | 23 | 10 | 9 | 29 | 11 | 17 |
| 4 | 4 | 4 | 4 | 6 | 4 | 4 | 4 | | | | |
| 9 | 9 | 9 | 4 | 9 | 9 | 9 | 9 | | | | |
| 13 | 13 | 17 | 13 | 17 | 13 | 17 | 9 | | | | |
| 22 | 22 | 26 | 17 | 30 | 27 | 35 | 17 | | | | |
| 35 | 28 | 43 | 35 | 48 | 48 | 48 | 28 | | | | |
| 52 | 35 | 82 | 43 | 61 | 56 | 78 | 38 | | | | |
| 112 | 52 | 86 | 48 | 91 | 113 | 113 | 52 | | | | |
| 38 | 37 | 52 | 37 | 51 | 37 | 48 | 31 | 32 | 26 | 38 | 35 |
| 30 | 27 | 44 | 30 | 35 | 29 | 40 | 18 | 18 | 13 | 15 | 20 |
| 58 | 37 | 59 | 53 | 53 | 52 | 48 | 33 | 35 | 38 | 43 | 35 |
| 43 | 18 | 39 | 20 | 24 | 43 | 33 | 19 | 8 | 2 | 20 | 37 |
| 28 | 30 | 61 | 15 | 63 | 50 | 53 | 32 | 27 | 80 | 13 | 36 |
| 4 | 6 | 10 | 2 | 31 | 11 | 31 | 7 | 1 | 65 | 0 | 15 |
| 260 | 195 | 247 | 143 | 161 | 170 | 234 | 113 | 71 | 364 | 78 | 113 |
| 368 | 195 | 257 | 143 | 273 | 340 | 397 | 156 | 91 | 429 | 130 | 227 |
| 38 | 33 | 47 | 34 | 45 | 37 | 43 | 30 | 28 | 35 | 29 | 28 |
| 29 | 23 | 28 | 23 | 28 | 30 | 28 | 17 | 10 | 56 | 13 | 21 |
| 13 | 13 | 13 | 13 | 13 | 13 | 13 | 13 | | | | |
| 26 | 23 | 26 | 13 | 26 | 13 | 26 | 13 | | | | |
| 26 | 26 | 39 | 26 | 39 | 26 | 39 | 26 | | | | |
| 46 | 39 | 65 | 52 | 52 | 46 | 52 | 39 | | | | |
| 78 | 65 | 78 | 65 | 78 | 78 | 78 | 52 | | | | |
| 85 | 78 | 113 | 78 | 92 | 113 | 104 | 69 | | | | |
| 184 | 104 | 127 | 130 | 156 | 170 | 156 | 78 | | | | |
| 260 | 195 | 130 | 130 | 161 | 170 | 227 | 113 | 65 | 338 | 52 | 113 |
| .8 | 0 | 0 | 0 | 0 | 0 | 0 | 3.0 | 0 | 2.6 | 0 | 0 |
| .5 | 0 | 0 | 0 | 0 | 0 | 0 | 1.1 | 0 | 4.0 | 0 | 0 |

TABLE 5.20.--PEANUT BUTTER COOKIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.4 | .7 | 2.4 | 2.9 | 5.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.9 | .7 | 1.6 | 2.2 | 4.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | 0 | .6 | .6 | .5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | .1 | .1 | .4 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER.... | 1.4 | 1.0 | 1.6 | 1.5 | 1.4 |
| STANDARD DEVIATION.....DO..... | .8 | .0 | .9 | .9 | 1.0 |
| MAXIMUM.....DO..... | 7.0 | 1.0 | 5.0 | 6.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 15 | 6 | 13 | 14 | 12 |
| STANDARD DEVIATION.....GM..... | 15 | 2 | 7 | 9 | 13 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 4 | | | 4 | 4 |
| 25TH.....GM..... | 6 | | | 8 | 4 |
| 50TH.....GM..... | 9 | | | 12 | 8 |
| 75TH.....GM..... | 17 | | | 19 | 15 |
| 90TH.....GM..... | 28 | | | 24 | 20 |
| 95TH.....GM..... | 40 | | | 28 | 46 |
| 99TH.....GM..... | 72 | | | 47 | 53 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 33 | 19 | 34 | 35 | 24 |
| STANDARD DEVIATION.....GM..... | 28 | 6 | 21 | 19 | 16 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 35 | 0 | 23 | 30 | 35 |
| STANDARD DEVIATION.....GM..... | 31 | 0 | 7 | 18 | 21 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 39 | 0 | 22 | 35 | 41 |
| STANDARD DEVIATION.....GM..... | 20 | 0 | 0 | 8 | 18 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 624 | 24 | 81 | 85 | 113 |
| MAXIMUM OVER 3 DAYS.....GM..... | 636 | 24 | 81 | 142 | 216 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 31 | 19 | 24 | 29 | 26 |
| STANDARD DEVIATION.....GM..... | 29 | 6 | 11 | 16 | 17 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 12 | | | 12 | 12 |
| 25TH.....GM..... | 12 | | | 12 | 12 |
| 50TH.....GM..... | 24 | | | 24 | 24 |
| 75TH.....GM..... | 36 | | | 36 | 36 |
| 90TH.....GM..... | 57 | | | 52 | 48 |
| 95TH.....GM..... | 72 | | | 60 | 52 |
| 99TH.....GM..... | 113 | | | 85 | 113 |
| MAXIMUM.....GM..... | 624 | 24 | 60 | 85 | 113 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER.... | 0 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 0 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.4 | 5.0 | 3.7 | 3.8 | 1.9 | 1.8 | 1.4 | 1.4 | 2.1 | 1.4 | .3 | 1.3 |
| 4.5 | 4.5 | 2.8 | 3.2 | 1.5 | 1.3 | 1.0 | 1.1 | 1.7 | .7 | .3 | .5 |
| .8 | .5 | .6 | .5 | .3 | .4 | .2 | .3 | .4 | .5 | 0 | .7 |
| .1 | 0 | .3 | (2) | .1 | .1 | .1 | .1 | 0 | .3 | 0 | .2 |
| 1.2 | 1.2 | 1.5 | 1.2 | 1.4 | 1.6 | 1.4 | 1.5 | 1.4 | 2.2 | 1.0 | 2.0 |
| .6 | .4 | .9 | .5 | .6 | 1.0 | .7 | .9 | .7 | 1.8 | .0 | .7 |
| 4.0 | 3.0 | 5.0 | 3.0 | 3.0 | 5.0 | 4.0 | 6.0 | 3.0 | 7.0 | 1.0 | 3.0 |
| 13 | 11 | 19 | 12 | 22 | 16 | 20 | 13 | 15 | 17 | 8 | 12 |
| 21 | 9 | 18 | 9 | 18 | 14 | 17 | 9 | 11 | 21 | 0 | 7 |
| 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | | | | |
| 4 | 4 | 8 | 4 | 8 | 4 | 8 | 8 | | | | |
| 8 | 8 | 12 | 8 | 16 | 12 | 12 | 8 | | | | |
| 15 | 14 | 24 | 16 | 29 | 24 | 20 | 16 | | | | |
| 24 | 20 | 51 | 24 | 36 | 38 | 52 | 24 | | | | |
| 24 | 33 | 63 | 33 | 64 | 44 | 64 | 38 | | | | |
| 48 | 48 | 83 | 44 | 89 | 80 | 66 | 56 | | | | |
| 27 | 29 | 42 | 31 | 55 | 37 | 40 | 32 | 32 | 17 | 24 | 30 |
| 17 | 23 | 38 | 24 | 53 | 32 | 29 | 22 | 15 | 6 | 0 | 10 |
| 49 | 35 | 44 | 32 | 50 | 37 | 37 | 26 | 51 | 21 | 0 | 13 |
| 71 | 18 | 27 | 15 | 19 | 28 | 20 | 14 | 15 | 7 | 0 | 2 |
| 35 | 0 | 54 | 24 | 41 | 26 | 62 | 24 | 0 | 52 | 0 | 28 |
| 7 | 0 | 22 | 0 | 10 | 12 | 6 | 15 | 0 | 24 | 0 | 0 |
| 624 | 170 | 227 | 106 | 268 | 168 | 156 | 113 | 72 | 120 | 24 | 36 |
| 636 | 170 | 248 | 132 | 268 | 240 | 197 | 168 | 120 | 216 | 24 | 85 |
| 32 | 28 | 40 | 30 | 48 | 30 | 41 | 26 | 32 | 24 | 24 | 18 |
| 53 | 20 | 32 | 21 | 42 | 23 | 28 | 16 | 14 | 11 | 0 | 9 |
| 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | | | | |
| 12 | 12 | 24 | 12 | 24 | 12 | 24 | 12 | | | | |
| 24 | 24 | 28 | 24 | 36 | 24 | 36 | 24 | | | | |
| 36 | 36 | 48 | 36 | 50 | 36 | 48 | 36 | | | | |
| 48 | 56 | 72 | 60 | 72 | 72 | 72 | 57 | | | | |
| 72 | 72 | 104 | 72 | 96 | 84 | 104 | 57 | | | | |
| 96 | 85 | 156 | 106 | 268 | 113 | 156 | 72 | | | | |
| 624 | 144 | 227 | 106 | 268 | 113 | 156 | 85 | 57 | 48 | 24 | 36 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 5.21.--SUGAR COOKIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.2 | .9 | 3.6 | 4.4 | 5.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.3 | .7 | 2.6 | 3.5 | 4.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .7 | 0 | .6 | .7 | 1.1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | .2 | .4 | .3 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.5 | 1.4 | 1.6 | 1.4 | 1.4 |
| STANDARD DEVIATION.....DO..... | .8 | .8 | 1.1 | .8 | .9 |
| MAXIMUM.....DO..... | 6.0 | 3.0 | 5.0 | 5.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 18 | 12 | 15 | 17 | 16 |
| STANDARD DEVIATION.....GM..... | 16 | 8 | 16 | 17 | 13 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 5 | | | 5 | 5 |
| 25TH.....GM..... | 8 | | | 9 | 5 |
| 50TH.....GM..... | 11 | | | 11 | 11 |
| 75TH.....GM..... | 21 | | | 21 | 21 |
| 90TH.....GM..... | 38 | | | 43 | 32 |
| 95TH.....GM..... | 53 | | | 57 | 48 |
| 99TH.....GM..... | 75 | | | 87 | 64 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 40 | 27 | 28 | 43 | 34 |
| STANDARD DEVIATION.....GM..... | 34 | 13 | 13 | 42 | 24 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 41 | 0 | 27 | 28 | 41 |
| STANDARD DEVIATION.....GM..... | 24 | 0 | 15 | 14 | 21 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 42 | 27 | 49 | 52 | 48 |
| STANDARD DEVIATION.....GM..... | 22 | 0 | 25 | 20 | 9 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 320 | 48 | 113 | 224 | 170 |
| MAXIMUM OVER 3 DAYS.....GM..... | 340 | 80 | 227 | 260 | 192 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 37 | 27 | 28 | 37 | 34 |
| STANDARD DEVIATION.....GM..... | 27 | 10 | 14 | 26 | 21 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 8 | | | 8 | 16 |
| 25TH.....GM..... | 16 | | | 16 | 16 |
| 50TH.....GM..... | 32 | | | 32 | 32 |
| 75TH.....GM..... | 48 | | | 48 | 48 |
| 90TH.....GM..... | 64 | | | 64 | 57 |
| 95TH.....GM..... | 81 | | | 96 | 64 |
| 99TH.....GM..... | 142 | | | 128 | 85 |
| MAXIMUM.....GM..... | 320 | 48 | 64 | 142 | 170 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 9.1 | 0 | 1.1 | 0 | 1.2 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .5 | 0 | 1.8 | 0 | .8 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.9 | 5.0 | 3.1 | 3.1 | 1.3 | 2.0 | 2.9 | 3.0 | 4.1 | 3.3 | 6.4 | 5.6 |
| 3.3 | 3.9 | 2.3 | 2.2 | .9 | 1.6 | 1.9 | 2.3 | 2.3 | 2.3 | 4.0 | 3.0 |
| .6 | .9 | .6 | .7 | .4 | .3 | .8 | .6 | .9 | .8 | 1.4 | 1.5 |
| .1 | .2 | .2 | .1 | (2) | .1 | .1 | .1 | .9 | .2 | 1.0 | 1.2 |
| 1.2 | 1.4 | 1.4 | 1.4 | 1.5 | 1.4 | 1.5 | 1.4 | 1.9 | 1.4 | 1.7 | 1.9 |
| .5 | .8 | .8 | .7 | .7 | .7 | .7 | .7 | 1.3 | .6 | 1.0 | 1.2 |
| 4.0 | 5.0 | 5.0 | 3.0 | 4.0 | 4.0 | 4.0 | 4.0 | 5.0 | 3.0 | 4.0 | 6.0 |
| 15 | 18 | 25 | 18 | 27 | 16 | 24 | 15 | 20 | 13 | 22 | 16 |
| 13 | 16 | 21 | 17 | 19 | 12 | 18 | 13 | 18 | 8 | 20 | 13 |
| 3 | 3 | | | 5 | 5 | 4 | 5 | | 5 | | 5 |
| 5 | 5 | | | 11 | 9 | 11 | 5 | | 9 | | 11 |
| 11 | 11 | | | 21 | 11 | 21 | 11 | | 11 | | 11 |
| 19 | 21 | | | 43 | 21 | 32 | 16 | | 18 | | 18 |
| 32 | 43 | | | 53 | 36 | 47 | 27 | | 27 | | 32 |
| 47 | 54 | | | 64 | 48 | 54 | 38 | | 37 | | 59 |
| 59 | 64 | | | 95 | 48 | 113 | 69 | | 37 | | 59 |
| 39 | 36 | 64 | 34 | 71 | 40 | 53 | 35 | 27 | 32 | 35 | 32 |
| 34 | 26 | 61 | 23 | 62 | 28 | 36 | 27 | 14 | 22 | 22 | 15 |
| 36 | 58 | 41 | 53 | 51 | 38 | 46 | 38 | 39 | 28 | 64 | 20 |
| 21 | 26 | 19 | 39 | 16 | 18 | 21 | 27 | 11 | 9 | 39 | 9 |
| 37 | 50 | 60 | 29 | 45 | 38 | 73 | 29 | 40 | 23 | 34 | 33 |
| 0 | 11 | 7 | 3 | 0 | 13 | 26 | 7 | 24 | 10 | 13 | 17 |
| 162 | 144 | 320 | 240 | 284 | 144 | 142 | 162 | 112 | 112 | 135 | 96 |
| 176 | 255 | 320 | 272 | 284 | 144 | 340 | 227 | 256 | 112 | 216 | 176 |
| 37 | 39 | 53 | 37 | 55 | 36 | 48 | 32 | 31 | 29 | 38 | 25 |
| 29 | 25 | 47 | 32 | 44 | 23 | 29 | 24 | 15 | 13 | 21 | 12 |
| 8 | 8 | | | 16 | 13 | 8 | 8 | | 16 | | 8 |
| 16 | 16 | | | 27 | 16 | 28 | 16 | | 16 | | 16 |
| 32 | 32 | | | 48 | 32 | 48 | 28 | | 28 | | 24 |
| 48 | 54 | | | 64 | 48 | 64 | 32 | | 32 | | 32 |
| 64 | 64 | | | 128 | 64 | 96 | 64 | | 48 | | 32 |
| 81 | 81 | | | 128 | 85 | 108 | 80 | | 54 | | 48 |
| 162 | 142 | | | 284 | 113 | 135 | 144 | | 64 | | 64 |
| 162 | 144 | 320 | 240 | 284 | 113 | 142 | 162 | 64 | 64 | 81 | 64 |
| 0 | .8 | 0 | 0 | 0 | 0 | 1.7 | 2.1 | 0 | 0 | 0 | 2.1 |
| 0 | .5 | 0 | 0 | 0 | 0 | .8 | .7 | 0 | 0 | 0 | 2.0 |

TABLE 5.22.--PIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 11.9 | .5 | 4.7 | 7.1 | 8.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 9.3 | .5 | 4.0 | 5.9 | 7.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.1 | 0 | .7 | 1.2 | 1.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .4 | 0 | 0 | (2) | (2) |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.3 | 1.2 | 1.3 | 1.3 | 1.2 |
| STANDARD DEVIATION.....DO..... | .6 | .4 | .6 | .5 | .4 |
| MAXIMUM.....DO..... | 5.0 | 2.0 | 4.0 | 4.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 57 | 22 | 37 | 44 | 45 |
| STANDARD DEVIATION.....GM..... | 38 | 15 | 24 | 24 | 31 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 19 | | | 15 | 11 |
| 25TH.....GM..... | 34 | | | 28 | 25 |
| 50TH.....GM..... | 45 | | | 39 | 39 |
| 75TH.....GM..... | 70 | | | 56 | 50 |
| 90TH.....GM..... | 102 | | | 79 | 77 |
| 95TH.....GM..... | 135 | | | 80 | 110 |
| 99TH.....GM..... | 197 | | | 133 | 180 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 136 | 65 | 103 | 117 | 113 |
| STANDARD DEVIATION.....GM..... | 70 | 46 | 56 | 57 | 58 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 144 | 0 | 84 | 102 | 134 |
| STANDARD DEVIATION.....GM..... | 68 | 0 | 57 | 40 | 73 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 134 | 0 | 0 | 133 | 133 |
| STANDARD DEVIATION.....GM..... | 51 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 960 | 140 | 228 | 275 | 450 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,283 | 140 | 405 | 480 | 675 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 129 | 53 | 88 | 106 | 116 |
| STANDARD DEVIATION.....GM..... | 60 | 30 | 50 | 48 | 58 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 57 | | | 43 | 32 |
| 25TH.....GM..... | 97 | | | 73 | 69 |
| 50TH.....GM..... | 120 | | | 101 | 114 |
| 75TH.....GM..... | 150 | | | 130 | 135 |
| 90TH.....GM..... | 195 | | | 158 | 187 |
| 95TH.....GM..... | 236 | | | 228 | 228 |
| 99TH.....GM..... | 360 | | | 264 | 270 |
| MAXIMUM.....GM..... | 896 | 102 | 228 | 264 | 342 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 1054.2 | 0 | 11.4 | 26.6 | 24.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 17.6 | 0 | 18.2 | 17.3 | 14.4 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 10.2 | 10.6 | 13.6 | 9.2 | 13.6 | 10.0 | 15.7 | 11.9 | 19.5 | 14.6 | 17.2 | 15.6 |
| 8.3 | 9.1 | 10.9 | 8.2 | 10.6 | 8.5 | 11.7 | 9.3 | 13.8 | 11.1 | 10.7 | 11.3 |
| 1.8 | 1.3 | 2.4 | .8 | 2.7 | 1.2 | 3.3 | 2.1 | 4.9 | 3.1 | 4.7 | 3.1 |
| .1 | .2 | .3 | .2 | .3 | .2 | .7 | .5 | .8 | .3 | 1.8 | 1.2 |
| 1.3 | 1.2 | 1.3 | 1.2 | 1.3 | 1.2 | 1.4 | 1.4 | 1.4 | 1.4 | 1.7 | 1.4 |
| .6 | .6 | .6 | .5 | .7 | .6 | .6 | .6 | .7 | .7 | .7 | .7 |
| 4.0 | 5.0 | 4.0 | 3.0 | 5.0 | 5.0 | 4.0 | 4.0 | 5.0 | 4.0 | 3.0 | 4.0 |
| 58 | 53 | 62 | 50 | 65 | 51 | 60 | 54 | 67 | 61 | 76 | 58 |
| 34 | 34 | 41 | 28 | 47 | 31 | 37 | 33 | 49 | 47 | 46 | 35 |
| 22 | 19 | 23 | 20 | 28 | 19 | 20 | 19 | 22 | 20 | 20 | 19 |
| 38 | 32 | 39 | 35 | 38 | 33 | 38 | 34 | 38 | 34 | 43 | 34 |
| 45 | 43 | 45 | 43 | 45 | 43 | 45 | 45 | 51 | 45 | 70 | 45 |
| 75 | 65 | 80 | 51 | 76 | 61 | 76 | 65 | 86 | 71 | 94 | 68 |
| 101 | 88 | 110 | 79 | 120 | 88 | 119 | 95 | 132 | 126 | 135 | 126 |
| 135 | 150 | 135 | 88 | 166 | 108 | 137 | 113 | 151 | 152 | 160 | 132 |
| 161 | 168 | 225 | 169 | 253 | 163 | 160 | 190 | 280 | 251 | 246 | 158 |
| 150 | 135 | 146 | 131 | 151 | 128 | 137 | 129 | 146 | 136 | 172 | 134 |
| 83 | 71 | 67 | 57 | 82 | 56 | 64 | 63 | 76 | 96 | 102 | 68 |
| 138 | 147 | 179 | 134 | 167 | 128 | 151 | 127 | 160 | 155 | 141 | 116 |
| 49 | 69 | 79 | 68 | 93 | 46 | 59 | 49 | 76 | 80 | 44 | 50 |
| 118 | 133 | 100 | 118 | 178 | 162 | 119 | 118 | 152 | 156 | 140 | 131 |
| 44 | 30 | 15 | 23 | 48 | 30 | 25 | 59 | 96 | 29 | 63 | 6 |
| 680 | 454 | 466 | 506 | 900 | 342 | 472 | 672 | 506 | 960 | 506 | 384 |
| 680 | 724 | 802 | 675 | 1,152 | 619 | 870 | 830 | 1,283 | 960 | 737 | 474 |
| 133 | 129 | 144 | 126 | 147 | 123 | 131 | 119 | 140 | 130 | 137 | 121 |
| 55 | 62 | 66 | 47 | 77 | 53 | 51 | 54 | 66 | 63 | 74 | 50 |
| 59 | 57 | 65 | 60 | 72 | 56 | 57 | 49 | 64 | 60 | 60 | 57 |
| 101 | 93 | 103 | 101 | 105 | 93 | 101 | 85 | 105 | 90 | 98 | 85 |
| 128 | 118 | 130 | 128 | 130 | 118 | 128 | 118 | 130 | 125 | 126 | 120 |
| 152 | 140 | 158 | 135 | 158 | 135 | 152 | 140 | 158 | 158 | 152 | 135 |
| 227 | 225 | 249 | 189 | 236 | 192 | 195 | 170 | 203 | 192 | 227 | 189 |
| 252 | 236 | 280 | 225 | 284 | 234 | 228 | 225 | 240 | 210 | 270 | 203 |
| 284 | 384 | 360 | 270 | 450 | 283 | 312 | 284 | 420 | 448 | 454 | 288 |
| 360 | 450 | 405 | 350 | 896 | 342 | 450 | 672 | 480 | 480 | 506 | 384 |
| 50.3 | 47.3 | 50.3 | 34.2 | 154.8 | 127.7 | 197.1 | 186.9 | 27.1 | 52.1 | 12.1 | 51.5 |
| 18.1 | 16.8 | 20.4 | 21.3 | 21.8 | 19.3 | 18.4 | 16.4 | 8.6 | 14.7 | 7.9 | 23.3 |

TABLE 5.23.--FRUIT PIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 7.1 | 0 | 2.1 | 4.0 | 4.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.8 | 0 | 1.8 | 3.4 | 4.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.1 | 0 | .3 | .6 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 0 | 1.3 | 1.2 | 1.1 |
| STANDARD DEVIATION.....DO..... | .6 | 0 | .7 | .5 | .3 |
| MAXIMUM.....DO..... | 5.0 | 0 | 4.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 57 | 0 | 35 | 43 | 42 |
| STANDARD DEVIATION.....GM..... | 39 | 0 | 27 | 25 | 29 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 21 | | | 16 | 15 |
| 25TH.....GM..... | 35 | | | 25 | 25 |
| 50TH.....GM..... | 45 | | | 35 | 39 |
| 75TH.....GM..... | 67 | | | 56 | 45 |
| 90TH.....GM..... | 104 | | | 79 | 76 |
| 95TH.....GM..... | 135 | | | 80 | 90 |
| 99TH.....GM..... | 208 | | | 160 | 180 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 138 | 0 | 81 | 111 | 114 |
| STANDARD DEVIATION.....GM..... | 72 | 0 | 40 | 56 | 53 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 152 | 0 | 125 | 116 | 201 |
| STANDARD DEVIATION.....GM..... | 70 | 0 | 56 | 46 | 82 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 150 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 55 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 960 | 0 | 203 | 270 | 450 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,283 | 0 | 405 | 480 | 675 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 133 | 0 | 79 | 106 | 120 |
| STANDARD DEVIATION.....GM..... | 60 | 0 | 31 | 48 | 56 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 57 | | | 48 | 50 |
| 25TH.....GM..... | 101 | | | 75 | 85 |
| 50TH.....GM..... | 128 | | | 101 | 118 |
| 75TH.....GM..... | 158 | | | 128 | 135 |
| 90TH.....GM..... | 200 | | | 158 | 225 |
| 95TH.....GM..... | 236 | | | 236 | 236 |
| 99TH.....GM..... | 384 | | | 240 | 270 |
| MAXIMUM.....GM..... | 540 | 0 | 158 | 240 | 270 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 531.4 | 0 | 4.2 | 5.9 | 16.2 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 15.4 | 0 | 14.5 | 7.1 | 17.0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 6.7 | 6.0 | 8.2 | 4.8 | 8.1 | 5.4 | 10.2 | 6.7 | 11.9 | 9.5 | 11.7 | 10.7 |
| 5.6 | 5.6 | 7.0 | 4.4 | 6.6 | 4.6 | 8.3 | 5.7 | 8.5 | 6.7 | 7.6 | 8.1 |
| 1.0 | .4 | 1.0 | .3 | 1.3 | .7 | 1.8 | .8 | 3.2 | 2.4 | 3.2 | 2.0 |
| .1 | .1 | .2 | .1 | .2 | .1 | .2 | .2 | .2 | .3 | .9 | .6 |
| 1.3 | 1.2 | 1.2 | 1.1 | 1.3 | 1.2 | 1.3 | 1.3 | 1.4 | 1.5 | 1.6 | 1.4 |
| .6 | .5 | .6 | .4 | .6 | .5 | .6 | .6 | .7 | .7 | .7 | .7 |
| 4.0 | 4.0 | 4.0 | 3.0 | 5.0 | 4.0 | 4.0 | 4.0 | 5.0 | 4.0 | 3.0 | 4.0 |
| 58 | 51 | 60 | 51 | 65 | 52 | 57 | 51 | 72 | 65 | 77 | 60 |
| 36 | 38 | 41 | 30 | 47 | 31 | 34 | 32 | 55 | 49 | 47 | 35 |
| 23 | 19 | 23 | 24 | 28 | 23 | 23 | 19 | 23 | 23 | 23 | 23 |
| 36 | 28 | 39 | 38 | 39 | 34 | 38 | 34 | 39 | 35 | 45 | 38 |
| 45 | 42 | 45 | 43 | 45 | 45 | 45 | 45 | 53 | 45 | 70 | 51 |
| 68 | 53 | 65 | 53 | 76 | 63 | 68 | 58 | 90 | 87 | 90 | 70 |
| 101 | 96 | 113 | 79 | 120 | 90 | 105 | 90 | 140 | 126 | 135 | 128 |
| 143 | 151 | 143 | 117 | 180 | 105 | 128 | 105 | 157 | 147 | 169 | 132 |
| 165 | 177 | 225 | 225 | 292 | 163 | 160 | 190 | 280 | 251 | 246 | 158 |
| 150 | 130 | 144 | 134 | 155 | 132 | 138 | 128 | 156 | 141 | 175 | 147 |
| 90 | 76 | 60 | 47 | 88 | 54 | 65 | 56 | 89 | 104 | 88 | 74 |
| 147 | 204 | 216 | 165 | 170 | 133 | 154 | 131 | 169 | 152 | 134 | 125 |
| 54 | 84 | 81 | 92 | 91 | 49 | 55 | 52 | 84 | 76 | 40 | 54 |
| 103 | 157 | 101 | 135 | 177 | 172 | 118 | 138 | 225 | 156 | 192 | 135 |
| 42 | 0 | 18 | 0 | 41 | 27 | 10 | 63 | 132 | 29 | 47 | 6 |
| 680 | 454 | 450 | 450 | 500 | 316 | 472 | 450 | 506 | 960 | 506 | 384 |
| 680 | 724 | 754 | 675 | 900 | 619 | 675 | 830 | 1,283 | 960 | 737 | 474 |
| 133 | 132 | 148 | 135 | 152 | 129 | 135 | 120 | 149 | 133 | 141 | 131 |
| 54 | 70 | 65 | 47 | 73 | 50 | 53 | 48 | 75 | 60 | 73 | 54 |
| 57 | 57 | 68 | 71 | 75 | 64 | 68 | 53 | 68 | 68 | 60 | 68 |
| 101 | 85 | 113 | 105 | 113 | 96 | 105 | 85 | 112 | 96 | 105 | 96 |
| 128 | 118 | 135 | 128 | 135 | 118 | 128 | 118 | 135 | 126 | 126 | 128 |
| 151 | 135 | 170 | 135 | 158 | 158 | 158 | 135 | 158 | 158 | 158 | 158 |
| 227 | 225 | 237 | 192 | 256 | 192 | 196 | 180 | 225 | 189 | 227 | 192 |
| 256 | 270 | 300 | 227 | 284 | 236 | 236 | 210 | 300 | 196 | 270 | 210 |
| 284 | 384 | 360 | 270 | 450 | 270 | 360 | 236 | 420 | 480 | 506 | 288 |
| 360 | 450 | 405 | 350 | 540 | 283 | 450 | 384 | 480 | 480 | 506 | 384 |
| 26.6 | 17.5 | 22.2 | 19.5 | 71.4 | 55.0 | 117.5 | 89.6 | 21.9 | 27.0 | 8.2 | 28.6 |
| 14.5 | 11.5 | 15.9 | 24.3 | 17.4 | 15.7 | 18.3 | 14.9 | 11.4 | 11.1 | 7.9 | 19.4 |

TABLE 5.24.--CREAM AND CUSTARD PIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.1 | .1 | 2.5 | 2.7 | 3.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.4 | .1 | 2.3 | 2.3 | 2.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .6 | 0 | .1 | .4 | .5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | 0 | (2) | (2) |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 2.0 | 1.2 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .5 | .0 | .4 | .5 | .5 |
| MAXIMUM.....DO..... | 4.0 | 2.0 | 2.0 | 4.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 52 | 47 | 38 | 45 | 45 |
| STANDARD DEVIATION.....GM..... | 31 | 0 | 20 | 24 | 26 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 19 | | | | 9 |
| 25TH.....GM..... | 33 | | | | 30 |
| 50TH.....GM..... | 43 | | | | 41 |
| 75TH.....GM..... | 63 | | | | 51 |
| 90TH.....GM..... | 87 | | | | 76 |
| 95TH.....GM..... | 113 | | | | 95 |
| 99TH.....GM..... | 163 | | | | 133 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 133 | 140 | 117 | 124 | 119 |
| STANDARD DEVIATION.....GM..... | 69 | 0 | 61 | 60 | 63 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 127 | 0 | 27 | 86 | 99 |
| STANDARD DEVIATION.....GM..... | 57 | 0 | 6 | 32 | 39 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 117 | 0 | 0 | 133 | 133 |
| STANDARD DEVIATION.....GM..... | 43 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 896 | 140 | 228 | 275 | 342 |
| MAXIMUM OVER 3 DAYS.....GM..... | 896 | 140 | 228 | 400 | 400 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 124 | 70 | 99 | 109 | 112 |
| STANDARD DEVIATION.....GM..... | 62 | 31 | 61 | 52 | 59 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 49 | | | | 28 |
| 25TH.....GM..... | 95 | | | | 65 |
| 50TH.....GM..... | 120 | | | | 114 |
| 75TH.....GM..... | 138 | | | | 130 |
| 90TH.....GM..... | 188 | | | | 171 |
| 95TH.....GM..... | 228 | | | | 228 |
| 99TH.....GM..... | 340 | | | | 342 |
| MAXIMUM.....GM..... | 896 | 102 | 228 | 264 | 342 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 379.7 | 0 | 6.2 | 14.0 | 8.7 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 19.5 | 0 | 20.9 | 24.5 | 12.5 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 2.8 | 4.0 | 4.3 | 3.6 | 4.4 | 3.7 | 4.7 | 4.5 | 7.3 | 4.7 | 4.2 | 5.2 |
| 2.4 | 3.4 | 3.4 | 3.3 | 3.6 | 3.3 | 3.8 | 3.7 | 5.7 | 4.2 | 2.3 | 4.1 |
| .5 | .5 | .9 | .1 | .8 | .3 | .7 | .8 | 1.6 | .5 | 1.3 | .7 |
| 0 | .1 | 0 | .1 | (2) | .1 | .2 | .1 | 0 | 0 | .5 | .4 |
| 1.2 | 1.2 | 1.3 | 1.1 | 1.3 | 1.2 | 1.3 | 1.3 | 1.3 | 1.2 | 1.7 | 1.3 |
| .5 | .6 | .5 | .5 | .6 | .5 | .6 | .5 | .5 | .4 | .8 | .6 |
| 3.0 | 4.0 | 3.0 | 3.0 | 4.0 | 4.0 | 4.0 | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| 56 | 52 | 57 | 46 | 60 | 47 | 57 | 51 | 53 | 49 | 69 | 45 |
| 26 | 27 | 37 | 22 | 43 | 25 | 32 | 30 | 27 | 36 | 40 | 28 |
| 22 | 17 | 11 | 19 | 22 | 15 | 22 | 19 | 21 | 14 | | 19 |
| 38 | 34 | 38 | 33 | 38 | 31 | 38 | 33 | 34 | 27 | | 33 |
| 51 | 43 | 43 | 42 | 43 | 43 | 43 | 43 | 43 | 43 | | 38 |
| 76 | 75 | 76 | 51 | 70 | 51 | 67 | 59 | 60 | 56 | | 43 |
| 87 | 87 | 101 | 75 | 102 | 87 | 101 | 87 | 93 | 77 | | 87 |
| 114 | 88 | 114 | 85 | 131 | 108 | 137 | 101 | 120 | 112 | | 130 |
| 130 | 150 | 267 | 132 | 253 | 114 | 152 | 183 | 135 | 193 | | 130 |
| 152 | 135 | 138 | 127 | 151 | 127 | 136 | 131 | 132 | 130 | 145 | 108 |
| 66 | 60 | 69 | 47 | 101 | 64 | 55 | 68 | 49 | 88 | 93 | 32 |
| 126 | 123 | 146 | 123 | 141 | 123 | 146 | 120 | 126 | 144 | 144 | 75 |
| 35 | 34 | 76 | 58 | 62 | 39 | 51 | 58 | 51 | 66 | 60 | 33 |
| 0 | 120 | 0 | 105 | 253 | 102 | 124 | 57 | 0 | 0 | 91 | 130 |
| 0 | 30 | 0 | 23 | 0 | 13 | 22 | 0 | 0 | 0 | 6 | (2) |
| 342 | 304 | 466 | 260 | 896 | 342 | 420 | 672 | 304 | 578 | 396 | 203 |
| 390 | 503 | 802 | 397 | 896 | 378 | 473 | 672 | 405 | 578 | 479 | 391 |
| 138 | 125 | 135 | 120 | 141 | 120 | 129 | 119 | 124 | 122 | 121 | 100 |
| 55 | 52 | 65 | 45 | 87 | 60 | 47 | 64 | 46 | 69 | 44 | 37 |
| 65 | 54 | 49 | 42 | 65 | 43 | 60 | 42 | 48 | 43 | | 33 |
| 112 | 95 | 100 | 98 | 100 | 88 | 100 | 85 | 100 | 65 | | 65 |
| 130 | 114 | 130 | 119 | 130 | 114 | 130 | 114 | 130 | 112 | | 101 |
| 152 | 152 | 152 | 130 | 152 | 130 | 152 | 138 | 152 | 152 | | 130 |
| 227 | 222 | 227 | 182 | 224 | 195 | 195 | 176 | 177 | 210 | | 130 |
| 252 | 228 | 284 | 224 | 260 | 252 | 228 | 228 | 203 | 228 | | 152 |
| 342 | 260 | 342 | 228 | 454 | 342 | 228 | 325 | 280 | 448 | | 203 |
| 342 | 264 | 342 | 228 | 896 | 342 | 420 | 672 | 280 | 448 | 240 | 203 |
| 18.0 | 23.1 | 20.9 | 8.6 | 55.8 | 53.9 | 49.7 | 72.6 | 4.2 | 19.3 | 3.9 | 20.5 |
| 24.8 | 21.5 | 27.8 | 14.4 | 25.2 | 23.6 | 16.2 | 17.8 | 4.0 | 19.6 | 10.2 | 29.8 |

TABLE 5.25.--DOUGHNUTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 9.9 | .8 | 6.6 | 8.6 | 10.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 7.3 | .8 | 5.6 | 7.0 | 9.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.0 | 0 | .6 | 1.4 | 1.6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .6 | 0 | .4 | .3 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1</u> /NUMBER..... | 1.4 | 1.2 | 1.3 | 1.3 | 1.3 |
| STANDARD DEVIATION.....DO..... | .7 | .4 | .8 | .6 | .5 |
| MAXIMUM.....DO..... | 7.0 | 2.0 | 5.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1</u> /GM..... | 30 | 15 | 20 | 24 | 25 |
| STANDARD DEVIATION.....GM..... | 23 | 8 | 13 | 15 | 16 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> / | | | | | |
| 5TH.....GM..... | 9 | | 4 | 8 | 9 |
| 25TH.....GM..... | 14 | | 14 | 14 | 14 |
| 50TH.....GM..... | 28 | | 14 | 17 | 19 |
| 75TH.....GM..... | 38 | | 28 | 28 | 29 |
| 90TH.....GM..... | 56 | | 38 | 42 | 42 |
| 95TH.....GM..... | 75 | | 47 | 56 | 56 |
| 99TH.....GM..... | 112 | | 57 | 84 | 84 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 69 | 44 | 50 | 61 | 65 |
| STANDARD DEVIATION.....GM..... | 49 | 25 | 30 | 40 | 37 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 69 | 0 | 50 | 53 | 66 |
| STANDARD DEVIATION.....GM..... | 38 | 0 | 9 | 23 | 27 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 61 | 0 | 48 | 48 | 49 |
| STANDARD DEVIATION.....GM..... | 27 | 0 | 9 | 10 | 7 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 914 | 84 | 142 | 294 | 336 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,276 | 84 | 171 | 294 | 336 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1</u> /GM..... | 64 | 36 | 47 | 54 | 60 |
| STANDARD DEVIATION.....GM..... | 40 | 22 | 26 | 28 | 30 |
| PERCENTILES: <u>1</u> / | | | | | |
| 5TH.....GM..... | 26 | | 13 | 25 | 26 |
| 25TH.....GM..... | 42 | | 28 | 42 | 42 |
| 50TH.....GM..... | 43 | | 42 | 42 | 43 |
| 75TH.....GM..... | 84 | | 52 | 57 | 84 |
| 90TH.....GM..... | 106 | | 84 | 85 | 88 |
| 95TH.....GM..... | 126 | | 112 | 126 | 126 |
| 99TH.....GM..... | 208 | | 142 | 168 | 168 |
| MAXIMUM.....GM..... | 851 | 84 | 142 | 171 | 171 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> / | | | | | |
| EATING OCCASIONS.....NUMBER..... | 20.8 | 0 | 0 | 2.9 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | 0 | 1.5 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 12.0 | 12.9 | 13.2 | 12.9 | 12.5 | 9.1 | 11.7 | 8.6 | 6.8 | 5.2 | 7.1 | 4.4 |
| 9.0 | 10.6 | 9.2 | 10.2 | 8.4 | 6.9 | 7.4 | 6.9 | 4.4 | 3.6 | 4.0 | 2.7 |
| 2.6 | 2.0 | 3.5 | 2.1 | 3.2 | 1.6 | 3.2 | 1.3 | 1.6 | 1.2 | 1.8 | .6 |
| .4 | .4 | .5 | .5 | .9 | .6 | 1.1 | .5 | .8 | .5 | 1.3 | 1.1 |
| 1.4 | 1.3 | 1.4 | 1.3 | 1.5 | 1.3 | 1.6 | 1.3 | 1.6 | 1.5 | 1.7 | 1.7 |
| .7 | .6 | .7 | .6 | .8 | .7 | .8 | .6 | .8 | .9 | .8 | 1.0 |
| 4.0 | 5.0 | 5.0 | 4.0 | 5.0 | 6.0 | 7.0 | 5.0 | 4.0 | 6.0 | 4.0 | 4.0 |
| 32 | 27 | 43 | 27 | 39 | 27 | 31 | 26 | 29 | 26 | 27 | 33 |
| 23 | 20 | 42 | 18 | 29 | 18 | 20 | 19 | 15 | 21 | 15 | 18 |
| 14 | 9 | 14 | 9 | 14 | 9 | 14 | 11 | 14 | 7 | | |
| 14 | 14 | 19 | 14 | 19 | 14 | 14 | 14 | 14 | 14 | | |
| 28 | 21 | 28 | 19 | 28 | 19 | 28 | 19 | 28 | 17 | | |
| 42 | 29 | 56 | 31 | 56 | 33 | 42 | 28 | 42 | 28 | | |
| 56 | 50 | 84 | 52 | 76 | 56 | 56 | 52 | 56 | 42 | | |
| 76 | 71 | 113 | 69 | 86 | 58 | 76 | 63 | 56 | 73 | | |
| 141 | 89 | 196 | 84 | 151 | 84 | 98 | 113 | 56 | 126 | | |
| 77 | 66 | 89 | 64 | 89 | 65 | 65 | 63 | 62 | 56 | 53 | 73 |
| 56 | 42 | 66 | 36 | 81 | 37 | 37 | 40 | 34 | 33 | 18 | 36 |
| 72 | 67 | 108 | 69 | 80 | 60 | 65 | 60 | 63 | 62 | 47 | 82 |
| 38 | 30 | 87 | 31 | 30 | 24 | 25 | 33 | 17 | 45 | 15 | 31 |
| 60 | 68 | 95 | 60 | 77 | 61 | 61 | 61 | 45 | 45 | 48 | 41 |
| 28 | 38 | 55 | 21 | 31 | 19 | 22 | 22 | 4 | 14 | 15 | 12 |
| 448 | 258 | 851 | 284 | 914 | 252 | 252 | 340 | 168 | 252 | 84 | 171 |
| 448 | 486 | 1,276 | 284 | 914 | 344 | 336 | 420 | 168 | 378 | 225 | 227 |
| 67 | 62 | 91 | 63 | 77 | 60 | 60 | 58 | 55 | 51 | 49 | 56 |
| 39 | 36 | 74 | 34 | 52 | 32 | 30 | 32 | 27 | 30 | 17 | 28 |
| 28 | 26 | 42 | 26 | 42 | 28 | 28 | 25 | 25 | 21 | | |
| 42 | 42 | 43 | 42 | 42 | 42 | 42 | 42 | 42 | 42 | | |
| 50 | 43 | 84 | 52 | 84 | 43 | 43 | 42 | 42 | 42 | | |
| 84 | 84 | 113 | 84 | 84 | 84 | 84 | 84 | 66 | 56 | | |
| 126 | 99 | 168 | 104 | 126 | 94 | 84 | 85 | 84 | 112 | | |
| 142 | 126 | 208 | 126 | 168 | 126 | 114 | 114 | 84 | 126 | | |
| 168 | 227 | 336 | 170 | 252 | 171 | 171 | 168 | 168 | 126 | | |
| 340 | 252 | 851 | 284 | 638 | 252 | 252 | 340 | 168 | 227 | 84 | 126 |
| 1.0 | 0 | 0 | 1.3 | 3.0 | 2.0 | 4.4 | 3.4 | 2.8 | 0 | 0 | 0 |
| .3 | 0 | 0 | .5 | .4 | .3 | .5 | .4 | 2.4 | 0 | 0 | 0 |

TABLE 5.26.--SWEET ROLLS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.5 | 0 | 2.0 | 2.7 | 4.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.3 | 0 | 1.8 | 2.2 | 4.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.0 | 0 | .2 | .3 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | 0 | .1 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 0 | 1.2 | 1.3 | 1.1 |
| STANDARD DEVIATION.....DO..... | .7 | 0 | .4 | .8 | .2 |
| MAXIMUM.....DO..... | 7.0 | 0 | 2.0 | 5.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 32 | 0 | 19 | 28 | 23 |
| STANDARD DEVIATION.....GM..... | 23 | 0 | 10 | 25 | 11 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 12 | | | | 12 |
| 25TH.....GM..... | 17 | | | | 17 |
| 50TH.....GM..... | 24 | | | | 17 |
| 75TH.....GM..... | 38 | | | | 28 |
| 90TH.....GM..... | 57 | | | | 38 |
| 95TH.....GM..... | 71 | | | | 50 |
| 99TH.....GM..... | 118 | | | | 57 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 76 | 0 | 57 | 65 | 68 |
| STANDARD DEVIATION.....GM..... | 46 | 0 | 30 | 39 | 33 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 73 | 0 | 35 | 57 | 48 |
| STANDARD DEVIATION.....GM..... | 32 | 0 | 14 | 15 | 7 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 72 | 0 | 0 | 133 | 0 |
| STANDARD DEVIATION.....GM..... | 39 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 510 | 0 | 113 | 200 | 170 |
| MAXIMUM OVER 3 DAYS.....GM..... | 780 | 0 | 113 | 400 | 170 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 72 | 0 | 50 | 62 | 65 |
| STANDARD DEVIATION.....GM..... | 39 | 0 | 29 | 32 | 31 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 35 | | | | 36 |
| 25TH.....GM..... | 50 | | | | 50 |
| 50TH.....GM..... | 56 | | | | 50 |
| 75TH.....GM..... | 85 | | | | 71 |
| 90TH.....GM..... | 113 | | | | 102 |
| 95TH.....GM..... | 142 | | | | 142 |
| 99TH.....GM..... | 227 | | | | 170 |
| MAXIMUM.....GM..... | 510 | 0 | 113 | 200 | 170 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 12.1 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.8 | 5.7 | 7.4 | 5.6 | 6.4 | 4.8 | 7.4 | 6.0 | 6.0 | 4.0 | 5.8 | 6.4 |
| 4.5 | 4.9 | 5.7 | 4.9 | 4.7 | 3.8 | 5.3 | 4.5 | 5.0 | 2.7 | 3.0 | 4.6 |
| 1.1 | .7 | 1.4 | .6 | 1.2 | .9 | 1.4 | 1.1 | .6 | .5 | 1.9 | 1.6 |
| .2 | .1 | .2 | .2 | .5 | .2 | .6 | .4 | .4 | .8 | .9 | .3 |
| 1.3 | 1.2 | 1.3 | 1.2 | 1.5 | 1.3 | 1.4 | 1.4 | 1.3 | 1.6 | 1.6 | 1.3 |
| .7 | .4 | .6 | .5 | .8 | .5 | .7 | .7 | .6 | .9 | .7 | .6 |
| 6.0 | 3.0 | 3.0 | 3.0 | 7.0 | 3.0 | 5.0 | 4.0 | 3.0 | 4.0 | 3.0 | 3.0 |
| 38 | 29 | 36 | 26 | 40 | 30 | 35 | 30 | 32 | 33 | 32 | 25 |
| 31 | 20 | 24 | 14 | 31 | 20 | 24 | 19 | 20 | 23 | 15 | 15 |
| 17 | 12 | 17 | 12 | 17 | 12 | 17 | 12 | 17 | 17 | | 12 |
| 17 | 17 | 17 | 17 | 19 | 17 | 17 | 17 | 17 | 17 | | 17 |
| 33 | 21 | 28 | 19 | 33 | 24 | 33 | 24 | 24 | 24 | | 17 |
| 47 | 33 | 50 | 33 | 47 | 33 | 47 | 33 | 38 | 38 | | 33 |
| 67 | 50 | 67 | 47 | 76 | 50 | 67 | 57 | 71 | 67 | | 47 |
| 113 | 57 | 76 | 52 | 113 | 67 | 76 | 68 | 76 | 83 | | 67 |
| 200 | 127 | 113 | 67 | 130 | 113 | 142 | 95 | 95 | 115 | | 67 |
| 91 | 74 | 90 | 68 | 91 | 69 | 81 | 68 | 85 | 72 | 64 | 51 |
| 61 | 50 | 60 | 29 | 59 | 33 | 43 | 40 | 53 | 46 | 25 | 12 |
| 83 | 77 | 89 | 63 | 83 | 74 | 72 | 68 | 75 | 59 | 54 | 63 |
| 38 | 34 | 34 | 13 | 41 | 27 | 33 | 23 | 35 | 18 | 13 | 23 |
| 117 | 55 | 54 | 63 | 84 | 85 | 74 | 68 | 55 | 57 | 57 | 51 |
| 76 | 28 | 34 | 13 | 54 | 33 | 37 | 22 | 13 | 28 | 7 | 5 |
| 425 | 510 | 340 | 200 | 425 | 213 | 250 | 400 | 227 | 340 | 100 | 100 |
| 695 | 510 | 484 | 263 | 780 | 400 | 542 | 400 | 284 | 344 | 200 | 200 |
| 85 | 72 | 83 | 64 | 82 | 71 | 74 | 65 | 75 | 62 | 58 | 56 |
| 51 | 47 | 46 | 23 | 52 | 32 | 37 | 30 | 45 | 36 | 19 | 20 |
| 35 | 35 | 35 | 35 | 50 | 35 | 35 | 28 | 50 | 18 | | 25 |
| 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | | 50 |
| 71 | 50 | 71 | 50 | 71 | 57 | 50 | 50 | 50 | 50 | | 50 |
| 100 | 85 | 100 | 71 | 100 | 100 | 100 | 85 | 100 | 71 | | 57 |
| 142 | 107 | 150 | 100 | 113 | 113 | 113 | 100 | 113 | 100 | | 100 |
| 170 | 150 | 170 | 100 | 150 | 140 | 150 | 113 | 211 | 100 | | 100 |
| 284 | 210 | 284 | 142 | 340 | 170 | 227 | 200 | 227 | 150 | | 100 |
| 400 | 510 | 284 | 150 | 340 | 213 | 227 | 284 | 227 | 340 | 100 | 100 |
| 0 | 6.0 | 2.2 | 0 | 2.1 | 1.8 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 4.2 | 1.6 | 0 | .6 | .6 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 5.27.--CRACKERS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 26.2 | 13.8 | 38.1 | 32.8 | 26.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 17.9 | 4.9 | 20.2 | 21.0 | 19.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 6.0 | 4.6 | 11.7 | 9.5 | 6.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.4 | 4.3 | 6.3 | 2.3 | 1.1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.5 | 2.6 | 1.9 | 1.6 | 1.4 |
| STANDARD DEVIATION.....DO..... | .9 | 1.9 | 1.3 | 1.0 | .7 |
| MAXIMUM.....DO..... | 9.0 | 9.0 | 9.0 | 9.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 11 | 9 | 9 | 10 | 10 |
| STANDARD DEVIATION.....GM..... | 13 | 10 | 10 | 10 | 11 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 2 | 2 | 1 | 2 | 2 |
| 25TH.....GM..... | 4 | 3 | 3 | 4 | 4 |
| 50TH.....GM..... | 8 | 6 | 6 | 6 | 6 |
| 75TH.....GM..... | 13 | 10 | 11 | 12 | 12 |
| 90TH.....GM..... | 23 | 19 | 18 | 21 | 20 |
| 95TH.....GM..... | 35 | 34 | 28 | 33 | 28 |
| 99TH.....GM..... | 65 | 56 | 56 | 42 | 56 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 23 | 10 | 16 | 23 | 20 |
| STANDARD DEVIATION.....GM..... | 23 | 5 | 18 | 28 | 20 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 24 | 11 | 15 | 19 | 23 |
| STANDARD DEVIATION.....GM..... | 21 | 5 | 14 | 14 | 16 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 26 | 17 | 19 | 21 | 29 |
| STANDARD DEVIATION.....GM..... | 25 | 14 | 13 | 11 | 28 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 364 | 57 | 170 | 239 | 255 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,036 | 168 | 227 | 239 | 454 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 22 | 10 | 14 | 18 | 20 |
| STANDARD DEVIATION.....GM..... | 21 | 9 | 14 | 20 | 19 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 6 | 3 | 3 | 6 | 6 |
| 25TH.....GM..... | 12 | 6 | 6 | 9 | 9 |
| 50TH.....GM..... | 15 | 7 | 12 | 12 | 14 |
| 75TH.....GM..... | 24 | 14 | 14 | 21 | 24 |
| 90TH.....GM..... | 42 | 15 | 28 | 30 | 42 |
| 95TH.....GM..... | 57 | 28 | 28 | 57 | 57 |
| 99TH.....GM..... | 113 | 56 | 85 | 113 | 113 |
| MAXIMUM.....GM..... | 340 | 57 | 113 | 227 | 170 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 123.1 | 3.3 | 4.8 | 2.2 | 8.4 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .8 | 1.8 | .6 | .2 | 1.2 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 22.1 | 22.1 | 18.0 | 19.6 | 18.5 | 23.1 | 26.5 | 29.9 | 37.4 | 34.9 | 31.6 | 36.0 |
| 16.7 | 17.3 | 14.1 | 15.0 | 13.7 | 16.5 | 18.3 | 20.0 | 24.3 | 21.1 | 18.3 | 20.2 |
| 4.7 | 3.8 | 2.7 | 3.5 | 3.9 | 5.4 | 5.7 | 7.2 | 7.9 | 8.7 | 5.0 | 8.6 |
| .7 | 1.0 | 1.3 | 1.0 | .9 | 1.3 | 2.4 | 2.7 | 5.2 | 5.1 | 8.3 | 7.3 |
| 1.4 | 1.3 | 1.4 | 1.4 | 1.4 | 1.4 | 1.6 | 1.6 | 1.7 | 1.6 | 2.0 | 1.8 |
| .7 | .6 | .8 | .7 | .8 | .8 | 1.0 | .9 | 1.2 | .9 | 1.4 | 1.2 |
| 6.0 | 4.0 | 6.0 | 5.0 | 6.0 | 9.0 | 9.0 | 6.0 | 9.0 | 6.0 | 7.0 | 8.0 |
| 11 | 9 | 14 | 11 | 15 | 10 | 13 | 11 | 12 | 9 | 15 | 11 |
| 11 | 8 | 15 | 11 | 19 | 11 | 14 | 14 | 13 | 9 | 16 | 13 |
| 2 | 2 | 2 | 2 | 3 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 4 | 4 | 5 | 4 | 5 | 4 | 5 | 4 | 5 | 4 | 5 | 4 |
| 8 | 6 | 9 | 7 | 8 | 7 | 8 | 7 | 8 | 6 | 8 | 6 |
| 14 | 10 | 19 | 12 | 19 | 12 | 16 | 12 | 12 | 11 | 18 | 12 |
| 22 | 19 | 29 | 22 | 37 | 19 | 28 | 20 | 28 | 18 | 37 | 22 |
| 33 | 26 | 50 | 30 | 47 | 30 | 41 | 28 | 36 | 23 | 48 | 30 |
| 57 | 38 | 76 | 56 | 85 | 57 | 76 | 59 | 70 | 57 | 67 | 66 |
| 26 | 21 | 32 | 23 | 33 | 22 | 27 | 20 | 22 | 17 | 21 | 16 |
| 27 | 18 | 33 | 21 | 34 | 20 | 25 | 19 | 18 | 14 | 20 | 16 |
| 28 | 21 | 39 | 28 | 34 | 24 | 30 | 24 | 21 | 17 | 33 | 19 |
| 20 | 15 | 28 | 26 | 29 | 19 | 25 | 21 | 11 | 14 | 28 | 16 |
| 24 | 20 | 32 | 23 | 51 | 25 | 32 | 25 | 31 | 19 | 27 | 24 |
| 14 | 12 | 24 | 14 | 43 | 25 | 21 | 32 | 22 | 14 | 18 | 20 |
| 340 | 175 | 227 | 227 | 340 | 224 | 245 | 364 | 143 | 113 | 113 | 132 |
| 340 | 204 | 336 | 340 | 579 | 421 | 361 | 1,036 | 255 | 232 | 227 | 340 |
| 24 | 20 | 32 | 23 | 33 | 21 | 26 | 20 | 21 | 17 | 22 | 18 |
| 24 | 16 | 29 | 21 | 35 | 20 | 23 | 20 | 15 | 14 | 19 | 16 |
| 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 |
| 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 9 |
| 18 | 14 | 24 | 18 | 24 | 15 | 18 | 14 | 18 | 12 | 18 | 12 |
| 30 | 24 | 41 | 28 | 40 | 28 | 30 | 24 | 28 | 18 | 24 | 18 |
| 45 | 39 | 57 | 42 | 57 | 42 | 56 | 32 | 36 | 28 | 44 | 30 |
| 57 | 56 | 85 | 57 | 113 | 56 | 60 | 56 | 48 | 44 | 57 | 56 |
| 120 | 85 | 170 | 113 | 170 | 110 | 113 | 113 | 84 | 85 | 113 | 85 |
| 340 | 113 | 227 | 170 | 340 | 224 | 245 | 196 | 135 | 113 | 113 | 113 |
| 8.1 | 3.5 | 8.5 | 4.7 | 10.7 | 22.0 | 15.4 | 12.4 | 5.0 | 8.0 | 1.4 | 4.8 |
| 1.3 | .6 | 2.5 | 1.2 | 1.0 | 1.2 | .8 | .4 | .7 | .8 | .4 | .7 |

TABLE 5.28.--GRAHAM CRACKERS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.1 | 5.2 | 8.2 | 6.2 | 3.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.4 | 2.9 | 4.7 | 4.4 | 2.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | 1.1 | 2.0 | 1.0 | .7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 1.2 | 1.5 | .8 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 1.6 | 2.5 | 1.8 | 1.5 | 1.4 |
| STANDARD DEVIATION.....DO..... | 1.0 | 2.3 | .9 | .8 | .7 |
| MAXIMUM.....DO..... | 9.0 | 9.0 | 5.0 | 4.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 16 | 12 | 11 | 14 | 14 |
| STANDARD DEVIATION.....GM..... | 15 | 13 | 9 | 13 | 11 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 5 | | 4 | 5 | 5 |
| 25TH.....GM..... | 5 | | 5 | 5 | 5 |
| 50TH.....GM..... | 9 | | 9 | 9 | 9 |
| 75TH.....GM..... | 19 | | 14 | 19 | 19 |
| 90TH.....GM..... | 35 | | 19 | 33 | 28 |
| 95TH.....GM..... | 47 | | 24 | 38 | 38 |
| 99TH.....GM..... | 75 | | 56 | 76 | 56 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 34 | 16 | 23 | 34 | 33 |
| STANDARD DEVIATION.....GM..... | 30 | 8 | 14 | 38 | 26 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 32 | 11 | 20 | 25 | 34 |
| STANDARD DEVIATION.....GM..... | 22 | 5 | 14 | 13 | 20 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 32 | 33 | 19 | 23 | 24 |
| STANDARD DEVIATION.....GM..... | 21 | 12 | 12 | 11 | 5 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 227 | 57 | 84 | 227 | 113 |
| MAXIMUM OVER 3 DAYS.....GM..... | 336 | 168 | 168 | 227 | 168 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 30 | 14 | 19 | 27 | 31 |
| STANDARD DEVIATION.....GM..... | 24 | 12 | 12 | 27 | 23 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 7 | | 7 | 7 | 14 |
| 25TH.....GM..... | 14 | | 14 | 14 | 14 |
| 50TH.....GM..... | 28 | | 14 | 21 | 28 |
| 75TH.....GM..... | 42 | | 28 | 28 | 42 |
| 90TH.....GM..... | 56 | | 28 | 42 | 56 |
| 95TH.....GM..... | 84 | | 28 | 64 | 84 |
| 99TH.....GM..... | 140 | | 71 | 151 | 113 |
| MAXIMUM.....GM..... | 227 | 57 | 84 | 227 | 113 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 4.1 | 1.0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 1.5 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 2.2 | 2.3 | 1.1 | 1.4 | .6 | 1.4 | 1.3 | 1.7 | 2.8 | 2.5 | 2.1 | 2.4 |
| 1.6 | 1.9 | 1.0 | 1.2 | .4 | 1.0 | .9 | 1.1 | 1.4 | 1.5 | 1.6 | 1.6 |
| .4 | .3 | .1 | .2 | .1 | .2 | .2 | .4 | .7 | .5 | .5 | .6 |
| .1 | .1 | (2) | .1 | (2) | .2 | .2 | .2 | .6 | .6 | 0 | .2 |
| 1.5 | 1.4 | 1.3 | 1.3 | 1.5 | 1.7 | 1.7 | 1.6 | 1.8 | 1.8 | 1.3 | 1.4 |
| 1.0 | .7 | 1.0 | .6 | .7 | 1.1 | 1.3 | 1.0 | .8 | 1.0 | .4 | .6 |
| 6.0 | 4.0 | 6.0 | 3.0 | 3.0 | 6.0 | 8.0 | 6.0 | 3.0 | 5.0 | 2.0 | 3.0 |
| 12 | 13 | 22 | 16 | 27 | 19 | 22 | 16 | 23 | 16 | 19 | 14 |
| 10 | 9 | 26 | 13 | 20 | 20 | 16 | 14 | 19 | 14 | 11 | 13 |
| | 2 | | | | 5 | 5 | 5 | | | | |
| | 5 | | | | 7 | 9 | 9 | | | | |
| | 9 | | | | 12 | 19 | 14 | | | | |
| | 19 | | | | 19 | 28 | 19 | | | | |
| | 28 | | | | 47 | 42 | 28 | | | | |
| | 28 | | | | 56 | 47 | 47 | | | | |
| | 38 | | | | 112 | 75 | 79 | | | | |
| 30 | 34 | 53 | 45 | 57 | 35 | 43 | 35 | 27 | 32 | 42 | 25 |
| 21 | 25 | 55 | 42 | 48 | 29 | 30 | 25 | 21 | 22 | 27 | 9 |
| 22 | 32 | 56 | 26 | 63 | 45 | 42 | 35 | 44 | 23 | 49 | 28 |
| 15 | 9 | 0 | 10 | 31 | 27 | 24 | 24 | 16 | 10 | 0 | 13 |
| 32 | 20 | 112 | 23 | 47 | 37 | 44 | 31 | 50 | 29 | 0 | 56 |
| 17 | 7 | 0 | 5 | 0 | 34 | 15 | 18 | 14 | 19 | 0 | 0 |
| 140 | 113 | 227 | 170 | 154 | 154 | 140 | 154 | 112 | 84 | 85 | 56 |
| 168 | 113 | 336 | 170 | 252 | 336 | 224 | 238 | 196 | 196 | 98 | 168 |
| 25 | 28 | 54 | 38 | 55 | 35 | 37 | 31 | 40 | 27 | 45 | 31 |
| 18 | 19 | 47 | 35 | 39 | 23 | 23 | 24 | 23 | 19 | 21 | 15 |
| | 14 | | | | 14 | 14 | 14 | | | | |
| | 14 | | | | 14 | 28 | 14 | | | | |
| | 28 | | | | 28 | 28 | 28 | | | | |
| | 28 | | | | 56 | 42 | 42 | | | | |
| | 56 | | | | 70 | 57 | 56 | | | | |
| | 56 | | | | 84 | 84 | 84 | | | | |
| | 113 | | | | 85 | 140 | 140 | | | | |
| 140 | 113 | 227 | 170 | 154 | 85 | 140 | 154 | 112 | 84 | 85 | 56 |
| 0 | 0 | 1.0 | .6 | 0 | 0 | .8 | 0 | 0 | 0 | 0 | .8 |
| 0 | 0 | 4.9 | 2.3 | 0 | 0 | .7 | 0 | 0 | 0 | 0 | 2.4 |

TABLE 5.29.--SALTINES AND SODA CRACKERS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 14.5 | 6.3 | 20.1 | 17.9 | 14.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 10.9 | 3.1 | 12.6 | 13.8 | 11.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.8 | 1.9 | 5.1 | 3.6 | 2.3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .8 | 1.3 | 2.5 | .4 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.4 | 2.0 | 1.6 | 1.4 | 1.2 |
| STANDARD DEVIATION.....DO..... | .7 | 1.2 | 1.0 | .6 | .5 |
| MAXIMUM.....DO..... | 9.0 | 5.0 | 6.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 8 | 5 | 6 | 6 | 7 |
| STANDARD DEVIATION.....GM..... | 9 | 4 | 8 | 6 | 7 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 2 | | 1 | 2 | 2 |
| 25TH.....GM..... | 4 | | 2 | 3 | 3 |
| 50TH.....GM..... | 6 | | 4 | 4 | 4 |
| 75TH.....GM..... | 9 | | 6 | 7 | 8 |
| 90TH.....GM..... | 16 | | 11 | 12 | 14 |
| 95TH.....GM..... | 22 | | 17 | 14 | 20 |
| 99TH.....GM..... | 48 | | 40 | 38 | 40 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 18 | 7 | 10 | 15 | 16 |
| STANDARD DEVIATION.....GM..... | 17 | 4 | 12 | 15 | 16 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 20 | 11 | 13 | 13 | 20 |
| STANDARD DEVIATION.....GM..... | 19 | 5 | 16 | 9 | 15 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 20 | 9 | 13 | 10 | 20 |
| STANDARD DEVIATION.....GM..... | 16 | 5 | 12 | 3 | 15 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 227 | 30 | 170 | 113 | 170 |
| MAXIMUM OVER 3 DAYS.....GM..... | 355 | 60 | 227 | 113 | 182 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 18 | 8 | 11 | 13 | 17 |
| STANDARD DEVIATION.....GM..... | 17 | 5 | 14 | 12 | 17 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 6 | | 3 | 6 | 6 |
| 25TH.....GM..... | 9 | | 6 | 6 | 6 |
| 50TH.....GM..... | 12 | | 6 | 12 | 12 |
| 75TH.....GM..... | 18 | | 12 | 12 | 18 |
| 90TH.....GM..... | 30 | | 18 | 21 | 30 |
| 95TH.....GM..... | 42 | | 24 | 28 | 36 |
| 99TH.....GM..... | 113 | | 113 | 57 | 60 |
| MAXIMUM.....GM..... | 227 | 30 | 113 | 113 | 170 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 31.7 | 1.0 | 2.6 | 1.3 | .9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 1.5 | .8 | .3 | .3 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 12.7 | 12.4 | 10.2 | 10.1 | 9.2 | 11.8 | 14.7 | 17.2 | 23.9 | 21.0 | 16.9 | 21.2 |
| 10.1 | 10.8 | 8.6 | 8.0 | 7.1 | 9.2 | 10.5 | 12.9 | 17.8 | 14.2 | 9.9 | 12.9 |
| 2.4 | 1.3 | 1.1 | 1.8 | 1.9 | 2.0 | 3.1 | 3.5 | 3.9 | 5.1 | 3.2 | 5.6 |
| .2 | .2 | .4 | .4 | .2 | .6 | 1.1 | .8 | 2.3 | 1.8 | 3.8 | 2.7 |
| 1.3 | 1.2 | 1.2 | 1.3 | 1.3 | 1.3 | 1.5 | 1.4 | 1.5 | 1.5 | 1.9 | 1.6 |
| .6 | .5 | .6 | .7 | .6 | .6 | .9 | .7 | .9 | .7 | 1.4 | .9 |
| 5.0 | 3.0 | 4.0 | 5.0 | 5.0 | 6.0 | 9.0 | 5.0 | 6.0 | 4.0 | 7.0 | 6.0 |
| 9 | 6 | 12 | 9 | 11 | 7 | 11 | 8 | 9 | 8 | 12 | 8 |
| 8 | 5 | 13 | 10 | 13 | 6 | 12 | 9 | 8 | 8 | 13 | 8 |
| 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 4 | 2 |
| 4 | 3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 5 | 4 |
| 6 | 4 | 8 | 6 | 6 | 6 | 8 | 6 | 7 | 6 | 6 | 6 |
| 10 | 8 | 12 | 10 | 10 | 8 | 12 | 8 | 10 | 9 | 12 | 10 |
| 18 | 12 | 20 | 19 | 21 | 12 | 22 | 14 | 16 | 15 | 31 | 16 |
| 26 | 16 | 40 | 24 | 38 | 16 | 32 | 19 | 28 | 18 | 47 | 19 |
| 41 | 26 | 76 | 76 | 76 | 28 | 57 | 40 | 48 | 47 | 76 | 30 |
| 20 | 16 | 26 | 22 | 25 | 16 | 23 | 16 | 20 | 15 | 18 | 14 |
| 17 | 13 | 23 | 25 | 26 | 11 | 22 | 13 | 17 | 11 | 14 | 7 |
| 25 | 18 | 42 | 22 | 23 | 16 | 26 | 20 | 16 | 19 | 25 | 18 |
| 18 | 10 | 38 | 19 | 18 | 9 | 24 | 20 | 6 | 17 | 26 | 18 |
| 23 | 12 | 27 | 19 | 51 | 14 | 24 | 21 | 24 | 15 | 23 | 14 |
| 15 | 3 | 19 | 13 | 44 | 13 | 16 | 16 | 13 | 6 | 16 | 4 |
| 132 | 131 | 227 | 227 | 227 | 113 | 227 | 227 | 143 | 113 | 113 | 125 |
| 227 | 131 | 247 | 227 | 355 | 251 | 340 | 312 | 174 | 170 | 227 | 239 |
| 21 | 15 | 28 | 20 | 24 | 15 | 22 | 17 | 18 | 16 | 18 | 14 |
| 18 | 12 | 26 | 19 | 25 | 11 | 20 | 15 | 12 | 12 | 16 | 11 |
| 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 3 |
| 12 | 9 | 12 | 12 | 12 | 9 | 12 | 12 | 12 | 12 | 12 | 9 |
| 15 | 12 | 20 | 15 | 18 | 12 | 18 | 12 | 15 | 12 | 12 | 12 |
| 27 | 18 | 30 | 24 | 24 | 18 | 24 | 18 | 24 | 18 | 24 | 18 |
| 36 | 24 | 57 | 36 | 48 | 24 | 36 | 28 | 30 | 24 | 28 | 21 |
| 48 | 30 | 113 | 48 | 85 | 30 | 57 | 30 | 36 | 30 | 28 | 28 |
| 113 | 60 | 113 | 113 | 113 | 57 | 113 | 113 | 60 | 57 | 113 | 57 |
| 120 | 113 | 140 | 140 | 227 | 113 | 227 | 170 | 135 | 113 | 113 | 113 |
| 3.8 | 1.6 | .9 | 1.3 | 1.6 | 4.6 | 7.0 | 2.0 | 2.1 | 0 | 0 | 1.0 |
| 1.1 | .5 | .5 | .7 | .3 | .6 | .7 | .1 | .5 | 0 | 0 | .3 |

TABLE 5.30.--BUTTER CRACKERS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.4 | 2.6 | 7.6 | 5.5 | 5.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.4 | 1.4 | 4.8 | 4.5 | 4.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .8 | 1.0 | 2.2 | .9 | 1.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | .2 | .6 | .1 | (2) |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.4 | 1.5 | 1.6 | 1.3 | 1.3 |
| STANDARD DEVIATION.....DO..... | .7 | .6 | .8 | .7 | .6 |
| MAXIMUM.....DO..... | 7.0 | 3.0 | 6.0 | 6.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 9 | 4 | 6 | 6 | 9 |
| STANDARD DEVIATION.....GM..... | 10 | 2 | 4 | 5 | 14 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 2 | | 1 | 2 | 2 |
| 25TH.....GM..... | 4 | | 3 | 3 | 4 |
| 50TH.....GM..... | 6 | | 6 | 5 | 6 |
| 75TH.....GM..... | 10 | | 8 | 8 | 9 |
| 90TH.....GM..... | 18 | | 12 | 12 | 13 |
| 95TH.....GM..... | 24 | | 15 | 16 | 18 |
| 99TH.....GM..... | 50 | | 23 | 28 | 30 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 21 | 7 | 14 | 17 | 19 |
| STANDARD DEVIATION.....GM..... | 19 | 3 | 11 | 13 | 11 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 21 | 8 | 12 | 12 | 19 |
| STANDARD DEVIATION.....GM..... | 14 | 2 | 4 | 4 | 11 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 24 | 6 | 13 | 24 | 151 |
| STANDARD DEVIATION.....GM..... | 22 | 0 | 5 | (2) | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 340 | 15 | 60 | 85 | 255 |
| MAXIMUM OVER 3 DAYS.....GM..... | 454 | 21 | 69 | 85 | 454 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 20 | 7 | 12 | 14 | 21 |
| STANDARD DEVIATION.....GM..... | 18 | 3 | 7 | 11 | 19 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 6 | | 3 | 4 | 4 |
| 25TH.....GM..... | 12 | | 6 | 8 | 12 |
| 50TH.....GM..... | 18 | | 12 | 12 | 18 |
| 75TH.....GM..... | 24 | | 15 | 16 | 24 |
| 90TH.....GM..... | 36 | | 18 | 27 | 36 |
| 95TH.....GM..... | 50 | | 28 | 30 | 57 |
| 99TH.....GM..... | 113 | | 30 | 57 | 113 |
| MAXIMUM.....GM..... | 340 | 15 | 30 | 85 | 142 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 7.2 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.4 | 3.4 | 3.0 | 3.7 | 3.6 | 4.2 | 4.9 | 4.5 | 4.0 | 4.8 | 4.6 | 7.3 |
| 2.9 | 2.8 | 2.4 | 2.8 | 2.9 | 3.6 | 3.7 | 3.4 | 2.4 | 3.6 | 2.9 | 3.9 |
| .5 | .5 | .4 | .7 | .5 | .4 | .9 | .9 | 1.4 | .3 | .2 | 1.7 |
| (2) | .1 | .2 | .3 | .1 | .1 | .3 | .2 | .2 | .9 | 1.5 | 1.7 |
| 1.2 | 1.2 | 1.3 | 1.4 | 1.3 | 1.2 | 1.3 | 1.3 | 1.6 | 1.5 | 1.8 | 2.0 |
| .5 | .5 | .7 | .7 | .6 | .4 | .6 | .6 | 1.0 | .9 | 1.1 | 1.4 |
| 3.0 | 3.0 | 4.0 | 4.0 | 4.0 | 3.0 | 5.0 | 4.0 | 7.0 | 4.0 | 4.0 | 6.0 |
| 11 | 7 | 13 | 10 | 14 | 8 | 10 | 8 | 11 | 7 | 14 | 12 |
| 12 | 4 | 10 | 7 | 14 | 6 | 11 | 8 | 10 | 4 | 10 | 17 |
| 3 | 2 | | 3 | 3 | 2 | 2 | 2 | | 2 | | 1 |
| 6 | 4 | | 4 | 6 | 4 | 4 | 4 | | 4 | | 4 |
| 8 | 6 | | 8 | 8 | 6 | 6 | 6 | | 5 | | 8 |
| 14 | 8 | | 14 | 16 | 10 | 12 | 10 | | 10 | | 12 |
| 20 | 12 | | 21 | 38 | 16 | 20 | 15 | | 12 | | 21 |
| 20 | 15 | | 27 | 47 | 19 | 24 | 20 | | 16 | | 50 |
| 113 | 25 | | 30 | 76 | 33 | 54 | 47 | | 22 | | 113 |
| 29 | 19 | 29 | 21 | 32 | 21 | 22 | 18 | 25 | 16 | 33 | 14 |
| 38 | 12 | 18 | 15 | 31 | 13 | 16 | 18 | 18 | 9 | 32 | 9 |
| 25 | 16 | 37 | 24 | 35 | 23 | 25 | 20 | 17 | 16 | 54 | 17 |
| 12 | 7 | 25 | 11 | 28 | 10 | 15 | 11 | 8 | 2 | 0 | 9 |
| 20 | 12 | 23 | 22 | 47 | 27 | 33 | 20 | 34 | 13 | 17 | 28 |
| 0 | 0 | 8 | 7 | 9 | 14 | 26 | 14 | 28 | 4 | 5 | 29 |
| 340 | 75 | 100 | 71 | 170 | 60 | 140 | 142 | 99 | 57 | 113 | 113 |
| 340 | 75 | 180 | 90 | 227 | 135 | 294 | 165 | 210 | 66 | 113 | 340 |
| 26 | 18 | 29 | 21 | 32 | 21 | 23 | 18 | 21 | 14 | 23 | 18 |
| 32 | 11 | 19 | 13 | 29 | 13 | 17 | 14 | 13 | 7 | 21 | 19 |
| 9 | 6 | | 6 | 5 | 6 | 6 | 6 | | 6 | | 3 |
| 15 | 12 | | 12 | 14 | 12 | 12 | 12 | | 12 | | 9 |
| 21 | 15 | | 18 | 24 | 18 | 18 | 15 | | 12 | | 12 |
| 30 | 18 | | 30 | 36 | 28 | 28 | 24 | | 18 | | 18 |
| 45 | 30 | | 36 | 57 | 42 | 45 | 30 | | 18 | | 30 |
| 50 | 45 | | 48 | 113 | 50 | 56 | 45 | | 24 | | 40 |
| 85 | 57 | | 71 | 142 | 60 | 85 | 60 | | 57 | | 113 |
| 340 | 75 | 100 | 71 | 170 | 60 | 113 | 142 | 84 | 57 | 113 | 113 |
| 0 | 0 | 0 | 0 | 0 | 2.9 | 1.3 | .8 | 0 | 0 | 0 | 2.2 |
| 0 | 0 | 0 | 0 | 0 | 1.1 | .4 | .2 | 0 | 0 | 0 | 1.5 |

TABLE 5.31.--POPCORN
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.6 | .1 | 5.7 | 8.5 | 9.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.9 | .1 | 4.9 | 7.6 | 7.9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .6 | 0 | .8 | .8 | 1.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | 0 | .1 | .4 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.2 | 1.0 | 1.2 | 1.1 | 1.2 |
| STANDARD DEVIATION.....DO..... | .5 | .0 | .4 | .4 | .5 |
| MAXIMUM.....DO..... | 4.0 | 1.0 | 2.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 7 | 24 | 3 | 5 | 6 |
| STANDARD DEVIATION.....GM..... | 9 | 0 | 4 | 4 | 4 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 2 | | 1 | 1 | 2 |
| 25TH.....GM..... | 3 | | 2 | 2 | 3 |
| 50TH.....GM..... | 6 | | 3 | 3 | 6 |
| 75TH.....GM..... | 9 | | 3 | 6 | 6 |
| 90TH.....GM..... | 12 | | 6 | 9 | 9 |
| 95TH.....GM..... | 18 | | 8 | 12 | 12 |
| 99TH.....GM..... | 43 | | 38 | 13 | 28 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 19 | 72 | 10 | 12 | 16 |
| STANDARD DEVIATION.....GM..... | 23 | 0 | 14 | 8 | 10 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 19 | 0 | 7 | 15 | 10 |
| STANDARD DEVIATION.....GM..... | 15 | 0 | 3 | 15 | 4 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 20 | 0 | 0 | 10 | 16 |
| STANDARD DEVIATION.....GM..... | 20 | 0 | 0 | 0 | 9 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 340 | 72 | 113 | 130 | 98 |
| MAXIMUM OVER 3 DAYS.....GM..... | 340 | 72 | 113 | 139 | 98 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 19 | 72 | 9 | 12 | 14 |
| STANDARD DEVIATION.....GM..... | 22 | 0 | 12 | 11 | 9 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 5 | | 2 | 3 | 5 |
| 25TH.....GM..... | 9 | | 5 | 6 | 9 |
| 50TH.....GM..... | 15 | | 6 | 9 | 12 |
| 75TH.....GM..... | 18 | | 9 | 18 | 18 |
| 90TH.....GM..... | 36 | | 14 | 18 | 27 |
| 95TH.....GM..... | 45 | | 24 | 27 | 28 |
| 99TH.....GM..... | 108 | | 113 | 36 | 36 |
| MAXIMUM.....GM..... | 340 | 72 | 113 | 130 | 98 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 66.2 | 0 | 0 | 6.2 | 12.3 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.7 | 0 | 0 | 3.7 | 5.8 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 9.6 | 9.1 | 6.1 | 7.8 | 6.5 | 7.1 | 4.1 | 3.8 | 1.5 | .9 | .3 | .8 |
| 8.7 | 7.7 | 4.9 | 6.6 | 5.8 | 6.2 | 3.5 | 3.4 | 1.4 | .9 | .2 | .6 |
| .8 | 1.1 | 1.1 | .9 | .5 | .7 | .5 | .3 | .1 | 0 | .1 | .2 |
| .1 | .3 | .1 | .3 | .1 | .2 | .1 | (2) | 0 | 0 | 0 | 0 |
| 1.1 | 1.2 | 1.3 | 1.2 | 1.1 | 1.2 | 1.2 | 1.1 | 1.5 | 1.0 | 1.5 | 1.2 |
| .4 | .6 | .5 | .5 | .4 | .5 | .5 | .4 | .9 | .0 | .5 | .4 |
| 3.0 | 4.0 | 3.0 | 3.0 | 4.0 | 3.0 | 4.0 | 3.0 | 4.0 | 1.0 | 2.0 | 2.0 |
| 7 | 7 | 8 | 7 | 9 | 7 | 11 | 6 | 9 | 4 | 12 | 5 |
| 6 | 6 | 10 | 9 | 12 | 9 | 15 | 6 | 9 | 1 | 6 | 1 |
| 2 | 2 | 2 | 2 | 1 | 1 | 1 | 2 | | | | |
| 3 | 3 | 4 | 3 | 3 | 3 | 4 | 3 | | | | |
| 6 | 6 | 6 | 6 | 6 | 6 | 8 | 6 | | | | |
| 9 | 9 | 9 | 6 | 12 | 9 | 12 | 7 | | | | |
| 12 | 12 | 15 | 12 | 19 | 12 | 18 | 12 | | | | |
| 15 | 18 | 19 | 24 | 24 | 15 | 32 | 18 | | | | |
| 28 | 28 | 84 | 72 | 90 | 38 | 95 | 43 | | | | |
| 19 | 18 | 20 | 16 | 25 | 20 | 28 | 16 | 18 | 11 | 18 | 14 |
| 18 | 14 | 15 | 12 | 34 | 26 | 41 | 11 | 6 | 4 | 0 | 3 |
| 15 | 17 | 23 | 21 | 26 | 19 | 22 | 23 | 54 | 0 | 27 | 9 |
| 7 | 15 | 29 | 17 | 11 | 12 | 13 | 15 | 0 | 0 | 0 | 0 |
| 6 | 14 | 9 | 32 | 22 | 10 | 56 | 38 | 0 | 0 | 0 | 0 |
| (2) | 5 | 0 | 27 | 30 | 2 | 12 | 23 | 0 | 0 | 0 | 0 |
| 170 | 112 | 144 | 180 | 320 | 340 | 284 | 112 | 54 | 18 | 36 | 18 |
| 170 | 130 | 252 | 216 | 320 | 340 | 284 | 168 | 108 | 18 | 54 | 18 |
| 18 | 17 | 20 | 18 | 24 | 19 | 27 | 18 | 18 | 11 | 24 | 12 |
| 17 | 15 | 20 | 20 | 29 | 24 | 36 | 13 | 11 | 4 | 8 | 3 |
| 5 | 5 | 5 | 5 | 3 | 5 | 5 | 5 | | | | |
| 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | | | | |
| 18 | 14 | 18 | 18 | 18 | 18 | 18 | 18 | | | | |
| 18 | 18 | 18 | 18 | 27 | 24 | 28 | 18 | | | | |
| 36 | 27 | 36 | 27 | 48 | 36 | 45 | 36 | | | | |
| 36 | 36 | 48 | 45 | 68 | 36 | 72 | 36 | | | | |
| 85 | 85 | 108 | 112 | 113 | 85 | 284 | 56 | | | | |
| 170 | 112 | 144 | 180 | 284 | 340 | 284 | 112 | 54 | 18 | 36 | 18 |
| 12.3 | 9.0 | 5.8 | 8.6 | 3.4 | 4.5 | 0 | 4.1 | 0 | 0 | 0 | 0 |
| 5.5 | 3.7 | 5.4 | 6.2 | 1.2 | 1.0 | 0 | 1.4 | 0 | 0 | 0 | 0 |

TABLE 5.32.--PRETZELS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.2 | .7 | 3.2 | 3.1 | 3.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.7 | .4 | 2.3 | 2.5 | 2.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | .3 | .9 | .6 | .3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | 0 | (2) | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 1.9 | 1.5 | 1.3 | 1.2 |
| STANDARD DEVIATION.....DO..... | .6 | 1.1 | 1.0 | .5 | .5 |
| MAXIMUM.....DO..... | 5.0 | 4.0 | 5.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 13 | 3 | 9 | 9 | 10 |
| STANDARD DEVIATION.....GM..... | 15 | 1 | 9 | 9 | 10 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 1 | | | (2) | 1 |
| 25TH.....GM..... | 5 | | | 2 | 4 |
| 50TH.....GM..... | 9 | | | 6 | 8 |
| 75TH.....GM..... | 16 | | | 11 | 11 |
| 90TH.....GM..... | 28 | | | 19 | 19 |
| 95TH.....GM..... | 42 | | | 30 | 30 |
| 99TH.....GM..... | 64 | | | 41 | 75 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 29 | 10 | 20 | 22 | 25 |
| STANDARD DEVIATION.....GM..... | 28 | 5 | 24 | 22 | 28 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 29 | 3 | 24 | 20 | 26 |
| STANDARD DEVIATION.....GM..... | 23 | 1 | 14 | 16 | 17 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 38 | 0 | 0 | 41 | 24 |
| STANDARD DEVIATION.....GM..... | 37 | 0 | 0 | 0 | 9 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 224 | 15 | 113 | 90 | 224 |
| MAXIMUM OVER 3 DAYS.....GM..... | 480 | 15 | 113 | 123 | 224 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 29 | 4 | 18 | 21 | 25 |
| STANDARD DEVIATION.....GM..... | 28 | 4 | 18 | 20 | 21 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 3 | | | 1 | 4 |
| 25TH.....GM..... | 12 | | | 5 | 12 |
| 50TH.....GM..... | 21 | | | 16 | 23 |
| 75TH.....GM..... | 36 | | | 32 | 28 |
| 90TH.....GM..... | 57 | | | 56 | 57 |
| 95TH.....GM..... | 85 | | | 57 | 68 |
| 99TH.....GM..... | 160 | | | 90 | 112 |
| MAXIMUM.....GM..... | 170 | 15 | 113 | 90 | 112 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 21.6 | .8 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.9 | 12.7 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.
2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 4.1 | 3.5 | 2.9 | 3.1 | 2.4 | 2.3 | 1.9 | 1.2 | 1.2 | .5 | 2.1 | .6 |
| 2.9 | 2.5 | 2.1 | 2.4 | 1.8 | 1.8 | 1.5 | 1.0 | .4 | .3 | 1.5 | .6 |
| .8 | .7 | .4 | .5 | .6 | .4 | .3 | .2 | .5 | .2 | .6 | 0 |
| .4 | .2 | .3 | .2 | .1 | .1 | 0 | .1 | .3 | 0 | 0 | 0 |
| 1.5 | 1.4 | 1.4 | 1.3 | 1.4 | 1.3 | 1.2 | 1.3 | 1.9 | 1.4 | 1.5 | 1.0 |
| .8 | .6 | .7 | .6 | .6 | .6 | .4 | .7 | .7 | .5 | .5 | .0 |
| 4.0 | 3.0 | 3.0 | 3.0 | 3.0 | 4.0 | 2.0 | 4.0 | 3.0 | 2.0 | 2.0 | 1.0 |
| 15 | 14 | 25 | 11 | 19 | 9 | 13 | 10 | 27 | 12 | 5 | 12 |
| 12 | 12 | 38 | 8 | 15 | 9 | 13 | 12 | 22 | 4 | 3 | 4 |
| 2 | 2 | | | 1 | 1 | 1 | 2 | | | | |
| 5 | 5 | | | 8 | 4 | 5 | 3 | | | | |
| 11 | 9 | | | 15 | 6 | 9 | 5 | | | | |
| 19 | 16 | | | 23 | 13 | 17 | 10 | | | | |
| 33 | 30 | | | 47 | 19 | 21 | 19 | | | | |
| 38 | 43 | | | 51 | 28 | 43 | 47 | | | | |
| 47 | 57 | | | 67 | 50 | 76 | 54 | | | | |
| 34 | 32 | 38 | 26 | 51 | 20 | 27 | 24 | 70 | 30 | 17 | 36 |
| 29 | 26 | 24 | 16 | 42 | 17 | 18 | 26 | 66 | 14 | 11 | 13 |
| 33 | 27 | 34 | 20 | 35 | 29 | 43 | 23 | 19 | 22 | 6 | 0 |
| 17 | 21 | 21 | 10 | 27 | 20 | 30 | 29 | 8 | 1 | (2) | 0 |
| 23 | 34 | 102 | 32 | 35 | 17 | 0 | 28 | 56 | 0 | 0 | 0 |
| 18 | 12 | 68 | 3 | 8 | 2 | 0 | 18 | 10 | 0 | 0 | 0 |
| 142 | 135 | 160 | 57 | 172 | 113 | 170 | 142 | 142 | 48 | 30 | 57 |
| 225 | 170 | 480 | 105 | 202 | 149 | 227 | 163 | 192 | 48 | 30 | 57 |
| 29 | 30 | 52 | 25 | 42 | 21 | 32 | 23 | 41 | 26 | 10 | 36 |
| 25 | 26 | 50 | 16 | 38 | 19 | 26 | 28 | 36 | 11 | 6 | 13 |
| 6 | 2 | | | 3 | 3 | 6 | 3 | | | | |
| 14 | 12 | | | 17 | 9 | 15 | 9 | | | | |
| 23 | 23 | | | 30 | 15 | 24 | 15 | | | | |
| 42 | 43 | | | 50 | 28 | 45 | 28 | | | | |
| 57 | 64 | | | 113 | 45 | 57 | 48 | | | | |
| 85 | 90 | | | 142 | 57 | 72 | 57 | | | | |
| 113 | 113 | | | 170 | 113 | 170 | 142 | | | | |
| 142 | 135 | 160 | 57 | 170 | 113 | 170 | 142 | 142 | 48 | 23 | 57 |
| 3.4 | 1.9 | .9 | 1.1 | 4.2 | 0 | 2.3 | 7.0 | 0 | 0 | 0 | 0 |
| 2.7 | 1.9 | 1.5 | 1.9 | 3.3 | 0 | 2.1 | 6.2 | 0 | 0 | 0 | 0 |

TABLE 5.33.--POTATO CHIPS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 17.0 | .8 | 19.3 | 21.7 | 26.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 12.1 | .6 | 13.7 | 15.7 | 17.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.7 | .2 | 4.7 | 4.9 | 7.1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.2 | 0 | 1.0 | 1.1 | 2.3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.4 | 1.3 | 1.4 | 1.4 | 1.5 |
| STANDARD DEVIATION.....DO..... | .7 | .4 | .7 | .7 | .8 |
| MAXIMUM.....DO..... | 6.0 | 2.0 | 5.0 | 5.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 11 | 4 | 7 | 8 | 11 |
| STANDARD DEVIATION.....GM..... | 11 | 1 | 8 | 7 | 11 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 2 | | 2 | 2 | 2 |
| 25TH.....GM..... | 5 | | 3 | 3 | 4 |
| 50TH.....GM..... | 7 | | 5 | 6 | 7 |
| 75TH.....GM..... | 13 | | 9 | 9 | 13 |
| 90TH.....GM..... | 24 | | 15 | 15 | 21 |
| 95TH.....GM..... | 33 | | 21 | 19 | 31 |
| 99TH.....GM..... | 57 | | 49 | 38 | 57 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 24 | 14 | 14 | 18 | 20 |
| STANDARD DEVIATION.....GM..... | 27 | 3 | 13 | 17 | 18 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 24 | 3 | 19 | 16 | 24 |
| STANDARD DEVIATION.....GM..... | 18 | 0 | 16 | 9 | 19 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 25 | 0 | 20 | 20 | 25 |
| STANDARD DEVIATION.....GM..... | 15 | 0 | 11 | 11 | 14 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 340 | 18 | 142 | 170 | 170 |
| MAXIMUM OVER 3 DAYS.....GM..... | 340 | 18 | 156 | 170 | 255 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 23 | 10 | 16 | 17 | 21 |
| STANDARD DEVIATION.....GM..... | 23 | 6 | 17 | 13 | 19 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 5 | | 5 | 5 | 5 |
| 25TH.....GM..... | 9 | | 9 | 9 | 9 |
| 50TH.....GM..... | 18 | | 10 | 14 | 18 |
| 75TH.....GM..... | 28 | | 18 | 18 | 23 |
| 90TH.....GM..... | 43 | | 28 | 28 | 36 |
| 95TH.....GM..... | 57 | | 43 | 40 | 57 |
| 99TH.....GM..... | 120 | | 113 | 57 | 128 |
| MAXIMUM.....GM..... | 340 | 18 | 142 | 170 | 170 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 240.0 | 0 | 6.9 | 6.7 | 21.7 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.6 | 0 | 2.4 | 1.3 | 2.9 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 29.0 | 29.0 | 28.7 | 25.0 | 23.5 | 17.7 | 12.4 | 10.2 | 3.8 | 3.5 | 2.7 | 4.2 |
| 20.0 | 20.0 | 19.4 | 18.8 | 15.1 | 13.5 | 8.8 | 8.4 | 2.8 | 2.8 | 1.6 | 3.6 |
| 6.9 | 6.5 | 6.9 | 5.0 | 6.0 | 3.4 | 2.7 | 1.4 | 1.0 | .5 | 1.1 | .2 |
| 2.1 | 2.5 | 2.4 | 1.2 | 2.4 | .9 | .9 | .4 | 0 | .1 | 0 | .4 |
| 1.5 | 1.5 | 1.5 | 1.4 | 1.6 | 1.4 | 1.4 | 1.3 | 1.3 | 1.2 | 1.4 | 1.3 |
| .7 | .8 | .8 | .7 | .8 | .7 | .7 | .6 | .4 | .5 | .5 | .8 |
| 4.0 | 6.0 | 6.0 | 6.0 | 6.0 | 5.0 | 4.0 | 5.0 | 2.0 | 3.0 | 2.0 | 4.0 |
| 12 | 12 | 14 | 11 | 14 | 10 | 11 | 9 | 8 | 6 | 5 | 8 |
| 12 | 11 | 14 | 12 | 14 | 9 | 9 | 10 | 8 | 5 | 3 | 6 |
| 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | | 2 | | |
| 6 | 6 | 6 | 5 | 6 | 5 | 5 | 3 | | 3 | | |
| 9 | 9 | 9 | 7 | 9 | 7 | 7 | 6 | | 6 | | |
| 14 | 14 | 18 | 12 | 18 | 12 | 12 | 9 | | 8 | | |
| 24 | 24 | 28 | 24 | 30 | 19 | 22 | 19 | | 9 | | |
| 38 | 33 | 46 | 38 | 42 | 27 | 31 | 24 | | 9 | | |
| 57 | 57 | 76 | 66 | 66 | 38 | 43 | 57 | | 30 | | |
| 28 | 27 | 33 | 26 | 29 | 24 | 24 | 22 | 22 | 17 | 12 | 19 |
| 29 | 27 | 38 | 26 | 35 | 25 | 21 | 27 | 26 | 8 | 5 | 12 |
| 27 | 26 | 27 | 26 | 27 | 22 | 23 | 21 | 18 | 8 | 10 | 13 |
| 21 | 19 | 20 | 20 | 19 | 12 | 15 | 16 | 9 | 3 | 5 | 0 |
| 21 | 20 | 30 | 31 | 31 | 21 | 23 | 24 | 0 | 30 | 0 | 20 |
| 9 | 10 | 18 | 24 | 18 | 8 | 11 | 12 | 0 | 0 | 0 | 8 |
| 284 | 227 | 227 | 227 | 340 | 340 | 128 | 284 | 142 | 36 | 18 | 73 |
| 312 | 255 | 245 | 263 | 340 | 340 | 170 | 284 | 142 | 90 | 27 | 78 |
| 25 | 24 | 28 | 25 | 27 | 21 | 23 | 21 | 20 | 16 | 11 | 18 |
| 25 | 22 | 29 | 24 | 28 | 20 | 18 | 24 | 22 | 9 | 6 | 16 |
| 5 | 6 | 5 | 5 | 5 | 5 | 5 | 5 | | 3 | | |
| 11 | 11 | 14 | 11 | 10 | 9 | 14 | 9 | | 8 | | |
| 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 | | 18 | | |
| 28 | 28 | 32 | 28 | 30 | 27 | 28 | 24 | | 20 | | |
| 54 | 43 | 57 | 53 | 57 | 36 | 43 | 36 | | 28 | | |
| 57 | 57 | 85 | 60 | 57 | 57 | 57 | 57 | | 36 | | |
| 128 | 128 | 170 | 113 | 142 | 85 | 90 | 128 | | 36 | | |
| 284 | 227 | 227 | 227 | 340 | 340 | 128 | 284 | 142 | 36 | 18 | 70 |
| 24.1 | 24.7 | 14.4 | 17.4 | 32.1 | 18.8 | 31.5 | 33.4 | 5.2 | 3.2 | 0 | 0 |
| 2.7 | 2.7 | 2.4 | 3.5 | 2.2 | 1.4 | 3.6 | 3.7 | 9.7 | 4.4 | 0 | 0 |

TABLE 5.34.--CORN-BASED SALTY SNACKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS 1/ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.9 | .6 | 6.6 | 8.6 | 10.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.5 | .4 | 5.2 | 6.8 | 7.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.1 | 0 | 1.3 | 1.3 | 1.9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | .2 | .1 | .4 | .8 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE 1/NUMBER..... | 1.4 | 1.7 | 1.3 | 1.3 | 1.4 |
| STANDARD DEVIATION.....DO..... | .7 | 1.0 | .5 | .6 | .8 |
| MAXIMUM.....DO..... | 8.0 | 3.0 | 3.0 | 4.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY 1/GM..... | 15 | 4 | 10 | 12 | 14 |
| STANDARD DEVIATION.....GM..... | 15 | 2 | 9 | 10 | 13 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: 1/ | | | | | |
| 5TH.....GM..... | 3 | | 1 | 3 | 3 |
| 25TH.....GM..... | 7 | | 4 | 5 | 7 |
| 50TH.....GM..... | 12 | | 7 | 7 | 9 |
| 75TH.....GM..... | 19 | | 14 | 13 | 17 |
| 90TH.....GM..... | 28 | | 22 | 27 | 30 |
| 95TH.....GM..... | 38 | | 27 | 38 | 38 |
| 99TH.....GM..... | 76 | | 38 | 57 | 57 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE 1/GM..... | 35 | 9 | 25 | 28 | 32 |
| STANDARD DEVIATION.....GM..... | 31 | 2 | 22 | 25 | 34 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE 1/GM..... | 34 | 0 | 28 | 24 | 30 |
| STANDARD DEVIATION.....GM..... | 25 | 0 | 13 | 12 | 14 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE 1/GM..... | 37 | 7 | 5 | 33 | 29 |
| STANDARD DEVIATION.....GM..... | 33 | 0 | 0 | 16 | 16 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 340 | 11 | 113 | 113 | 340 |
| MAXIMUM OVER 3 DAYS.....GM..... | 542 | 21 | 120 | 170 | 340 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE 1/GM..... | 33 | 8 | 24 | 27 | 29 |
| STANDARD DEVIATION.....GM..... | 30 | 2 | 20 | 22 | 26 |
| PERCENTILES: 1/ | | | | | |
| 5TH.....GM..... | 9 | | 4 | 10 | 10 |
| 25TH.....GM..... | 18 | | 10 | 13 | 16 |
| 50TH.....GM..... | 21 | | 18 | 20 | 21 |
| 75TH.....GM..... | 40 | | 36 | 36 | 35 |
| 90TH.....GM..... | 60 | | 54 | 43 | 57 |
| 95TH.....GM..... | 80 | | 57 | 72 | 80 |
| 99TH.....GM..... | 156 | | 85 | 113 | 113 |
| MAXIMUM.....GM..... | 340 | 11 | 113 | 113 | 340 |
| QUANTITIES NOT SPECIFIED BY USERS: 1/ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 67.2 | 1.0 | 4.0 | 2.8 | 4.0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.2 | 20.2 | 4.6 | 1.5 | 1.5 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 9.9 | 11.3 | 8.3 | 10.7 | 8.2 | 6.3 | 3.8 | 3.1 | 1.2 | .5 | 1.4 | 1.7 |
| 6.9 | 8.0 | 6.5 | 8.5 | 6.0 | 5.1 | 2.8 | 2.6 | 1.2 | .5 | 1.2 | 1.3 |
| 2.5 | 2.7 | 1.4 | 1.7 | 1.8 | .9 | .7 | .4 | 0 | 0 | .2 | .2 |
| .5 | .6 | .4 | .6 | .4 | .3 | .3 | .1 | 0 | 0 | 0 | .2 |
| 1.4 | 1.4 | 1.4 | 1.4 | 1.3 | 1.2 | 1.4 | 1.3 | 1.2 | 1.0 | 1.1 | 1.4 |
| .7 | .7 | .8 | .7 | .6 | .5 | .7 | .9 | .4 | .0 | .3 | .7 |
| 4.0 | 4.0 | 5.0 | 5.0 | 5.0 | 3.0 | 4.0 | 8.0 | 2.0 | 1.0 | 2.0 | 3.0 |
| 16 | 15 | 21 | 16 | 18 | 14 | 16 | 14 | 8 | 6 | 6 | 10 |
| 15 | 13 | 22 | 11 | 19 | 13 | 13 | 19 | 6 | 2 | 4 | 8 |
| 3 | 3 | 5 | 3 | 3 | 4 | 3 | 2 | | | | |
| 7 | 7 | 7 | 7 | 7 | 7 | 7 | 6 | | | | |
| 13 | 13 | 13 | 13 | 13 | 12 | 13 | 9 | | | | |
| 19 | 20 | 27 | 20 | 20 | 14 | 23 | 13 | | | | |
| 30 | 28 | 52 | 28 | 30 | 27 | 28 | 27 | | | | |
| 41 | 40 | 71 | 38 | 40 | 38 | 33 | 40 | | | | |
| 95 | 57 | 95 | 53 | 88 | 75 | 61 | 151 | | | | |
| 37 | 37 | 50 | 39 | 39 | 32 | 39 | 33 | 25 | 19 | 19 | 21 |
| 34 | 37 | 48 | 26 | 26 | 20 | 31 | 35 | 19 | 7 | 13 | 15 |
| 33 | 32 | 52 | 34 | 38 | 46 | 25 | 27 | 0 | 0 | 5 | 18 |
| 27 | 18 | 48 | 20 | 20 | 36 | 14 | 14 | 0 | 0 | 0 | 0 |
| 36 | 26 | 45 | 34 | 56 | 26 | 35 | 74 | 0 | 0 | 0 | 28 |
| 16 | 9 | 25 | 8 | 59 | 8 | 21 | 65 | 0 | 0 | 0 | 0 |
| 284 | 340 | 227 | 128 | 340 | 252 | 184 | 240 | 57 | 28 | 40 | 56 |
| 288 | 340 | 397 | 213 | 542 | 280 | 280 | 454 | 57 | 28 | 40 | 85 |
| 33 | 32 | 46 | 34 | 39 | 34 | 33 | 32 | 21 | 19 | 15 | 23 |
| 29 | 30 | 44 | 22 | 36 | 28 | 26 | 30 | 11 | 7 | 13 | 12 |
| 7 | 9 | 14 | 10 | 10 | 10 | 10 | 7 | | | | |
| 20 | 20 | 20 | 20 | 20 | 20 | 20 | 14 | | | | |
| 22 | 21 | 30 | 28 | 28 | 28 | 21 | 21 | | | | |
| 40 | 40 | 54 | 40 | 44 | 40 | 40 | 40 | | | | |
| 60 | 57 | 85 | 60 | 80 | 57 | 72 | 57 | | | | |
| 80 | 80 | 160 | 85 | 85 | 80 | 80 | 85 | | | | |
| 128 | 170 | 213 | 113 | 170 | 156 | 120 | 170 | | | | |
| 284 | 340 | 227 | 128 | 340 | 252 | 184 | 240 | 40 | 28 | 40 | 56 |
| 5.4 | 5.3 | 4.2 | 7.9 | 8.6 | 9.0 | 8.8 | 4.1 | 0 | 0 | 0 | 2.2 |
| 1.8 | 1.5 | 2.6 | 3.7 | 2.0 | 2.2 | 3.3 | 1.4 | 0 | 0 | 0 | 9.2 |

TABLE 5.35.--PASTA
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 11.4 | 3.4 | 14.1 | 14.7 | 14.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 10.1 | 2.4 | 11.5 | 12.3 | 12.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.3 | .7 | 2.5 | 2.3 | 1.8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | .3 | .1 | (2) | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 1.5 | 1.2 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .4 | .7 | .5 | .4 | .4 |
| MAXIMUM.....DO..... | 6.0 | 3.0 | 4.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 59 | 28 | 33 | 39 | 45 |
| STANDARD DEVIATION.....GM..... | 46 | 32 | 26 | 26 | 31 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 12 | | 5 | 12 | 9 |
| 25TH.....GM..... | 27 | | 13 | 23 | 23 |
| 50TH.....GM..... | 47 | | 27 | 27 | 40 |
| 75TH.....GM..... | 80 | | 47 | 47 | 53 |
| 90TH.....GM..... | 107 | | 70 | 70 | 93 |
| 95TH.....GM..... | 140 | | 93 | 93 | 103 |
| 99TH.....GM..... | 233 | | 123 | 123 | 160 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 156 | 50 | 83 | 100 | 119 |
| STANDARD DEVIATION.....GM..... | 110 | 30 | 58 | 61 | 75 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 159 | 52 | 82 | 100 | 122 |
| STANDARD DEVIATION.....GM..... | 106 | 42 | 45 | 41 | 61 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 141 | 98 | 192 | 157 | 0 |
| STANDARD DEVIATION.....GM..... | 61 | 48 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 980 | 160 | 490 | 420 | 480 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,960 | 413 | 577 | 472 | 620 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 153 | 58 | 82 | 99 | 116 |
| STANDARD DEVIATION.....GM..... | 108 | 42 | 59 | 58 | 74 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 35 | | 14 | 35 | 27 |
| 25TH.....GM..... | 70 | | 40 | 70 | 70 |
| 50TH.....GM..... | 140 | | 70 | 80 | 105 |
| 75TH.....GM..... | 210 | | 105 | 140 | 140 |
| 90TH.....GM..... | 280 | | 160 | 160 | 210 |
| 95TH.....GM..... | 320 | | 210 | 210 | 280 |
| 99TH.....GM..... | 560 | | 280 | 280 | 320 |
| MAXIMUM.....GM..... | 980 | 160 | 350 | 400 | 480 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 22.6 | 0 | .7 | .7 | 3.4 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .5 | 0 | .4 | .2 | 1.1 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 14.0 | 14.5 | 11.2 | 10.8 | 12.3 | 12.6 | 10.8 | 9.9 | 8.9 | 6.7 | 7.5 | 7.3 |
| 12.7 | 13.1 | 10.3 | 9.8 | 10.8 | 11.0 | 9.8 | 8.7 | 7.8 | 5.8 | 6.8 | 5.7 |
| 1.2 | 1.4 | .9 | 1.0 | 1.4 | 1.5 | 1.0 | 1.1 | .9 | .9 | .7 | 1.4 |
| .1 | (2) | 0 | 0 | .1 | (2) | 0 | (2) | .3 | 0 | 0 | .2 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.2 | 1.1 | 1.3 |
| .4 | .4 | .4 | .3 | .4 | .4 | .3 | .4 | .8 | .5 | .3 | .5 |
| 3.0 | 4.0 | 4.0 | 2.0 | 3.0 | 3.0 | 3.0 | 3.0 | 6.0 | 4.0 | 2.0 | 3.0 |
| 60 | 54 | 75 | 59 | 76 | 61 | 70 | 54 | 57 | 50 | 51 | 51 |
| 42 | 38 | 58 | 43 | 60 | 45 | 52 | 38 | 47 | 48 | 28 | 31 |
| 13 | 12 | 20 | 12 | 13 | 13 | 17 | 12 | 18 | 13 | | 7 |
| 27 | 27 | 38 | 27 | 41 | 27 | 35 | 27 | 27 | 26 | | 27 |
| 47 | 47 | 53 | 47 | 53 | 47 | 53 | 47 | 47 | 40 | | 47 |
| 93 | 70 | 93 | 80 | 93 | 80 | 93 | 70 | 76 | 53 | | 70 |
| 107 | 105 | 151 | 107 | 147 | 107 | 135 | 104 | 107 | 93 | | 100 |
| 140 | 117 | 210 | 157 | 187 | 140 | 187 | 117 | 140 | 107 | | 107 |
| 213 | 187 | 280 | 207 | 280 | 280 | 280 | 210 | 245 | 373 | | 107 |
| 164 | 142 | 198 | 160 | 201 | 158 | 192 | 147 | 146 | 126 | 145 | 131 |
| 104 | 89 | 144 | 114 | 141 | 103 | 129 | 95 | 91 | 75 | 82 | 83 |
| 169 | 173 | 250 | 177 | 208 | 179 | 200 | 138 | 143 | 145 | 113 | 118 |
| 97 | 75 | 118 | 66 | 141 | 102 | 128 | 84 | 112 | 151 | 22 | 39 |
| 140 | 110 | 0 | 0 | 171 | 80 | 0 | 210 | 165 | 0 | 0 | 80 |
| 0 | 0 | 0 | 0 | 47 | 0 | 0 | 0 | 87 | 0 | 0 | 0 |
| 640 | 560 | 720 | 800 | 980 | 640 | 840 | 700 | 613 | 560 | 320 | 320 |
| 840 | 700 | 1,040 | 800 | 1,960 | 980 | 1,120 | 840 | 900 | 1,120 | 320 | 320 |
| 162 | 145 | 198 | 158 | 198 | 160 | 189 | 142 | 138 | 127 | 133 | 121 |
| 102 | 89 | 133 | 99 | 141 | 101 | 123 | 95 | 108 | 84 | 69 | 71 |
| 40 | 35 | 61 | 35 | 40 | 40 | 40 | 35 | 35 | 35 | | 20 |
| 80 | 80 | 120 | 80 | 105 | 80 | 105 | 70 | 70 | 70 | | 80 |
| 140 | 140 | 140 | 140 | 160 | 140 | 160 | 140 | 105 | 85 | | 99 |
| 210 | 175 | 280 | 210 | 280 | 210 | 280 | 160 | 160 | 160 | | 160 |
| 280 | 280 | 320 | 280 | 350 | 280 | 320 | 280 | 280 | 280 | | 210 |
| 320 | 280 | 454 | 320 | 480 | 350 | 420 | 320 | 420 | 320 | | 280 |
| 560 | 420 | 701 | 480 | 700 | 420 | 640 | 480 | 560 | 341 | | 320 |
| 640 | 560 | 720 | 560 | 980 | 640 | 840 | 700 | 560 | 480 | 320 | 320 |
| .8 | 0 | 1.4 | .9 | 1.1 | 5.3 | 4.0 | 2.2 | 0 | 0 | 2.1 | 0 |
| .2 | 0 | .8 | .5 | .2 | .7 | .7 | .3 | 0 | 0 | 4.5 | 0 |

TABLE 5.36.--SPAGHETTI
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.6 | 1.6 | 5.8 | 7.1 | 7.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.2 | 1.3 | 5.2 | 6.2 | 7.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | .3 | .7 | .9 | .6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 1.3 | 1.1 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .3 | .5 | .3 | .3 | .3 |
| MAXIMUM.....DO..... | 4.0 | 2.0 | 2.0 | 2.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 63 | 22 | 31 | 40 | 43 |
| STANDARD DEVIATION.....GM..... | 48 | 25 | 21 | 26 | 25 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 12 | | 9 | 12 | 9 |
| 25TH.....GM..... | 35 | | 18 | 23 | 23 |
| 50TH.....GM..... | 47 | | 23 | 35 | 47 |
| 75TH.....GM..... | 93 | | 47 | 47 | 47 |
| 90TH.....GM..... | 107 | | 47 | 93 | 82 |
| 95TH.....GM..... | 140 | | 82 | 93 | 93 |
| 99TH.....GM..... | 245 | | 93 | 117 | 117 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 173 | 42 | 85 | 104 | 124 |
| STANDARD DEVIATION.....GM..... | 118 | 33 | 58 | 64 | 69 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 185 | 82 | 72 | 111 | 105 |
| STANDARD DEVIATION.....GM..... | 124 | 52 | 40 | 44 | 52 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 245 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 0 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 980 | 140 | 280 | 280 | 280 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,960 | 253 | 280 | 350 | 350 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 172 | 53 | 83 | 104 | 118 |
| STANDARD DEVIATION.....GM..... | 117 | 39 | 55 | 59 | 66 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 35 | | 26 | 35 | 28 |
| 25TH.....GM..... | 87 | | 35 | 70 | 70 |
| 50TH.....GM..... | 140 | | 70 | 105 | 140 |
| 75TH.....GM..... | 210 | | 140 | 140 | 140 |
| 90TH.....GM..... | 280 | | 140 | 140 | 210 |
| 95TH.....GM..... | 420 | | 140 | 210 | 280 |
| 99TH.....GM..... | 561 | | 280 | 280 | 280 |
| MAXIMUM.....GM..... | 980 | 130 | 280 | 280 | 280 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 10.1 | 0 | .7 | .7 | 2.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | 1.0 | .5 | 1.6 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 7.1 | 7.4 | 6.4 | 5.4 | 6.0 | 6.4 | 5.0 | 4.9 | 4.2 | 2.6 | 3.4 | 2.6 |
| 6.5 | 7.0 | 5.9 | 5.1 | 5.7 | 5.8 | 4.7 | 4.5 | 4.0 | 2.2 | 3.4 | 2.5 |
| . | . | .5 | .3 | .3 | .6 | .2 | .4 | .1 | .3 | 0 | .1 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | .1 | 0 | 0 | 0 |
| 1.1 | 1.1 | 1.2 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.2 | 1.1 | 1.0 |
| .3 | .3 | .5 | .3 | .2 | .3 | .2 | .3 | .6 | .6 | .3 | .2 |
| 3.0 | 3.0 | 4.0 | 2.0 | 2.0 | 2.0 | 3.0 | 2.0 | 4.0 | 4.0 | 2.0 | 2.0 |
| 64 | 59 | 82 | 58 | 83 | 67 | 74 | 58 | 66 | 59 | 44 | 41 |
| 43 | 37 | 58 | 36 | 67 | 47 | 51 | 39 | 50 | 63 | 22 | 24 |
| 18 | 12 | 23 | 12 | 18 | 18 | 23 | 12 | | | | |
| 38 | 38 | 47 | 35 | 47 | 47 | 38 | 35 | | | | |
| 47 | 47 | 70 | 47 | 70 | 47 | 65 | 47 | | | | |
| 93 | 70 | 93 | 70 | 93 | 93 | 93 | 93 | | | | |
| 117 | 93 | 187 | 107 | 187 | 117 | 140 | 93 | | | | |
| 140 | 117 | 213 | 140 | 187 | 140 | 187 | 140 | | | | |
| 233 | 187 | 280 | 163 | 280 | 280 | 280 | 187 | | | | |
| 172 | 165 | 222 | 164 | 233 | 177 | 212 | 162 | 175 | 139 | 133 | 121 |
| 96 | 94 | 150 | 97 | 163 | 103 | 138 | 102 | 109 | 82 | 66 | 72 |
| 207 | 194 | 269 | 174 | 267 | 222 | 218 | 166 | 175 | 214 | 0 | 92 |
| 104 | 77 | 95 | 61 | 215 | 111 | 145 | 80 | 0 | 199 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 245 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 560 | 560 | 700 | 490 | 980 | 640 | 840 | 700 | 613 | 560 | 280 | 280 |
| 840 | 700 | 840 | 490 | 1,960 | 840 | 980 | 840 | 736 | 1,120 | 280 | 280 |
| 176 | 166 | 213 | 162 | 236 | 183 | 210 | 161 | 170 | 148 | 119 | 118 |
| 97 | 93 | 129 | 87 | 170 | 108 | 135 | 99 | 124 | 86 | 38 | 70 |
| 53 | 35 | 70 | 53 | 53 | 53 | 70 | 35 | | | | |
| 113 | 112 | 140 | 105 | 140 | 105 | 112 | 105 | | | | |
| 140 | 140 | 140 | 140 | 210 | 140 | 158 | 140 | | | | |
| 210 | 210 | 280 | 210 | 280 | 280 | 280 | 210 | | | | |
| 280 | 280 | 420 | 280 | 448 | 280 | 350 | 280 | | | | |
| 280 | 280 | 454 | 320 | 561 | 420 | 420 | 280 | | | | |
| 560 | 420 | 641 | 420 | 841 | 560 | 701 | 560 | | | | |
| 560 | 560 | 700 | 420 | 980 | 640 | 840 | 700 | 560 | 341 | 210 | 280 |
| .8 | 0 | 1.4 | .9 | 0 | 1.4 | 1.6 | 0 | 0 | 0 | 0 | 0 |
| .5 | 0 | 1.4 | 1.0 | 0 | .4 | .6 | 0 | 0 | 0 | 0 | 0 |

TABLE 5.37.--NOODLES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.7 | 1.5 | 5.3 | 4.8 | 4.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.4 | .9 | 4.4 | 4.3 | 4.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .3 | .5 | .9 | .5 | .3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | .1 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.1 | 1.6 | 1.2 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .3 | .7 | .4 | .3 | .3 |
| MAXIMUM.....DO..... | 3.0 | 3.0 | 2.0 | 2.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 52 | 34 | 29 | 34 | 42 |
| STANDARD DEVIATION.....GM..... | 41 | 20 | 18 | 21 | 32 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 13 | | 3 | 13 | 13 |
| 25TH.....GM..... | 27 | | 13 | 27 | 27 |
| 50TH.....GM..... | 53 | | 27 | 27 | 27 |
| 75TH.....GM..... | 53 | | 40 | 53 | 53 |
| 90TH.....GM..... | 107 | | 53 | 53 | 80 |
| 95TH.....GM..... | 107 | | 60 | 67 | 107 |
| 99TH.....GM..... | 213 | | 80 | 133 | 160 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 143 | 89 | 77 | 97 | 121 |
| STANDARD DEVIATION.....GM..... | 103 | 41 | 50 | 59 | 86 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 141 | 60 | 64 | 80 | 118 |
| STANDARD DEVIATION.....GM..... | 105 | 44 | 26 | 30 | 80 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 119 | 40 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 60 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 800 | 160 | 240 | 400 | 480 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,120 | 240 | 240 | 400 | 480 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 140 | 65 | 71 | 93 | 117 |
| STANDARD DEVIATION.....GM..... | 98 | 47 | 49 | 56 | 86 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 40 | | 10 | 40 | 40 |
| 25TH.....GM..... | 80 | | 40 | 61 | 80 |
| 50TH.....GM..... | 120 | | 61 | 80 | 80 |
| 75TH.....GM..... | 160 | | 80 | 120 | 160 |
| 90TH.....GM..... | 320 | | 160 | 160 | 200 |
| 95TH.....GM..... | 320 | | 160 | 160 | 320 |
| 99TH.....GM..... | 480 | | 240 | 400 | 480 |
| MAXIMUM.....GM..... | 720 | 160 | 240 | 400 | 480 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 6.7 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.9 | 4.6 | 3.0 | 3.0 | 4.3 | 3.7 | 3.5 | 3.2 | 3.6 | 3.0 | 2.4 | 4.2 |
| 3.6 | 4.3 | 2.9 | 2.8 | 4.0 | 3.4 | 3.2 | 3.1 | 3.3 | 2.9 | 2.4 | 3.2 |
| .3 | .3 | .1 | .2 | .3 | .4 | .3 | .1 | .3 | .1 | 0 | .8 |
| 0 | 0 | 0 | 0 | .1 | (2) | 0 | 0 | 0 | 0 | 0 | .2 |
| 1.1 | 1.1 | 1.0 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.0 | 1.0 | 1.3 |
| .3 | .3 | .2 | .3 | .4 | .4 | .4 | .3 | .3 | .2 | .0 | .6 |
| 2.0 | 2.0 | 2.0 | 2.0 | 3.0 | 3.0 | 3.0 | 3.0 | 2.0 | 2.0 | 1.0 | 3.0 |
| 52 | 45 | 64 | 58 | 63 | 52 | 72 | 47 | 44 | 41 | 49 | 54 |
| 35 | 34 | 60 | 51 | 45 | 36 | 58 | 31 | 30 | 28 | 31 | 30 |
| 13 | 13 | | | 13 | 13 | 13 | 7 | | 7 | | |
| 27 | 27 | | | 27 | 27 | 40 | 27 | | 27 | | |
| 53 | 27 | | | 53 | 53 | 53 | 53 | | 27 | | |
| 53 | 53 | | | 80 | 53 | 80 | 53 | | 53 | | |
| 107 | 107 | | | 107 | 107 | 133 | 107 | | 57 | | |
| 107 | 107 | | | 160 | 120 | 187 | 107 | | 107 | | |
| 213 | 187 | | | 213 | 160 | 373 | 160 | | 160 | | |
| 152 | 124 | 171 | 168 | 165 | 144 | 186 | 140 | 118 | 122 | 148 | 144 |
| 107 | 87 | 124 | 155 | 105 | 97 | 124 | 91 | 82 | 86 | 92 | 87 |
| 96 | 142 | 520 | 139 | 205 | 126 | 252 | 93 | 129 | 60 | 0 | 105 |
| 38 | 72 | 0 | 42 | 75 | 72 | 150 | 47 | 39 | 0 | 0 | 42 |
| 0 | 0 | 0 | 0 | 171 | 80 | 0 | 0 | 0 | 0 | 0 | 80 |
| 0 | 0 | 0 | 0 | 47 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 640 | 480 | 720 | 800 | 480 | 480 | 800 | 480 | 480 | 480 | 320 | 320 |
| 640 | 560 | 1,040 | 800 | 640 | 640 | 1,120 | 480 | 480 | 480 | 320 | 320 |
| 142 | 126 | 186 | 159 | 168 | 136 | 193 | 133 | 120 | 118 | 148 | 121 |
| 103 | 85 | 148 | 122 | 103 | 79 | 118 | 88 | 81 | 85 | 92 | 75 |
| 40 | 40 | | | 40 | 40 | 40 | 20 | | 20 | | |
| 80 | 80 | | | 80 | 80 | 113 | 80 | | 80 | | |
| 120 | 80 | | | 160 | 120 | 160 | 150 | | 80 | | |
| 160 | 160 | | | 240 | 160 | 240 | 160 | | 160 | | |
| 320 | 240 | | | 320 | 240 | 320 | 240 | | 170 | | |
| 320 | 320 | | | 320 | 320 | 400 | 320 | | 320 | | |
| 640 | 480 | | | 480 | 320 | 640 | 480 | | 480 | | |
| 640 | 480 | 720 | 480 | 480 | 360 | 640 | 480 | 480 | 480 | 320 | 320 |
| 0 | 0 | 0 | 0 | 1.1 | 2.6 | 1.0 | 0 | 0 | 0 | 2.1 | 0 |
| 0 | 0 | 0 | 0 | .6 | 1.1 | .5 | 0 | 0 | 0 | 16.4 | 0 |

TABLE 5.38.--MACARONI
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.5 | .4 | 3.7 | 3.3 | 2.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.3 | .4 | 3.2 | 2.9 | 2.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .2 | 0 | .5 | .5 | .1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.1 | 1.0 | 1.2 | 1.2 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .0 | .4 | .4 | .3 |
| MAXIMUM.....DO..... | 6.0 | 1.0 | 2.0 | 3.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 49 | 15 | 37 | 37 | 39 |
| STANDARD DEVIATION.....GM..... | 37 | 7 | 28 | 26 | 30 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 12 | | | 12 | 6 |
| 25TH.....GM..... | 23 | | | 23 | 23 |
| 50TH.....GM..... | 47 | | | 23 | 23 |
| 75TH.....GM..... | 58 | | | 47 | 47 |
| 90TH.....GM..... | 93 | | | 70 | 93 |
| 95TH.....GM..... | 104 | | | 93 | 103 |
| 99TH.....GM..... | 187 | | | 140 | 140 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 134 | 46 | 101 | 98 | 111 |
| STANDARD DEVIATION.....GM..... | 94 | 22 | 81 | 67 | 84 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 139 | 0 | 82 | 100 | 135 |
| STANDARD DEVIATION.....GM..... | 91 | 0 | 41 | 43 | 24 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 129 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 69 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 700 | 70 | 490 | 420 | 420 |
| MAXIMUM OVER 3 DAYS.....GM..... | 980 | 70 | 490 | 420 | 420 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 131 | 46 | 93 | 96 | 109 |
| STANDARD DEVIATION.....GM..... | 93 | 22 | 67 | 57 | 73 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 35 | | | 35 | 18 |
| 25TH.....GM..... | 70 | | | 70 | 70 |
| 50TH.....GM..... | 140 | | | 70 | 70 |
| 75TH.....GM..... | 140 | | | 140 | 140 |
| 90TH.....GM..... | 280 | | | 140 | 210 |
| 95TH.....GM..... | 280 | | | 210 | 280 |
| 99TH.....GM..... | 420 | | | 280 | 420 |
| MAXIMUM.....GM..... | 700 | 70 | 350 | 310 | 420 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 5.8 | 0 | 0 | 0 | .8 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | 0 | 0 | 0 | 1.4 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.4 | 3.1 | 1.8 | 2.8 | 2.4 | 2.7 | 2.5 | 2.2 | 1.6 | 1.3 | 2.4 | .9 |
| 3.3 | 2.9 | 1.7 | 2.7 | 2.2 | 2.5 | 2.3 | 1.9 | 1.5 | 1.1 | 2.4 | .9 |
| .1 | .2 | .1 | .1 | .2 | .3 | .2 | .2 | 0 | .2 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | (2) | .1 | 0 | 0 | 0 |
| 1.0 | 1.1 | 1.1 | 1.0 | 1.1 | 1.1 | 1.1 | 1.2 | 1.4 | 1.2 | 1.0 | 1.0 |
| .2 | .2 | .3 | .2 | .3 | .3 | .3 | .4 | 1.3 | .4 | .0 | .0 |
| 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 3.0 | 6.0 | 2.0 | 1.0 | 1.0 |
| 54 | 44 | 57 | 52 | 63 | 51 | 53 | 45 | 46 | 43 | 47 | 43 |
| 41 | 28 | 48 | 33 | 42 | 45 | 33 | 38 | 31 | 32 | 26 | 16 |
| 12 | 6 | | | 6 | 17 | 12 | 12 | | | | |
| 23 | 23 | | | 47 | 23 | 23 | 23 | | | | |
| 47 | 47 | | | 47 | 47 | 47 | 35 | | | | |
| 93 | 47 | | | 93 | 47 | 93 | 47 | | | | |
| 93 | 93 | | | 98 | 93 | 93 | 93 | | | | |
| 140 | 93 | | | 140 | 140 | 93 | 93 | | | | |
| 187 | 105 | | | 187 | 327 | 187 | 233 | | | | |
| 153 | 124 | 155 | 155 | 166 | 129 | 158 | 117 | 132 | 112 | 140 | 129 |
| 112 | 79 | 127 | 100 | 97 | 92 | 99 | 83 | 95 | 73 | 78 | 47 |
| 247 | 127 | 181 | 105 | 204 | 185 | 98 | 117 | 0 | 118 | 0 | 0 |
| 35 | 29 | 96 | 0 | 75 | 121 | 51 | 91 | 0 | 69 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 210 | 70 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 560 | 280 | 700 | 560 | 420 | 560 | 560 | 560 | 420 | 280 | 280 | 210 |
| 560 | 315 | 700 | 560 | 560 | 980 | 560 | 700 | 420 | 350 | 280 | 210 |
| 156 | 124 | 159 | 149 | 168 | 139 | 144 | 115 | 100 | 108 | 140 | 129 |
| 107 | 76 | 124 | 96 | 100 | 101 | 93 | 87 | 90 | 74 | 78 | 47 |
| 35 | 18 | | | 17 | 52 | 35 | 26 | | | | |
| 70 | 70 | | | 105 | 70 | 70 | 70 | | | | |
| 140 | 140 | | | 140 | 140 | 140 | 85 | | | | |
| 280 | 140 | | | 280 | 140 | 210 | 140 | | | | |
| 280 | 280 | | | 280 | 280 | 280 | 210 | | | | |
| 420 | 280 | | | 310 | 420 | 280 | 280 | | | | |
| 420 | 280 | | | 420 | 560 | 560 | 560 | | | | |
| 420 | 280 | 700 | 560 | 420 | 560 | 560 | 560 | 420 | 280 | 280 | 210 |
| 0 | 0 | 0 | 0 | 0 | 1.4 | 1.4 | 2.2 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | .9 | 1.0 | 1.2 | 0 | 0 | 0 | 0 |

TABLE 5.39.--SPAGHETTI WITH TOMATO SAUCE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 10.6 | 4.0 | 16.8 | 17.2 | 17.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 9.6 | 3.1 | 14.4 | 14.8 | 16.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.0 | .9 | 2.3 | 2.2 | 1.4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | .1 | .2 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 1.2 | 1.2 | 1.2 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .4 | .4 | .4 | .4 |
| MAXIMUM.....DO..... | 4.0 | 2.0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 132 | 69 | 71 | 90 | 98 |
| STANDARD DEVIATION.....GM..... | 93 | 32 | 44 | 53 | 64 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 42 | | 21 | 21 | 42 |
| 25TH.....GM..... | 83 | | 42 | 42 | 63 |
| 50TH.....GM..... | 104 | | 63 | 83 | 83 |
| 75TH.....GM..... | 167 | | 83 | 125 | 125 |
| 90TH.....GM..... | 250 | | 146 | 167 | 167 |
| 95TH.....GM..... | 333 | | 167 | 167 | 208 |
| 99TH.....GM..... | 458 | | 167 | 250 | 333 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 361 | 178 | 194 | 240 | 261 |
| STANDARD DEVIATION.....GM..... | 233 | 85 | 120 | 136 | 143 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 355 | 153 | 161 | 225 | 321 |
| STANDARD DEVIATION.....GM..... | 202 | 26 | 74 | 81 | 137 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 273 | 0 | 125 | 195 | 260 |
| STANDARD DEVIATION.....GM..... | 196 | 0 | 0 | 39 | 15 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 2,500 | 250 | 750 | 1,000 | 1,500 |
| MAXIMUM OVER 3 DAYS.....GM..... | 4,500 | 375 | 750 | 1,000 | 1,500 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 351 | 169 | 184 | 227 | 264 |
| STANDARD DEVIATION.....GM..... | 225 | 75 | 104 | 121 | 142 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 95 | | 47 | 63 | 125 |
| 25TH.....GM..... | 250 | | 125 | 125 | 188 |
| 50TH.....GM..... | 250 | | 155 | 250 | 250 |
| 75TH.....GM..... | 500 | | 250 | 250 | 250 |
| 90TH.....GM..... | 500 | | 250 | 375 | 500 |
| 95TH.....GM..... | 751 | | 375 | 500 | 500 |
| 99TH.....GM..... | 1,001 | | 500 | 500 | 750 |
| MAXIMUM.....GM..... | 2,500 | 250 | 500 | 1,000 | 1,500 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 67.5 | 0 | 2.1 | 6.1 | 2.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.5 | 0 | 1.0 | 1.7 | .8 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 19.1 | 17.4 | 13.2 | 12.6 | 8.9 | 9.9 | 8.8 | 7.5 | 5.2 | 5.6 | 4.4 | 3.4 |
| 16.8 | 15.6 | 12.0 | 11.5 | 8.1 | 9.2 | 8.2 | 6.9 | 4.3 | 4.7 | 2.8 | 2.8 |
| 2.1 | 1.6 | 1.1 | 1.1 | .6 | .7 | .6 | .6 | .8 | .8 | 1.6 | .6 |
| .1 | .1 | .1 | 0 | .1 | .1 | 0 | (2) | .1 | .2 | 0 | 0 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.3 | 1.4 | 1.2 |
| .4 | .4 | .4 | .3 | .4 | .3 | .4 | .3 | .5 | .7 | .5 | .4 |
| 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 4.0 | 4.0 | 3.0 | 4.0 | 2.0 | 2.0 |
| 145 | 118 | 167 | 129 | 182 | 133 | 164 | 125 | 155 | 127 | 152 | 149 |
| 119 | 76 | 111 | 89 | 107 | 76 | 101 | 88 | 93 | 89 | 97 | 61 |
| 42 | 32 | 42 | 42 | 63 | 42 | 42 | 42 | 38 | 42 | | |
| 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | | |
| 125 | 83 | 167 | 83 | 167 | 125 | 167 | 83 | 139 | 83 | | |
| 167 | 167 | 167 | 167 | 250 | 167 | 208 | 167 | 250 | 167 | | |
| 250 | 208 | 292 | 250 | 333 | 250 | 292 | 208 | 292 | 250 | | |
| 333 | 250 | 417 | 258 | 375 | 250 | 333 | 319 | 333 | 333 | | |
| 500 | 417 | 584 | 500 | 583 | 417 | 500 | 417 | 417 | 417 | | |
| 380 | 323 | 460 | 342 | 500 | 372 | 470 | 337 | 406 | 328 | 374 | 390 |
| 243 | 198 | 291 | 200 | 278 | 192 | 297 | 191 | 250 | 199 | 147 | 127 |
| 432 | 317 | 452 | 422 | 523 | 351 | 403 | 387 | 377 | 377 | 300 | 353 |
| 343 | 138 | 220 | 203 | 163 | 170 | 97 | 165 | 119 | 160 | 201 | 87 |
| 212 | 253 | 333 | 0 | 439 | 317 | 0 | 1,167 | 208 | 42 | 0 | 0 |
| 44 | 47 | 0 | 0 | 13 | 62 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2,500 | 1,500 | 2,000 | 1,750 | 2,000 | 1,250 | 2,500 | 2,000 | 1,000 | 1,250 | 750 | 500 |
| 4,500 | 1,500 | 2,000 | 2,000 | 2,000 | 1,500 | 2,500 | 3,500 | 1,250 | 1,250 | 1,000 | 1,000 |
| 381 | 314 | 446 | 348 | 491 | 363 | 442 | 342 | 375 | 298 | 334 | 378 |
| 268 | 185 | 280 | 198 | 258 | 190 | 279 | 192 | 198 | 189 | 196 | 121 |
| 125 | 95 | 125 | 125 | 188 | 125 | 125 | 125 | 125 | 31 | | |
| 250 | 250 | 250 | 250 | 250 | 250 | 250 | 250 | 250 | 125 | | |
| 250 | 250 | 500 | 250 | 500 | 375 | 375 | 250 | 375 | 250 | | |
| 500 | 375 | 500 | 500 | 563 | 500 | 500 | 500 | 500 | 470 | | |
| 625 | 500 | 751 | 500 | 751 | 500 | 751 | 500 | 750 | 500 | | |
| 751 | 625 | 1,000 | 750 | 1,000 | 750 | 1,000 | 750 | 751 | 750 | | |
| 1,501 | 1,000 | 1,501 | 1,001 | 1,500 | 1,000 | 1,501 | 1,124 | 1,000 | 750 | | |
| 2,500 | 1,500 | 2,000 | 1,500 | 2,000 | 1,250 | 2,500 | 1,500 | 1,000 | 750 | 750 | 500 |
| 3.5 | 2.7 | 1.9 | 1.4 | 5.7 | 19.0 | 13.5 | 5.1 | 2.0 | 1.6 | 0 | 0 |
| .8 | .6 | .9 | .7 | 1.5 | 3.3 | 2.8 | .9 | 2.8 | 1.3 | 0 | 0 |

TABLE 5.40.--MACARONI AND CHEESE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 8.7 | 3.0 | 14.5 | 14.7 | 13.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 7.9 | 2.5 | 12.8 | 13.5 | 11.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .7 | .4 | 1.7 | 1.3 | 1.8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 1.2 | 1.1 | 1.1 | 1.2 |
| STANDARD DEVIATION.....DO..... | .3 | .4 | .3 | .3 | .4 |
| MAXIMUM.....DO..... | 3.0 | 2.0 | 2.0 | 2.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 76 | 37 | 46 | 58 | 69 |
| STANDARD DEVIATION.....GM..... | 55 | 19 | 31 | 47 | 48 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 15 | | 5 | 10 | 17 |
| 25TH.....GM..... | 33 | | 25 | 33 | 33 |
| 50TH.....GM..... | 67 | | 40 | 40 | 67 |
| 75TH.....GM..... | 83 | | 67 | 80 | 80 |
| 90TH.....GM..... | 133 | | 80 | 100 | 133 |
| 95TH.....GM..... | 167 | | 100 | 160 | 160 |
| 99TH.....GM..... | 267 | | 160 | 267 | 240 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 209 | 107 | 132 | 154 | 190 |
| STANDARD DEVIATION.....GM..... | 139 | 53 | 85 | 100 | 129 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 199 | 70 | 95 | 195 | 144 |
| STANDARD DEVIATION.....GM..... | 125 | 31 | 62 | 132 | 86 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 208 | 0 | 0 | 0 | 136 |
| STANDARD DEVIATION.....GM..... | 106 | 0 | 0 | 0 | 67 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,200 | 240 | 480 | 720 | 800 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,494 | 240 | 480 | 960 | 800 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 203 | 93 | 121 | 157 | 175 |
| STANDARD DEVIATION.....GM..... | 136 | 51 | 81 | 109 | 118 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 38 | | 13 | 25 | 30 |
| 25TH.....GM..... | 100 | | 60 | 100 | 100 |
| 50TH.....GM..... | 200 | | 100 | 120 | 150 |
| 75TH.....GM..... | 240 | | 200 | 200 | 240 |
| 90TH.....GM..... | 400 | | 240 | 240 | 300 |
| 95TH.....GM..... | 480 | | 240 | 400 | 400 |
| 99TH.....GM..... | 720 | | 360 | 480 | 720 |
| MAXIMUM.....GM..... | 1,200 | 240 | 480 | 720 | 720 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 13.7 | 0 | 2.2 | 0 | 2.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | 1.3 | 0 | 1.0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 11.2 | 11.4 | 10.2 | 8.7 | 8.1 | 8.9 | 6.6 | 7.2 | 5.3 | 6.4 | 3.9 | 5.4 |
| 10.3 | 10.5 | 9.3 | 8.0 | 7.3 | 8.1 | 6.0 | 6.7 | 5.1 | 5.8 | 3.7 | 5.0 |
| .9 | .8 | .7 | .6 | .8 | .7 | .6 | .5 | .2 | .6 | .2 | .4 |
| .1 | (2) | .1 | .1 | 0 | .1 | 0 | (2) | 0 | 0 | 0 | 0 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 |
| .3 | .3 | .4 | .4 | .3 | .4 | .3 | .3 | .2 | .3 | .4 | .3 |
| 3.0 | 3.0 | 3.0 | 3.0 | 2.0 | 3.0 | 3.0 | 3.0 | 2.0 | 2.0 | 2.0 | 2.0 |
| 79 | 73 | 108 | 78 | 107 | 75 | 85 | 65 | 62 | 84 | 69 | 61 |
| 48 | 53 | 82 | 52 | 73 | 49 | 53 | 40 | 49 | 57 | 58 | 37 |
| 22 | 17 | 33 | 20 | 33 | 17 | 17 | 8 | 8 | 8 | | 1 |
| 40 | 33 | 67 | 50 | 67 | 40 | 50 | 33 | 27 | 33 | | 33 |
| 67 | 67 | 80 | 67 | 80 | 67 | 67 | 67 | 60 | 67 | | 66 |
| 100 | 80 | 133 | 80 | 133 | 100 | 120 | 80 | 67 | 133 | | 80 |
| 133 | 133 | 200 | 133 | 200 | 133 | 160 | 120 | 120 | 160 | | 120 |
| 167 | 167 | 333 | 160 | 267 | 160 | 200 | 133 | 160 | 200 | | 133 |
| 267 | 267 | 360 | 300 | 400 | 240 | 240 | 180 | 220 | 267 | | 133 |
| 214 | 204 | 289 | 216 | 287 | 210 | 237 | 182 | 182 | 231 | 191 | 186 |
| 118 | 132 | 200 | 125 | 178 | 128 | 144 | 102 | 148 | 158 | 165 | 111 |
| 239 | 204 | 302 | 173 | 303 | 192 | 225 | 161 | 150 | 224 | 237 | 67 |
| 100 | 145 | 147 | 116 | 153 | 101 | 80 | 104 | 0 | 91 | 0 | 45 |
| 113 | 200 | 358 | 300 | 0 | 217 | 0 | 183 | 0 | 0 | 0 | 0 |
| 0 | 0 | 175 | 0 | 0 | 17 | 0 | 37 | 0 | 0 | 0 | 0 |
| 800 | 960 | 1,000 | 960 | 1,200 | 720 | 720 | 600 | 660 | 800 | 480 | 400 |
| 900 | 960 | 1,494 | 960 | 1,440 | 800 | 1,000 | 900 | 660 | 800 | 480 | 400 |
| 213 | 203 | 284 | 207 | 285 | 202 | 229 | 178 | 175 | 219 | 180 | 167 |
| 115 | 134 | 196 | 128 | 176 | 119 | 137 | 101 | 142 | 145 | 139 | 107 |
| 60 | 25 | 60 | 50 | 76 | 50 | 50 | 25 | 25 | 50 | | 4 |
| 120 | 100 | 175 | 120 | 200 | 100 | 120 | 100 | 80 | 100 | | 100 |
| 200 | 200 | 200 | 200 | 240 | 200 | 200 | 200 | 150 | 200 | | 180 |
| 240 | 240 | 400 | 240 | 400 | 240 | 300 | 240 | 200 | 300 | | 240 |
| 400 | 400 | 498 | 400 | 480 | 360 | 400 | 300 | 360 | 400 | | 360 |
| 400 | 480 | 720 | 420 | 601 | 400 | 480 | 400 | 480 | 480 | | 400 |
| 600 | 601 | 1,000 | 601 | 961 | 720 | 720 | 480 | 660 | 800 | | 400 |
| 720 | 960 | 1,000 | 960 | 1,200 | 720 | 720 | 600 | 660 | 800 | 480 | 400 |
| 3.3 | 0 | 0 | 1.5 | .9 | 0 | 1.5 | 1.4 | 0 | 0 | 0 | 0 |
| 1.3 | 0 | 0 | 1.1 | .3 | 0 | .4 | .3 | 0 | 0 | 0 | 0 |

TABLE 5.41.--LASAGNE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 1.9 | .2 | 1.6 | 1.5 | 2.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.8 | 0 | 1.2 | 1.5 | 2.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .1 | .2 | .3 | 0 | .1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 2.0 | 1.2 | 1.0 | 1.0 |
| STANDARD DEVIATION.....DO..... | .3 | .0 | .4 | .0 | .2 |
| MAXIMUM.....DO..... | 3.0 | 2.0 | 2.0 | 1.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 121 | 83 | 56 | 56 | 82 |
| STANDARD DEVIATION.....GM..... | 96 | 0 | 46 | 24 | 39 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 28 | | | | 34 |
| 25TH.....GM..... | 63 | | | | 63 |
| 50TH.....GM..... | 83 | | | | 83 |
| 75TH.....GM..... | 167 | | | | 83 |
| 90TH.....GM..... | 230 | | | | 136 |
| 95TH.....GM..... | 272 | | | | 167 |
| 99TH.....GM..... | 584 | | | | 230 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 319 | 0 | 101 | 167 | 235 |
| STANDARD DEVIATION.....GM..... | 195 | 0 | 46 | 71 | 100 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 420 | 125 | 208 | 0 | 279 |
| STANDARD DEVIATION.....GM..... | 266 | 0 | 26 | 0 | 58 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 514 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 125 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,632 | 125 | 250 | 250 | 500 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,040 | 250 | 500 | 250 | 689 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 333 | 125 | 139 | 167 | 238 |
| STANDARD DEVIATION.....GM..... | 218 | 0 | 68 | 71 | 100 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 85 | | | | 102 |
| 25TH.....GM..... | 188 | | | | 188 |
| 50TH.....GM..... | 250 | | | | 250 |
| 75TH.....GM..... | 500 | | | | 250 |
| 90TH.....GM..... | 510 | | | | 375 |
| 95TH.....GM..... | 751 | | | | 500 |
| 99TH.....GM..... | 1,001 | | | | 500 |
| MAXIMUM.....GM..... | 1,632 | 125 | 250 | 250 | 500 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 19.8 | 0 | 2.0 | 2.0 | 1.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.5 | 0 | 10.1 | 7.7 | 3.0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.9 | 2.6 | 2.8 | 2.7 | 2.8 | 1.9 | 1.7 | 1.4 | .5 | .7 | 1.4 | .6 |
| 3.7 | 2.4 | 2.6 | 2.7 | 2.5 | 1.7 | 1.5 | 1.3 | .3 | .7 | 1.4 | .6 |
| .2 | .2 | .3 | 0 | .3 | .2 | .2 | .1 | .2 | 0 | 0 | 0 |
| (2) | 0 | 0 | 0 | .1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1.1 | 1.1 | 1.1 | 1.0 | 1.1 | 1.1 | 1.1 | 1.1 | 1.3 | 1.0 | 1.0 | 1.0 |
| .3 | .3 | .3 | .0 | .4 | .3 | .3 | .3 | .5 | .0 | .0 | .0 |
| 3.0 | 2.0 | 2.0 | 1.0 | 3.0 | 2.0 | 2.0 | 2.0 | 2.0 | 1.0 | 1.0 | 1.0 |
| 102 | 93 | 132 | 89 | 188 | 126 | 153 | 111 | 137 | 103 | 101 | 59 |
| 72 | 67 | 74 | 48 | 140 | 92 | 114 | 66 | 71 | 47 | 50 | 36 |
| 28 | 32 | | | 63 | 38 | 42 | 28 | | | | |
| 63 | 42 | | | 104 | 83 | 83 | 63 | | | | |
| 83 | 83 | | | 167 | 95 | 125 | 83 | | | | |
| 125 | 95 | | | 230 | 166 | 167 | 167 | | | | |
| 167 | 167 | | | 333 | 208 | 333 | 167 | | | | |
| 272 | 250 | | | 596 | 255 | 408 | 250 | | | | |
| 442 | 375 | | | 681 | 544 | 584 | 281 | | | | |
| 280 | 237 | 376 | 267 | 454 | 320 | 390 | 320 | 460 | 310 | 304 | 176 |
| 168 | 129 | 215 | 144 | 217 | 163 | 239 | 184 | 248 | 140 | 149 | 109 |
| 387 | 389 | 294 | 0 | 679 | 425 | 524 | 244 | 156 | 0 | 0 | 0 |
| 191 | 114 | 105 | 0 | 318 | 231 | 240 | 150 | 0 | 0 | 0 | 0 |
| 323 | 0 | 0 | 0 | 595 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 816 | 750 | 1,250 | 750 | 1,632 | 1,000 | 1,250 | 816 | 750 | 500 | 500 | 250 |
| 1,326 | 1,125 | 1,250 | 750 | 2,040 | 1,632 | 1,750 | 1,000 | 750 | 500 | 500 | 250 |
| 289 | 259 | 355 | 267 | 498 | 332 | 416 | 305 | 308 | 310 | 304 | 176 |
| 171 | 145 | 218 | 144 | 287 | 174 | 245 | 180 | 241 | 140 | 149 | 109 |
| 85 | 95 | | | 188 | 113 | 125 | 85 | | | | |
| 188 | 125 | | | 313 | 230 | 250 | 159 | | | | |
| 250 | 250 | | | 500 | 250 | 375 | 250 | | | | |
| 375 | 250 | | | 562 | 459 | 500 | 425 | | | | |
| 500 | 500 | | | 817 | 500 | 817 | 500 | | | | |
| 816 | 582 | | | 1,001 | 626 | 876 | 681 | | | | |
| 816 | 750 | | | 1,632 | 875 | 1,250 | 816 | | | | |
| 816 | 750 | 1,250 | 750 | 1,632 | 875 | 1,250 | 816 | 750 | 500 | 500 | 250 |
| 4.1 | 0 | 2.8 | 1.0 | 1.3 | 2.6 | 0 | 2.3 | 0 | 0 | 0 | 0 |
| 4.8 | 0 | 6.5 | 2.6 | 1.0 | 2.2 | 0 | 2.2 | 0 | 0 | 0 | 0 |

TABLE 5.42.--PIZZA
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 9.8 | .5 | 6.9 | 10.1 | 15.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 8.9 | .5 | 6.4 | 9.4 | 14.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .8 | 0 | .6 | .6 | 1.1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | 0 | (2) | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.1 | 1.0 | 1.1 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .0 | .3 | .6 | .4 |
| MAXIMUM.....DO..... | 8.0 | 1.0 | 2.0 | 8.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 64 | 10 | 28 | 38 | 42 |
| STANDARD DEVIATION.....GM..... | 52 | 7 | 21 | 35 | 29 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 17 | | 9 | 9 | 13 |
| 25TH.....GM..... | 30 | | 13 | 20 | 23 |
| 50TH.....GM..... | 45 | | 23 | 28 | 36 |
| 75TH.....GM..... | 81 | | 37 | 45 | 45 |
| 90TH.....GM..... | 123 | | 68 | 72 | 78 |
| 95TH.....GM..... | 158 | | 76 | 84 | 94 |
| 99TH.....GM..... | 272 | | 102 | 113 | 149 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 175 | 31 | 80 | 106 | 116 |
| STANDARD DEVIATION.....GM..... | 139 | 20 | 60 | 69 | 70 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 164 | 0 | 71 | 80 | 112 |
| STANDARD DEVIATION.....GM..... | 101 | 0 | 35 | 35 | 54 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 187 | 0 | 0 | 407 | 125 |
| STANDARD DEVIATION.....GM..... | 103 | 0 | 0 | 0 | 121 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,945 | 57 | 305 | 477 | 461 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,945 | 57 | 305 | 1,222 | 840 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 171 | 31 | 78 | 102 | 115 |
| STANDARD DEVIATION.....GM..... | 133 | 20 | 59 | 62 | 70 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 43 | | 26 | 28 | 38 |
| 25TH.....GM..... | 85 | | 34 | 60 | 61 |
| 50TH.....GM..... | 136 | | 68 | 85 | 108 |
| 75TH.....GM..... | 227 | | 91 | 136 | 136 |
| 90TH.....GM..... | 338 | | 128 | 181 | 216 |
| 95TH.....GM..... | 408 | | 213 | 227 | 255 |
| 99TH.....GM..... | 639 | | 305 | 268 | 377 |
| MAXIMUM.....GM..... | 1,945 | | 305 | 457 | 461 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 1008.3 | 0 | 31.9 | 51.7 | 72.4 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 24.4 | 0 | 40.3 | 26.6 | 22.9 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 19.9 | 17.8 | 17.5 | 17.2 | 12.5 | 12.0 | 5.0 | 5.8 | 1.8 | 2.8 | 1.0 | 2.2 |
| 17.5 | 15.7 | 14.9 | 15.0 | 11.6 | 11.4 | 4.5 | 5.3 | 1.7 | 2.5 | .8 | 2.0 |
| 2.2 | 1.9 | 2.3 | 2.0 | .9 | .6 | .5 | .4 | .1 | .2 | .2 | .2 |
| .2 | .2 | .3 | .1 | (2) | (2) | 0 | (2) | 0 | 0 | 0 | 0 |
| 1.2 | 1.2 | 1.2 | 1.2 | 1.1 | 1.1 | 1.1 | 1.1 | 1.0 | 1.1 | 1.2 | 1.1 |
| .4 | .5 | .4 | .5 | .3 | .3 | .3 | .3 | .2 | .3 | .4 | .3 |
| 4.0 | 5.0 | 3.0 | 7.0 | 3.0 | 4.0 | 3.0 | 3.0 | 2.0 | 2.0 | 2.0 | 2.0 |
| 61 | 52 | 84 | 57 | 93 | 62 | 87 | 55 | 59 | 50 | 62 | 60 |
| 44 | 40 | 68 | 39 | 74 | 44 | 59 | 37 | 42 | 30 | 13 | 35 |
| 19 | 14 | 20 | 18 | 20 | 20 | 22 | 14 | | | | |
| 33 | 27 | 37 | 32 | 45 | 35 | 43 | 28 | | | | |
| 45 | 40 | 69 | 45 | 76 | 54 | 81 | 45 | | | | |
| 81 | 68 | 112 | 74 | 122 | 80 | 113 | 68 | | | | |
| 122 | 102 | 163 | 102 | 170 | 113 | 181 | 110 | | | | |
| 151 | 136 | 205 | 122 | 221 | 136 | 213 | 132 | | | | |
| 221 | 204 | 321 | 181 | 362 | 279 | 288 | 181 | | | | |
| 166 | 140 | 213 | 155 | 270 | 175 | 236 | 154 | 176 | 144 | 187 | 156 |
| 118 | 101 | 151 | 101 | 222 | 111 | 145 | 96 | 129 | 87 | 44 | 80 |
| 144 | 127 | 226 | 128 | 209 | 191 | 244 | 160 | 113 | 112 | 94 | 200 |
| 77 | 53 | 148 | 62 | 87 | 107 | 125 | 79 | 0 | 37 | 0 | 0 |
| 151 | 172 | 221 | 194 | 172 | 279 | 0 | 117 | 0 | 0 | 0 | 0 |
| 54 | 92 | 107 | 102 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 816 | 682 | 1,418 | 680 | 1,945 | 880 | 865 | 610 | 486 | 370 | 251 | 284 |
| 816 | 840 | 1,758 | 984 | 1,945 | 907 | 1,092 | 636 | 486 | 370 | 251 | 401 |
| 158 | 137 | 211 | 148 | 260 | 174 | 235 | 154 | 171 | 140 | 155 | 164 |
| 115 | 93 | 144 | 96 | 211 | 103 | 145 | 101 | 125 | 83 | 57 | 75 |
| 42 | 43 | 61 | 56 | 61 | 60 | 61 | 31 | | | | |
| 81 | 69 | 108 | 85 | 122 | 94 | 122 | 84 | | | | |
| 122 | 112 | 183 | 122 | 227 | 152 | 227 | 136 | | | | |
| 216 | 170 | 272 | 183 | 338 | 227 | 305 | 204 | | | | |
| 305 | 244 | 380 | 258 | 476 | 284 | 424 | 272 | | | | |
| 377 | 324 | 525 | 340 | 567 | 377 | 510 | 340 | | | | |
| 550 | 486 | 742 | 543 | 849 | 486 | 680 | 543 | | | | |
| 816 | 682 | 851 | 680 | 1,945 | 785 | 865 | 610 | 486 | 370 | 251 | 284 |
| 92.7 | 107.2 | 65.7 | 79.7 | 111.0 | 196.0 | 67.2 | 120.6 | 1.7 | 5.9 | 0 | 4.6 |
| 19.4 | 24.2 | 22.6 | 27.4 | 21.0 | 28.4 | 24.7 | 27.3 | 8.0 | 11.3 | 0 | 19.2 |

TABLE 5.43.--RICE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 18.5 | 4.3 | 20.9 | 22.2 | 23.4 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 13.4 | 3.4 | 14.0 | 15.3 | 17.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.6 | .8 | 4.6 | 4.9 | 4.4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.5 | .1 | 2.3 | 2.0 | 1.8 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.5 | 1.5 | 1.6 | 1.5 | 1.4 |
| STANDARD DEVIATION.....DO..... | .9 | 1.0 | 1.0 | .8 | .8 |
| MAXIMUM.....DO..... | 9.0 | 6.0 | 7.0 | 5.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 71 | 26 | 42 | 47 | 57 |
| STANDARD DEVIATION.....GM..... | 73 | 22 | 38 | 42 | 52 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 14 | | 5 | 7 | 11 |
| 25TH.....GM..... | 29 | | 15 | 19 | 29 |
| 50TH.....GM..... | 58 | | 29 | 29 | 29 |
| 75TH.....GM..... | 88 | | 58 | 58 | 58 |
| 90TH.....GM..... | 146 | | 97 | 117 | 117 |
| 95TH.....GM..... | 229 | | 131 | 117 | 175 |
| 99TH.....GM..... | 350 | | 190 | 204 | 233 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 133 | 57 | 76 | 87 | 108 |
| STANDARD DEVIATION.....GM..... | 84 | 52 | 49 | 57 | 76 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 169 | 69 | 102 | 105 | 167 |
| STANDARD DEVIATION.....GM..... | 103 | 20 | 65 | 63 | 100 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 208 | 87 | 95 | 124 | 120 |
| STANDARD DEVIATION.....GM..... | 129 | 0 | 47 | 57 | 53 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,400 | 175 | 350 | 525 | 875 |
| MAXIMUM OVER 3 DAYS.....GM..... | 3,500 | 263 | 613 | 755 | 1,225 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 147 | 53 | 81 | 95 | 120 |
| STANDARD DEVIATION.....GM..... | 91 | 42 | 50 | 58 | 77 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 41 | | 11 | 21 | 41 |
| 25TH.....GM..... | 88 | | 44 | 44 | 88 |
| 50TH.....GM..... | 165 | | 88 | 88 | 88 |
| 75TH.....GM..... | 175 | | 88 | 88 | 175 |
| 90TH.....GM..... | 263 | | 175 | 175 | 175 |
| 95TH.....GM..... | 350 | | 175 | 175 | 263 |
| 99TH.....GM..... | 438 | | 263 | 263 | 350 |
| MAXIMUM.....GM..... | 1,400 | 175 | 350 | 350 | 875 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 67.2 | 0 | 0 | 1.5 | .7 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .7 | 0 | 0 | .3 | .1 |

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 18.9 | 22.4 | 20.9 | 19.0 | 18.5 | 18.3 | 17.7 | 18.4 | 15.4 | 14.1 | 18.8 | 12.0 |
| 13.9 | 16.7 | 15.4 | 14.1 | 13.2 | 13.0 | 13.3 | 13.1 | 10.4 | 10.7 | 14.6 | 8.6 |
| 4.3 | 3.6 | 3.2 | 3.1 | 3.6 | 3.7 | 3.0 | 3.9 | 3.1 | 2.3 | 2.8 | 2.8 |
| .7 | 2.0 | 2.3 | 1.7 | 1.7 | 1.6 | 1.5 | 1.3 | 1.9 | 1.1 | 1.4 | .6 |
| 1.4 | 1.4 | 1.5 | 1.4 | 1.5 | 1.5 | 1.4 | 1.5 | 1.6 | 1.5 | 1.6 | 1.5 |
| .7 | .9 | .8 | .8 | .8 | .9 | .9 | .8 | 1.0 | 1.0 | 1.2 | .9 |
| 5.0 | 6.0 | 6.0 | 5.0 | 7.0 | 9.0 | 7.0 | 9.0 | 5.0 | 7.0 | 7.0 | 5.0 |
| 68 | 66 | 95 | 76 | 88 | 73 | 82 | 69 | 79 | 61 | 74 | 61 |
| 56 | 59 | 103 | 69 | 81 | 76 | 86 | 70 | 89 | 63 | 73 | 50 |
| 15 | 15 | 22 | 15 | 15 | 15 | 15 | 9 | 15 | 14 | 19 | 9 |
| 29 | 29 | 44 | 29 | 36 | 29 | 29 | 29 | 29 | 29 | 29 | 29 |
| 58 | 55 | 58 | 58 | 58 | 58 | 58 | 55 | 58 | 44 | 58 | 55 |
| 88 | 83 | 117 | 88 | 117 | 66 | 88 | 83 | 88 | 58 | 58 | 58 |
| 117 | 146 | 194 | 175 | 193 | 175 | 175 | 146 | 175 | 117 | 146 | 139 |
| 175 | 175 | 233 | 233 | 233 | 233 | 233 | 233 | 182 | 175 | 230 | 175 |
| 277 | 292 | 467 | 350 | 467 | 350 | 350 | 350 | 526 | 408 | 408 | 233 |
| 140 | 129 | 180 | 147 | 165 | 126 | 155 | 126 | 138 | 115 | 148 | 128 |
| 86 | 80 | 107 | 96 | 101 | 70 | 85 | 78 | 76 | 64 | 84 | 74 |
| 173 | 163 | 200 | 184 | 210 | 180 | 191 | 168 | 136 | 134 | 159 | 147 |
| 99 | 86 | 95 | 100 | 106 | 122 | 94 | 106 | 75 | 53 | 77 | 82 |
| 189 | 177 | 272 | 214 | 238 | 219 | 262 | 211 | 239 | 222 | 268 | 157 |
| 61 | 85 | 199 | 91 | 128 | 110 | 166 | 118 | 153 | 102 | 107 | 79 |
| 700 | 700 | 1,400 | 700 | 875 | 1,400 | 875 | 875 | 700 | 525 | 525 | 350 |
| 1,185 | 1,488 | 3,500 | 1,225 | 1,575 | 1,575 | 2,538 | 1,925 | 1,750 | 1,225 | 1,225 | 700 |
| 149 | 138 | 195 | 160 | 180 | 151 | 170 | 142 | 149 | 125 | 142 | 120 |
| 86 | 77 | 117 | 89 | 105 | 98 | 95 | 83 | 90 | 56 | 61 | 58 |
| 44 | 41 | 67 | 44 | 44 | 41 | 44 | 28 | 44 | 44 | 54 | 11 |
| 88 | 88 | 124 | 88 | 88 | 88 | 88 | 88 | 88 | 88 | 88 | 88 |
| 175 | 131 | 175 | 175 | 175 | 175 | 175 | 131 | 131 | 131 | 175 | 131 |
| 175 | 175 | 248 | 175 | 175 | 175 | 175 | 175 | 175 | 175 | 175 | 175 |
| 263 | 262 | 350 | 350 | 350 | 263 | 350 | 263 | 284 | 175 | 175 | 175 |
| 350 | 263 | 350 | 350 | 350 | 350 | 350 | 350 | 350 | 175 | 175 | 175 |
| 350 | 350 | 700 | 350 | 526 | 350 | 438 | 350 | 350 | 263 | 350 | 263 |
| 700 | 525 | 700 | 525 | 700 | 1,400 | 525 | 525 | 438 | 350 | 350 | 263 |
| 4.9 | .8 | 2.2 | 4.2 | 5.3 | 14.5 | 13.0 | 8.8 | 3.1 | 2.0 | 2.8 | 3.6 |
| .9 | .1 | .5 | 1.1 | .5 | 1.0 | 1.0 | .5 | 1.1 | .6 | 1.8 | 2.0 |

TABLE 5.44.--COOKED CEREALS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 17.4 | 16.3 | 33.1 | 26.0 | 21.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 10.3 | 5.8 | 16.2 | 15.3 | 13.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 4.3 | 4.9 | 11.1 | 7.2 | 6.3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.7 | 5.5 | 5.8 | 3.6 | 1.9 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 1.6 | 2.1 | 1.7 | 1.6 | 1.5 |
| STANDARD DEVIATION.....DO..... | .8 | 1.1 | .8 | .8 | .7 |
| MAXIMUM.....DO..... | 6.0 | 6.0 | 5.0 | 4.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 107 | 82 | 85 | 92 | 98 |
| STANDARD DEVIATION.....GM..... | 81 | 74 | 64 | 68 | 66 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 20 | 5 | 10 | 13 | 20 |
| 25TH.....GM..... | 50 | 26 | 40 | 41 | 41 |
| 50TH.....GM..... | 82 | 62 | 80 | 80 | 82 |
| 75TH.....GM..... | 160 | 109 | 120 | 123 | 140 |
| 90TH.....GM..... | 240 | 210 | 163 | 184 | 172 |
| 95TH.....GM..... | 245 | 245 | 220 | 243 | 240 |
| 99TH.....GM..... | 388 | 320 | 280 | 320 | 320 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 202 | 98 | 142 | 167 | 202 |
| STANDARD DEVIATION.....GM..... | 109 | 75 | 100 | 95 | 114 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 208 | 112 | 155 | 192 | 198 |
| STANDARD DEVIATION.....GM..... | 100 | 71 | 82 | 91 | 84 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 208 | 142 | 157 | 178 | 196 |
| STANDARD DEVIATION.....GM..... | 108 | 86 | 71 | 82 | 91 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,200 | 480 | 660 | 735 | 960 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,640 | 960 | 980 | 980 | 1,470 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 203 | 116 | 149 | 177 | 198 |
| STANDARD DEVIATION.....GM..... | 110 | 82 | 87 | 97 | 104 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 31 | 14 | 28 | 31 | 38 |
| 25TH.....GM..... | 123 | 60 | 81 | 120 | 123 |
| 50TH.....GM..... | 240 | 105 | 123 | 180 | 240 |
| 75TH.....GM..... | 245 | 123 | 240 | 245 | 245 |
| 90TH.....GM..... | 360 | 245 | 245 | 245 | 263 |
| 95TH.....GM..... | 480 | 263 | 245 | 263 | 368 |
| 99TH.....GM..... | 490 | 360 | 480 | 480 | 490 |
| MAXIMUM.....GM..... | 960 | 462 | 540 | 735 | 960 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 45.4 | .7 | 3.7 | 3.7 | 3.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | .4 | .6 | .5 | .6 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 19.5 | 17.3 | 14.3 | 12.1 | 10.5 | 12.2 | 16.0 | 15.0 | 26.4 | 24.7 | 33.1 | 33.6 |
| 12.2 | 11.9 | 9.8 | 9.2 | 6.8 | 9.0 | 8.4 | 9.5 | 13.4 | 13.1 | 12.3 | 15.8 |
| 5.1 | 4.0 | 2.6 | 1.7 | 2.7 | 2.5 | 4.2 | 3.1 | 5.3 | 6.8 | 10.3 | 8.1 |
| 2.2 | 1.5 | 1.9 | 1.1 | 1.1 | .8 | 3.3 | 2.4 | 7.7 | 4.7 | 10.5 | 9.7 |
| 1.5 | 1.4 | 1.5 | 1.3 | 1.5 | 1.3 | 1.7 | 1.5 | 1.8 | 1.7 | 2.0 | 1.8 |
| .7 | .6 | .7 | .6 | .7 | .6 | .8 | .8 | .9 | .8 | .8 | .9 |
| 3.0 | 3.0 | 4.0 | 3.0 | 4.0 | 3.0 | 4.0 | 4.0 | 4.0 | 4.0 | 3.0 | 3.0 |
| 111 | 99 | 127 | 102 | 118 | 88 | 130 | 97 | 135 | 108 | 140 | 120 |
| 78 | 70 | 107 | 67 | 91 | 63 | 93 | 70 | 92 | 84 | 102 | 88 |
| 20 | 30 | 31 | 40 | 20 | 17 | 21 | 15 | 40 | 21 | 28 | 20 |
| 60 | 54 | 80 | 80 | 73 | 41 | 80 | 41 | 80 | 41 | 80 | 54 |
| 82 | 82 | 82 | 82 | 82 | 80 | 88 | 80 | 99 | 82 | 120 | 82 |
| 160 | 123 | 160 | 123 | 162 | 88 | 163 | 123 | 163 | 160 | 200 | 163 |
| 220 | 163 | 283 | 163 | 240 | 163 | 245 | 179 | 263 | 240 | 243 | 242 |
| 280 | 240 | 351 | 245 | 286 | 245 | 323 | 245 | 327 | 245 | 286 | 245 |
| 360 | 368 | 490 | 360 | 490 | 325 | 480 | 327 | 480 | 480 | 551 | 435 |
| 223 | 217 | 246 | 238 | 232 | 190 | 230 | 188 | 222 | 184 | 220 | 183 |
| 125 | 99 | 111 | 114 | 126 | 100 | 128 | 93 | 91 | 97 | 93 | 79 |
| 244 | 211 | 247 | 204 | 237 | 217 | 237 | 195 | 231 | 189 | 234 | 183 |
| 121 | 86 | 129 | 74 | 101 | 110 | 102 | 89 | 76 | 85 | 93 | 98 |
| 193 | 206 | 308 | 233 | 277 | 188 | 228 | 191 | 226 | 213 | 201 | 215 |
| 98 | 133 | 152 | 100 | 139 | 77 | 116 | 86 | 103 | 106 | 136 | 87 |
| 960 | 960 | 1,200 | 735 | 735 | 960 | 789 | 526 | 490 | 608 | 551 | 490 |
| 1,440 | 1,920 | 2,640 | 1,470 | 1,960 | 1,103 | 2,160 | 1,440 | 1,470 | 1,470 | 1,654 | 1,450 |
| 223 | 212 | 259 | 229 | 242 | 197 | 230 | 188 | 224 | 192 | 213 | 198 |
| 126 | 107 | 132 | 106 | 126 | 104 | 122 | 94 | 102 | 100 | 119 | 95 |
| 35 | 43 | 61 | 110 | 61 | 43 | 41 | 31 | 60 | 46 | 28 | 29 |
| 123 | 123 | 184 | 123 | 161 | 123 | 123 | 123 | 123 | 123 | 123 | 123 |
| 240 | 240 | 245 | 240 | 245 | 240 | 240 | 220 | 240 | 184 | 240 | 240 |
| 245 | 245 | 306 | 245 | 245 | 245 | 245 | 245 | 245 | 245 | 245 | 245 |
| 368 | 360 | 490 | 368 | 480 | 263 | 368 | 245 | 360 | 245 | 360 | 245 |
| 490 | 420 | 490 | 480 | 490 | 368 | 490 | 360 | 480 | 368 | 480 | 360 |
| 612 | 490 | 527 | 490 | 700 | 490 | 601 | 490 | 490 | 490 | 551 | 490 |
| 960 | 960 | 960 | 735 | 735 | 960 | 789 | 526 | 490 | 490 | 551 | 490 |
| 2.6 | 4.9 | 1.8 | 2.0 | 2.6 | 2.8 | 8.2 | 5.7 | 1.8 | 0 | 0 | 1.3 |
| .4 | .9 | .6 | .8 | .4 | .3 | .6 | .4 | .3 | 0 | 0 | .2 |

TABLE 5.45.--GRITS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.7 | 2.4 | 9.2 | 8.4 | 5.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.4 | 1.3 | 3.9 | 5.3 | 3.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.4 | .6 | 3.3 | 2.0 | 1.7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .9 | .5 | 2.0 | 1.1 | .4 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.6 | 1.7 | 1.9 | 1.5 | 1.5 |
| STANDARD DEVIATION.....DO..... | .8 | .9 | .9 | .8 | .6 |
| MAXIMUM.....DO..... | 5.0 | 4.0 | 5.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 97 | 67 | 77 | 78 | 89 |
| STANDARD DEVIATION.....GM..... | 78 | 58 | 63 | 62 | 69 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 15 | | 8 | 10 | 20 |
| 25TH.....GM..... | 41 | | 28 | 41 | 41 |
| 50TH.....GM..... | 82 | | 55 | 61 | 82 |
| 75TH.....GM..... | 123 | | 114 | 82 | 123 |
| 90TH.....GM..... | 204 | | 163 | 163 | 163 |
| 95TH.....GM..... | 245 | | 218 | 245 | 204 |
| 99TH.....GM..... | 388 | | 245 | 305 | 490 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 173 | 100 | 136 | 141 | 159 |
| STANDARD DEVIATION.....GM..... | 93 | 82 | 126 | 83 | 71 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 190 | 119 | 115 | 163 | 181 |
| STANDARD DEVIATION.....GM..... | 94 | 36 | 69 | 73 | 72 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 200 | 143 | 137 | 164 | 225 |
| STANDARD DEVIATION.....GM..... | 105 | 70 | 76 | 84 | 111 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 980 | 245 | 660 | 490 | 490 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,960 | 735 | 980 | 980 | 1,470 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 184 | 117 | 123 | 152 | 179 |
| STANDARD DEVIATION.....GM..... | 103 | 76 | 84 | 87 | 87 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 31 | | 15 | 31 | 55 |
| 25TH.....GM..... | 123 | | 61 | 110 | 123 |
| 50TH.....GM..... | 184 | | 123 | 123 | 184 |
| 75TH.....GM..... | 245 | | 163 | 245 | 245 |
| 90TH.....GM..... | 245 | | 245 | 245 | 245 |
| 95TH.....GM..... | 368 | | 245 | 245 | 245 |
| 99TH.....GM..... | 490 | | 368 | 430 | 490 |
| MAXIMUM.....GM..... | 735 | 245 | 490 | 490 | 490 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 15.4 | 0 | 1.5 | .8 | 1.4 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .5 | 0 | .8 | .4 | 1.0 |

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 6.8 | 4.1 | 6.2 | 4.9 | 4.8 | 4.6 | 6.1 | 5.0 | 7.0 | 7.0 | 7.7 | 7.7 |
| 4.5 | 2.6 | 3.7 | 3.1 | 3.1 | 3.3 | 3.1 | 3.4 | 3.5 | 3.7 | 2.5 | 4.4 |
| 1.7 | 1.0 | 1.4 | 1.1 | 1.1 | .9 | 1.8 | .9 | 1.3 | 1.8 | 3.5 | 1.6 |
| .6 | .5 | 1.1 | .8 | .6 | .3 | 1.2 | .8 | 2.2 | 1.5 | 1.8 | 1.6 |
| 1.4 | 1.5 | 1.6 | 1.5 | 1.5 | 1.4 | 1.7 | 1.5 | 1.8 | 1.7 | 1.9 | 1.7 |
| .7 | .7 | .8 | .8 | .7 | .6 | .8 | .8 | .9 | .9 | .7 | .8 |
| 3.0 | 3.0 | 4.0 | 3.0 | 4.0 | 3.0 | 4.0 | 4.0 | 3.0 | 4.0 | 3.0 | 3.0 |
| 88 | 94 | 121 | 95 | 114 | 86 | 116 | 88 | 117 | 95 | 105 | 96 |
| 66 | 67 | 102 | 62 | 103 | 68 | 85 | 67 | 89 | 79 | 58 | 76 |
| 18 | 27 | 20 | 37 | 20 | 15 | 19 | 10 | 20 | 15 | | 10 |
| 41 | 41 | 55 | 41 | 41 | 41 | 61 | 41 | 41 | 41 | | 41 |
| 82 | 82 | 82 | 82 | 82 | 82 | 82 | 82 | 82 | 73 | | 82 |
| 95 | 136 | 163 | 123 | 163 | 82 | 163 | 110 | 163 | 123 | | 123 |
| 163 | 163 | 245 | 204 | 245 | 163 | 245 | 184 | 245 | 204 | | 245 |
| 204 | 204 | 327 | 245 | 327 | 245 | 286 | 245 | 327 | 245 | | 245 |
| 408 | 368 | 490 | 245 | 490 | 368 | 408 | 286 | 408 | 327 | | 368 |
| 173 | 175 | 196 | 192 | 198 | 169 | 195 | 169 | 174 | 138 | 209 | 181 |
| 80 | 87 | 90 | 92 | 98 | 86 | 102 | 91 | 91 | 75 | 47 | 94 |
| 206 | 202 | 225 | 176 | 218 | 233 | 203 | 169 | 201 | 177 | 226 | 134 |
| 122 | 71 | 94 | 66 | 96 | 111 | 94 | 78 | 69 | 89 | 79 | 78 |
| 173 | 198 | 269 | 187 | 313 | 188 | 217 | 194 | 200 | 186 | 68 | 200 |
| 58 | 84 | 132 | 70 | 149 | 82 | 101 | 74 | 101 | 93 | 40 | 90 |
| 613 | 490 | 980 | 490 | 735 | 613 | 735 | 490 | 490 | 490 | 490 | 490 |
| 1,225 | 1,103 | 1,470 | 735 | 1,960 | 1,103 | 1,470 | 980 | 1,225 | 1,470 | 858 | 1,103 |
| 184 | 189 | 227 | 186 | 230 | 191 | 204 | 175 | 193 | 164 | 166 | 174 |
| 105 | 86 | 117 | 81 | 124 | 101 | 106 | 91 | 103 | 92 | 101 | 97 |
| 26 | 46 | 61 | 55 | 61 | 61 | 31 | 31 | 41 | 28 | | 28 |
| 123 | 123 | 123 | 123 | 123 | 123 | 123 | 123 | 123 | 123 | | 123 |
| 184 | 245 | 245 | 220 | 245 | 220 | 245 | 184 | 184 | 123 | | 184 |
| 245 | 245 | 245 | 245 | 245 | 245 | 245 | 245 | 245 | 245 | | 245 |
| 245 | 245 | 368 | 245 | 490 | 245 | 245 | 245 | 368 | 245 | | 245 |
| 368 | 368 | 490 | 245 | 490 | 368 | 490 | 245 | 368 | 245 | | 245 |
| 612 | 490 | 490 | 490 | 490 | 490 | 490 | 490 | 490 | 490 | | 490 |
| 613 | 490 | 490 | 490 | 735 | 613 | 490 | 490 | 490 | 490 | 490 | 490 |
| 1.6 | 1.4 | .8 | 2.0 | 0 | 0 | 2.1 | 2.0 | 1.8 | 0 | 0 | 0 |
| .8 | 1.1 | .6 | 1.8 | 0 | 0 | .4 | .4 | 1.3 | 0 | 0 | 0 |

TABLE 5.46.--OATMEAL
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 8.3 | 9.0 | 18.1 | 13.1 | 10.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.7 | 3.4 | 11.5 | 9.0 | 7.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.7 | 3.3 | 4.8 | 3.0 | 2.8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.0 | 2.3 | 1.8 | 1.1 | .6 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.4 | 2.0 | 1.5 | 1.4 | 1.4 |
| STANDARD DEVIATION.....DO..... | .7 | .9 | .7 | .7 | .6 |
| MAXIMUM.....DO..... | 4.0 | 4.0 | 3.0 | 4.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 102 | 79 | 79 | 87 | 95 |
| STANDARD DEVIATION.....GM..... | 76 | 82 | 59 | 62 | 64 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 20 | | 13 | 20 | 26 |
| 25TH.....GM..... | 54 | | 40 | 40 | 40 |
| 50TH.....GM..... | 80 | | 80 | 80 | 80 |
| 75TH.....GM..... | 120 | | 100 | 97 | 120 |
| 90TH.....GM..... | 200 | | 160 | 160 | 172 |
| 95TH.....GM..... | 240 | | 195 | 240 | 240 |
| 99TH.....GM..... | 360 | | 295 | 320 | 335 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 216 | 82 | 154 | 184 | 212 |
| STANDARD DEVIATION.....GM..... | 112 | 51 | 89 | 97 | 118 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 209 | 106 | 174 | 192 | 206 |
| STANDARD DEVIATION.....GM..... | 99 | 70 | 91 | 101 | 92 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 215 | 166 | 154 | 184 | 196 |
| STANDARD DEVIATION.....GM..... | 117 | 106 | 76 | 77 | 112 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,200 | 480 | 540 | 480 | 960 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,640 | 960 | 960 | 960 | 1,080 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 213 | 121 | 161 | 185 | 206 |
| STANDARD DEVIATION.....GM..... | 113 | 93 | 93 | 96 | 115 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 43 | | 43 | 38 | 43 |
| 25TH.....GM..... | 120 | | 120 | 120 | 120 |
| 50TH.....GM..... | 240 | | 120 | 180 | 240 |
| 75TH.....GM..... | 240 | | 240 | 240 | 240 |
| 90TH.....GM..... | 360 | | 240 | 263 | 360 |
| 95TH.....GM..... | 480 | | 360 | 360 | 462 |
| 99TH.....GM..... | 480 | | 480 | 480 | 526 |
| MAXIMUM.....GM..... | 960 | 462 | 540 | 480 | 960 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 19.1 | .7 | 0 | 2.9 | 2.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | .8 | 0 | .9 | .8 |

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY, 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 8.7 | 8.7 | 6.4 | 4.6 | 3.8 | 5.3 | 7.3 | 7.4 | 15.0 | 11.4 | 17.3 | 19.6 |
| 6.2 | 6.8 | 5.3 | 4.1 | 3.1 | 4.5 | 4.7 | 5.0 | 8.5 | 7.4 | 6.6 | 10.3 |
| 1.7 | 1.3 | .8 | .3 | .4 | .7 | 1.6 | 1.5 | 3.2 | 2.6 | 5.3 | 4.6 |
| .8 | .7 | .3 | .2 | .3 | .2 | 1.1 | .8 | 3.3 | 1.4 | 5.3 | 4.7 |
| 1.4 | 1.3 | 1.2 | 1.2 | 1.3 | 1.2 | 1.5 | 1.4 | 1.7 | 1.5 | 2.0 | 1.7 |
| .6 | .6 | .5 | .5 | .6 | .5 | .7 | .7 | .8 | .7 | .9 | .8 |
| 3.0 | 3.0 | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| 115 | 94 | 115 | 102 | 106 | 79 | 126 | 93 | 129 | 104 | 149 | 115 |
| 80 | 75 | 97 | 65 | 62 | 55 | 93 | 64 | 82 | 85 | 88 | 79 |
| 19 | 30 | 40 | 40 | 40 | 19 | 20 | 19 | 40 | 26 | 40 | 28 |
| 80 | 53 | 80 | 80 | 80 | 40 | 80 | 40 | 80 | 54 | 80 | 54 |
| 80 | 80 | 80 | 80 | 80 | 80 | 80 | 80 | 88 | 80 | 140 | 80 |
| 160 | 100 | 120 | 120 | 120 | 80 | 160 | 120 | 160 | 120 | 240 | 160 |
| 220 | 160 | 165 | 160 | 200 | 160 | 240 | 160 | 240 | 200 | 240 | 240 |
| 263 | 200 | 240 | 240 | 240 | 170 | 280 | 200 | 280 | 240 | 240 | 240 |
| 408 | 400 | 480 | 360 | 320 | 320 | 480 | 360 | 480 | 480 | 480 | 320 |
| 255 | 223 | 272 | 253 | 260 | 198 | 253 | 196 | 252 | 200 | 275 | 195 |
| 135 | 96 | 109 | 102 | 113 | 105 | 138 | 96 | 82 | 85 | 116 | 86 |
| 245 | 199 | 256 | 275 | 319 | 218 | 238 | 204 | 228 | 185 | 213 | 200 |
| 123 | 103 | 148 | 72 | 102 | 131 | 86 | 69 | 87 | 77 | 84 | 96 |
| 239 | 228 | 400 | 302 | 182 | 154 | 256 | 185 | 221 | 260 | 226 | 207 |
| 112 | 172 | 227 | 60 | 73 | 78 | 138 | 103 | 105 | 126 | 101 | 78 |
| 960 | 960 | 1,200 | 480 | 720 | 960 | 789 | 526 | 480 | 480 | 480 | 480 |
| 1,440 | 1,920 | 2,640 | 1,080 | 960 | 1,080 | 2,160 | 1,440 | 1,440 | 1,440 | 1,440 | 1,306 |
| 249 | 218 | 282 | 262 | 254 | 198 | 250 | 195 | 233 | 211 | 228 | 202 |
| 131 | 119 | 137 | 98 | 120 | 113 | 129 | 95 | 100 | 103 | 101 | 89 |
| 35 | 35 | 88 | 120 | 60 | 35 | 60 | 43 | 60 | 60 | 43 | 60 |
| 180 | 120 | 240 | 240 | 180 | 120 | 180 | 120 | 180 | 120 | 180 | 120 |
| 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 |
| 263 | 240 | 360 | 263 | 277 | 240 | 263 | 240 | 240 | 240 | 240 | 240 |
| 480 | 360 | 480 | 480 | 480 | 263 | 480 | 263 | 360 | 360 | 360 | 240 |
| 480 | 480 | 526 | 480 | 480 | 360 | 480 | 360 | 480 | 480 | 480 | 346 |
| 527 | 480 | 721 | 480 | 480 | 480 | 721 | 480 | 480 | 480 | 480 | 480 |
| 960 | 960 | 960 | 480 | 720 | 960 | 789 | 526 | 480 | 480 | 480 | 480 |
| 1.0 | 1.5 | 1.0 | 0 | 0 | 2.8 | 6.1 | 1.0 | 0 | 0 | 0 | 0 |
| .4 | .6 | .9 | 0 | 0 | .8 | 1.1 | .1 | 0 | 0 | 0 | 0 |

TABLE 5.47.--COOKED WHEAT CEREALS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.2 | 6.4 | 7.9 | 6.2 | 6.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.0 | 3.0 | 5.1 | 4.1 | 4.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .8 | 2.2 | 1.8 | 1.6 | 1.1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .4 | 1.1 | .9 | .5 | .3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.4 | 1.8 | 1.5 | 1.4 | 1.3 |
| STANDARD DEVIATION.....DO..... | .7 | .8 | .7 | .6 | .6 |
| MAXIMUM.....DO..... | 5.0 | 5.0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 99 | 65 | 78 | 94 | 88 |
| STANDARD DEVIATION.....GM..... | 71 | 55 | 59 | 70 | 51 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 20 | | 10 | 20 | 20 |
| 25TH.....GM..... | 54 | | 41 | 41 | 41 |
| 50TH.....GM..... | 82 | | 61 | 82 | 82 |
| 75TH.....GM..... | 123 | | 96 | 116 | 102 |
| 90TH.....GM..... | 163 | | 163 | 184 | 163 |
| 95TH.....GM..... | 245 | | 204 | 245 | 194 |
| 99TH.....GM..... | 327 | | 245 | 327 | 255 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 221 | 102 | 140 | 183 | 211 |
| STANDARD DEVIATION.....GM..... | 112 | 65 | 75 | 89 | 101 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 202 | 101 | 163 | 196 | 214 |
| STANDARD DEVIATION.....GM..... | 98 | 68 | 55 | 104 | 82 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 217 | 143 | 201 | 245 | 159 |
| STANDARD DEVIATION.....GM..... | 113 | 55 | 45 | 50 | 63 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 735 | 245 | 306 | 735 | 490 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,654 | 735 | 735 | 980 | 766 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 214 | 110 | 160 | 198 | 204 |
| STANDARD DEVIATION.....GM..... | 112 | 68 | 71 | 103 | 97 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 31 | | 31 | 57 | 38 |
| 25TH.....GM..... | 123 | | 123 | 123 | 123 |
| 50TH.....GM..... | 245 | | 163 | 245 | 245 |
| 75TH.....GM..... | 245 | | 245 | 245 | 245 |
| 90TH.....GM..... | 368 | | 245 | 245 | 245 |
| 95TH.....GM..... | 490 | | 245 | 368 | 368 |
| 99TH.....GM..... | 490 | | 245 | 490 | 490 |
| MAXIMUM.....GM..... | 735 | 245 | 306 | 735 | 490 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 10.9 | 0 | 2.2 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .5 | 0 | 1.8 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 4.9 | 5.3 | 2.4 | 2.7 | 2.5 | 2.5 | 3.3 | 3.2 | 5.8 | 8.3 | 9.7 | 9.5 |
| 3.4 | 4.4 | 1.9 | 2.4 | 2.0 | 1.7 | 2.2 | 2.4 | 4.2 | 5.6 | 6.0 | 5.5 |
| 1.3 | .8 | .3 | .3 | .4 | .6 | .7 | .6 | .7 | 1.6 | 1.1 | 2.6 |
| .3 | .1 | .2 | .1 | .2 | .2 | .4 | .3 | .9 | 1.0 | 2.5 | 1.4 |
| 1.4 | 1.2 | 1.3 | 1.1 | 1.3 | 1.4 | 1.5 | 1.3 | 1.4 | 1.4 | 1.6 | 1.6 |
| .6 | .4 | .6 | .4 | .6 | .6 | .8 | .6 | .7 | .7 | .9 | .7 |
| 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 4.0 | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| 106 | 89 | 131 | 107 | 108 | 98 | 120 | 87 | 119 | 91 | 122 | 110 |
| 69 | 49 | 117 | 72 | 87 | 57 | 70 | 56 | 87 | 64 | 124 | 73 |
| 31 | 30 | | | 9 | 15 | 27 | 19 | 41 | 20 | 15 | 25 |
| 49 | 61 | | | 82 | 71 | 82 | 41 | 82 | 41 | 41 | 61 |
| 82 | 82 | | | 82 | 82 | 102 | 82 | 82 | 82 | 82 | 82 |
| 163 | 115 | | | 123 | 123 | 163 | 82 | 163 | 123 | 163 | 163 |
| 190 | 163 | | | 184 | 175 | 204 | 163 | 163 | 163 | 245 | 245 |
| 245 | 204 | | | 306 | 245 | 245 | 163 | 368 | 245 | 551 | 245 |
| 327 | 245 | | | 490 | 245 | 327 | 327 | 490 | 286 | 552 | 327 |
| 244 | 233 | 244 | 278 | 231 | 217 | 293 | 213 | 243 | 188 | 205 | 227 |
| 126 | 99 | 100 | 127 | 104 | 96 | 140 | 96 | 92 | 102 | 107 | 114 |
| 239 | 216 | 365 | 248 | 245 | 209 | 208 | 176 | 230 | 190 | 254 | 186 |
| 126 | 102 | 122 | 84 | 112 | 77 | 119 | 68 | 40 | 72 | 108 | 78 |
| 174 | 180 | 455 | 490 | 338 | 219 | 207 | 174 | 265 | 186 | 232 | 224 |
| 86 | 61 | 55 | 0 | 116 | 30 | 78 | 113 | 122 | 85 | 186 | 88 |
| 613 | 490 | 490 | 735 | 571 | 490 | 613 | 490 | 490 | 490 | 551 | 490 |
| 980 | 735 | 1,470 | 1,470 | 1,470 | 735 | 980 | 1,225 | 1,470 | 858 | 1,654 | 980 |
| 234 | 227 | 306 | 284 | 252 | 214 | 243 | 195 | 248 | 188 | 225 | 212 |
| 126 | 102 | 130 | 129 | 123 | 87 | 129 | 96 | 101 | 94 | 155 | 102 |
| 57 | 56 | | | 31 | 61 | 29 | 29 | 123 | 46 | 28 | 31 |
| 123 | 173 | | | 216 | 173 | 173 | 123 | 164 | 123 | 123 | 123 |
| 245 | 245 | | | 245 | 245 | 245 | 245 | 245 | 245 | 245 | 245 |
| 245 | 245 | | | 306 | 245 | 245 | 245 | 245 | 245 | 245 | 245 |
| 490 | 368 | | | 490 | 306 | 490 | 245 | 368 | 245 | 490 | 306 |
| 490 | 368 | | | 490 | 368 | 490 | 368 | 490 | 368 | 551 | 490 |
| 612 | 490 | | | 571 | 490 | 612 | 490 | 490 | 490 | 551 | 490 |
| 613 | 490 | 490 | 735 | 571 | 490 | 613 | 490 | 490 | 490 | 551 | 490 |
| 0 | 2.0 | 0 | 0 | 2.6 | 0 | 0 | 2.8 | 0 | 0 | 0 | 1.3 |
| 0 | 1.5 | 0 | 0 | 2.1 | 0 | 0 | .9 | 0 | 0 | 0 | .9 |

TABLE 5.48.--READY-TO-EAT CEREALS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 43.4 | 68.7 | 68.0 | 75.8 | 76.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 17.4 | 7.5 | 23.5 | 25.5 | 24.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 13.5 | 10.4 | 23.8 | 27.0 | 26.7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 12.4 | 50.7 | 20.8 | 23.3 | 25.9 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 2.0 | 3.8 | 2.1 | 2.1 | 2.1 |
| STANDARD DEVIATION.....DO..... | 1.0 | 2.1 | 1.0 | 1.0 | .9 |
| MAXIMUM.....DO..... | 15.0 | 15.0 | 9.0 | 10.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 23 | 17 | 16 | 20 | 23 |
| STANDARD DEVIATION.....GM..... | 20 | 15 | 12 | 14 | 16 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 4 | 2 | 3 | 5 | 5 |
| 25TH.....GM..... | 10 | 7 | 8 | 9 | 10 |
| 50TH.....GM..... | 18 | 13 | 13 | 17 | 19 |
| 75TH.....GM..... | 30 | 20 | 21 | 27 | 30 |
| 90TH.....GM..... | 46 | 40 | 31 | 38 | 43 |
| 95TH.....GM..... | 58 | 43 | 38 | 47 | 54 |
| 99TH.....GM..... | 100 | 66 | 55 | 67 | 78 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 37 | 18 | 22 | 30 | 34 |
| STANDARD DEVIATION.....GM..... | 25 | 11 | 14 | 17 | 18 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 36 | 18 | 24 | 30 | 32 |
| STANDARD DEVIATION.....GM..... | 22 | 12 | 13 | 16 | 18 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 38 | 19 | 25 | 32 | 35 |
| STANDARD DEVIATION.....GM..... | 25 | 16 | 13 | 16 | 18 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 495 | 158 | 150 | 200 | 224 |
| MAXIMUM OVER 3 DAYS.....GM..... | 935 | 342 | 261 | 400 | 383 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 36 | 13 | 23 | 29 | 33 |
| STANDARD DEVIATION.....GM..... | 25 | 11 | 14 | 17 | 19 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 8 | 2 | 6 | 10 | 9 |
| 25TH.....GM..... | 22 | 5 | 13 | 19 | 25 |
| 50TH.....GM..... | 29 | 10 | 21 | 27 | 28 |
| 75TH.....GM..... | 45 | 20 | 28 | 35 | 40 |
| 90TH.....GM..... | 60 | 28 | 40 | 50 | 56 |
| 95TH.....GM..... | 84 | 40 | 50 | 60 | 65 |
| 99TH.....GM..... | 120 | 46 | 70 | 90 | 100 |
| MAXIMUM.....GM..... | 440 | 80 | 110 | 170 | 224 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 130.9 | 21.5 | 8.7 | 4.8 | 7.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 1.7 | .6 | .2 | .3 |

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 69.8 | 64.0 | 50.4 | 43.7 | 28.8 | 30.5 | 30.3 | 30.4 | 47.5 | 46.6 | 48.2 | 47.4 |
| 24.0 | 25.5 | 19.5 | 22.7 | 14.1 | 16.5 | 12.8 | 14.6 | 14.4 | 16.5 | 15.3 | 13.9 |
| 24.3 | 21.4 | 15.0 | 13.0 | 8.5 | 9.5 | 8.4 | 8.8 | 13.4 | 14.8 | 16.0 | 13.5 |
| 21.6 | 17.2 | 15.9 | 8.0 | 6.2 | 4.4 | 9.1 | 7.0 | 19.7 | 15.4 | 17.0 | 20.0 |
| 2.1 | 1.9 | 2.0 | 1.7 | 1.8 | 1.6 | 1.9 | 1.8 | 2.2 | 2.0 | 2.1 | 2.2 |
| 1.0 | .9 | 1.0 | .8 | .9 | .8 | .9 | .9 | 1.0 | .9 | 1.0 | 1.0 |
| 7.0 | 6.0 | 6.0 | 6.0 | 6.0 | 5.0 | 5.0 | 6.0 | 6.0 | 6.0 | 6.0 | 7.0 |
| 28 | 23 | 33 | 21 | 29 | 20 | 27 | 20 | 27 | 22 | 24 | 24 |
| 24 | 17 | 27 | 16 | 27 | 16 | 22 | 16 | 23 | 17 | 18 | 20 |
| 6 | 5 | 7 | 5 | 5 | 4 | 4 | 4 | 7 | 4 | 5 | 4 |
| 13 | 10 | 15 | 10 | 12 | 9 | 11 | 9 | 12 | 10 | 12 | 10 |
| 23 | 19 | 27 | 17 | 20 | 17 | 20 | 17 | 22 | 17 | 20 | 20 |
| 37 | 30 | 45 | 28 | 37 | 25 | 33 | 27 | 35 | 28 | 30 | 30 |
| 53 | 43 | 67 | 41 | 57 | 38 | 54 | 40 | 50 | 43 | 42 | 43 |
| 67 | 51 | 83 | 50 | 80 | 52 | 70 | 50 | 65 | 52 | 57 | 52 |
| 127 | 85 | 129 | 79 | 136 | 81 | 100 | 84 | 133 | 100 | 110 | 83 |
| 41 | 37 | 51 | 38 | 48 | 36 | 40 | 34 | 36 | 32 | 33 | 31 |
| 28 | 24 | 43 | 22 | 33 | 24 | 26 | 22 | 19 | 20 | 18 | 18 |
| 42 | 36 | 47 | 37 | 44 | 37 | 43 | 35 | 35 | 33 | 34 | 32 |
| 26 | 19 | 27 | 20 | 29 | 22 | 24 | 20 | 20 | 20 | 18 | 18 |
| 46 | 37 | 55 | 38 | 58 | 40 | 44 | 35 | 41 | 34 | 37 | 34 |
| 29 | 21 | 30 | 20 | 37 | 20 | 27 | 20 | 28 | 20 | 22 | 24 |
| 440 | 275 | 495 | 220 | 320 | 220 | 226 | 220 | 240 | 180 | 168 | 300 |
| 935 | 500 | 688 | 385 | 581 | 352 | 678 | 380 | 530 | 398 | 371 | 758 |
| 41 | 36 | 49 | 37 | 49 | 37 | 42 | 34 | 37 | 32 | 34 | 32 |
| 28 | 21 | 31 | 22 | 35 | 23 | 28 | 22 | 25 | 21 | 22 | 20 |
| 13 | 11 | 15 | 13 | 10 | 10 | 10 | 10 | 10 | 8 | 9 | 8 |
| 25 | 25 | 28 | 25 | 25 | 25 | 25 | 21 | 25 | 20 | 20 | 21 |
| 35 | 30 | 43 | 30 | 42 | 30 | 37 | 28 | 30 | 28 | 28 | 28 |
| 50 | 45 | 60 | 45 | 60 | 50 | 50 | 45 | 50 | 40 | 40 | 38 |
| 74 | 57 | 86 | 60 | 90 | 60 | 75 | 60 | 68 | 55 | 57 | 54 |
| 90 | 75 | 110 | 80 | 113 | 83 | 100 | 75 | 84 | 60 | 70 | 64 |
| 128 | 111 | 167 | 110 | 172 | 112 | 131 | 112 | 120 | 120 | 120 | 113 |
| 440 | 226 | 330 | 220 | 280 | 220 | 226 | 220 | 224 | 180 | 168 | 150 |
| 7.2 | 3.0 | 8.2 | 0 | 24.9 | 10.2 | 12.0 | 11.1 | 3.4 | 3.3 | 1.5 | 3.7 |
| .2 | .1 | .6 | 0 | 1.2 | .4 | .4 | .3 | .3 | .2 | .3 | .4 |

TABLE 5.49.--READY-TO-EAT CEREALS, 15-21 GM PER CUP, ALL SUGAR LEVELS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.6 | .3 | 3.4 | 4.7 | 4.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.2 | .3 | 2.5 | 3.6 | 3.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .9 | 0 | .7 | .8 | 1.1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .5 | 0 | .2 | .3 | .5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.6 | 1.0 | 1.3 | 1.4 | 1.4 |
| STANDARD DEVIATION.....CO..... | .8 | .0 | .6 | .7 | .7 |
| MAXIMUM.....DO..... | 5.0 | 1.0 | 3.0 | 4.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 11 | 2 | 8 | 9 | 9 |
| STANDARD DEVIATION.....GM..... | 7 | 0 | 4 | 6 | 6 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 3 | | | 2 | 2 |
| 25TH.....GM..... | 5 | | | 5 | 5 |
| 50TH.....GM..... | 8 | | | 7 | 7 |
| 75TH.....GM..... | 13 | | | 11 | 13 |
| 90TH.....GM..... | 20 | | | 17 | 19 |
| 95TH.....GM..... | 25 | | | 23 | 21 |
| 99TH.....GM..... | 38 | | | 30 | 35 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 21 | 7 | 19 | 20 | 21 |
| STANDARD DEVIATION.....GM..... | 12 | 0 | 10 | 11 | 11 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 20 | 0 | 15 | 19 | 15 |
| STANDARD DEVIATION.....GM..... | 10 | 0 | 6 | 9 | 9 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 21 | 0 | 14 | 21 | 22 |
| STANDARD DEVIATION.....GM..... | 9 | 0 | 6 | 7 | 6 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 84 | 8 | 40 | 57 | 63 |
| MAXIMUM OVER 3 DAYS.....GM..... | 180 | 8 | 60 | 90 | 105 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 20 | 7 | 17 | 19 | 19 |
| STANDARD DEVIATION.....GM..... | 11 | 0 | 9 | 10 | 11 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 5 | | | 4 | 5 |
| 25TH.....GM..... | 15 | | | 11 | 11 |
| 50TH.....GM..... | 20 | | | 20 | 20 |
| 75TH.....GM..... | 21 | | | 23 | 21 |
| 90TH.....GM..... | 40 | | | 32 | 32 |
| 95TH.....GM..... | 42 | | | 42 | 42 |
| 99TH.....GM..... | 56 | | | 43 | 45 |
| MAXIMUM.....GM..... | 80 | 8 | 40 | 43 | 63 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 2.5 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .1 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.5 | 4.7 | 4.2 | 2.8 | 1.4 | 2.0 | 2.7 | 3.2 | 6.9 | 7.8 | 8.5 | 6.5 |
| 3.6 | 3.0 | 2.7 | 1.8 | .8 | 1.5 | 1.5 | 2.0 | 3.3 | 3.6 | 4.6 | 2.6 |
| 1.3 | 1.3 | 1.0 | 1.0 | .5 | .5 | .7 | .7 | 1.8 | 2.3 | 2.0 | 1.6 |
| .7 | .5 | .5 | 0 | .1 | 0 | .6 | .5 | 1.7 | 1.9 | 1.9 | 2.2 |
| 1.5 | 1.5 | 1.5 | 1.4 | 1.4 | 1.3 | 1.7 | 1.5 | 1.9 | 1.8 | 1.7 | 1.9 |
| .7 | .7 | .7 | .5 | .6 | .4 | .8 | .8 | 1.0 | .9 | .8 | .9 |
| 4.0 | 3.0 | 3.0 | 2.0 | 3.0 | 2.0 | 3.0 | 3.0 | 5.0 | 4.0 | 3.0 | 3.0 |
| 13 | 11 | 14 | 11 | 11 | 9 | 12 | 10 | 12 | 11 | 10 | 11 |
| 9 | 7 | 9 | 5 | 9 | 5 | 8 | 7 | 8 | 8 | 9 | 7 |
| 3 | 3 | 4 | | 3 | 2 | 3 | 2 | 3 | 3 | | 3 |
| 7 | 7 | 7 | | 5 | 7 | 7 | 5 | 7 | 5 | | 5 |
| 10 | 10 | 13 | | 9 | 7 | 10 | 7 | 10 | 8 | | 8 |
| 18 | 14 | 20 | | 14 | 13 | 13 | 13 | 15 | 15 | | 16 |
| 25 | 20 | 27 | | 20 | 14 | 20 | 19 | 25 | 20 | | 20 |
| 28 | 20 | 30 | | 28 | 19 | 28 | 20 | 28 | 27 | | 20 |
| 40 | 30 | 47 | | 53 | 28 | 40 | 38 | 37 | 40 | | 40 |
| 27 | 24 | 29 | 29 | 21 | 22 | 20 | 18 | 22 | 16 | 15 | 16 |
| 14 | 12 | 17 | 14 | 11 | 12 | 11 | 8 | 8 | 8 | 8 | 4 |
| 22 | 20 | 27 | 18 | 27 | 21 | 20 | 22 | 19 | 19 | 15 | 19 |
| 12 | 9 | 9 | 8 | 18 | 9 | 11 | 9 | 11 | 7 | 6 | 4 |
| 27 | 23 | 29 | 0 | 21 | 0 | 22 | 18 | 20 | 19 | 23 | 17 |
| 11 | 8 | 10 | 0 | 2 | 0 | 8 | 7 | 8 | 9 | 10 | 8 |
| 84 | 48 | 80 | 63 | 80 | 60 | 80 | 57 | 45 | 57 | 40 | 40 |
| 180 | 126 | 140 | 70 | 160 | 85 | 120 | 120 | 112 | 120 | 120 | 120 |
| 25 | 22 | 28 | 24 | 24 | 21 | 21 | 19 | 19 | 18 | 18 | 17 |
| 12 | 11 | 14 | 12 | 15 | 10 | 10 | 8 | 10 | 9 | 9 | 7 |
| 5 | 5 | 15 | | 8 | 7 | 8 | 8 | 5 | 5 | | 7 |
| 16 | 16 | 20 | | 15 | 15 | 15 | 15 | 15 | 10 | | 11 |
| 21 | 20 | 21 | | 20 | 20 | 20 | 20 | 20 | 20 | | 20 |
| 40 | 30 | 40 | | 28 | 23 | 23 | 20 | 20 | 20 | | 20 |
| 42 | 42 | 42 | | 42 | 40 | 30 | 28 | 36 | 30 | | 28 |
| 42 | 42 | 50 | | 43 | 43 | 40 | 30 | 42 | 40 | | 30 |
| 60 | 43 | 80 | | 80 | 43 | 56 | 56 | 42 | 40 | | 40 |
| 63 | 43 | 80 | 63 | 80 | 43 | 80 | 57 | 42 | 57 | 40 | 40 |
| 0 | 0 | 1.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1.5 | 0 |
| 0 | 0 | 1.2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1.9 | 0 |

TABLE 5.50.--READY-TO-EAT CEREALS, 25-32 GM PER CUP, 30 PCT OR MORE SUGAR
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 6.1 | 1.1 | 18.5 | 23.3 | 22.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.5 | .5 | 10.2 | 13.5 | 10.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.7 | .6 | 4.9 | 5.6 | 7.8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .9 | 0 | 3.5 | 4.2 | 3.9 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.6 | 1.6 | 1.7 | 1.7 | 1.8 |
| STANDARD DEVIATION.....DO..... | .8 | .5 | 1.0 | .9 | .8 |
| MAXIMUM.....DO..... | 6.0 | 2.0 | 6.0 | 6.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 17 | 6 | 13 | 15 | 18 |
| STANDARD DEVIATION.....GM..... | 13 | 4 | 10 | 12 | 12 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 5 | | 3 | 4 | 5 |
| 25TH.....GM..... | 9 | | 5 | 9 | 9 |
| 50TH.....GM..... | 14 | | 9 | 10 | 14 |
| 75TH.....GM..... | 20 | | 18 | 20 | 25 |
| 90TH.....GM..... | 33 | | 21 | 30 | 37 |
| 95TH.....GM..... | 40 | | 28 | 37 | 43 |
| 99TH.....GM..... | 60 | | 60 | 60 | 57 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 32 | 4 | 22 | 26 | 30 |
| STANDARD DEVIATION.....GM..... | 18 | 3 | 12 | 12 | 16 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 33 | 14 | 22 | 30 | 33 |
| STANDARD DEVIATION.....GM..... | 18 | 3 | 9 | 14 | 17 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 33 | 0 | 25 | 31 | 32 |
| STANDARD DEVIATION.....GM..... | 15 | 0 | 15 | 15 | 13 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 168 | 28 | 96 | 150 | 168 |
| MAXIMUM OVER 3 DAYS.....GM..... | 336 | 42 | 244 | 276 | 224 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 31 | 12 | 22 | 28 | 31 |
| STANDARD DEVIATION.....GM..... | 17 | 6 | 13 | 14 | 15 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 8 | | 7 | 8 | 8 |
| 25TH.....GM..... | 21 | | 14 | 16 | 25 |
| 50TH.....GM..... | 28 | | 16 | 28 | 28 |
| 75TH.....GM..... | 32 | | 28 | 30 | 32 |
| 90TH.....GM..... | 56 | | 30 | 45 | 56 |
| 95TH.....GM..... | 60 | | 56 | 57 | 57 |
| 99TH.....GM..... | 90 | | 64 | 90 | 64 |
| MAXIMUM.....GM..... | 168 | 28 | 96 | 96 | 112 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 3.4 | 0 | .9 | .9 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .1 | 0 | .3 | .1 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 17.5 | 16.9 | 7.6 | 7.8 | 2.3 | 2.1 | .8 | .9 | .4 | 1.4 | 1.3 | 1.6 |
| 10.4 | 9.3 | 4.0 | 5.4 | 1.6 | 1.6 | .5 | .6 | .4 | .8 | .8 | 1.0 |
| 4.9 | 4.9 | 2.2 | 1.5 | .5 | .5 | .2 | .2 | 0 | .3 | .2 | .3 |
| 2.2 | 2.7 | 1.3 | .9 | .2 | .1 | .1 | .1 | 0 | .3 | .3 | .3 |
| 1.6 | 1.6 | 1.7 | 1.4 | 1.4 | 1.3 | 1.4 | 1.4 | 1.0 | 1.7 | 1.5 | 1.6 |
| .8 | .8 | .8 | .8 | .6 | .5 | .6 | .7 | .0 | .9 | .8 | .8 |
| 4.0 | 5.0 | 6.0 | 6.0 | 3.0 | 3.0 | 3.0 | 3.0 | 1.0 | 3.0 | 3.0 | 3.0 |
| 19 | 18 | 25 | 15 | 18 | 14 | 14 | 13 | 11 | 16 | 12 | 14 |
| 14 | 12 | 18 | 9 | 17 | 9 | 8 | 8 | 2 | 9 | 4 | 12 |
| 5 | 5 | 5 | 5 | 3 | 4 | | 2 | | | | |
| 9 | 9 | 9 | 9 | 9 | 9 | | 9 | | | | |
| 19 | 14 | 20 | 10 | 10 | 10 | | 10 | | | | |
| 24 | 21 | 35 | 20 | 20 | 19 | | 17 | | | | |
| 37 | 33 | 56 | 28 | 30 | 28 | | 28 | | | | |
| 47 | 40 | 59 | 30 | 56 | 33 | | 30 | | | | |
| 63 | 56 | 84 | 56 | 103 | 37 | | 37 | | | | |
| 39 | 33 | 44 | 33 | 33 | 33 | 37 | 30 | 33 | 30 | 38 | 23 |
| 23 | 17 | 25 | 14 | 21 | 17 | 17 | 10 | 7 | 15 | 13 | 14 |
| 38 | 32 | 44 | 28 | 44 | 34 | 27 | 26 | 0 | 28 | 14 | 15 |
| 23 | 13 | 20 | 11 | 38 | 14 | 15 | 10 | 0 | 13 | 0 | (2) |
| 36 | 34 | 51 | 32 | 43 | 33 | 9 | 25 | 0 | 27 | 9 | 36 |
| 14 | 13 | 19 | 10 | 16 | 5 | 4 | 13 | 0 | 3 | 0 | 7 |
| 168 | 120 | 126 | 84 | 168 | 84 | 56 | 56 | 42 | 60 | 56 | 57 |
| 336 | 252 | 294 | 176 | 308 | 112 | 112 | 112 | 42 | 90 | 56 | 126 |
| 36 | 32 | 45 | 31 | 38 | 33 | 28 | 28 | 33 | 27 | 22 | 26 |
| 20 | 15 | 22 | 13 | 28 | 15 | 18 | 12 | 7 | 11 | 16 | 13 |
| 14 | 11 | 14 | 14 | 8 | 11 | | 6 | | | | |
| 28 | 28 | 28 | 28 | 24 | 28 | | 28 | | | | |
| 28 | 28 | 42 | 28 | 30 | 28 | | 28 | | | | |
| 45 | 32 | 60 | 30 | 53 | 42 | | 30 | | | | |
| 60 | 56 | 70 | 56 | 75 | 56 | | 42 | | | | |
| 64 | 60 | 84 | 60 | 90 | 60 | | 45 | | | | |
| 112 | 84 | 120 | 84 | 168 | 84 | | 56 | | | | |
| 168 | 120 | 120 | 84 | 168 | 84 | 56 | 56 | 42 | 45 | 56 | 57 |
| 0 | 0 | 0 | 0 | 0 | 1.6 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 1.1 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 5.51.--READY-TO-EAT CEREALS, 25-32 GM PER CUP, LESS THAN 15 PCT SUGAR
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 19.4 | 6.2 | 31.8 | 36.1 | 35.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 10.9 | 2.0 | 16.6 | 18.4 | 17.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 5.5 | 2.7 | 10.0 | 12.0 | 11.3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 3.1 | 1.5 | 5.2 | 5.7 | 6.4 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.6 | 2.0 | 1.7 | 1.7 | 1.7 |
| STANDARD DEVIATION.....DO..... | .8 | .9 | .8 | .9 | .8 |
| MAXIMUM.....DO..... | 10.0 | 4.0 | 6.0 | 10.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 16 | 9 | 11 | 14 | 16 |
| STANDARD DEVIATION.....GM..... | 12 | 9 | 8 | 10 | 11 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 4 | | 2 | 4 | 4 |
| 25TH.....GM..... | 8 | | 5 | 8 | 8 |
| 50TH.....GM..... | 13 | | 8 | 10 | 13 |
| 75TH.....GM..... | 20 | | 15 | 17 | 20 |
| 90TH.....GM..... | 30 | | 21 | 25 | 30 |
| 95TH.....GM..... | 38 | | 27 | 33 | 38 |
| 99TH.....GM..... | 58 | | 33 | 46 | 50 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 29 | 13 | 19 | 25 | 27 |
| STANDARD DEVIATION.....GM..... | 16 | 8 | 11 | 12 | 12 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 29 | 13 | 20 | 24 | 28 |
| STANDARD DEVIATION.....GM..... | 15 | 11 | 9 | 11 | 15 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 31 | 15 | 20 | 27 | 28 |
| STANDARD DEVIATION.....GM..... | 16 | 13 | 10 | 14 | 11 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 275 | 60 | 75 | 200 | 125 |
| MAXIMUM OVER 3 DAYS.....GM..... | 510 | 120 | 180 | 400 | 200 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 29 | 13 | 19 | 24 | 27 |
| STANDARD DEVIATION.....GM..... | 16 | 13 | 10 | 12 | 13 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 8 | | 6 | 10 | 8 |
| 25TH.....GM..... | 25 | | 13 | 15 | 25 |
| 50TH.....GM..... | 25 | | 15 | 25 | 25 |
| 75TH.....GM..... | 30 | | 25 | 30 | 30 |
| 90TH.....GM..... | 50 | | 30 | 38 | 50 |
| 95TH.....GM..... | 57 | | 30 | 50 | 50 |
| 99TH.....GM..... | 85 | | 50 | 60 | 75 |
| MAXIMUM.....GM..... | 275 | 60 | 75 | 112 | 100 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 52.1 | 0 | 4.0 | 1.6 | 5.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | .7 | .1 | .5 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 34.0 | 28.6 | 24.5 | 19.1 | 12.5 | 13.4 | 13.8 | 13.1 | 24.0 | 20.0 | 25.2 | 21.8 |
| 17.9 | 16.4 | 13.8 | 11.9 | 7.3 | 9.0 | 7.7 | 8.4 | 10.6 | 10.3 | 9.5 | 11.3 |
| 11.1 | 8.4 | 5.6 | 5.2 | 3.4 | 3.5 | 2.9 | 3.3 | 6.9 | 5.8 | 9.0 | 5.9 |
| 5.0 | 3.8 | 5.1 | 2.0 | 1.8 | .9 | 3.2 | 1.4 | 6.5 | 3.9 | 6.7 | 4.6 |
| 1.7 | 1.6 | 1.7 | 1.5 | 1.6 | 1.4 | 1.7 | 1.5 | 1.9 | 1.7 | 2.0 | 1.8 |
| .8 | .8 | .9 | .8 | .7 | .6 | .8 | .7 | .9 | .8 | 1.0 | .9 |
| 6.0 | 5.0 | 6.0 | 5.0 | 4.0 | 5.0 | 4.0 | 5.0 | 4.0 | 4.0 | 5.0 | 7.0 |
| 19 | 15 | 23 | 15 | 19 | 13 | 18 | 13 | 17 | 14 | 18 | 14 |
| 15 | 10 | 19 | 10 | 14 | 8 | 15 | 9 | 11 | 11 | 12 | 10 |
| 4 | 4 | 5 | 4 | 4 | 4 | 4 | 2 | 4 | 4 | 5 | 3 |
| 8 | 8 | 9 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 |
| 17 | 13 | 17 | 13 | 17 | 10 | 13 | 9 | 15 | 13 | 15 | 10 |
| 25 | 20 | 30 | 20 | 25 | 17 | 24 | 17 | 25 | 18 | 25 | 18 |
| 38 | 27 | 50 | 30 | 38 | 21 | 33 | 25 | 33 | 25 | 30 | 25 |
| 50 | 33 | 53 | 33 | 50 | 27 | 45 | 30 | 35 | 30 | 45 | 30 |
| 67 | 50 | 95 | 49 | 70 | 42 | 70 | 45 | 50 | 50 | 60 | 60 |
| 33 | 30 | 39 | 32 | 34 | 29 | 29 | 26 | 28 | 25 | 27 | 25 |
| 23 | 14 | 25 | 16 | 18 | 14 | 17 | 13 | 11 | 12 | 11 | 12 |
| 36 | 29 | 39 | 32 | 37 | 28 | 31 | 27 | 26 | 25 | 30 | 23 |
| 18 | 16 | 17 | 14 | 20 | 13 | 11 | 12 | 12 | 7 | 13 | 11 |
| 38 | 29 | 46 | 27 | 39 | 28 | 33 | 26 | 30 | 27 | 28 | 26 |
| 19 | 9 | 23 | 10 | 17 | 8 | 20 | 12 | 10 | 16 | 14 | 13 |
| 275 | 180 | 188 | 125 | 168 | 170 | 180 | 120 | 75 | 170 | 125 | 90 |
| 375 | 330 | 510 | 150 | 300 | 270 | 380 | 240 | 180 | 365 | 200 | 210 |
| 35 | 29 | 40 | 30 | 36 | 28 | 31 | 26 | 28 | 26 | 27 | 23 |
| 20 | 13 | 21 | 14 | 18 | 12 | 18 | 13 | 12 | 13 | 14 | 11 |
| 11 | 10 | 13 | 13 | 11 | 10 | 6 | 6 | 11 | 8 | 13 | 6 |
| 25 | 25 | 25 | 25 | 25 | 25 | 25 | 19 | 25 | 19 | 19 | 15 |
| 30 | 25 | 30 | 25 | 30 | 25 | 28 | 25 | 25 | 25 | 25 | 25 |
| 45 | 30 | 50 | 38 | 50 | 30 | 38 | 30 | 30 | 30 | 30 | 30 |
| 60 | 50 | 60 | 50 | 60 | 50 | 50 | 45 | 50 | 38 | 45 | 30 |
| 60 | 50 | 75 | 57 | 75 | 50 | 60 | 50 | 50 | 47 | 50 | 38 |
| 100 | 75 | 120 | 84 | 100 | 60 | 120 | 60 | 60 | 60 | 75 | 60 |
| 275 | 100 | 150 | 99 | 120 | 125 | 180 | 100 | 75 | 170 | 125 | 60 |
| 2.9 | .9 | 1.7 | 0 | 16.6 | 4.2 | 10.2 | 2.9 | 0 | 2.0 | 0 | 0 |
| .2 | .1 | .3 | 0 | 2.2 | .4 | .9 | .2 | 0 | .3 | 0 | 0 |

TABLE 5.52.--READY-TO-EAT CEREALS, 35-45 GM PER CUP, 30 PCT OR MORE SUGAR
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.7 | .7 | 11.6 | 15.5 | 16.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.5 | .7 | 6.8 | 9.3 | 9.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.5 | 0 | 2.7 | 4.6 | 4.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .7 | 0 | 2.1 | 1.5 | 2.5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.6 | 1.0 | 1.7 | 1.5 | 1.6 |
| STANDARD DEVIATION.....DO..... | .8 | .0 | .9 | .7 | .8 |
| MAXIMUM.....DO..... | 7.0 | 1.0 | 5.0 | 4.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 22 | 8 | 16 | 18 | 21 |
| STANDARD DEVIATION.....GM..... | 16 | 5 | 11 | 11 | 14 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 6 | | 3 | 5 | 6 |
| 25TH.....GM..... | 12 | | 7 | 12 | 12 |
| 50TH.....GM..... | 15 | | 12 | 13 | 18 |
| 75TH.....GM..... | 27 | | 23 | 25 | 26 |
| 90TH.....GM..... | 42 | | 33 | 37 | 40 |
| 95TH.....GM..... | 53 | | 40 | 40 | 49 |
| 99TH.....GM..... | 80 | | 47 | 53 | 74 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 43 | 24 | 29 | 36 | 40 |
| STANDARD DEVIATION.....GM..... | 24 | 14 | 23 | 19 | 20 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 43 | 0 | 29 | 35 | 40 |
| STANDARD DEVIATION.....GM..... | 22 | 0 | 11 | 15 | 19 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 44 | 0 | 31 | 38 | 37 |
| STANDARD DEVIATION.....GM..... | 22 | 0 | 10 | 9 | 16 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 320 | 37 | 140 | 140 | 148 |
| MAXIMUM OVER 3 DAYS.....GM..... | 520 | 37 | 148 | 200 | 240 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 42 | 24 | 29 | 36 | 39 |
| STANDARD DEVIATION.....GM..... | 23 | 14 | 15 | 16 | 19 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 12 | | 9 | 12 | 10 |
| 25TH.....GM..... | 28 | | 18 | 23 | 28 |
| 50TH.....GM..... | 37 | | 28 | 35 | 37 |
| 75TH.....GM..... | 50 | | 37 | 40 | 40 |
| 90TH.....GM..... | 74 | | 40 | 57 | 70 |
| 95TH.....GM..... | 80 | | 53 | 70 | 80 |
| 99TH.....GM..... | 120 | | 80 | 80 | 111 |
| MAXIMUM.....GM..... | 175 | 37 | 90 | 120 | 148 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 2.2 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .1 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 17.4 | 13.2 | 10.0 | 8.5 | 3.7 | 2.7 | 1.3 | 1.5 | 1.9 | 1.9 | 1.3 | 2.0 |
| 10.5 | 8.0 | 6.0 | 5.6 | 2.2 | 2.0 | .8 | 1.1 | 1.1 | 1.1 | .3 | 1.1 |
| 4.7 | 3.4 | 2.7 | 2.0 | 1.0 | .6 | .3 | .2 | .6 | .5 | .8 | .3 |
| 2.2 | 1.8 | 1.3 | .9 | .5 | .1 | .2 | .2 | .2 | .3 | .2 | .6 |
| 1.6 | 1.6 | 1.6 | 1.5 | 1.5 | 1.3 | 1.6 | 1.5 | 1.7 | 1.6 | 1.9 | 1.7 |
| .8 | .8 | .8 | .7 | .8 | .6 | .8 | .8 | .9 | .8 | .6 | .9 |
| 7.0 | 5.0 | 5.0 | 3.0 | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| 25 | 22 | 30 | 21 | 25 | 19 | 22 | 18 | 23 | 20 | 30 | 16 |
| 19 | 17 | 20 | 14 | 20 | 14 | 17 | 13 | 18 | 12 | 20 | 6 |
| 6 | 6 | 6 | 9 | 5 | 6 | 3 | 6 | | | | |
| 12 | 12 | 13 | 12 | 12 | 12 | 12 | 12 | | | | |
| 20 | 16 | 25 | 13 | 20 | 13 | 13 | 13 | | | | |
| 30 | 27 | 41 | 26 | 30 | 23 | 27 | 23 | | | | |
| 49 | 41 | 58 | 38 | 57 | 35 | 40 | 40 | | | | |
| 58 | 53 | 67 | 53 | 67 | 49 | 70 | 43 | | | | |
| 89 | 93 | 93 | 80 | 100 | 75 | 80 | 60 | | | | |
| 48 | 42 | 65 | 44 | 50 | 45 | 40 | 38 | 31 | 33 | 35 | 41 |
| 23 | 20 | 39 | 18 | 28 | 24 | 23 | 14 | 8 | 8 | 0 | 14 |
| 48 | 45 | 53 | 42 | 52 | 38 | 38 | 39 | 54 | 39 | 40 | 40 |
| 23 | 25 | 27 | 22 | 31 | 18 | 13 | 14 | 16 | 11 | 0 | 0 |
| 52 | 44 | 56 | 44 | 46 | 57 | 50 | 40 | 45 | 42 | 80 | 16 |
| 31 | 23 | 21 | 16 | 28 | 19 | 21 | 14 | 24 | 3 | 0 | 5 |
| 320 | 170 | 225 | 140 | 170 | 180 | 100 | 80 | 80 | 60 | 80 | 80 |
| 520 | 383 | 320 | 280 | 300 | 225 | 240 | 200 | 240 | 140 | 240 | 80 |
| 47 | 43 | 57 | 43 | 49 | 44 | 42 | 38 | 39 | 38 | 47 | 29 |
| 23 | 23 | 31 | 19 | 32 | 21 | 21 | 14 | 15 | 10 | 16 | 15 |
| 14 | 13 | 18 | 18 | 5 | 13 | 10 | 19 | | | | |
| 35 | 35 | 37 | 35 | 28 | 35 | 30 | 30 | | | | |
| 40 | 37 | 53 | 40 | 40 | 40 | 40 | 40 | | | | |
| 60 | 53 | 74 | 45 | 70 | 53 | 40 | 40 | | | | |
| 80 | 70 | 90 | 74 | 80 | 70 | 80 | 60 | | | | |
| 80 | 80 | 120 | 80 | 120 | 80 | 80 | 70 | | | | |
| 111 | 140 | 167 | 120 | 170 | 120 | 100 | 80 | | | | |
| 175 | 170 | 167 | 140 | 170 | 120 | 100 | 80 | 80 | 60 | 80 | 80 |
| 0 | 1.3 | 0 | 0 | 0 | 0 | .9 | 0 | 0 | 0 | 0 | 0 |
| 0 | .3 | 0 | 0 | 0 | 0 | .9 | 0 | 0 | 0 | 0 | 0 |

TABLE 5.53.--READY-TO-EAT CEREALS, 35-45 GM PER CUP, LESS THAN 20 PCT SUGAR
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.9 | .8 | 3.4 | 4.8 | 4.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.8 | .5 | 1.8 | 3.0 | 3.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.3 | 0 | .8 | 1.3 | 1.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .8 | .3 | .7 | .5 | .5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.6 | 1.8 | 1.7 | 1.5 | 1.5 |
| STANDARD DEVIATION.....DO..... | .8 | 1.0 | .8 | .7 | .8 |
| MAXIMUM.....DO..... | 6.0 | 3.0 | 3.0 | 4.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 20 | 9 | 19 | 17 | 20 |
| STANDARD DEVIATION.....GM..... | 16 | 6 | 13 | 13 | 14 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 4 | | | 3 | 5 |
| 25TH.....GM..... | 12 | | | 9 | 12 |
| 50TH.....GM..... | 15 | | | 13 | 17 |
| 75TH.....GM..... | 27 | | | 25 | 22 |
| 90TH.....GM..... | 38 | | | 29 | 37 |
| 95TH.....GM..... | 47 | | | 39 | 55 |
| 99TH.....GM..... | 83 | | | 62 | 72 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 39 | 14 | 34 | 33 | 43 |
| STANDARD DEVIATION.....GM..... | 22 | 4 | 21 | 18 | 23 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 38 | 0 | 30 | 34 | 35 |
| STANDARD DEVIATION.....GM..... | 19 | 0 | 12 | 18 | 11 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 39 | 16 | 35 | 36 | 46 |
| STANDARD DEVIATION.....GM..... | 23 | 5 | 12 | 16 | 22 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 215 | 22 | 86 | 138 | 129 |
| MAXIMUM OVER 3 DAYS.....GM..... | 409 | 65 | 165 | 193 | 215 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 38 | 15 | 34 | 33 | 40 |
| STANDARD DEVIATION.....GM..... | 21 | 5 | 17 | 19 | 20 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 9 | | | 5 | 11 |
| 25TH.....GM..... | 23 | | | 21 | 22 |
| 50TH.....GM..... | 35 | | | 32 | 38 |
| 75TH.....GM..... | 43 | | | 43 | 55 |
| 90TH.....GM..... | 65 | | | 55 | 65 |
| 95TH.....GM..... | 83 | | | 65 | 74 |
| 99TH.....GM..... | 110 | | | 86 | 86 |
| MAXIMUM.....GM..... | 172 | 22 | 86 | 138 | 111 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 10.1 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.1 | 4.4 | 4.8 | 3.5 | 4.0 | 4.1 | 4.9 | 4.7 | 8.7 | 9.1 | 7.2 | 8.7 |
| 3.2 | 2.9 | 2.6 | 2.7 | 2.7 | 2.4 | 2.2 | 2.6 | 5.5 | 5.1 | 4.1 | 3.0 |
| 1.4 | .9 | 1.2 | .5 | 1.0 | 1.3 | 1.7 | 1.4 | .8 | 1.8 | 2.2 | 2.5 |
| .4 | .6 | 1.0 | .3 | .3 | .4 | .9 | .7 | 2.4 | 2.2 | .9 | 3.1 |
| 1.5 | 1.5 | 1.7 | 1.3 | 1.4 | 1.5 | 1.8 | 1.6 | 1.6 | 1.7 | 1.6 | 2.1 |
| .7 | .8 | .9 | .6 | .8 | .7 | .8 | .7 | .9 | .9 | .7 | 1.0 |
| 4.0 | 3.0 | 4.0 | 3.0 | 6.0 | 5.0 | 5.0 | 3.0 | 3.0 | 4.0 | 3.0 | 5.0 |
| 24 | 22 | 37 | 20 | 26 | 19 | 23 | 17 | 19 | 17 | 17 | 21 |
| 20 | 13 | 28 | 13 | 22 | 12 | 18 | 11 | 13 | 10 | 12 | 14 |
| 4 | 6 | 9 | 5 | 4 | 4 | 4 | 3 | 5 | 2 | | 3 |
| 12 | 13 | 18 | 12 | 13 | 9 | 12 | 9 | 9 | 9 | | 12 |
| 19 | 22 | 28 | 14 | 22 | 14 | 18 | 13 | 13 | 13 | | 13 |
| 29 | 29 | 49 | 27 | 31 | 25 | 31 | 23 | 25 | 23 | | 27 |
| 46 | 37 | 72 | 37 | 46 | 37 | 43 | 35 | 38 | 33 | | 43 |
| 55 | 50 | 110 | 43 | 57 | 43 | 70 | 38 | 41 | 37 | | 53 |
| 128 | 57 | 129 | 70 | 136 | 55 | 83 | 47 | 73 | 43 | | 58 |
| 46 | 48 | 58 | 43 | 53 | 36 | 38 | 33 | 34 | 29 | 33 | 28 |
| 25 | 23 | 24 | 23 | 28 | 19 | 19 | 17 | 16 | 16 | 14 | 14 |
| 48 | 42 | 74 | 50 | 45 | 37 | 37 | 32 | 34 | 35 | 28 | 35 |
| 15 | 19 | 38 | 12 | 16 | 18 | 17 | 15 | 4 | 12 | 9 | 19 |
| 67 | 38 | 68 | 45 | 78 | 39 | 45 | 29 | 34 | 27 | 37 | 29 |
| 36 | 12 | 31 | 13 | 34 | 7 | 23 | 13 | 14 | 9 | 19 | 14 |
| 165 | 111 | 215 | 110 | 175 | 110 | 123 | 86 | 75 | 75 | 70 | 140 |
| 387 | 185 | 387 | 210 | 409 | 172 | 258 | 180 | 219 | 129 | 210 | 175 |
| 50 | 43 | 64 | 45 | 54 | 37 | 39 | 31 | 34 | 29 | 32 | 29 |
| 25 | 19 | 27 | 20 | 28 | 18 | 21 | 16 | 16 | 13 | 15 | 17 |
| 11 | 18 | 28 | 14 | 13 | 11 | 9 | 4 | 12 | 4 | | 4 |
| 35 | 28 | 43 | 35 | 35 | 28 | 26 | 20 | 19 | 18 | | 19 |
| 43 | 40 | 57 | 43 | 43 | 35 | 37 | 35 | 35 | 30 | | 28 |
| 65 | 55 | 86 | 53 | 70 | 43 | 53 | 40 | 38 | 37 | | 39 |
| 83 | 70 | 110 | 70 | 86 | 60 | 70 | 43 | 57 | 43 | | 43 |
| 105 | 86 | 110 | 85 | 110 | 74 | 83 | 57 | 70 | 53 | | 53 |
| 129 | 95 | 129 | 110 | 129 | 86 | 86 | 84 | 75 | 57 | | 105 |
| 129 | 110 | 129 | 110 | 172 | 110 | 110 | 86 | 75 | 70 | 70 | 105 |
| .7 | 0 | 1.3 | 0 | 2.5 | 0 | 0 | 1.8 | 0 | 0 | 0 | 3.7 |
| .5 | 0 | 1.1 | 0 | 1.1 | 0 | 0 | .3 | 0 | 0 | 0 | 2.1 |

TABLE 5.54.--READY-TO-EAT CEREALS, 50-75 GM PER CUP, ALL SUGAR LEVELS
 CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 7.8 | .2 | 6.2 | 7.8 | 7.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.5 | 0 | 3.6 | 4.6 | 5.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.9 | .2 | 1.6 | 2.1 | 2.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.4 | 0 | 1.0 | 1.1 | .6 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.6 | 2.0 | 1.6 | 1.6 | 1.4 |
| STANDARD DEVIATION.....DO..... | .8 | .0 | .8 | .8 | .7 |
| MAXIMUM.....DO..... | 6.0 | 2.0 | 4.0 | 5.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 25 | 17 | 18 | 21 | 22 |
| STANDARD DEVIATION.....GM..... | 20 | 0 | 13 | 14 | 17 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 6 | | 4 | 8 | 4 |
| 25TH.....GM..... | 13 | | 8 | 9 | 17 |
| 50TH.....GM..... | 17 | | 17 | 17 | 17 |
| 75TH.....GM..... | 33 | | 25 | 25 | 30 |
| 90TH.....GM..... | 50 | | 38 | 47 | 40 |
| 95TH.....GM..... | 60 | | 42 | 50 | 60 |
| 99TH.....GM..... | 100 | | 50 | 67 | 85 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 48 | 0 | 30 | 39 | 46 |
| STANDARD DEVIATION.....GM..... | 28 | 0 | 14 | 17 | 21 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 47 | 25 | 38 | 43 | 43 |
| STANDARD DEVIATION.....GM..... | 23 | 0 | 13 | 21 | 22 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 47 | 0 | 37 | 39 | 61 |
| STANDARD DEVIATION.....GM..... | 28 | 0 | 10 | 16 | 19 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 400 | 25 | 80 | 100 | 120 |
| MAXIMUM OVER 3 DAYS.....GM..... | 600 | 50 | 150 | 200 | 270 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 46 | 25 | 35 | 39 | 47 |
| STANDARD DEVIATION.....GM..... | 26 | 0 | 14 | 18 | 22 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 13 | | 13 | 13 | 13 |
| 25TH.....GM..... | 25 | | 25 | 25 | 25 |
| 50TH.....GM..... | 50 | | 30 | 38 | 50 |
| 75TH.....GM..... | 50 | | 50 | 50 | 50 |
| 90TH.....GM..... | 75 | | 50 | 60 | 75 |
| 95TH.....GM..... | 100 | | 50 | 73 | 100 |
| 99TH.....GM..... | 142 | | 60 | 100 | 120 |
| MAXIMUM.....GM..... | 400 | 25 | 80 | 100 | 120 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 12.5 | 0 | 0 | 1.0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 0 | 0 | .5 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 7.0 | 6.8 | 6.8 | 5.5 | 5.8 | 6.4 | 8.4 | 8.0 | 13.9 | 13.8 | 13.4 | 13.6 |
| 4.9 | 4.7 | 4.2 | 3.9 | 3.4 | 3.8 | 4.8 | 4.6 | 6.0 | 6.7 | 6.1 | 5.4 |
| 1.4 | 1.6 | 1.5 | .9 | 1.5 | 1.6 | 2.2 | 1.8 | 3.3 | 4.5 | 2.4 | 3.3 |
| .6 | .6 | 1.2 | .7 | .9 | .9 | 1.4 | 1.6 | 4.6 | 2.7 | 4.8 | 4.8 |
| 1.5 | 1.4 | 1.6 | 1.4 | 1.6 | 1.6 | 1.6 | 1.6 | 2.0 | 1.7 | 1.9 | 2.0 |
| .8 | .7 | .9 | .7 | .8 | .8 | .8 | .8 | 1.1 | .8 | 1.0 | 1.0 |
| 5.0 | 4.0 | 5.0 | 3.0 | 4.0 | 4.0 | 4.0 | 5.0 | 6.0 | 4.0 | 5.0 | 4.0 |
| 31 | 26 | 36 | 24 | 32 | 23 | 27 | 23 | 28 | 21 | 24 | 26 |
| 26 | 18 | 30 | 15 | 25 | 15 | 19 | 16 | 27 | 17 | 17 | 23 |
| 8 | 8 | 9 | 4 | 10 | 7 | 6 | 5 | 5 | 4 | 5 | 5 |
| 17 | 17 | 17 | 17 | 17 | 13 | 15 | 13 | 9 | 10 | 13 | 13 |
| 25 | 17 | 25 | 17 | 25 | 17 | 20 | 17 | 19 | 17 | 17 | 17 |
| 42 | 33 | 40 | 33 | 37 | 33 | 33 | 30 | 33 | 25 | 33 | 33 |
| 60 | 50 | 75 | 50 | 58 | 50 | 50 | 46 | 60 | 40 | 50 | 45 |
| 67 | 60 | 111 | 50 | 75 | 50 | 67 | 55 | 75 | 50 | 63 | 60 |
| 133 | 100 | 133 | 67 | 150 | 67 | 100 | 83 | 120 | 100 | 80 | 83 |
| 66 | 57 | 61 | 54 | 66 | 46 | 50 | 43 | 37 | 37 | 40 | 37 |
| 48 | 36 | 26 | 29 | 34 | 23 | 27 | 23 | 17 | 16 | 17 | 18 |
| 67 | 52 | 57 | 54 | 52 | 48 | 51 | 43 | 46 | 37 | 46 | 38 |
| 34 | 21 | 21 | 21 | 24 | 18 | 22 | 21 | 25 | 22 | 29 | 17 |
| 74 | 59 | 88 | 41 | 67 | 40 | 51 | 41 | 46 | 37 | 34 | 41 |
| 37 | 25 | 33 | 17 | 37 | 18 | 22 | 20 | 36 | 24 | 17 | 31 |
| 400 | 275 | 200 | 168 | 200 | 150 | 200 | 200 | 240 | 180 | 120 | 300 |
| 510 | 350 | 540 | 225 | 450 | 225 | 300 | 338 | 530 | 360 | 240 | 600 |
| 65 | 54 | 67 | 50 | 61 | 44 | 50 | 42 | 42 | 37 | 38 | 38 |
| 40 | 26 | 31 | 25 | 33 | 21 | 25 | 22 | 26 | 22 | 21 | 24 |
| 25 | 25 | 25 | 15 | 17 | 14 | 13 | 13 | 9 | 7 | 8 | 6 |
| 50 | 50 | 50 | 30 | 40 | 25 | 30 | 25 | 25 | 25 | 19 | 25 |
| 50 | 50 | 50 | 50 | 50 | 50 | 50 | 43 | 38 | 30 | 38 | 30 |
| 75 | 56 | 90 | 50 | 75 | 50 | 60 | 50 | 50 | 50 | 50 | 50 |
| 100 | 100 | 100 | 75 | 100 | 75 | 83 | 60 | 75 | 60 | 60 | 60 |
| 125 | 100 | 120 | 100 | 142 | 75 | 100 | 75 | 100 | 60 | 63 | 75 |
| 200 | 150 | 180 | 120 | 150 | 120 | 113 | 110 | 120 | 120 | 120 | 150 |
| 400 | 200 | 200 | 168 | 150 | 150 | 200 | 200 | 170 | 180 | 120 | 150 |
| 3.6 | .8 | 0 | 0 | 0 | 0 | 0 | 2.4 | 3.4 | 1.3 | 0 | 0 |
| 1.7 | .4 | 0 | 0 | 0 | 0 | 0 | .3 | 1.1 | .3 | 0 | 0 |

TABLE 5.55.--READY-TO-EAT CEREALS, 80 GM OR MORE PER CUP, ALL SUGAR LEVELS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.5 | .8 | 2.4 | 2.8 | 2.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.0 | .2 | 1.6 | 2.0 | 2.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .9 | .4 | .7 | .6 | .5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .6 | .2 | .1 | .2 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.6 | 2.1 | 1.4 | 1.4 | 1.4 |
| STANDARD DEVIATION.....DO..... | .8 | .7 | .6 | .9 | .7 |
| MAXIMUM.....DO..... | 6.0 | 3.0 | 3.0 | 6.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 32 | 29 | 14 | 28 | 28 |
| STANDARD DEVIATION.....GM..... | 29 | 14 | 10 | 23 | 25 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 2 | | | | 2 |
| 25TH.....GM..... | 12 | | | | 14 |
| 50TH.....GM..... | 28 | | | | 18 |
| 75TH.....GM..... | 39 | | | | 37 |
| 90TH.....GM..... | 73 | | | | 73 |
| 95TH.....GM..... | 92 | | | | 75 |
| 99TH.....GM..... | 131 | | | | 112 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 64 | 27 | 41 | 53 | 58 |
| STANDARD DEVIATION.....GM..... | 49 | 0 | 30 | 33 | 42 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 61 | 47 | 23 | 73 | 71 |
| STANDARD DEVIATION.....GM..... | 45 | 16 | 17 | 30 | 46 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 58 | 42 | 23 | 68 | 69 |
| STANDARD DEVIATION.....GM..... | 43 | 0 | 0 | 35 | 27 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 495 | 65 | 110 | 170 | 224 |
| MAXIMUM OVER 3 DAYS.....GM..... | 678 | 129 | 110 | 339 | 337 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 60 | 42 | 31 | 59 | 61 |
| STANDARD DEVIATION.....GM..... | 46 | 12 | 26 | 35 | 42 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 5 | | | | 7 |
| 25TH.....GM..... | 28 | | | | 28 |
| 50TH.....GM..... | 55 | | | | 55 |
| 75TH.....GM..... | 84 | | | | 84 |
| 90TH.....GM..... | 112 | | | | 112 |
| 95TH.....GM..... | 126 | | | | 113 |
| 99TH.....GM..... | 220 | | | | 224 |
| MAXIMUM.....GM..... | 440 | 65 | 110 | 170 | 224 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 20.4 | 0 | 2.3 | 1.3 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.0 | 0 | 6.6 | 1.9 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 2.2 | 2.3 | 2.2 | 2.3 | 3.5 | 3.6 | 3.7 | 3.9 | 5.6 | 4.8 | 3.9 | 6.2 |
| 1.2 | 1.6 | 1.7 | 1.7 | 2.1 | 2.2 | 2.0 | 2.3 | 2.4 | 2.5 | 1.8 | 3.1 |
| .7 | .4 | .3 | .3 | .9 | 1.0 | 1.0 | 1.0 | 1.0 | 1.2 | 1.7 | 1.4 |
| .2 | .4 | .2 | .2 | .5 | .4 | .7 | .7 | 2.2 | 1.1 | .4 | 1.7 |
| 1.6 | 1.5 | 1.4 | 1.4 | 1.6 | 1.5 | 1.7 | 1.6 | 2.0 | 1.7 | 1.6 | 1.9 |
| .8 | .8 | .7 | .7 | .8 | .7 | .8 | .8 | .9 | .8 | .7 | 1.1 |
| 4.0 | 3.0 | 3.0 | 3.0 | 4.0 | 4.0 | 4.0 | 4.0 | 4.0 | 3.0 | 3.0 | 5.0 |
| 51 | 36 | 51 | 32 | 42 | 29 | 38 | 27 | 34 | 28 | 31 | 21 |
| 42 | 27 | 33 | 26 | 38 | 26 | 34 | 22 | 32 | 23 | 35 | 19 |
| | 5 | | | 3 | 2 | 4 | 2 | 2 | 2 | | 2 |
| | 18 | | | 9 | 9 | 14 | 9 | 14 | 9 | | 9 |
| | 28 | | | 37 | 19 | 28 | 19 | 28 | 21 | | 19 |
| | 55 | | | 56 | 37 | 56 | 37 | 37 | 37 | | 28 |
| | 73 | | | 85 | 73 | 84 | 57 | 84 | 56 | | 37 |
| | 73 | | | 113 | 84 | 110 | 75 | 120 | 84 | | 37 |
| | 151 | | | 194 | 112 | 147 | 85 | 149 | 121 | | 120 |
| 101 | 76 | 128 | 72 | 75 | 55 | 66 | 55 | 67 | 69 | 34 | 59 |
| 68 | 45 | 89 | 39 | 60 | 44 | 42 | 42 | 28 | 41 | 21 | 30 |
| 100 | 67 | 91 | 67 | 83 | 59 | 73 | 53 | 41 | 34 | 60 | 21 |
| 80 | 36 | 26 | 48 | 53 | 40 | 42 | 29 | 24 | 26 | 54 | 16 |
| 94 | 74 | 98 | 77 | 84 | 58 | 70 | 44 | 49 | 44 | 79 | 30 |
| 36 | 33 | 30 | 37 | 50 | 33 | 51 | 30 | 44 | 34 | 40 | 29 |
| 440 | 226 | 495 | 220 | 280 | 220 | 226 | 220 | 224 | 180 | 168 | 120 |
| 660 | 452 | 495 | 385 | 581 | 336 | 678 | 360 | 448 | 363 | 336 | 360 |
| 95 | 73 | 111 | 72 | 79 | 56 | 69 | 50 | 51 | 49 | 56 | 33 |
| 71 | 41 | 57 | 43 | 57 | 41 | 47 | 36 | 41 | 39 | 48 | 31 |
| | 28 | | | 10 | 5 | 7 | 7 | 2 | 2 | | 2 |
| | 55 | | | 28 | 28 | 28 | 28 | 21 | 21 | | 7 |
| | 56 | | | 63 | 43 | 63 | 43 | 36 | 42 | | 28 |
| | 110 | | | 110 | 84 | 110 | 78 | 84 | 57 | | 49 |
| | 112 | | | 165 | 112 | 112 | 105 | 112 | 112 | | 84 |
| | 113 | | | 220 | 112 | 149 | 113 | 120 | 120 | | 110 |
| | 226 | | | 280 | 220 | 226 | 128 | 121 | 126 | | 120 |
| 440 | 226 | 330 | 220 | 280 | 220 | 226 | 220 | 224 | 180 | 168 | 120 |
| 0 | 0 | 0 | 0 | 7.2 | 4.5 | 0 | 3.2 | 0 | 0 | 0 | 1.9 |
| 0 | 0 | 0 | 0 | 3.3 | 1.5 | 0 | .7 | 0 | 0 | 0 | 1.6 |

TABLE 6.01.--COOKED DRIED BEANS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 9.8 | 2.8 | 11.7 | 13.4 | 12.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 6.9 | 2.3 | 7.6 | 9.3 | 8.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.2 | .5 | 3.2 | 2.6 | 2.9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .7 | 0 | .8 | 1.4 | .7 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.5 | 1.3 | 1.5 | 1.5 | 1.5 |
| STANDARD DEVIATION.....DO..... | .8 | .8 | .7 | .9 | .8 |
| MAXIMUM.....DO..... | 6.0 | 4.0 | 4.0 | 6.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 78 | 24 | 40 | 55 | 58 |
| STANDARD DEVIATION.....GM..... | 65 | 29 | 28 | 53 | 45 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 15 | | 8 | 8 | 12 |
| 25TH.....GM..... | 32 | | 16 | 24 | 32 |
| 50TH.....GM..... | 63 | | 32 | 32 | 48 |
| 75TH.....GM..... | 118 | | 62 | 63 | 63 |
| 90TH.....GM..... | 154 | | 85 | 123 | 123 |
| 95TH.....GM..... | 190 | | 95 | 157 | 127 |
| 99TH.....GM..... | 317 | | 123 | 247 | 247 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 167 | 39 | 86 | 110 | 132 |
| STANDARD DEVIATION.....GM..... | 118 | 30 | 61 | 78 | 86 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 174 | 113 | 93 | 125 | 125 |
| STANDARD DEVIATION.....GM..... | 105 | 55 | 45 | 79 | 80 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 175 | 0 | 64 | 126 | 126 |
| STANDARD DEVIATION.....GM..... | 102 | 0 | 22 | 93 | 60 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,140 | 185 | 370 | 740 | 555 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,420 | 370 | 555 | 1,758 | 925 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 156 | 53 | 81 | 108 | 119 |
| STANDARD DEVIATION.....GM..... | 99 | 45 | 54 | 74 | 80 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 36 | | 24 | 24 | 24 |
| 25TH.....GM..... | 93 | | 48 | 48 | 70 |
| 50TH.....GM..... | 185 | | 70 | 95 | 95 |
| 75TH.....GM..... | 190 | | 95 | 139 | 185 |
| 90TH.....GM..... | 285 | | 185 | 190 | 190 |
| 95TH.....GM..... | 380 | | 190 | 190 | 231 |
| 99TH.....GM..... | 475 | | 285 | 380 | 380 |
| MAXIMUM.....GM..... | 760 | 185 | 370 | 380 | 555 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 67.4 | 0 | 3.9 | 9.6 | 8.8 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.2 | 0 | 2.1 | 2.7 | 2.7 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 12.1 | 12.4 | 11.1 | 10.9 | 9.1 | 9.3 | 8.5 | 9.4 | 10.1 | 6.6 | 9.8 | 8.6 |
| 8.1 | 8.6 | 8.0 | 7.5 | 6.7 | 6.5 | 5.7 | 6.5 | 7.8 | 5.0 | 7.1 | 6.1 |
| 3.1 | 2.4 | 2.6 | 2.5 | 1.6 | 2.2 | 2.3 | 2.2 | 1.8 | 1.4 | 2.0 | 2.1 |
| .9 | 1.3 | .5 | .9 | .8 | .6 | .5 | .7 | .5 | .2 | .7 | .5 |
| 1.5 | 1.5 | 1.4 | 1.5 | 1.5 | 1.5 | 1.6 | 1.5 | 1.5 | 1.4 | 1.6 | 1.5 |
| .8 | .9 | .7 | .8 | .9 | .8 | .9 | .9 | .8 | .8 | .9 | .8 |
| 5.0 | 5.0 | 4.0 | 6.0 | 6.0 | 6.0 | 6.0 | 6.0 | 4.0 | 4.0 | 4.0 | 4.0 |
| 80 | 75 | 101 | 79 | 96 | 78 | 98 | 76 | 79 | 65 | 85 | 67 |
| 60 | 60 | 75 | 53 | 74 | 66 | 82 | 62 | 71 | 51 | 75 | 44 |
| 16 | 8 | 19 | 16 | 15 | 15 | 24 | 15 | 12 | 12 | 8 | 15 |
| 32 | 32 | 62 | 32 | 42 | 32 | 59 | 32 | 40 | 31 | 32 | 31 |
| 63 | 63 | 63 | 63 | 63 | 62 | 63 | 62 | 62 | 62 | 63 | 63 |
| 100 | 95 | 127 | 123 | 127 | 123 | 127 | 123 | 93 | 95 | 123 | 94 |
| 135 | 127 | 190 | 158 | 190 | 143 | 185 | 135 | 127 | 127 | 190 | 127 |
| 190 | 187 | 247 | 185 | 253 | 190 | 247 | 190 | 190 | 127 | 190 | 143 |
| 317 | 317 | 377 | 278 | 317 | 380 | 380 | 315 | 507 | 317 | 370 | 190 |
| 170 | 159 | 217 | 167 | 211 | 161 | 210 | 163 | 188 | 145 | 156 | 159 |
| 106 | 97 | 135 | 99 | 150 | 110 | 128 | 118 | 114 | 101 | 101 | 101 |
| 160 | 156 | 239 | 178 | 218 | 184 | 199 | 173 | 217 | 164 | 202 | 136 |
| 84 | 83 | 110 | 58 | 111 | 106 | 105 | 110 | 188 | 91 | 76 | 67 |
| 196 | 168 | 246 | 165 | 205 | 184 | 250 | 157 | 98 | 179 | 266 | 142 |
| 76 | 89 | 118 | 67 | 85 | 96 | 166 | 66 | 51 | 14 | 89 | 42 |
| 760 | 555 | 760 | 570 | 855 | 760 | 925 | 1,140 | 760 | 570 | 555 | 380 |
| 950 | 1,130 | 1,425 | 855 | 1,213 | 1,500 | 2,420 | 1,520 | 1,520 | 950 | 1,110 | 570 |
| 156 | 150 | 213 | 161 | 194 | 159 | 190 | 149 | 161 | 136 | 160 | 130 |
| 89 | 89 | 119 | 82 | 126 | 88 | 109 | 90 | 106 | 85 | 93 | 64 |
| 46 | 24 | 69 | 48 | 38 | 35 | 46 | 28 | 24 | 46 | 23 | 36 |
| 93 | 95 | 185 | 95 | 95 | 95 | 95 | 93 | 93 | 95 | 93 | 93 |
| 185 | 139 | 190 | 185 | 190 | 185 | 185 | 143 | 185 | 95 | 185 | 95 |
| 190 | 190 | 278 | 190 | 190 | 190 | 190 | 190 | 190 | 190 | 190 | 190 |
| 278 | 190 | 380 | 190 | 380 | 278 | 370 | 238 | 370 | 190 | 285 | 190 |
| 370 | 380 | 380 | 370 | 380 | 370 | 380 | 370 | 380 | 380 | 370 | 190 |
| 380 | 380 | 571 | 380 | 571 | 380 | 570 | 380 | 380 | 380 | 416 | 278 |
| 570 | 555 | 740 | 570 | 760 | 570 | 740 | 760 | 555 | 380 | 416 | 380 |
| .8 | 9.9 | .9 | 2.6 | 3.8 | 11.2 | 4.6 | 11.3 | 0 | 0 | 0 | 0 |
| .2 | 2.4 | .4 | 1.1 | .7 | 1.5 | .7 | 1.1 | 0 | 0 | 0 | 0 |

TABLE 6.02.--BAKED BEANS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 10.4 | 2.4 | 10.9 | 13.4 | 14.0 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 9.3 | 1.8 | 9.7 | 11.8 | 12.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.0 | .2 | 1.1 | 1.5 | 1.5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | .4 | .1 | .1 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 1.4 | 1.1 | 1.1 | 1.2 |
| STANDARD DEVIATION.....DO..... | .4 | .8 | .4 | .4 | .4 |
| MAXIMUM.....DO..... | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 67 | 32 | 37 | 42 | 53 |
| STANDARD DEVIATION.....GM..... | 53 | 37 | 35 | 29 | 39 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 11 | | 5 | 5 | 11 |
| 25TH.....GM..... | 43 | | 16 | 21 | 32 |
| 50TH.....GM..... | 43 | | 39 | 43 | 43 |
| 75TH.....GM..... | 85 | | 43 | 53 | 79 |
| 90TH.....GM..... | 128 | | 76 | 85 | 85 |
| 95TH.....GM..... | 170 | | 85 | 85 | 128 |
| 99TH.....GM..... | 298 | | 255 | 128 | 170 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 179 | 60 | 100 | 117 | 141 |
| STANDARD DEVIATION.....GM..... | 121 | 43 | 100 | 81 | 84 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 172 | 48 | 95 | 93 | 132 |
| STANDARD DEVIATION.....GM..... | 112 | 0 | 51 | 48 | 70 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 223 | 84 | 75 | 82 | 167 |
| STANDARD DEVIATION.....GM..... | 145 | 57 | 0 | 64 | 102 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,140 | 190 | 765 | 510 | 510 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,710 | 475 | 765 | 510 | 1,140 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 177 | 68 | 97 | 110 | 138 |
| STANDARD DEVIATION.....GM..... | 119 | 52 | 92 | 78 | 84 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 32 | | 16 | 16 | 31 |
| 25TH.....GM..... | 128 | | 43 | 64 | 80 |
| 50TH.....GM..... | 128 | | 84 | 127 | 128 |
| 75TH.....GM..... | 255 | | 128 | 128 | 158 |
| 90TH.....GM..... | 255 | | 128 | 255 | 255 |
| 95TH.....GM..... | 383 | | 255 | 255 | 255 |
| 99TH.....GM..... | 510 | | 765 | 383 | 383 |
| MAXIMUM.....GM..... | 1,020 | 190 | 765 | 510 | 510 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 28.0 | 0 | 0 | 1.6 | 4.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | 0 | 0 | .6 | 1.4 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 14.7 | 13.5 | 12.2 | 9.9 | 11.2 | 9.6 | 12.2 | 8.0 | 8.6 | 5.5 | 9.2 | 5.3 |
| 12.7 | 12.1 | 11.3 | 9.2 | 9.7 | 8.6 | 10.5 | 7.5 | 8.1 | 4.6 | 8.4 | 5.1 |
| 1.9 | 1.1 | .7 | .7 | 1.3 | .9 | 1.5 | .5 | .4 | .9 | .8 | .2 |
| .2 | .3 | .2 | 0 | .1 | .1 | .2 | (2) | .1 | 0 | 0 | 0 |
| 1.2 | 1.1 | 1.1 | 1.1 | 1.2 | 1.1 | 1.2 | 1.1 | 1.1 | 1.2 | 1.1 | 1.0 |
| .4 | .4 | .4 | .3 | .4 | .4 | .4 | .3 | .3 | .4 | .4 | .2 |
| 3.0 | 4.0 | 4.0 | 3.0 | 4.0 | 3.0 | 4.0 | 3.0 | 3.0 | 2.0 | 3.0 | 2.0 |
| 75 | 64 | 75 | 59 | 86 | 60 | 89 | 54 | 74 | 67 | 88 | 53 |
| 60 | 57 | 59 | 36 | 62 | 42 | 67 | 36 | 44 | 34 | 49 | 35 |
| 21 | 11 | 11 | 11 | 21 | 11 | 21 | 11 | 16 | 11 | | 5 |
| 43 | 43 | 43 | 43 | 43 | 43 | 43 | 32 | 43 | 43 | | 43 |
| 64 | 43 | 64 | 43 | 85 | 43 | 85 | 43 | 85 | 85 | | 43 |
| 85 | 85 | 85 | 85 | 85 | 85 | 85 | 85 | 85 | 85 | | 75 |
| 128 | 127 | 128 | 106 | 170 | 85 | 170 | 85 | 128 | 92 | | 85 |
| 170 | 170 | 170 | 128 | 170 | 128 | 234 | 127 | 170 | 127 | | 85 |
| 340 | 340 | 298 | 170 | 340 | 221 | 340 | 170 | 213 | 190 | | 170 |
| 194 | 171 | 202 | 163 | 230 | 166 | 227 | 152 | 213 | 183 | 233 | 144 |
| 127 | 124 | 125 | 98 | 141 | 114 | 138 | 96 | 116 | 85 | 107 | 76 |
| 180 | 136 | 191 | 172 | 222 | 141 | 221 | 148 | 168 | 148 | 291 | 255 |
| 98 | 75 | 96 | 55 | 155 | 84 | 130 | 72 | 111 | 64 | 49 | 0 |
| 359 | 238 | 337 | 0 | 174 | 116 | 340 | 223 | 212 | 0 | 0 | 0 |
| 160 | 167 | 175 | 0 | 60 | 50 | 93 | 32 | 0 | 0 | 0 | 0 |
| 1,020 | 1,020 | 765 | 510 | 1,140 | 1,020 | 1,020 | 765 | 510 | 383 | 510 | 255 |
| 1,710 | 1,520 | 1,710 | 638 | 1,520 | 1,020 | 1,290 | 765 | 638 | 570 | 765 | 510 |
| 194 | 168 | 203 | 164 | 222 | 158 | 226 | 151 | 207 | 171 | 238 | 153 |
| 129 | 123 | 129 | 94 | 137 | 98 | 137 | 92 | 120 | 82 | 100 | 79 |
| 64 | 32 | 32 | 32 | 64 | 32 | 64 | 32 | 32 | 32 | | 16 |
| 128 | 84 | 128 | 128 | 128 | 120 | 128 | 84 | 128 | 128 | | 128 |
| 128 | 128 | 190 | 128 | 255 | 128 | 255 | 128 | 191 | 143 | | 128 |
| 255 | 255 | 255 | 255 | 255 | 255 | 255 | 255 | 255 | 255 | | 255 |
| 255 | 255 | 383 | 255 | 383 | 255 | 383 | 255 | 383 | 255 | | 255 |
| 383 | 383 | 510 | 319 | 510 | 281 | 510 | 255 | 510 | 255 | | 255 |
| 638 | 570 | 571 | 510 | 637 | 510 | 571 | 510 | 510 | 380 | | 255 |
| 1,020 | 1,020 | 765 | 510 | 1,020 | 663 | 1,020 | 765 | 510 | 383 | 510 | 255 |
| .6 | 3.5 | 0 | 2.1 | 3.0 | 6.7 | 5.0 | 1.3 | 0 | 0 | 0 | 0 |
| .2 | 1.0 | 0 | 1.3 | .6 | 1.2 | .7 | .2 | 0 | 0 | 0 | 0 |

TABLE 6.03.--NUTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.0 | 0 | 1.9 | 3.2 | 5.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.9 | 0 | 1.3 | 2.4 | 4.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .8 | 0 | .5 | .8 | .8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | .1 | 0 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 0 | 1.3 | 1.3 | 1.3 |
| STANDARD DEVIATION.....DO..... | .7 | 0 | .5 | .5 | .6 |
| MAXIMUM.....DO..... | 8.0 | 0 | 3.0 | 3.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 19 | 0 | 8 | 17 | 18 |
| STANDARD DEVIATION.....GM..... | 21 | 0 | 8 | 19 | 23 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 2 | | | 1 | 2 |
| 25TH.....GM..... | 6 | | | 6 | 6 |
| 50TH.....GM..... | 12 | | | 11 | 12 |
| 75TH.....GM..... | 24 | | | 24 | 24 |
| 90TH.....GM..... | 48 | | | 37 | 43 |
| 95TH.....GM..... | 57 | | | 57 | 52 |
| 99TH.....GM..... | 109 | | | 117 | 121 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 44 | 0 | 20 | 33 | 47 |
| STANDARD DEVIATION.....GM..... | 46 | 0 | 18 | 33 | 60 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 52 | 0 | 15 | 51 | 44 |
| STANDARD DEVIATION.....GM..... | 44 | 0 | 9 | 39 | 49 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 43 | 0 | 36 | 0 | 42 |
| STANDARD DEVIATION.....GM..... | 30 | 0 | 0 | 0 | 9 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 432 | 0 | 57 | 288 | 363 |
| MAXIMUM OVER 3 DAYS.....GM..... | 495 | 0 | 108 | 351 | 363 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 43 | 0 | 19 | 39 | 44 |
| STANDARD DEVIATION.....GM..... | 43 | 0 | 16 | 36 | 48 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 5 | | | 4 | 6 |
| 25TH.....GM..... | 16 | | | 14 | 17 |
| 50TH.....GM..... | 32 | | | 28 | 28 |
| 75TH.....GM..... | 57 | | | 57 | 47 |
| 90TH.....GM..... | 85 | | | 85 | 85 |
| 95TH.....GM..... | 144 | | | 113 | 144 |
| 99TH.....GM..... | 189 | | | 144 | 252 |
| MAXIMUM.....GM..... | 426 | 0 | 57 | 144 | 288 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 114.2 | 0 | .8 | .9 | 8.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 4.5 | 0 | 2.8 | 1.3 | 6.7 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.9 | 4.1 | 3.2 | 3.2 | 6.7 | 5.1 | 6.8 | 5.8 | 4.6 | 4.6 | 2.0 | 3.0 |
| 3.3 | 3.6 | 2.8 | 2.8 | 5.3 | 4.2 | 5.2 | 4.4 | 3.7 | 2.9 | 1.5 | 2.2 |
| .5 | .4 | .3 | .4 | 1.1 | .7 | 1.1 | 1.0 | .7 | 1.2 | 0 | .7 |
| .1 | (2) | 0 | .1 | .3 | .2 | .4 | .4 | .2 | .5 | .5 | 0 |
| 1.2 | 1.2 | 1.1 | 1.2 | 1.3 | 1.3 | 1.3 | 1.4 | 1.2 | 1.6 | 2.0 | 1.2 |
| .5 | .5 | .3 | .5 | .6 | .6 | .7 | .9 | .5 | .9 | 1.8 | .4 |
| 4.0 | 4.0 | 2.0 | 3.0 | 4.0 | 4.0 | 5.0 | 8.0 | 3.0 | 5.0 | 6.0 | 2.0 |
| 22 | 16 | 22 | 18 | 24 | 18 | 23 | 17 | 13 | 13 | 22 | 12 |
| 30 | 19 | 19 | 18 | 24 | 19 | 24 | 19 | 9 | 12 | 25 | 22 |
| 3 | 2 | | | 2 | 2 | 2 | 1 | 2 | 2 | | |
| 6 | 6 | | | 9 | 6 | 7 | 6 | 6 | 4 | | |
| 12 | 11 | | | 15 | 12 | 16 | 12 | 12 | 9 | | |
| 24 | 23 | | | 30 | 24 | 24 | 24 | 18 | 19 | | |
| 48 | 42 | | | 48 | 48 | 57 | 44 | 28 | 30 | | |
| 96 | 48 | | | 57 | 57 | 72 | 48 | 30 | 33 | | |
| 144 | 120 | | | 142 | 78 | 109 | 93 | 36 | 62 | | |
| 45 | 38 | 59 | 41 | 58 | 49 | 44 | 39 | 31 | 31 | 24 | 17 |
| 51 | 35 | 58 | 35 | 60 | 54 | 33 | 40 | 24 | 29 | 16 | 14 |
| 97 | 71 | 59 | 58 | 62 | 38 | 77 | 39 | 28 | 19 | 0 | 49 |
| 71 | 55 | 11 | 39 | 50 | 21 | 48 | 26 | 14 | 14 | 0 | 55 |
| 62 | 31 | 0 | 72 | 39 | 38 | 54 | 41 | 30 | 28 | 64 | 0 |
| 35 | 0 | 0 | 0 | 24 | 24 | 28 | 36 | 2 | 18 | 6 | 0 |
| 432 | 216 | 340 | 221 | 426 | 324 | 288 | 280 | 108 | 145 | 85 | 144 |
| 495 | 360 | 340 | 246 | 432 | 324 | 435 | 424 | 108 | 185 | 218 | 288 |
| 54 | 43 | 59 | 44 | 55 | 44 | 51 | 36 | 30 | 24 | 34 | 30 |
| 57 | 41 | 53 | 36 | 54 | 43 | 41 | 36 | 21 | 23 | 19 | 39 |
| 6 | 6 | | | 6 | 5 | 7 | 5 | 5 | 4 | | |
| 18 | 18 | | | 18 | 14 | 24 | 9 | 17 | 6 | | |
| 36 | 32 | | | 36 | 32 | 36 | 28 | 27 | 14 | | |
| 63 | 57 | | | 72 | 57 | 72 | 48 | 36 | 33 | | |
| 144 | 85 | | | 144 | 85 | 113 | 72 | 57 | 57 | | |
| 144 | 144 | | | 144 | 144 | 144 | 112 | 72 | 72 | | |
| 288 | 216 | | | 227 | 216 | 168 | 168 | 108 | 136 | | |
| 288 | 216 | 340 | 166 | 426 | 233 | 288 | 240 | 108 | 136 | 73 | 144 |
| 5.7 | 8.4 | 2.8 | 3.4 | 15.1 | 14.5 | 19.7 | 26.4 | 1.1 | 6.3 | 0 | 0 |
| 5.6 | 8.3 | 5.8 | 5.9 | 4.5 | 4.2 | 4.4 | 4.5 | 1.7 | 4.8 | 0 | 0 |

TABLE 6.04.--PEANUTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.8 | 0 | 1.1 | 2.2 | 4.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.3 | 0 | .8 | 1.8 | 3.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | 0 | .2 | .4 | .5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | .1 | 0 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.3 | 0 | 1.3 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .6 | 0 | .6 | .5 | .5 |
| MAXIMUM.....DO..... | 4.0 | 0 | 3.0 | 3.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 22 | 0 | 11 | 15 | 20 |
| STANDARD DEVIATION.....GM..... | 23 | 0 | 9 | 20 | 24 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 3 | | | | 5 |
| 25TH.....GM..... | 9 | | | | 6 |
| 50TH.....GM..... | 14 | | | | 12 |
| 75TH.....GM..... | 24 | | | | 24 |
| 90TH.....GM..... | 48 | | | | 48 |
| 95TH.....GM..... | 63 | | | | 84 |
| 99TH.....GM..... | 120 | | | | 121 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 52 | 0 | 26 | 34 | 54 |
| STANDARD DEVIATION.....GM..... | 46 | 0 | 17 | 34 | 64 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 62 | 0 | 21 | 53 | 47 |
| STANDARD DEVIATION.....GM..... | 51 | 0 | 10 | 54 | 58 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 55 | 0 | 36 | 0 | 34 |
| STANDARD DEVIATION.....GM..... | 35 | 0 | 0 | 0 | 3 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 432 | 0 | 57 | 288 | 363 |
| MAXIMUM OVER 3 DAYS.....GM..... | 495 | 0 | 108 | 351 | 363 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 53 | 0 | 26 | 38 | 49 |
| STANDARD DEVIATION.....GM..... | 44 | 0 | 16 | 37 | 53 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 9 | | | | 9 |
| 25TH.....GM..... | 20 | | | | 17 |
| 50TH.....GM..... | 36 | | | | 36 |
| 75TH.....GM..... | 72 | | | | 63 |
| 90TH.....GM..... | 113 | | | | 111 |
| 95TH.....GM..... | 144 | | | | 144 |
| 99TH.....GM..... | 216 | | | | 288 |
| MAXIMUM.....GM..... | 340 | 0 | 57 | 144 | 288 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 40.0 | 0 | .8 | 0 | 3.4 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 3.1 | 0 | 5.1 | 0 | 3.5 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 2.7 | 2.7 | 1.7 | 1.6 | 3.6 | 2.6 | 3.8 | 2.9 | 3.1 | 1.7 | .9 | 1.5 |
| 2.2 | 2.3 | 1.4 | 1.3 | 2.9 | 2.4 | 3.0 | 2.4 | 2.2 | 1.2 | .9 | 1.0 |
| .5 | .3 | .3 | .2 | .6 | .1 | .7 | .3 | .7 | .4 | 0 | .5 |
| 0 | (2) | 0 | 0 | .1 | (2) | .2 | .2 | .2 | .1 | 0 | 0 |
| 1.2 | 1.2 | 1.2 | 1.2 | 1.3 | 1.1 | 1.3 | 1.3 | 1.4 | 1.4 | 1.0 | 1.3 |
| .5 | .5 | .4 | .5 | .5 | .4 | .6 | .7 | .6 | .7 | .0 | .5 |
| 3.0 | 4.0 | 2.0 | 3.0 | 3.0 | 4.0 | 4.0 | 4.0 | 3.0 | 4.0 | 1.0 | 2.0 |
| 27 | 18 | 29 | 18 | 27 | 21 | 26 | 20 | 15 | 17 | 12 | 17 |
| 32 | 21 | 22 | 16 | 23 | 20 | 24 | 21 | 9 | 15 | 4 | 30 |
| 3 | 2 | | | 6 | 3 | 5 | 3 | | | | |
| 9 | 6 | | | 12 | 6 | 12 | 6 | | | | |
| 12 | 12 | | | 23 | 12 | 21 | 12 | | | | |
| 32 | 24 | | | 36 | 24 | 25 | 24 | | | | |
| 64 | 48 | | | 48 | 48 | 57 | 48 | | | | |
| 96 | 48 | | | 63 | 60 | 72 | 48 | | | | |
| 165 | 120 | | | 144 | 108 | 144 | 141 | | | | |
| 53 | 42 | 81 | 44 | 61 | 58 | 53 | 48 | 38 | 37 | 35 | 18 |
| 54 | 36 | 68 | 32 | 41 | 59 | 31 | 41 | 25 | 37 | 13 | 20 |
| 97 | 73 | 60 | 63 | 79 | 56 | 80 | 39 | 28 | 30 | 0 | 63 |
| 71 | 61 | 12 | 32 | 52 | 16 | 51 | 24 | 14 | 13 | 0 | 65 |
| 0 | 31 | 0 | 0 | 60 | 70 | 55 | 60 | 30 | 61 | 0 | 0 |
| 0 | 0 | 0 | 0 | 15 | 4 | 35 | 45 | 2 | 0 | 0 | 0 |
| 432 | 216 | 340 | 144 | 288 | 324 | 288 | 280 | 108 | 145 | 57 | 144 |
| 495 | 360 | 340 | 216 | 432 | 324 | 435 | 424 | 108 | 185 | 57 | 288 |
| 64 | 47 | 75 | 47 | 64 | 56 | 59 | 47 | 33 | 34 | 35 | 39 |
| 61 | 44 | 59 | 34 | 42 | 47 | 41 | 39 | 21 | 30 | 13 | 52 |
| 12 | 6 | | | 17 | 8 | 16 | 9 | | | | |
| 28 | 18 | | | 36 | 19 | 28 | 18 | | | | |
| 48 | 32 | | | 57 | 36 | 48 | 36 | | | | |
| 72 | 48 | | | 72 | 72 | 72 | 72 | | | | |
| 144 | 126 | | | 144 | 126 | 113 | 108 | | | | |
| 144 | 144 | | | 144 | 144 | 144 | 144 | | | | |
| 288 | 216 | | | 189 | 233 | 168 | 168 | | | | |
| 288 | 216 | 340 | 144 | 189 | 233 | 288 | 170 | 108 | 136 | 57 | 144 |
| 4.7 | 6.0 | 1.1 | 1.0 | 3.5 | 1.8 | 5.6 | 11.0 | 1.1 | 0 | 0 | 0 |
| 6.8 | 8.9 | 4.1 | 3.5 | 2.0 | 1.1 | 2.3 | 4.2 | 2.3 | 0 | 0 | 0 |

TABLE 6.05.--PEANUT BUTTER
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 15.2 | 2.5 | 24.7 | 32.4 | 32.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 10.5 | 2.1 | 15.8 | 20.7 | 22.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.3 | .3 | 7.1 | 8.3 | 7.1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.4 | .2 | 1.8 | 3.4 | 3.2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1</u> /NUMBER..... | 1.5 | 1.2 | 1.5 | 1.5 | 1.5 |
| STANDARD DEVIATION.....DO..... | .8 | .6 | .7 | .8 | .8 |
| MAXIMUM.....DO..... | 8.0 | 3.0 | 5.0 | 6.0 | 8.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1</u> /GM..... | 13 | 8 | 10 | 11 | 13 |
| STANDARD DEVIATION.....GM..... | 13 | 4 | 10 | 10 | 11 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> / | | | | | |
| 5TH.....GM..... | 2 | | 2 | 2 | 2 |
| 25TH.....GM..... | 5 | | 4 | 5 | 5 |
| 50TH.....GM..... | 11 | | 5 | 11 | 11 |
| 75TH.....GM..... | 16 | | 11 | 14 | 16 |
| 90TH.....GM..... | 27 | | 22 | 22 | 27 |
| 95TH.....GM..... | 34 | | 27 | 32 | 32 |
| 99TH.....GM..... | 65 | | 51 | 43 | 57 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 29 | 21 | 20 | 24 | 26 |
| STANDARD DEVIATION.....GM..... | 25 | 8 | 17 | 16 | 20 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 29 | 24 | 20 | 23 | 28 |
| STANDARD DEVIATION.....GM..... | 20 | 0 | 13 | 14 | 17 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 29 | 9 | 21 | 25 | 26 |
| STANDARD DEVIATION.....GM..... | 20 | 0 | 21 | 17 | 16 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 454 | 32 | 194 | 194 | 161 |
| MAXIMUM OVER 3 DAYS.....GM..... | 454 | 48 | 274 | 323 | 242 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1</u> /GM..... | 27 | 19 | 20 | 23 | 25 |
| STANDARD DEVIATION.....GM..... | 22 | 9 | 18 | 16 | 19 |
| PERCENTILES: <u>1</u> / | | | | | |
| 5TH.....GM..... | 5 | | 5 | 5 | 5 |
| 25TH.....GM..... | 16 | | 8 | 16 | 16 |
| 50TH.....GM..... | 16 | | 16 | 16 | 16 |
| 75TH.....GM..... | 32 | | 32 | 32 | 32 |
| 90TH.....GM..... | 48 | | 32 | 32 | 48 |
| 95TH.....GM..... | 65 | | 48 | 48 | 65 |
| 99TH.....GM..... | 129 | | 98 | 65 | 97 |
| MAXIMUM.....GM..... | 454 | 32 | 194 | 194 | 161 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> / | | | | | |
| EATING OCCASIONS.....NUMBER..... | 229.7 | 0 | 6.1 | 13.0 | 32.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.7 | 0 | 1.6 | 1.5 | 3.7 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 27.0 | 24.1 | 19.3 | 12.5 | 12.8 | 12.2 | 10.6 | 9.2 | 12.9 | 9.3 | 10.1 | 10.7 |
| 18.7 | 16.9 | 12.4 | 9.1 | 8.6 | 9.2 | 6.8 | 7.3 | 8.2 | 6.3 | 5.4 | 7.3 |
| 5.9 | 5.4 | 5.4 | 2.4 | 2.6 | 2.4 | 2.3 | 1.4 | 2.7 | 2.5 | 2.6 | 2.0 |
| 2.4 | 1.8 | 1.6 | 1.0 | 1.6 | .6 | 1.4 | .5 | 1.9 | .6 | 2.1 | 1.4 |
| 1.5 | 1.4 | 1.5 | 1.4 | 1.6 | 1.3 | 1.6 | 1.3 | 1.6 | 1.4 | 1.7 | 1.5 |
| .8 | .7 | .8 | .8 | .9 | .6 | .9 | .6 | 1.1 | .6 | .8 | .8 |
| 6.0 | 4.0 | 5.0 | 4.0 | 6.0 | 4.0 | 8.0 | 5.0 | 6.0 | 3.0 | 3.0 | 4.0 |
| 15 | 13 | 19 | 12 | 20 | 11 | 16 | 10 | 16 | 9 | 15 | 11 |
| 12 | 10 | 18 | 12 | 21 | 9 | 14 | 8 | 12 | 7 | 13 | 11 |
| 3 | 3 | 4 | 2 | 4 | 2 | 4 | 2 | 4 | 2 | 5 | 2 |
| 5 | 5 | 11 | 5 | 9 | 5 | 5 | 5 | 5 | 5 | 7 | 5 |
| 11 | 11 | 16 | 11 | 11 | 9 | 11 | 7 | 11 | 5 | 11 | 8 |
| 22 | 16 | 22 | 16 | 22 | 13 | 22 | 11 | 22 | 11 | 16 | 13 |
| 32 | 22 | 40 | 22 | 43 | 22 | 32 | 22 | 32 | 18 | 32 | 27 |
| 43 | 32 | 48 | 32 | 59 | 32 | 43 | 22 | 43 | 22 | 48 | 32 |
| 65 | 59 | 97 | 65 | 108 | 43 | 65 | 38 | 54 | 43 | 57 | 57 |
| 32 | 30 | 39 | 27 | 40 | 26 | 32 | 25 | 34 | 21 | 26 | 22 |
| 25 | 24 | 29 | 20 | 43 | 20 | 24 | 20 | 29 | 18 | 11 | 14 |
| 32 | 27 | 41 | 32 | 44 | 25 | 33 | 19 | 31 | 19 | 25 | 28 |
| 18 | 17 | 26 | 26 | 28 | 15 | 27 | 11 | 15 | 12 | 15 | 20 |
| 31 | 24 | 45 | 25 | 42 | 26 | 28 | 22 | 26 | 12 | 32 | 21 |
| 18 | 12 | 32 | 12 | 31 | 17 | 15 | 15 | 15 | 6 | 16 | 16 |
| 194 | 258 | 290 | 145 | 454 | 129 | 194 | 145 | 145 | 145 | 97 | 98 |
| 258 | 258 | 387 | 258 | 454 | 199 | 290 | 274 | 161 | 145 | 170 | 170 |
| 30 | 27 | 38 | 26 | 38 | 25 | 30 | 23 | 29 | 19 | 28 | 22 |
| 22 | 20 | 26 | 21 | 35 | 19 | 23 | 18 | 22 | 16 | 14 | 18 |
| 5 | 5 | 11 | 5 | 11 | 5 | 5 | 5 | 5 | 5 | 11 | 5 |
| 16 | 16 | 16 | 16 | 16 | 16 | 16 | 16 | 16 | 11 | 16 | 8 |
| 32 | 24 | 32 | 16 | 32 | 16 | 32 | 16 | 32 | 16 | 28 | 16 |
| 32 | 32 | 48 | 32 | 48 | 32 | 32 | 32 | 32 | 31 | 32 | 32 |
| 65 | 48 | 65 | 48 | 65 | 48 | 57 | 32 | 52 | 32 | 48 | 48 |
| 65 | 65 | 97 | 65 | 93 | 65 | 65 | 65 | 65 | 48 | 57 | 57 |
| 129 | 97 | 129 | 129 | 145 | 97 | 129 | 97 | 129 | 65 | 57 | 98 |
| 194 | 258 | 129 | 145 | 454 | 129 | 194 | 145 | 145 | 145 | 65 | 98 |
| 22.6 | 23.1 | 10.1 | 10.7 | 27.7 | 14.3 | 28.7 | 28.6 | 5.1 | 1.5 | 1.7 | 3.7 |
| 2.7 | 3.1 | 2.5 | 4.1 | 3.5 | 1.7 | 3.5 | 3.4 | 2.2 | .6 | 1.8 | 2.3 |

TABLE 7.01.--MEAT, POULTRY, FISH, AND THEIR MIXTURES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1: | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 98.6 | 55.0 | 98.5 | 99.3 | 99.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.8 | 13.0 | 4.4 | 3.3 | 1.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 14.2 | 13.2 | 20.5 | 19.0 | 16.6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 81.5 | 28.8 | 73.7 | 77.0 | 81.4 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 4.8 | 3.3 | 4.3 | 4.5 | 4.7 |
| STANDARD DEVIATION.....DO..... | 1.7 | 1.8 | 1.7 | 1.6 | 1.5 |
| MAXIMUM.....DO..... | 12.0 | 10.0 | 9.0 | 9.0 | 10.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 193 | 87 | 96 | 114 | 146 |
| STANDARD DEVIATION.....GM..... | 114 | 77 | 59 | 64 | 75 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 51 | 8 | 23 | 33 | 48 |
| 25TH.....GM..... | 113 | 37 | 54 | 70 | 93 |
| 50TH.....GM..... | 172 | 64 | 82 | 102 | 133 |
| 75TH.....GM..... | 249 | 128 | 125 | 145 | 185 |
| 90TH.....GM..... | 338 | 191 | 172 | 199 | 244 |
| 95TH.....GM..... | 405 | 255 | 216 | 233 | 283 |
| 99TH.....GM..... | 562 | 336 | 283 | 309 | 379 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 146 | 84 | 73 | 125 | 129 |
| STANDARD DEVIATION.....GM..... | 133 | 62 | 57 | 109 | 135 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 173 | 104 | 102 | 111 | 136 |
| STANDARD DEVIATION.....GM..... | 113 | 64 | 69 | 64 | 72 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 211 | 122 | 108 | 127 | 159 |
| STANDARD DEVIATION.....GM..... | 112 | 85 | 58 | 63 | 74 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 2,350 | 638 | 980 | 745 | 992 |
| MAXIMUM OVER 3 DAYS.....GM..... | 4,010 | 1,696 | 1,116 | 1,435 | 1,720 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 121 | 80 | 66 | 77 | 93 |
| STANDARD DEVIATION.....GM..... | 107 | 53 | 57 | 64 | 76 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 16 | 14 | 8 | 14 | 16 |
| 25TH.....GM..... | 56 | 40 | 28 | 30 | 44 |
| 50TH.....GM..... | 97 | 63 | 56 | 57 | 84 |
| 75TH.....GM..... | 162 | 115 | 85 | 97 | 112 |
| 90TH.....GM..... | 235 | 133 | 125 | 142 | 180 |
| 95TH.....GM..... | 324 | 184 | 170 | 205 | 240 |
| 99TH.....GM..... | 510 | 224 | 268 | 336 | 383 |
| MAXIMUM.....GM..... | 1,792 | 425 | 528 | 720 | 960 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 2704.5 | 5.9 | 52.3 | 103.2 | 127.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.5 | .7 | 1.2 | 1.4 | 1.5 |

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 99.7 | 99.6 | 99.8 | 99.5 | 99.3 | 98.3 | 99.5 | 99.3 | 98.6 | 98.3 | 98.8 | 98.4 |
| 1.9 | 2.5 | 1.4 | 3.5 | 1.8 | 3.9 | 1.0 | 2.9 | 2.6 | 4.3 | 3.7 | 5.1 |
| 13.9 | 16.9 | 11.4 | 17.5 | 10.6 | 17.3 | 7.4 | 14.8 | 10.7 | 16.9 | 11.1 | 18.0 |
| 83.8 | 80.2 | 87.0 | 78.5 | 86.9 | 77.0 | 91.0 | 81.6 | 85.3 | 77.0 | 84.0 | 75.3 |
| 4.9 | 4.6 | 5.3 | 4.6 | 5.3 | 4.5 | 5.5 | 4.6 | 5.0 | 4.2 | 4.7 | 4.1 |
| 1.6 | 1.6 | 1.7 | 1.6 | 1.7 | 1.6 | 1.7 | 1.6 | 1.8 | 1.6 | 1.8 | 1.6 |
| 10.0 | 11.0 | 12.0 | 10.0 | 12.0 | 10.0 | 12.0 | 10.0 | 12.0 | 9.0 | 10.0 | 9.0 |
| 191 | 162 | 253 | 175 | 277 | 179 | 261 | 177 | 207 | 153 | 182 | 136 |
| 92 | 86 | 126 | 89 | 143 | 96 | 122 | 93 | 108 | 87 | 100 | 79 |
| 65 | 49 | 84 | 56 | 94 | 52 | 94 | 53 | 64 | 43 | 51 | 35 |
| 124 | 104 | 170 | 112 | 180 | 110 | 174 | 113 | 131 | 93 | 112 | 85 |
| 178 | 148 | 235 | 162 | 254 | 164 | 244 | 163 | 189 | 141 | 166 | 124 |
| 243 | 204 | 311 | 222 | 343 | 228 | 323 | 225 | 266 | 196 | 230 | 173 |
| 310 | 273 | 417 | 289 | 453 | 304 | 423 | 301 | 343 | 258 | 332 | 229 |
| 363 | 319 | 498 | 343 | 537 | 353 | 489 | 354 | 407 | 312 | 374 | 297 |
| 460 | 417 | 647 | 462 | 755 | 473 | 639 | 456 | 557 | 440 | 490 | 410 |
| 171 | 111 | 215 | 141 | 192 | 173 | 225 | 129 | 144 | 129 | 233 | 127 |
| 138 | 89 | 191 | 124 | 119 | 140 | 156 | 96 | 119 | 121 | 391 | 96 |
| 184 | 154 | 225 | 171 | 269 | 170 | 237 | 174 | 182 | 145 | 169 | 138 |
| 110 | 86 | 125 | 93 | 167 | 93 | 143 | 105 | 102 | 84 | 130 | 89 |
| 205 | 178 | 270 | 194 | 294 | 199 | 271 | 193 | 223 | 172 | 195 | 152 |
| 88 | 83 | 123 | 85 | 139 | 93 | 119 | 89 | 104 | 85 | 94 | 76 |
| 1,536 | 1,376 | 1,708 | 1,225 | 2,350 | 1,329 | 1,957 | 1,501 | 1,360 | 1,869 | 1,502 | 1,128 |
| 2,018 | 2,791 | 2,957 | 1,879 | 4,010 | 2,260 | 3,401 | 2,695 | 2,627 | 2,360 | 1,665 | 1,540 |
| 117 | 105 | 143 | 115 | 158 | 120 | 143 | 117 | 125 | 108 | 116 | 99 |
| 97 | 87 | 123 | 92 | 135 | 99 | 123 | 99 | 106 | 92 | 107 | 83 |
| 17 | 20 | 24 | 19 | 24 | 16 | 18 | 16 | 16 | 16 | 16 | 14 |
| 56 | 54 | 57 | 56 | 64 | 56 | 57 | 56 | 56 | 54 | 44 | 43 |
| 90 | 85 | 112 | 97 | 113 | 97 | 112 | 92 | 108 | 88 | 90 | 85 |
| 148 | 119 | 180 | 142 | 220 | 160 | 184 | 150 | 168 | 135 | 150 | 113 |
| 227 | 210 | 273 | 224 | 314 | 227 | 278 | 227 | 235 | 224 | 227 | 218 |
| 291 | 250 | 368 | 270 | 442 | 304 | 368 | 288 | 340 | 252 | 294 | 245 |
| 490 | 480 | 612 | 480 | 705 | 490 | 600 | 490 | 490 | 480 | 490 | 460 |
| 1,020 | 960 | 1,596 | 1,225 | 1,792 | 1,275 | 1,596 | 1,440 | 1,020 | 1,588 | 1,470 | 750 |
| 178.7 | 121.3 | 101.2 | 119.3 | 295.9 | 325.6 | 495.3 | 514.0 | 69.2 | 115.8 | 30.6 | 48.7 |
| 1.7 | 1.2 | 1.4 | 1.8 | 1.4 | 1.4 | 1.8 | 1.6 | 1.3 | 1.6 | 1.2 | 1.2 |

TABLE 7.02.--MEAT, POULTRY, AND FISH
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 96.9 | 35.5 | 95.8 | 97.7 | 98.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 8.8 | 12.7 | 12.1 | 9.5 | 9.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 27.2 | 7.5 | 29.8 | 31.6 | 32.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 61.0 | 15.3 | 53.9 | 56.6 | 56.7 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 3.9 | 2.7 | 3.6 | 3.7 | 3.7 |
| STANDARD DEVIATION.....DO..... | 1.7 | 1.7 | 1.7 | 1.6 | 1.6 |
| MAXIMUM.....DO..... | 12.0 | 10.0 | 9.0 | 9.0 | 10.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 134 | 51 | 65 | 80 | 96 |
| STANDARD DEVIATION.....GM..... | 87 | 55 | 41 | 47 | 55 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 28 | 4 | 13 | 19 | 23 |
| 25TH.....GM..... | 73 | 16 | 33 | 47 | 56 |
| 50TH.....GM..... | 117 | 38 | 57 | 73 | 85 |
| 75TH.....GM..... | 176 | 61 | 87 | 103 | 127 |
| 90TH.....GM..... | 245 | 118 | 122 | 138 | 168 |
| 95TH.....GM..... | 296 | 151 | 142 | 165 | 200 |
| 99TH.....GM..... | 413 | 283 | 183 | 232 | 269 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 126 | 59 | 64 | 84 | 99 |
| STANDARD DEVIATION.....GM..... | 97 | 49 | 46 | 60 | 65 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 142 | 59 | 74 | 94 | 109 |
| STANDARD DEVIATION.....GM..... | 83 | 41 | 39 | 54 | 57 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 165 | 83 | 83 | 98 | 120 |
| STANDARD DEVIATION.....GM..... | 87 | 67 | 41 | 45 | 54 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 2,350 | 383 | 651 | 559 | 992 |
| MAXIMUM OVER 3 DAYS.....GM..... | 3,480 | 1,021 | 889 | 1,030 | 1,178 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 102 | 56 | 54 | 64 | 77 |
| STANDARD DEVIATION.....GM..... | 82 | 39 | 39 | 46 | 54 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 16 | 9 | 8 | 13 | 16 |
| 25TH.....GM..... | 44 | 28 | 28 | 28 | 40 |
| 50TH.....GM..... | 85 | 47 | 44 | 56 | 61 |
| 75TH.....GM..... | 129 | 78 | 71 | 85 | 110 |
| 90TH.....GM..... | 209 | 128 | 112 | 112 | 140 |
| 95TH.....GM..... | 234 | 128 | 113 | 144 | 170 |
| 99TH.....GM..... | 393 | 135 | 176 | 224 | 276 |
| MAXIMUM.....GM..... | 1,792 | 255 | 336 | 476 | 907 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 2152.3 | 4.0 | 43.2 | 90.4 | 97.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.5 | .8 | 1.2 | 1.4 | 1.4 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 98.5 | 98.1 | 97.8 | 97.2 | 98.4 | 96.3 | 98.9 | 97.9 | 98.1 | 97.2 | 97.4 | 96.1 |
| 8.7 | 9.3 | 7.3 | 12.3 | 7.7 | 10.8 | 4.4 | 8.9 | 8.3 | 9.3 | 5.4 | 10.7 |
| 31.1 | 33.6 | 27.4 | 29.8 | 25.6 | 29.4 | 19.9 | 27.5 | 21.0 | 28.1 | 22.4 | 26.2 |
| 58.7 | 55.2 | 63.1 | 55.2 | 65.0 | 56.0 | 74.6 | 61.5 | 68.9 | 59.8 | 69.6 | 59.2 |
| 3.9 | 3.7 | 4.2 | 3.6 | 4.2 | 3.6 | 4.6 | 3.8 | 4.3 | 3.7 | 4.2 | 3.6 |
| 1.7 | 1.6 | 1.8 | 1.6 | 1.8 | 1.6 | 1.8 | 1.7 | 1.8 | 1.6 | 1.8 | 1.6 |
| 10.0 | 9.0 | 12.0 | 9.0 | 11.0 | 9.0 | 12.0 | 10.0 | 12.0 | 9.0 | 10.0 | 9.0 |
| 125 | 106 | 164 | 117 | 186 | 122 | 187 | 127 | 150 | 112 | 138 | 102 |
| 71 | 62 | 95 | 71 | 108 | 73 | 99 | 75 | 85 | 65 | 83 | 61 |
| 29 | 26 | 38 | 24 | 46 | 28 | 54 | 28 | 37 | 28 | 28 | 24 |
| 74 | 62 | 96 | 65 | 113 | 68 | 117 | 75 | 88 | 65 | 76 | 56 |
| 113 | 98 | 149 | 107 | 168 | 110 | 173 | 115 | 140 | 103 | 123 | 95 |
| 163 | 139 | 211 | 154 | 240 | 160 | 238 | 166 | 194 | 150 | 192 | 135 |
| 219 | 181 | 283 | 208 | 320 | 217 | 313 | 223 | 260 | 201 | 241 | 176 |
| 259 | 207 | 328 | 245 | 375 | 257 | 366 | 264 | 295 | 233 | 295 | 212 |
| 340 | 288 | 475 | 371 | 509 | 345 | 490 | 372 | 426 | 321 | 364 | 335 |
| 126 | 107 | 159 | 128 | 181 | 129 | 178 | 121 | 130 | 107 | 131 | 111 |
| 94 | 75 | 103 | 112 | 143 | 89 | 109 | 87 | 78 | 61 | 165 | 75 |
| 137 | 120 | 175 | 126 | 208 | 137 | 186 | 141 | 159 | 120 | 124 | 110 |
| 69 | 60 | 95 | 66 | 107 | 70 | 89 | 79 | 89 | 58 | 55 | 56 |
| 154 | 134 | 197 | 151 | 219 | 153 | 211 | 155 | 176 | 140 | 163 | 127 |
| 69 | 62 | 93 | 67 | 108 | 73 | 97 | 72 | 80 | 63 | 82 | 60 |
| 952 | 1,376 | 1,411 | 952 | 2,350 | 1,072 | 1,438 | 1,501 | 937 | 688 | 897 | 1,128 |
| 1,455 | 2,605 | 2,957 | 1,427 | 3,480 | 2,028 | 2,208 | 2,005 | 1,854 | 1,274 | 1,563 | 1,311 |
| 96 | 86 | 117 | 96 | 132 | 100 | 122 | 99 | 105 | 92 | 98 | 85 |
| 70 | 64 | 90 | 72 | 106 | 75 | 95 | 78 | 80 | 67 | 78 | 63 |
| 16 | 16 | 16 | 16 | 24 | 16 | 16 | 16 | 16 | 14 | 14 | 13 |
| 46 | 44 | 56 | 48 | 56 | 48 | 56 | 45 | 50 | 44 | 36 | 38 |
| 85 | 84 | 97 | 85 | 112 | 85 | 112 | 85 | 86 | 84 | 85 | 82 |
| 113 | 112 | 168 | 113 | 170 | 124 | 168 | 123 | 140 | 113 | 126 | 112 |
| 180 | 168 | 224 | 170 | 255 | 194 | 227 | 184 | 224 | 170 | 210 | 168 |
| 224 | 196 | 280 | 224 | 336 | 227 | 288 | 225 | 227 | 224 | 227 | 218 |
| 336 | 302 | 448 | 344 | 480 | 368 | 448 | 368 | 392 | 336 | 368 | 304 |
| 896 | 960 | 1,344 | 896 | 1,792 | 960 | 1,131 | 1,440 | 768 | 624 | 678 | 564 |
| 148.8 | 94.8 | 79.4 | 93.9 | 225.3 | 265.9 | 393.7 | 406.4 | 47.1 | 85.9 | 29.2 | 46.6 |
| 1.9 | 1.2 | 1.4 | 1.8 | 1.4 | 1.4 | 1.8 | 1.5 | 1.0 | 1.4 | 1.3 | 1.3 |

TABLE 7.03.--BEEF, PORK, LAMB, AND VEAL
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.... | 37,900 | 524 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 84.6 | 23.2 | 78.2 | 82.8 | 84.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 29.8 | 12.7 | 33.3 | 34.5 | 34.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 33.8 | 4.7 | 31.2 | 33.0 | 35.6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 21.0 | 5.8 | 13.7 | 15.2 | 14.3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER.... | 2.4 | 2.0 | 2.1 | 2.1 | 2.1 |
| STANDARD DEVIATION.....DO..... | 1.3 | 1.4 | 1.2 | 1.1 | 1.1 |
| MAXIMUM.....DO..... | 10.0 | 8.0 | 8.0 | 7.0 | 9.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 85 | 39 | 37 | 46 | 57 |
| STANDARD DEVIATION.....GM..... | 67 | 49 | 28 | 34 | 41 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 11 | 3 | 5 | 5 | 8 |
| 25TH.....GM..... | 37 | 9 | 17 | 21 | 28 |
| 50TH.....GM..... | 69 | 24 | 29 | 37 | 48 |
| 75TH.....GM..... | 112 | 46 | 52 | 62 | 75 |
| 90TH.....GM..... | 168 | 85 | 77 | 88 | 111 |
| 95TH.....GM..... | 211 | 135 | 89 | 108 | 131 |
| 99TH.....GM..... | 317 | 340 | 121 | 161 | 206 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 128 | 56 | 61 | 75 | 96 |
| STANDARD DEVIATION.....GM..... | 98 | 47 | 45 | 50 | 74 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 133 | 53 | 64 | 82 | 99 |
| STANDARD DEVIATION.....GM..... | 79 | 28 | 37 | 47 | 52 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 139 | 88 | 63 | 75 | 95 |
| STANDARD DEVIATION.....GM..... | 83 | 74 | 33 | 38 | 48 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,792 | 383 | 418 | 559 | 624 |
| MAXIMUM OVER 3 DAYS.....GM..... | 3,328 | 1,021 | 579 | 856 | 972 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 107 | 58 | 53 | 66 | 82 |
| STANDARD DEVIATION.....GM..... | 85 | 42 | 40 | 46 | 55 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 16 | 9 | 8 | 8 | 14 |
| 25TH.....GM..... | 46 | 28 | 23 | 28 | 43 |
| 50TH.....GM..... | 86 | 47 | 46 | 56 | 84 |
| 75TH.....GM..... | 140 | 85 | 80 | 85 | 112 |
| 90TH.....GM..... | 224 | 128 | 112 | 112 | 142 |
| 95TH.....GM..... | 252 | 128 | 115 | 140 | 171 |
| 99TH.....GM..... | 432 | 135 | 170 | 224 | 280 |
| MAXIMUM.....GM..... | 1,792 | 255 | 280 | 448 | 504 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER.... | 1097.1 | 1.6 | 29.4 | 52.6 | 42.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.4 | .6 | 1.7 | 1.8 | 1.3 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 87.1 | 84.2 | 87.9 | 82.6 | 87.5 | 82.1 | 90.7 | 86.1 | 85.7 | 84.5 | 87.5 | 80.7 |
| 33.4 | 33.9 | 27.0 | 32.2 | 27.8 | 32.5 | 24.6 | 30.1 | 24.2 | 30.6 | 21.6 | 28.8 |
| 35.5 | 34.5 | 37.1 | 32.5 | 34.8 | 32.3 | 35.2 | 34.7 | 33.5 | 33.6 | 32.4 | 33.0 |
| 18.2 | 15.7 | 23.8 | 17.9 | 24.9 | 17.2 | 31.0 | 21.3 | 28.0 | 20.4 | 33.4 | 18.8 |
| 2.2 | 2.1 | 2.6 | 2.2 | 2.6 | 2.2 | 2.8 | 2.4 | 2.7 | 2.3 | 2.9 | 2.3 |
| 1.2 | 1.1 | 1.4 | 1.2 | 1.4 | 1.2 | 1.5 | 1.3 | 1.5 | 1.2 | 1.6 | 1.3 |
| 9.0 | 8.0 | 9.0 | 8.0 | 8.0 | 9.0 | 10.0 | 9.0 | 9.0 | 9.0 | 8.0 | 7.0 |
| 76 | 67 | 105 | 76 | 119 | 78 | 117 | 80 | 92 | 68 | 93 | 65 |
| 55 | 52 | 73 | 56 | 87 | 58 | 82 | 58 | 63 | 49 | 69 | 50 |
| 13 | 11 | 17 | 11 | 16 | 13 | 20 | 11 | 16 | 9 | 11 | 5 |
| 37 | 32 | 56 | 37 | 57 | 37 | 58 | 37 | 46 | 35 | 42 | 35 |
| 65 | 56 | 92 | 61 | 102 | 65 | 99 | 67 | 76 | 56 | 75 | 56 |
| 103 | 90 | 140 | 105 | 160 | 106 | 157 | 108 | 130 | 92 | 129 | 85 |
| 149 | 127 | 197 | 143 | 224 | 149 | 219 | 155 | 169 | 134 | 187 | 136 |
| 181 | 151 | 241 | 184 | 272 | 181 | 270 | 193 | 208 | 168 | 224 | 156 |
| 266 | 233 | 340 | 276 | 405 | 285 | 409 | 278 | 302 | 234 | 303 | 224 |
| 120 | 109 | 152 | 113 | 177 | 129 | 172 | 128 | 142 | 108 | 138 | 96 |
| 76 | 81 | 99 | 88 | 134 | 97 | 106 | 97 | 101 | 66 | 110 | 63 |
| 129 | 115 | 158 | 127 | 180 | 129 | 167 | 126 | 130 | 112 | 122 | 111 |
| 69 | 59 | 81 | 67 | 95 | 66 | 95 | 70 | 70 | 62 | 70 | 63 |
| 123 | 113 | 164 | 128 | 185 | 131 | 171 | 126 | 136 | 106 | 135 | 102 |
| 65 | 70 | 84 | 62 | 103 | 70 | 90 | 68 | 71 | 58 | 79 | 62 |
| 896 | 1,376 | 1,025 | 896 | 1,792 | 960 | 1,344 | 972 | 720 | 688 | 840 | 1,128 |
| 1,308 | 2,605 | 2,375 | 1,176 | 3,328 | 1,512 | 2,118 | 1,527 | 1,420 | 965 | 1,204 | 1,249 |
| 103 | 94 | 123 | 102 | 140 | 107 | 126 | 102 | 103 | 89 | 97 | 86 |
| 71 | 69 | 90 | 73 | 109 | 80 | 99 | 79 | 79 | 66 | 81 | 66 |
| 16 | 16 | 16 | 16 | 16 | 16 | 16 | 14 | 16 | 10 | 10 | 8 |
| 56 | 56 | 56 | 56 | 56 | 56 | 56 | 43 | 40 | 35 | 24 | 32 |
| 86 | 84 | 112 | 88 | 112 | 92 | 112 | 86 | 86 | 84 | 84 | 84 |
| 138 | 112 | 168 | 129 | 194 | 140 | 168 | 138 | 140 | 112 | 138 | 112 |
| 184 | 168 | 224 | 170 | 276 | 224 | 226 | 188 | 224 | 168 | 224 | 168 |
| 224 | 224 | 280 | 224 | 336 | 232 | 291 | 224 | 226 | 224 | 226 | 224 |
| 336 | 336 | 448 | 344 | 468 | 368 | 448 | 384 | 368 | 280 | 368 | 322 |
| 896 | 960 | 896 | 896 | 1,792 | 960 | 1,080 | 972 | 552 | 540 | 602 | 564 |
| 62.0 | 50.1 | 48.5 | 47.6 | 119.7 | 144.1 | 194.2 | 191.3 | 30.1 | 40.1 | 18.8 | 24.5 |
| 1.5 | 1.3 | 1.5 | 1.8 | 1.4 | 1.5 | 1.6 | 1.3 | 1.2 | 1.2 | 1.4 | 1.3 |

TABLE 7.04.--BEEF
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 67.3 | 15.6 | 60.1 | 65.5 | 67.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 38.5 | 9.4 | 38.6 | 41.6 | 40.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 22.4 | 4.0 | 17.0 | 19.7 | 22.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 6.4 | 2.2 | 4.4 | 4.2 | 4.5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.7 | 1.7 | 1.5 | 1.5 | 1.6 |
| STANDARD DEVIATION.....DO..... | .9 | 1.1 | .8 | .8 | .8 |
| MAXIMUM.....DO..... | 8.0 | 6.0 | 5.0 | 6.0 | 8.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 75 | 32 | 33 | 40 | 51 |
| STANDARD DEVIATION.....GM..... | 59 | 32 | 24 | 30 | 35 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 19 | 5 | 7 | 9 | 14 |
| 25TH.....GM..... | 37 | 9 | 18 | 19 | 28 |
| 50TH.....GM..... | 57 | 19 | 28 | 34 | 38 |
| 75TH.....GM..... | 95 | 43 | 43 | 55 | 65 |
| 90TH.....GM..... | 149 | 85 | 67 | 75 | 96 |
| 95TH.....GM..... | 187 | 100 | 81 | 93 | 114 |
| 99TH.....GM..... | 283 | 151 | 108 | 143 | 173 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 142 | 53 | 67 | 82 | 104 |
| STANDARD DEVIATION.....GM..... | 95 | 40 | 40 | 46 | 61 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 147 | 65 | 71 | 86 | 105 |
| STANDARD DEVIATION.....GM..... | 80 | 37 | 36 | 46 | 46 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 159 | 72 | 72 | 90 | 107 |
| STANDARD DEVIATION.....GM..... | 83 | 49 | 29 | 39 | 48 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,792 | 228 | 418 | 448 | 561 |
| MAXIMUM OVER 3 DAYS.....GM..... | 3,312 | 454 | 579 | 816 | 928 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 133 | 56 | 64 | 79 | 97 |
| STANDARD DEVIATION.....GM..... | 85 | 41 | 38 | 43 | 52 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 41 | 9 | 14 | 28 | 28 |
| 25TH.....GM..... | 84 | 27 | 35 | 56 | 56 |
| 50TH.....GM..... | 112 | 46 | 56 | 80 | 85 |
| 75TH.....GM..... | 168 | 84 | 84 | 112 | 112 |
| 90TH.....GM..... | 224 | 128 | 112 | 112 | 162 |
| 95TH.....GM..... | 280 | 128 | 126 | 149 | 184 |
| 99TH.....GM..... | 448 | 135 | 170 | 224 | 288 |
| MAXIMUM.....GM..... | 1,792 | 198 | 280 | 448 | 504 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 683.2 | 1.6 | 13.4 | 31.2 | 29.2 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.6 | 1.2 | 1.4 | 1.8 | 1.5 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 69.0 | 68.2 | 70.3 | 65.9 | 72.2 | 65.1 | 75.2 | 68.1 | 64.2 | 63.4 | 65.2 | 57.3 |
| 39.2 | 40.3 | 36.4 | 38.7 | 37.5 | 40.1 | 38.1 | 39.1 | 34.6 | 40.7 | 39.5 | 34.3 |
| 24.5 | 23.2 | 25.2 | 21.0 | 26.6 | 19.9 | 26.9 | 22.8 | 21.4 | 17.6 | 18.3 | 18.4 |
| 5.3 | 4.7 | 8.7 | 6.1 | 8.0 | 5.0 | 10.2 | 6.2 | 8.2 | 5.1 | 7.4 | 4.6 |
| 1.7 | 1.6 | 1.8 | 1.7 | 1.8 | 1.6 | 1.8 | 1.7 | 1.8 | 1.6 | 1.7 | 1.6 |
| .9 | .8 | 1.0 | .9 | .9 | .8 | 1.0 | .8 | 1.0 | .8 | 1.0 | .8 |
| 8.0 | 5.0 | 7.0 | 5.0 | 7.0 | 6.0 | 7.0 | 8.0 | 6.0 | 6.0 | 6.0 | 5.0 |
| 69 | 60 | 93 | 68 | 105 | 69 | 99 | 71 | 79 | 59 | 81 | 61 |
| 50 | 47 | 65 | 49 | 78 | 49 | 71 | 51 | 53 | 42 | 58 | 44 |
| 19 | 19 | 27 | 19 | 27 | 19 | 24 | 19 | 19 | 19 | 28 | 14 |
| 37 | 29 | 47 | 37 | 56 | 37 | 47 | 37 | 37 | 29 | 37 | 37 |
| 56 | 47 | 75 | 56 | 84 | 56 | 75 | 56 | 66 | 47 | 61 | 47 |
| 92 | 75 | 121 | 90 | 138 | 93 | 131 | 93 | 112 | 75 | 93 | 75 |
| 131 | 112 | 171 | 127 | 203 | 131 | 191 | 137 | 149 | 112 | 150 | 121 |
| 164 | 138 | 211 | 167 | 243 | 160 | 233 | 173 | 177 | 149 | 187 | 159 |
| 246 | 227 | 311 | 240 | 357 | 241 | 336 | 252 | 271 | 215 | 303 | 224 |
| 129 | 121 | 175 | 133 | 196 | 141 | 175 | 139 | 149 | 121 | 173 | 117 |
| 72 | 77 | 100 | 93 | 128 | 89 | 103 | 91 | 85 | 69 | 115 | 57 |
| 138 | 121 | 171 | 131 | 192 | 144 | 181 | 142 | 145 | 122 | 148 | 125 |
| 62 | 81 | 77 | 54 | 94 | 67 | 91 | 72 | 66 | 59 | 77 | 63 |
| 161 | 128 | 179 | 147 | 212 | 141 | 190 | 141 | 154 | 133 | 158 | 132 |
| 70 | 50 | 88 | 61 | 115 | 60 | 86 | 57 | 57 | 49 | 76 | 61 |
| 896 | 1,376 | 896 | 896 | 1,792 | 960 | 1,344 | 972 | 560 | 672 | 840 | 448 |
| 1,120 | 2,421 | 2,246 | 1,176 | 3,312 | 1,460 | 1,891 | 1,221 | 948 | 896 | 1,120 | 840 |
| 124 | 111 | 152 | 123 | 174 | 131 | 160 | 128 | 133 | 113 | 140 | 112 |
| 66 | 70 | 87 | 73 | 110 | 79 | 96 | 79 | 70 | 62 | 77 | 60 |
| 43 | 42 | 56 | 43 | 56 | 43 | 56 | 43 | 56 | 40 | 43 | 35 |
| 84 | 84 | 92 | 84 | 112 | 84 | 92 | 84 | 85 | 84 | 85 | 81 |
| 112 | 112 | 113 | 112 | 161 | 112 | 138 | 112 | 112 | 112 | 112 | 112 |
| 162 | 113 | 184 | 140 | 224 | 168 | 224 | 168 | 168 | 138 | 170 | 138 |
| 224 | 170 | 230 | 192 | 288 | 224 | 276 | 224 | 224 | 184 | 224 | 210 |
| 224 | 224 | 312 | 224 | 368 | 276 | 336 | 270 | 227 | 224 | 270 | 224 |
| 336 | 368 | 448 | 392 | 553 | 448 | 504 | 448 | 368 | 345 | 420 | 336 |
| 896 | 960 | 896 | 896 | 1,792 | 960 | 896 | 972 | 552 | 540 | 602 | 336 |
| 37.2 | 30.2 | 26.2 | 25.7 | 85.7 | 80.3 | 127.8 | 132.7 | 16.3 | 17.3 | 13.5 | 14.9 |
| 1.5 | 1.3 | 1.5 | 1.6 | 1.7 | 1.5 | 1.9 | 1.7 | 1.3 | 1.0 | 2.2 | 1.6 |

TABLE 7.05.--BEEF STEAKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 27.6 | 2.3 | 19.5 | 24.2 | 25.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 23.4 | 2.1 | 16.8 | 21.6 | 23.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.9 | .1 | 2.3 | 2.4 | 1.8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .4 | .1 | .4 | .2 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 1.2 | 1.2 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .5 | .5 | .5 | .4 | .3 |
| MAXIMUM.....DO..... | 6.0 | 3.0 | 3.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 65 | 17 | 24 | 29 | 39 |
| STANDARD DEVIATION.....GM..... | 51 | 10 | 16 | 24 | 29 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 15 | | 5 | 8 | 12 |
| 25TH.....GM..... | 31 | | 12 | 15 | 23 |
| 50TH.....GM..... | 56 | | 19 | 27 | 31 |
| 75TH.....GM..... | 75 | | 31 | 37 | 46 |
| 90TH.....GM..... | 123 | | 46 | 51 | 68 |
| 95TH.....GM..... | 151 | | 56 | 61 | 95 |
| 99TH.....GM..... | 245 | | 75 | 112 | 153 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 163 | 44 | 62 | 75 | 106 |
| STANDARD DEVIATION.....GM..... | 107 | 22 | 41 | 43 | 63 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 172 | 69 | 61 | 86 | 126 |
| STANDARD DEVIATION.....GM..... | 88 | 0 | 28 | 72 | 87 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 200 | 28 | 54 | 91 | 87 |
| STANDARD DEVIATION.....GM..... | 149 | 0 | 11 | 38 | 12 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,792 | 85 | 255 | 448 | 504 |
| MAXIMUM OVER 3 DAYS.....GM..... | 3,312 | 138 | 269 | 816 | 816 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 162 | 45 | 60 | 77 | 107 |
| STANDARD DEVIATION.....GM..... | 104 | 21 | 37 | 50 | 67 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 46 | | 14 | 23 | 30 |
| 25TH.....GM..... | 92 | | 28 | 46 | 69 |
| 50TH.....GM..... | 138 | | 56 | 70 | 92 |
| 75TH.....GM..... | 216 | | 84 | 92 | 115 |
| 90TH.....GM..... | 276 | | 112 | 112 | 184 |
| 95TH.....GM..... | 368 | | 122 | 168 | 224 |
| 99TH.....GM..... | 552 | | 168 | 252 | 368 |
| MAXIMUM.....GM..... | 1,792 | 85 | 255 | 448 | 504 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 498.5 | 1.6 | 9.8 | 22.8 | 19.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 4.0 | 11.6 | 4.1 | 4.8 | 3.9 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 24.5 | 25.9 | 27.3 | 23.7 | 31.8 | 27.1 | 35.0 | 30.3 | 25.0 | 23.3 | 24.8 | 18.1 |
| 21.8 | 23.0 | 24.0 | 21.0 | 26.2 | 23.4 | 27.4 | 25.2 | 20.6 | 19.8 | 21.9 | 14.9 |
| 2.6 | 2.6 | 3.1 | 2.6 | 5.0 | 3.3 | 6.6 | 4.8 | 3.9 | 3.3 | 2.0 | 3.0 |
| 0 | .2 | .2 | .1 | .6 | .4 | 1.0 | .3 | .5 | .1 | .8 | .1 |
| 1.1 | 1.1 | 1.2 | 1.1 | 1.2 | 1.2 | 1.3 | 1.2 | 1.3 | 1.2 | 1.3 | 1.2 |
| .3 | .4 | .4 | .4 | .5 | .4 | .6 | .5 | .6 | .5 | .6 | .5 |
| 3.0 | 3.0 | 3.0 | 3.0 | 6.0 | 4.0 | 6.0 | 5.0 | 5.0 | 4.0 | 4.0 | 3.0 |
| 51 | 47 | 61 | 56 | 90 | 63 | 84 | 64 | 67 | 53 | 71 | 56 |
| 34 | 31 | 51 | 40 | 74 | 41 | 56 | 45 | 40 | 34 | 43 | 41 |
| 15 | 14 | 19 | 18 | 24 | 19 | 28 | 19 | 23 | 18 | 23 | 14 |
| 31 | 28 | 37 | 31 | 54 | 37 | 47 | 37 | 37 | 31 | 37 | 28 |
| 38 | 37 | 60 | 46 | 75 | 56 | 75 | 56 | 56 | 46 | 61 | 46 |
| 61 | 56 | 84 | 65 | 112 | 75 | 112 | 75 | 77 | 67 | 82 | 75 |
| 85 | 75 | 117 | 107 | 160 | 120 | 149 | 123 | 122 | 87 | 123 | 107 |
| 112 | 112 | 149 | 131 | 216 | 141 | 187 | 149 | 155 | 123 | 165 | 136 |
| 168 | 157 | 262 | 224 | 328 | 205 | 276 | 240 | 204 | 184 | 224 | 224 |
| 135 | 126 | 181 | 149 | 223 | 165 | 202 | 163 | 166 | 140 | 193 | 135 |
| 78 | 73 | 96 | 99 | 139 | 95 | 110 | 105 | 83 | 87 | 116 | 79 |
| 148 | 126 | 172 | 159 | 217 | 157 | 206 | 164 | 169 | 134 | 168 | 165 |
| 71 | 55 | 90 | 71 | 103 | 62 | 94 | 76 | 70 | 55 | 56 | 83 |
| 0 | 149 | 341 | 178 | 309 | 184 | 209 | 176 | 162 | 65 | 131 | 103 |
| 0 | 81 | 212 | 24 | 267 | 101 | 82 | 77 | 39 | 0 | 56 | 0 |
| 896 | 504 | 896 | 693 | 1,792 | 736 | 1,344 | 972 | 552 | 672 | 672 | 448 |
| 896 | 800 | 1,798 | 740 | 3,312 | 1,460 | 1,540 | 1,040 | 644 | 672 | 672 | 672 |
| 137 | 125 | 178 | 147 | 217 | 161 | 196 | 159 | 157 | 132 | 169 | 137 |
| 78 | 71 | 105 | 91 | 140 | 91 | 105 | 97 | 82 | 72 | 85 | 76 |
| 46 | 43 | 56 | 54 | 68 | 56 | 72 | 56 | 56 | 54 | 56 | 45 |
| 85 | 84 | 112 | 92 | 113 | 105 | 112 | 92 | 92 | 85 | 96 | 84 |
| 112 | 112 | 168 | 115 | 184 | 140 | 170 | 138 | 138 | 112 | 161 | 112 |
| 168 | 149 | 224 | 168 | 276 | 184 | 224 | 184 | 210 | 168 | 224 | 184 |
| 224 | 224 | 300 | 224 | 368 | 276 | 336 | 271 | 230 | 224 | 276 | 252 |
| 280 | 252 | 368 | 324 | 448 | 360 | 414 | 336 | 306 | 224 | 368 | 306 |
| 368 | 448 | 576 | 553 | 673 | 540 | 553 | 540 | 448 | 448 | 368 | 336 |
| 896 | 504 | 896 | 693 | 1,792 | 736 | 720 | 972 | 552 | 460 | 448 | 336 |
| 23.1 | 23.4 | 17.5 | 18.0 | 66.8 | 61.7 | 93.3 | 91.3 | 10.6 | 14.0 | 11.5 | 13.2 |
| 4.0 | 3.7 | 4.0 | 4.5 | 4.3 | 3.6 | 4.2 | 3.5 | 3.0 | 2.9 | 6.9 | 6.0 |

TABLE 7.06.--BEEF ROASTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 19.1 | 2.1 | 12.6 | 14.8 | 16.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 15.6 | 2.1 | 11.4 | 13.1 | 14.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.2 | 0 | 1.2 | 1.6 | 1.5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .4 | 0 | 0 | .1 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 1.0 | 1.1 | 1.2 | 1.1 |
| STANDARD DEVIATION.....DO..... | .6 | .0 | .3 | .4 | .4 |
| MAXIMUM.....DO..... | 7.0 | 1.0 | 2.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 52 | 9 | 20 | 26 | 33 |
| STANDARD DEVIATION.....GM..... | 39 | 7 | 12 | 20 | 22 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 14 | | 5 | 9 | 9 |
| 25TH.....GM..... | 28 | | 9 | 14 | 19 |
| 50TH.....GM..... | 37 | | 19 | 19 | 28 |
| 75TH.....GM..... | 68 | | 27 | 30 | 37 |
| 90TH.....GM..... | 101 | | 37 | 56 | 57 |
| 95TH.....GM..... | 131 | | 47 | 65 | 75 |
| 99TH.....GM..... | 205 | | 56 | 112 | 112 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 130 | 26 | 54 | 70 | 90 |
| STANDARD DEVIATION.....GM..... | 90 | 21 | 31 | 48 | 55 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 129 | 0 | 54 | 67 | 100 |
| STANDARD DEVIATION.....GM..... | 71 | 0 | 19 | 43 | 44 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 138 | 0 | 0 | 113 | 21 |
| STANDARD DEVIATION.....GM..... | 60 | 0 | 0 | 19 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 896 | 70 | 168 | 420 | 336 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,312 | 70 | 168 | 560 | 396 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 124 | 26 | 54 | 68 | 89 |
| STANDARD DEVIATION.....GM..... | 82 | 21 | 29 | 46 | 53 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 35 | | 14 | 28 | 28 |
| 25TH.....GM..... | 72 | | 28 | 43 | 56 |
| 50TH.....GM..... | 112 | | 56 | 56 | 84 |
| 75TH.....GM..... | 168 | | 70 | 84 | 112 |
| 90TH.....GM..... | 224 | | 84 | 112 | 168 |
| 95TH.....GM..... | 280 | | 112 | 144 | 168 |
| 99TH.....GM..... | 448 | | 140 | 280 | 288 |
| MAXIMUM.....GM..... | 896 | 70 | 144 | 336 | 336 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 55.0 | 0 | 1.7 | 1.4 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | 0 | 1.2 | .5 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 17.6 | 18.0 | 22.5 | 18.6 | 21.2 | 16.5 | 25.7 | 20.0 | 21.8 | 18.2 | 20.2 | 13.8 |
| 15.1 | 16.0 | 19.1 | 15.3 | 17.5 | 13.8 | 19.9 | 16.0 | 16.7 | 14.2 | 13.2 | 9.4 |
| 2.4 | 2.0 | 3.1 | 3.0 | 3.4 | 2.4 | 5.1 | 3.5 | 4.4 | 3.7 | 6.1 | 3.5 |
| .1 | .1 | .3 | .3 | .4 | .2 | .7 | .5 | .6 | .3 | .9 | .8 |
| 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.3 | 1.3 | 1.3 | 1.3 | 1.5 | 1.5 |
| .5 | .4 | .5 | .5 | .5 | .5 | .6 | .6 | .7 | .6 | .8 | .9 |
| 5.0 | 3.0 | 4.0 | 5.0 | 4.0 | 4.0 | 7.0 | 5.0 | 5.0 | 4.0 | 4.0 | 5.0 |
| 50 | 45 | 61 | 48 | 63 | 49 | 66 | 49 | 54 | 47 | 65 | 53 |
| 30 | 38 | 37 | 34 | 45 | 34 | 49 | 33 | 38 | 37 | 49 | 42 |
| 14 | 14 | 19 | 14 | 19 | 14 | 19 | 14 | 14 | 14 | 14 | 14 |
| 29 | 28 | 37 | 28 | 37 | 28 | 37 | 28 | 28 | 24 | 28 | 27 |
| 43 | 37 | 56 | 37 | 47 | 37 | 56 | 37 | 47 | 37 | 48 | 37 |
| 56 | 56 | 75 | 56 | 75 | 59 | 75 | 56 | 75 | 56 | 75 | 68 |
| 93 | 75 | 108 | 93 | 115 | 93 | 131 | 93 | 108 | 93 | 149 | 112 |
| 112 | 108 | 132 | 103 | 149 | 112 | 149 | 112 | 113 | 115 | 149 | 149 |
| 149 | 240 | 216 | 172 | 224 | 174 | 233 | 159 | 205 | 187 | 201 | 205 |
| 134 | 118 | 167 | 121 | 154 | 129 | 161 | 123 | 130 | 107 | 148 | 106 |
| 73 | 92 | 95 | 73 | 93 | 84 | 112 | 78 | 75 | 71 | 111 | 60 |
| 122 | 131 | 134 | 111 | 169 | 119 | 153 | 112 | 119 | 122 | 135 | 114 |
| 50 | 86 | 62 | 40 | 88 | 58 | 84 | 54 | 55 | 57 | 79 | 50 |
| 137 | 87 | 131 | 197 | 155 | 112 | 147 | 114 | 175 | 162 | 135 | 156 |
| 79 | 65 | 51 | 60 | 50 | 50 | 64 | 45 | 43 | 67 | 50 | 56 |
| 560 | 896 | 720 | 720 | 864 | 718 | 896 | 896 | 448 | 540 | 602 | 336 |
| 645 | 1,120 | 720 | 792 | 972 | 718 | 1,312 | 896 | 688 | 730 | 602 | 616 |
| 127 | 116 | 151 | 117 | 153 | 120 | 149 | 114 | 121 | 108 | 127 | 104 |
| 66 | 86 | 88 | 64 | 95 | 75 | 100 | 67 | 66 | 68 | 81 | 59 |
| 43 | 36 | 56 | 43 | 43 | 28 | 43 | 42 | 28 | 43 | 28 | 28 |
| 84 | 70 | 86 | 84 | 86 | 70 | 84 | 72 | 84 | 56 | 56 | 56 |
| 112 | 112 | 129 | 112 | 114 | 112 | 112 | 112 | 112 | 86 | 112 | 86 |
| 168 | 129 | 196 | 140 | 196 | 162 | 168 | 140 | 168 | 129 | 168 | 135 |
| 224 | 216 | 243 | 172 | 243 | 224 | 252 | 172 | 224 | 168 | 224 | 210 |
| 224 | 224 | 288 | 224 | 336 | 280 | 336 | 224 | 224 | 224 | 224 | 224 |
| 336 | 448 | 448 | 336 | 448 | 336 | 540 | 336 | 336 | 450 | 420 | 224 |
| 448 | 896 | 720 | 448 | 864 | 576 | 896 | 896 | 448 | 540 | 602 | 336 |
| 4.0 | .8 | 4.0 | 4.5 | 3.5 | 4.4 | 16.0 | 8.9 | 3.5 | 1.4 | 0 | .8 |
| .9 | .2 | 1.1 | 1.3 | .3 | .4 | 1.0 | .5 | 1.1 | .3 | 0 | .4 |

TABLE 7.07.--GROUND BEEF
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 36.3 | 3.3 | 35.8 | 39.2 | 41.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 28.8 | 2.9 | 29.3 | 31.2 | 33.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 6.6 | .5 | 5.5 | 7.4 | 7.6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.0 | 0 | 1.0 | .6 | .7 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 1.1 | 1.3 | 1.3 | 1.3 |
| STANDARD DEVIATION.....DO..... | .6 | .4 | .5 | .6 | .5 |
| MAXIMUM.....DO..... | 6.0 | 2.0 | 4.0 | 5.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 53 | 18 | 29 | 36 | 41 |
| STANDARD DEVIATION.....GM..... | 37 | 11 | 18 | 23 | 23 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 19 | | 7 | 9 | 19 |
| 25TH.....GM..... | 28 | | 19 | 19 | 28 |
| 50TH.....GM..... | 37 | | 25 | 28 | 37 |
| 75TH.....GM..... | 75 | | 37 | 42 | 51 |
| 90TH.....GM..... | 93 | | 56 | 63 | 75 |
| 95TH.....GM..... | 117 | | 67 | 75 | 80 |
| 99TH.....GM..... | 187 | | 80 | 112 | 131 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 127 | 48 | 74 | 88 | 101 |
| STANDARD DEVIATION.....GM..... | 71 | 27 | 42 | 44 | 47 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 130 | 41 | 70 | 85 | 100 |
| STANDARD DEVIATION.....GM..... | 63 | 21 | 31 | 29 | 37 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 138 | 0 | 63 | 132 | 98 |
| STANDARD DEVIATION.....GM..... | 58 | 0 | 16 | 53 | 29 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,045 | 112 | 280 | 448 | 399 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,525 | 136 | 308 | 560 | 560 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 123 | 46 | 70 | 84 | 97 |
| STANDARD DEVIATION.....GM..... | 66 | 26 | 36 | 36 | 41 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 56 | | 19 | 28 | 42 |
| 25TH.....GM..... | 84 | | 44 | 56 | 71 |
| 50TH.....GM..... | 112 | | 56 | 84 | 85 |
| 75TH.....GM..... | 149 | | 85 | 112 | 112 |
| 90TH.....GM..... | 224 | | 112 | 113 | 142 |
| 95TH.....GM..... | 224 | | 113 | 148 | 170 |
| 99TH.....GM..... | 339 | | 170 | 224 | 224 |
| MAXIMUM.....GM..... | 960 | 112 | 280 | 224 | 336 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 82.7 | 0 | 1.8 | 5.4 | 7.3 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .5 | 0 | .4 | .6 | .8 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 45.2 | 41.4 | 43.5 | 38.3 | 40.1 | 34.4 | 37.1 | 32.9 | 31.3 | 29.5 | 32.8 | 30.4 |
| 35.1 | 33.5 | 32.6 | 27.6 | 29.8 | 27.6 | 29.6 | 27.2 | 25.2 | 25.2 | 27.7 | 23.0 |
| 9.0 | 7.5 | 8.5 | 9.0 | 8.8 | 6.2 | 6.4 | 5.1 | 5.5 | 3.3 | 4.5 | 6.2 |
| 1.1 | .5 | 2.4 | 1.6 | 1.5 | .7 | 1.1 | .7 | .7 | 1.0 | .6 | 1.2 |
| 1.3 | 1.3 | 1.4 | 1.4 | 1.4 | 1.3 | 1.3 | 1.2 | 1.3 | 1.2 | 1.2 | 1.3 |
| .6 | .5 | .7 | .7 | .7 | .5 | .6 | .5 | .6 | .6 | .5 | .6 |
| 5.0 | 4.0 | 5.0 | 5.0 | 4.0 | 4.0 | 5.0 | 4.0 | 5.0 | 6.0 | 4.0 | 4.0 |
| 53 | 44 | 67 | 53 | 74 | 50 | 63 | 48 | 55 | 45 | 54 | 49 |
| 36 | 32 | 44 | 38 | 50 | 30 | 41 | 29 | 31 | 26 | 27 | 29 |
| 19 | 19 | 19 | 19 | 24 | 19 | 24 | 19 | 24 | 19 | 28 | 19 |
| 28 | 28 | 37 | 28 | 37 | 28 | 37 | 28 | 37 | 28 | 37 | 37 |
| 37 | 37 | 56 | 37 | 57 | 37 | 56 | 37 | 47 | 37 | 47 | 37 |
| 65 | 56 | 84 | 66 | 85 | 65 | 75 | 56 | 75 | 56 | 75 | 56 |
| 95 | 75 | 131 | 99 | 149 | 84 | 112 | 75 | 93 | 75 | 93 | 75 |
| 112 | 93 | 149 | 112 | 187 | 103 | 149 | 112 | 112 | 93 | 112 | 107 |
| 187 | 149 | 242 | 207 | 243 | 156 | 224 | 149 | 179 | 149 | 149 | 163 |
| 121 | 109 | 155 | 116 | 172 | 122 | 152 | 121 | 136 | 114 | 141 | 114 |
| 59 | 59 | 86 | 69 | 100 | 59 | 79 | 63 | 54 | 45 | 58 | 39 |
| 133 | 115 | 156 | 124 | 172 | 121 | 155 | 115 | 129 | 110 | 136 | 118 |
| 52 | 74 | 64 | 60 | 72 | 44 | 80 | 44 | 45 | 45 | 44 | 56 |
| 151 | 108 | 145 | 127 | 178 | 127 | 145 | 132 | 163 | 121 | 114 | 105 |
| 74 | 28 | 54 | 39 | 70 | 57 | 48 | 39 | 40 | 45 | 2 | 42 |
| 480 | 1,045 | 725 | 896 | 896 | 454 | 672 | 800 | 336 | 448 | 336 | 448 |
| 1,014 | 1,525 | 924 | 1,008 | 1,120 | 922 | 1,344 | 800 | 616 | 672 | 448 | 616 |
| 120 | 106 | 142 | 114 | 162 | 118 | 146 | 116 | 130 | 109 | 132 | 111 |
| 59 | 62 | 74 | 64 | 82 | 57 | 74 | 57 | 51 | 45 | 49 | 46 |
| 56 | 56 | 56 | 42 | 56 | 56 | 56 | 56 | 56 | 50 | 84 | 56 |
| 84 | 84 | 85 | 84 | 112 | 84 | 112 | 84 | 98 | 84 | 112 | 84 |
| 112 | 112 | 112 | 112 | 142 | 112 | 112 | 112 | 112 | 112 | 112 | 112 |
| 142 | 112 | 170 | 112 | 224 | 113 | 170 | 113 | 168 | 112 | 150 | 112 |
| 224 | 168 | 224 | 168 | 226 | 170 | 224 | 168 | 224 | 168 | 224 | 168 |
| 224 | 213 | 255 | 224 | 336 | 224 | 255 | 224 | 224 | 224 | 224 | 224 |
| 336 | 336 | 448 | 320 | 448 | 336 | 340 | 336 | 227 | 224 | 336 | 255 |
| 480 | 960 | 640 | 896 | 672 | 454 | 672 | 800 | 336 | 448 | 336 | 336 |
| 6.3 | 4.7 | 3.1 | 3.1 | 13.9 | 9.0 | 9.0 | 14.1 | 2.2 | 2.0 | 0 | .9 |
| .5 | .4 | .4 | .4 | .6 | .4 | .4 | .5 | .5 | .3 | 0 | .2 |

TABLE 7.08.--PORK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 49.9 | 10.1 | 44.2 | 46.0 | 46.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 30.5 | 9.2 | 29.2 | 31.3 | 33.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 13.5 | .6 | 11.6 | 11.6 | 10.4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 5.9 | .3 | 3.4 | 3.2 | 2.7 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.7 | 1.3 | 1.5 | 1.5 | 1.5 |
| STANDARD DEVIATION.....DO..... | 1.0 | 1.1 | .8 | .8 | .8 |
| MAXIMUM.....DO..... | 8.0 | 8.0 | 6.0 | 6.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 39 | 28 | 19 | 24 | 28 |
| STANDARD DEVIATION.....GM..... | 39 | 53 | 17 | 21 | 24 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 5 | 3 | 3 | 3 | 3 |
| 25TH.....GM..... | 11 | 9 | 5 | 8 | 9 |
| 50TH.....GM..... | 28 | 19 | 13 | 19 | 23 |
| 75TH.....GM..... | 54 | 33 | 28 | 32 | 37 |
| 90TH.....GM..... | 84 | 43 | 42 | 49 | 56 |
| 95TH.....GM..... | 111 | 45 | 54 | 63 | 75 |
| 99TH.....GM..... | 178 | 340 | 77 | 97 | 119 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 84 | 55 | 46 | 56 | 68 |
| STANDARD DEVIATION.....GM..... | 77 | 40 | 43 | 46 | 60 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 76 | 64 | 35 | 48 | 56 |
| STANDARD DEVIATION.....GM..... | 62 | 41 | 25 | 34 | 35 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 70 | 340 | 36 | 46 | 50 |
| STANDARD DEVIATION.....GM..... | 63 | 0 | 26 | 37 | 39 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,128 | 383 | 243 | 389 | 512 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,564 | 1,021 | 312 | 543 | 512 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 69 | 66 | 37 | 47 | 57 |
| STANDARD DEVIATION.....GM..... | 69 | 44 | 36 | 44 | 49 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 8 | 8 | 5 | 8 | 8 |
| 25TH.....GM..... | 16 | 28 | 8 | 16 | 16 |
| 50TH.....GM..... | 44 | 57 | 24 | 30 | 43 |
| 75TH.....GM..... | 92 | 116 | 56 | 70 | 85 |
| 90TH.....GM..... | 160 | 128 | 86 | 97 | 112 |
| 95TH.....GM..... | 194 | 128 | 112 | 135 | 142 |
| 99TH.....GM..... | 320 | 135 | 160 | 188 | 224 |
| MAXIMUM.....GM..... | 1,080 | 135 | 243 | 389 | 432 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 379.0 | 0 | 16.0 | 17.6 | 12.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.2 | 0 | 2.2 | 1.5 | 1.0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 48.8 | 47.0 | 56.1 | 46.2 | 52.6 | 47.0 | 57.6 | 51.2 | 53.8 | 47.9 | 59.2 | 45.4 |
| 33.0 | 34.4 | 34.4 | 30.8 | 31.4 | 31.2 | 29.4 | 31.5 | 25.1 | 25.7 | 25.0 | 26.2 |
| 12.0 | 9.7 | 15.3 | 11.2 | 14.8 | 11.3 | 18.6 | 13.6 | 17.7 | 15.3 | 20.1 | 12.1 |
| 3.8 | 3.0 | 6.4 | 4.2 | 6.4 | 4.5 | 9.5 | 6.2 | 10.9 | 6.9 | 14.1 | 7.0 |
| 1.6 | 1.5 | 1.7 | 1.6 | 1.7 | 1.6 | 1.9 | 1.7 | 2.0 | 1.8 | 2.2 | 1.9 |
| .9 | .8 | 1.0 | .8 | 1.0 | .9 | 1.1 | 1.0 | 1.1 | 1.0 | 1.3 | 1.1 |
| 7.0 | 6.0 | 7.0 | 5.0 | 6.0 | 6.0 | 8.0 | 7.0 | 7.0 | 7.0 | 7.0 | 6.0 |
| 35 | 31 | 45 | 35 | 51 | 37 | 51 | 36 | 45 | 34 | 44 | 33 |
| 33 | 28 | 39 | 31 | 47 | 39 | 48 | 35 | 50 | 32 | 43 | 35 |
| 4 | 3 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 4 |
| 11 | 11 | 15 | 12 | 16 | 11 | 16 | 11 | 16 | 11 | 15 | 9 |
| 28 | 27 | 36 | 28 | 37 | 28 | 38 | 28 | 32 | 27 | 33 | 27 |
| 47 | 41 | 61 | 47 | 71 | 49 | 72 | 48 | 57 | 47 | 56 | 47 |
| 75 | 65 | 94 | 75 | 111 | 76 | 104 | 79 | 95 | 75 | 92 | 73 |
| 96 | 81 | 120 | 103 | 143 | 100 | 149 | 99 | 130 | 96 | 141 | 79 |
| 165 | 124 | 168 | 145 | 224 | 166 | 221 | 158 | 230 | 134 | 235 | 124 |
| 82 | 75 | 97 | 76 | 109 | 83 | 105 | 78 | 95 | 74 | 85 | 67 |
| 73 | 69 | 78 | 66 | 98 | 70 | 90 | 71 | 92 | 67 | 81 | 58 |
| 76 | 67 | 82 | 77 | 101 | 72 | 90 | 72 | 75 | 61 | 75 | 69 |
| 59 | 44 | 56 | 55 | 77 | 52 | 77 | 54 | 65 | 50 | 55 | 44 |
| 58 | 61 | 86 | 66 | 83 | 77 | 83 | 59 | 67 | 55 | 64 | 52 |
| 44 | 41 | 59 | 36 | 65 | 79 | 66 | 54 | 77 | 44 | 60 | 64 |
| 648 | 945 | 988 | 630 | 816 | 648 | 1,112 | 864 | 720 | 464 | 468 | 1,128 |
| 689 | 945 | 1,020 | 673 | 1,238 | 1,512 | 1,564 | 1,347 | 1,420 | 965 | 784 | 1,249 |
| 68 | 64 | 79 | 68 | 88 | 70 | 79 | 64 | 67 | 56 | 59 | 53 |
| 65 | 57 | 75 | 60 | 84 | 67 | 81 | 62 | 72 | 56 | 67 | 57 |
| 8 | 8 | 14 | 8 | 16 | 8 | 12 | 8 | 8 | 8 | 8 | 8 |
| 16 | 24 | 24 | 21 | 24 | 20 | 24 | 16 | 16 | 16 | 16 | 16 |
| 53 | 53 | 56 | 52 | 56 | 56 | 56 | 43 | 40 | 32 | 28 | 35 |
| 92 | 87 | 112 | 92 | 113 | 92 | 112 | 92 | 86 | 85 | 84 | 84 |
| 158 | 135 | 170 | 144 | 194 | 160 | 184 | 142 | 162 | 117 | 144 | 112 |
| 194 | 170 | 224 | 184 | 264 | 194 | 227 | 184 | 194 | 172 | 194 | 140 |
| 291 | 272 | 336 | 280 | 408 | 312 | 368 | 276 | 368 | 230 | 288 | 258 |
| 648 | 630 | 648 | 426 | 680 | 576 | 1,080 | 504 | 460 | 432 | 454 | 564 |
| 20.8 | 19.9 | 21.3 | 21.0 | 34.0 | 63.8 | 61.8 | 52.0 | 13.8 | 14.4 | 5.3 | 4.7 |
| 1.3 | 1.3 | 1.6 | 2.0 | .9 | 1.6 | 1.1 | .8 | 1.1 | 1.0 | .7 | .6 |

TABLE 7.09.--PORK CHOPS AND STEAKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 14.3 | 1.6 | 13.7 | 16.8 | 15.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 13.4 | 1.6 | 13.0 | 16.1 | 15.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .9 | 0 | .6 | .6 | .4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | 0 | (2) | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 1.0 | 1.1 | 1.1 | 1.0 |
| STANDARD DEVIATION.....DO..... | .3 | .0 | .3 | .2 | .2 |
| MAXIMUM.....DO..... | 4.0 | 1.0 | 2.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 46 | 16 | 24 | 29 | 33 |
| STANDARD DEVIATION.....GM..... | 28 | 12 | 14 | 17 | 17 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 15 | | 8 | 9 | 15 |
| 25TH.....GM..... | 27 | | 15 | 18 | 23 |
| 50TH.....GM..... | 38 | | 23 | 27 | 31 |
| 75TH.....GM..... | 61 | | 32 | 32 | 37 |
| 90TH.....GM..... | 78 | | 41 | 50 | 53 |
| 95TH.....GM..... | 97 | | 47 | 61 | 65 |
| 99TH.....GM..... | 149 | | 74 | 97 | 97 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 129 | 48 | 69 | 81 | 94 |
| STANDARD DEVIATION.....GM..... | 72 | 36 | 38 | 39 | 44 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 132 | 0 | 68 | 110 | 110 |
| STANDARD DEVIATION.....GM..... | 60 | 0 | 25 | 34 | 45 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 131 | 0 | 0 | 112 | 77 |
| STANDARD DEVIATION.....GM..... | 65 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 552 | 116 | 243 | 360 | 291 |
| MAXIMUM OVER 3 DAYS.....GM..... | 895 | 116 | 243 | 403 | 368 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 128 | 48 | 67 | 83 | 95 |
| STANDARD DEVIATION.....GM..... | 70 | 36 | 37 | 41 | 45 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 45 | | 23 | 28 | 44 |
| 25TH.....GM..... | 80 | | 43 | 53 | 69 |
| 50TH.....GM..... | 104 | | 58 | 80 | 92 |
| 75TH.....GM..... | 160 | | 92 | 97 | 106 |
| 90TH.....GM..... | 203 | | 116 | 138 | 156 |
| 95TH.....GM..... | 276 | | 138 | 176 | 184 |
| 99TH.....GM..... | 368 | | 188 | 194 | 276 |
| MAXIMUM.....GM..... | 552 | 116 | 243 | 360 | 291 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 83.2 | 0 | 5.5 | 3.5 | 1.7 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.4 | 0 | 3.6 | 1.2 | .6 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 14.5 | 15.3 | 16.4 | 14.7 | 14.6 | 14.5 | 15.0 | 13.9 | 14.4 | 10.1 | 17.4 | 10.2 |
| 13.8 | 14.8 | 15.7 | 14.1 | 13.6 | 13.8 | 13.5 | 12.8 | 12.5 | 8.7 | 16.2 | 9.6 |
| .7 | .4 | .7 | .7 | 1.0 | .6 | 1.4 | .9 | 1.6 | 1.2 | 1.2 | .6 |
| 0 | .1 | .1 | 0 | 0 | (2) | .1 | .1 | .3 | .1 | 0 | 0 |
| 1.1 | 1.0 | 1.1 | 1.0 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.2 | 1.1 | 1.1 |
| .2 | .2 | .3 | .2 | .3 | .2 | .4 | .4 | .5 | .5 | .3 | .3 |
| 2.0 | 3.0 | 3.0 | 2.0 | 2.0 | 3.0 | 3.0 | 4.0 | 3.0 | 4.0 | 3.0 | 2.0 |
| 41 | 36 | 50 | 42 | 63 | 43 | 62 | 44 | 54 | 46 | 47 | 37 |
| 23 | 18 | 25 | 23 | 35 | 26 | 32 | 24 | 33 | 26 | 31 | 22 |
| 15 | 15 | 20 | 15 | 23 | 15 | 27 | 15 | 19 | 20 | 19 | 9 |
| 27 | 27 | 31 | 27 | 36 | 27 | 39 | 27 | 31 | 31 | 28 | 23 |
| 32 | 32 | 46 | 32 | 61 | 34 | 53 | 38 | 46 | 32 | 32 | 32 |
| 53 | 46 | 61 | 53 | 77 | 53 | 77 | 53 | 61 | 59 | 56 | 47 |
| 65 | 61 | 84 | 65 | 97 | 65 | 97 | 69 | 92 | 85 | 92 | 63 |
| 86 | 65 | 96 | 80 | 129 | 92 | 124 | 93 | 123 | 107 | 129 | 65 |
| 129 | 107 | 129 | 138 | 184 | 142 | 184 | 123 | 186 | 123 | 156 | 120 |
| 118 | 106 | 144 | 117 | 176 | 124 | 173 | 122 | 145 | 118 | 126 | 108 |
| 61 | 49 | 66 | 56 | 87 | 65 | 80 | 58 | 84 | 57 | 78 | 65 |
| 121 | 105 | 129 | 143 | 189 | 108 | 138 | 128 | 120 | 124 | 163 | 85 |
| 49 | 51 | 44 | 47 | 66 | 47 | 68 | 54 | 43 | 43 | 39 | 20 |
| 0 | 38 | 135 | 0 | 0 | 259 | 189 | 76 | 135 | 108 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 27 | 5 | 22 | 49 | 1 | 0 | 0 |
| 473 | 368 | 368 | 426 | 552 | 552 | 464 | 426 | 485 | 368 | 468 | 360 |
| 473 | 381 | 414 | 426 | 650 | 895 | 582 | 596 | 558 | 434 | 468 | 360 |
| 116 | 105 | 140 | 119 | 175 | 124 | 163 | 120 | 137 | 115 | 125 | 104 |
| 56 | 50 | 64 | 57 | 87 | 66 | 79 | 57 | 74 | 49 | 55 | 62 |
| 46 | 44 | 59 | 46 | 69 | 46 | 69 | 46 | 53 | 56 | 57 | 20 |
| 80 | 80 | 92 | 80 | 97 | 80 | 92 | 80 | 85 | 88 | 85 | 69 |
| 97 | 97 | 138 | 97 | 180 | 97 | 160 | 97 | 122 | 97 | 106 | 92 |
| 145 | 117 | 184 | 160 | 224 | 160 | 194 | 154 | 184 | 138 | 160 | 116 |
| 194 | 184 | 234 | 194 | 284 | 194 | 276 | 184 | 194 | 184 | 194 | 184 |
| 232 | 194 | 276 | 225 | 320 | 240 | 291 | 224 | 291 | 194 | 234 | 194 |
| 291 | 284 | 348 | 368 | 468 | 368 | 448 | 300 | 368 | 284 | 276 | 360 |
| 322 | 368 | 368 | 426 | 552 | 552 | 464 | 368 | 460 | 368 | 276 | 360 |
| 3.2 | 3.7 | 5.5 | 3.2 | 13.0 | 12.8 | 15.5 | 5.7 | 4.2 | 3.4 | 2.1 | 0 |
| 1.0 | 1.1 | 2.3 | 1.4 | 2.1 | 1.6 | 1.8 | .5 | 2.2 | 1.6 | 2.0 | 0 |

TABLE 7.10.--HAM
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 16.3 | 5.4 | 9.5 | 11.7 | 12.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 12.6 | 4.5 | 7.7 | 9.8 | 10.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.2 | .6 | 1.6 | 2.0 | 1.8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .5 | .3 | .2 | 0 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.4 | 1.5 | 1.2 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .7 | 1.5 | .5 | .5 | .4 |
| MAXIMUM.....DO..... | 8.0 | 8.0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 39 | 40 | 20 | 23 | 27 |
| STANDARD DEVIATION.....GM..... | 38 | 69 | 14 | 15 | 16 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 9 | | 5 | 7 | 9 |
| 25TH.....GM..... | 19 | | 9 | 12 | 14 |
| 50TH.....GM..... | 29 | | 19 | 19 | 23 |
| 75TH.....GM..... | 48 | | 28 | 28 | 37 |
| 90TH.....GM..... | 75 | | 37 | 38 | 47 |
| 95TH.....GM..... | 103 | | 56 | 56 | 56 |
| 99TH.....GM..... | 192 | | 75 | 75 | 75 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 90 | 70 | 55 | 61 | 72 |
| STANDARD DEVIATION.....GM..... | 69 | 35 | 37 | 36 | 40 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 99 | 64 | 39 | 53 | 65 |
| STANDARD DEVIATION.....GM..... | 76 | 41 | 25 | 31 | 26 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 104 | 340 | 46 | 0 | 24 |
| STANDARD DEVIATION.....GM..... | 79 | 0 | 20 | 0 | 3 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,080 | 383 | 224 | 194 | 224 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,512 | 1,021 | 224 | 280 | 252 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 87 | 80 | 50 | 55 | 66 |
| STANDARD DEVIATION.....GM..... | 67 | 40 | 35 | 33 | 37 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 28 | | 9 | 14 | 22 |
| 25TH.....GM..... | 43 | | 28 | 28 | 42 |
| 50TH.....GM..... | 70 | | 43 | 56 | 56 |
| 75TH.....GM..... | 112 | | 56 | 70 | 86 |
| 90TH.....GM..... | 168 | | 86 | 112 | 112 |
| 95TH.....GM..... | 224 | | 112 | 112 | 135 |
| 99TH.....GM..... | 336 | | 224 | 170 | 180 |
| MAXIMUM.....GM..... | 1,080 | 135 | 224 | 189 | 224 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 64.7 | 0 | 2.2 | 3.0 | 1.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .8 | 0 | 1.8 | 1.2 | .5 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 14.6 | 13.4 | 19.1 | 14.9 | 18.8 | 15.2 | 22.1 | 16.4 | 17.0 | 16.2 | 14.8 | 17.3 |
| 11.9 | 10.9 | 14.7 | 11.9 | 13.2 | 12.4 | 15.7 | 13.3 | 12.4 | 12.2 | 10.6 | 12.0 |
| 2.5 | 2.1 | 3.8 | 2.3 | 4.8 | 2.6 | 5.5 | 2.5 | 3.3 | 3.4 | 4.2 | 4.4 |
| .2 | .4 | .6 | .7 | .8 | .2 | .9 | .5 | 1.3 | .7 | 0 | .8 |
| 1.3 | 1.3 | 1.3 | 1.3 | 1.5 | 1.3 | 1.4 | 1.3 | 1.5 | 1.4 | 1.4 | 1.5 |
| .5 | .6 | .6 | .6 | .7 | .6 | .8 | .7 | .8 | .7 | .7 | .8 |
| 4.0 | 4.0 | 5.0 | 4.0 | 5.0 | 5.0 | 8.0 | 5.0 | 4.0 | 5.0 | 4.0 | 5.0 |
| 37 | 32 | 46 | 32 | 50 | 37 | 48 | 35 | 47 | 39 | 41 | 34 |
| 33 | 29 | 38 | 26 | 44 | 42 | 45 | 35 | 44 | 30 | 26 | 21 |
| 9 | 9 | 12 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 |
| 19 | 14 | 23 | 14 | 19 | 19 | 19 | 15 | 19 | 19 | 18 | 19 |
| 28 | 24 | 37 | 28 | 37 | 28 | 37 | 28 | 36 | 28 | 38 | 29 |
| 47 | 37 | 56 | 38 | 57 | 46 | 57 | 38 | 56 | 56 | 56 | 47 |
| 75 | 65 | 85 | 63 | 102 | 70 | 93 | 69 | 93 | 75 | 75 | 70 |
| 96 | 76 | 112 | 84 | 137 | 86 | 142 | 85 | 115 | 96 | 84 | 75 |
| 187 | 149 | 196 | 132 | 227 | 164 | 224 | 159 | 230 | 131 | 125 | 93 |
| 91 | 82 | 114 | 80 | 110 | 86 | 102 | 81 | 104 | 90 | 96 | 76 |
| 70 | 85 | 92 | 69 | 87 | 65 | 73 | 54 | 80 | 64 | 62 | 39 |
| 101 | 74 | 99 | 74 | 116 | 86 | 118 | 102 | 115 | 89 | 96 | 80 |
| 75 | 34 | 70 | 37 | 78 | 50 | 89 | 96 | 101 | 44 | 33 | 28 |
| 69 | 63 | 100 | 69 | 108 | 233 | 112 | 90 | 83 | 98 | 0 | 69 |
| 24 | 20 | 40 | 27 | 67 | 183 | 65 | 40 | 46 | 54 | 0 | 21 |
| 560 | 945 | 972 | 630 | 680 | 648 | 1,080 | 864 | 560 | 336 | 342 | 224 |
| 689 | 945 | 972 | 630 | 1,238 | 1,512 | 1,123 | 1,339 | 1,064 | 648 | 374 | 280 |
| 88 | 75 | 103 | 73 | 103 | 86 | 100 | 81 | 96 | 84 | 88 | 67 |
| 66 | 60 | 79 | 51 | 78 | 66 | 78 | 59 | 70 | 55 | 54 | 33 |
| 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 |
| 56 | 43 | 56 | 43 | 56 | 43 | 56 | 43 | 56 | 43 | 45 | 43 |
| 72 | 56 | 84 | 56 | 85 | 67 | 84 | 57 | 84 | 72 | 84 | 56 |
| 112 | 86 | 113 | 86 | 113 | 112 | 112 | 112 | 112 | 112 | 112 | 86 |
| 168 | 113 | 172 | 129 | 180 | 168 | 196 | 168 | 170 | 168 | 162 | 112 |
| 224 | 168 | 224 | 168 | 245 | 224 | 224 | 170 | 224 | 216 | 196 | 114 |
| 340 | 336 | 448 | 315 | 448 | 432 | 425 | 315 | 360 | 270 | 288 | 168 |
| 560 | 630 | 648 | 410 | 680 | 576 | 1,080 | 448 | 432 | 288 | 288 | 170 |
| 2.8 | 3.8 | 4.4 | 2.6 | .9 | 9.7 | 11.8 | 10.9 | 4.4 | 3.8 | 0 | 2.9 |
| .7 | 1.0 | 1.3 | .9 | .1 | .9 | .8 | .7 | 1.6 | 1.0 | 0 | 1.1 |

TABLE 7.11.--PORK ROASTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.5 | .3 | 2.1 | 1.6 | 2.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.2 | .3 | 1.9 | 1.5 | 2.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .3 | 0 | .2 | .1 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 1.2 | 1.0 | 1.1 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .5 | .0 | .4 | .3 | .2 |
| MAXIMUM.....DO..... | 5.0 | 1.0 | 3.0 | 2.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 47 | 23 | 18 | 21 | 30 |
| STANDARD DEVIATION.....GM..... | 37 | 14 | 10 | 12 | 24 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 9 | | | | |
| 25TH.....GM..... | 27 | | | | |
| 50TH.....GM..... | 37 | | | | |
| 75TH.....GM..... | 56 | | | | |
| 90TH.....GM..... | 84 | | | | |
| 95TH.....GM..... | 112 | | | | |
| 99TH.....GM..... | 205 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 121 | 68 | 47 | 63 | 84 |
| STANDARD DEVIATION.....GM..... | 78 | 42 | 24 | 36 | 69 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 122 | 0 | 56 | 23 | 88 |
| STANDARD DEVIATION.....GM..... | 76 | 0 | (2) | 0 | 21 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 178 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 79 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,080 | 112 | 112 | 144 | 432 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,116 | 112 | 113 | 144 | 432 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 117 | 68 | 47 | 58 | 84 |
| STANDARD DEVIATION.....GM..... | 78 | 42 | 22 | 33 | 65 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 28 | | | | |
| 25TH.....GM..... | 56 | | | | |
| 50TH.....GM..... | 112 | | | | |
| 75TH.....GM..... | 162 | | | | |
| 90TH.....GM..... | 224 | | | | |
| 95TH.....GM..... | 227 | | | | |
| 99TH.....GM..... | 432 | | | | |
| MAXIMUM.....GM..... | 648 | 112 | 112 | 140 | 432 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 16.2 | 0 | 0 | 0 | 1.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.4 | 0 | 0 | 0 | 2.9 |

1/ SEE TABLE NOTES IN APPENDIX.
2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 2.8 | 2.1 | 3.0 | 2.4 | 1.9 | 1.8 | 3.3 | 3.1 | 2.7 | 2.6 | 4.2 | 2.3 |
| 2.7 | 1.9 | 2.9 | 2.0 | 1.6 | 1.7 | 2.7 | 2.6 | 1.8 | 2.0 | 3.1 | 1.7 |
| (2) | .2 | .2 | .4 | .3 | .1 | .6 | .4 | .9 | .6 | .7 | .5 |
| 0 | 0 | 0 | 0 | (2) | 0 | .1 | (2) | 0 | 0 | .4 | .2 |
| 1.1 | 1.1 | 1.1 | 1.2 | 1.3 | 1.1 | 1.2 | 1.2 | 1.4 | 1.3 | 1.7 | 1.6 |
| .3 | .4 | .4 | .4 | .6 | .3 | .5 | .5 | .5 | .5 | 1.2 | .9 |
| 3.0 | 3.0 | 3.0 | 3.0 | 4.0 | 2.0 | 3.0 | 4.0 | 2.0 | 2.0 | 5.0 | 4.0 |
| 43 | 38 | 48 | 50 | 65 | 35 | 59 | 46 | 65 | 38 | 82 | 42 |
| 32 | 31 | 25 | 33 | 45 | 19 | 46 | 29 | 63 | 27 | 55 | 27 |
| 9 | | | | 19 | 9 | 19 | 14 | | | | |
| 28 | | | | 37 | 19 | 29 | 28 | | | | |
| 37 | | | | 56 | 37 | 46 | 37 | | | | |
| 47 | | | | 75 | 38 | 75 | 56 | | | | |
| 75 | | | | 103 | 64 | 112 | 75 | | | | |
| 93 | | | | 149 | 75 | 149 | 108 | | | | |
| 216 | | | | 299 | 93 | 205 | 168 | | | | |
| 130 | 97 | 139 | 126 | 167 | 101 | 136 | 124 | 111 | 100 | 217 | 106 |
| 94 | 61 | 74 | 71 | 78 | 51 | 75 | 74 | 55 | 82 | 126 | 80 |
| 9 | 131 | 107 | 140 | 124 | 92 | 160 | 108 | 180 | 79 | 86 | 101 |
| 0 | 80 | 16 | 61 | 54 | 38 | 94 | 44 | 122 | 30 | 58 | 22 |
| 0 | 0 | 0 | 0 | 299 | 0 | 205 | 149 | 0 | 0 | 184 | 43 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 44 | 0 |
| 648 | 504 | 392 | 420 | 560 | 227 | 1,080 | 504 | 360 | 432 | 432 | 258 |
| 648 | 585 | 392 | 448 | 896 | 280 | 1,116 | 504 | 720 | 432 | 672 | 258 |
| 120 | 102 | 129 | 127 | 155 | 98 | 141 | 116 | 139 | 88 | 140 | 81 |
| 89 | 63 | 66 | 75 | 82 | 52 | 80 | 70 | 111 | 68 | 112 | 54 |
| 28 | | | | 36 | 28 | 43 | 35 | | | | |
| 72 | | | | 90 | 56 | 86 | 70 | | | | |
| 105 | | | | 168 | 86 | 113 | 112 | | | | |
| 140 | | | | 224 | 112 | 170 | 140 | | | | |
| 196 | | | | 280 | 184 | 224 | 203 | | | | |
| 224 | | | | 280 | 224 | 258 | 224 | | | | |
| 648 | | | | 448 | 224 | 448 | 454 | | | | |
| 648 | 420 | 280 | 420 | 448 | 227 | 540 | 504 | 360 | 432 | 432 | 258 |
| 1.8 | 0 | 2.8 | 0 | 1.0 | 0 | 5.4 | 3.7 | 0 | 0 | 0 | 0 |
| 2.9 | 0 | 5.9 | 0 | 1.0 | 0 | 2.7 | 1.4 | 0 | 0 | 0 | 0 |

TABLE 7.12.--BACON
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 25.3 | 1.7 | 23.5 | 22.0 | 20.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 16.7 | 1.7 | 15.9 | 16.2 | 15.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 5.8 | 0 | 5.9 | 4.1 | 4.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.9 | 0 | 1.7 | 1.6 | 1.2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.5 | 1.0 | 1.4 | 1.4 | 1.3 |
| STANDARD DEVIATION.....DO..... | .7 | .0 | .6 | .6 | .6 |
| MAXIMUM.....DO..... | 6.0 | 1.0 | 3.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 12 | 4 | 7 | 7 | 9 |
| STANDARD DEVIATION.....GM..... | 11 | 3 | 7 | 7 | 7 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 3 | | 2 | 3 | 3 |
| 25TH.....GM..... | 5 | | 3 | 3 | 5 |
| 50TH.....GM..... | 8 | | 5 | 5 | 5 |
| 75TH.....GM..... | 13 | | 8 | 9 | 11 |
| 90TH.....GM..... | 21 | | 11 | 13 | 16 |
| 95TH.....GM..... | 32 | | 16 | 19 | 21 |
| 99TH.....GM..... | 57 | | 38 | 53 | 43 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 25 | 12 | 14 | 17 | 20 |
| STANDARD DEVIATION.....GM..... | 22 | 8 | 10 | 11 | 16 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 23 | 0 | 16 | 17 | 20 |
| STANDARD DEVIATION.....GM..... | 18 | 0 | 15 | 16 | 12 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 23 | 0 | 12 | 16 | 17 |
| STANDARD DEVIATION.....GM..... | 17 | 0 | 5 | 11 | 9 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 454 | 28 | 112 | 112 | 142 |
| MAXIMUM OVER 3 DAYS.....GM..... | 624 | 28 | 168 | 196 | 142 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 24 | 12 | 14 | 16 | 20 |
| STANDARD DEVIATION.....GM..... | 20 | 8 | 13 | 13 | 15 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 8 | | 4 | 5 | 8 |
| 25TH.....GM..... | 16 | | 8 | 8 | 12 |
| 50TH.....GM..... | 16 | | 8 | 16 | 16 |
| 75TH.....GM..... | 24 | | 16 | 16 | 24 |
| 90TH.....GM..... | 40 | | 24 | 24 | 32 |
| 95TH.....GM..... | 57 | | 28 | 36 | 48 |
| 99TH.....GM..... | 113 | | 85 | 84 | 85 |
| MAXIMUM.....GM..... | 454 | 28 | 112 | 112 | 142 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 125.2 | 0 | 6.9 | 4.1 | 3.2 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .9 | 0 | 2.0 | .8 | .6 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 23.4 | 20.7 | 28.6 | 21.2 | 26.9 | 21.6 | 31.7 | 26.7 | 32.2 | 26.5 | 40.1 | 24.0 |
| 17.4 | 16.5 | 18.4 | 15.8 | 17.8 | 14.9 | 18.7 | 17.7 | 16.9 | 14.9 | 18.8 | 14.4 |
| 4.3 | 3.4 | 7.7 | 4.3 | 6.1 | 4.6 | 8.2 | 5.8 | 8.6 | 7.8 | 11.4 | 5.5 |
| 1.7 | .8 | 2.6 | 1.1 | 3.0 | 2.1 | 4.8 | 3.2 | 6.8 | 3.8 | 9.9 | 4.0 |
| 1.4 | 1.3 | 1.5 | 1.3 | 1.5 | 1.4 | 1.6 | 1.5 | 1.7 | 1.7 | 1.9 | 1.6 |
| .6 | .6 | .7 | .6 | .7 | .7 | .8 | .7 | .9 | .9 | .9 | .8 |
| 4.0 | 5.0 | 4.0 | 4.0 | 4.0 | 5.0 | 5.0 | 5.0 | 4.0 | 6.0 | 6.0 | 5.0 |
| 10 | 10 | 13 | 10 | 16 | 11 | 14 | 10 | 14 | 11 | 14 | 10 |
| 8 | 9 | 11 | 8 | 14 | 10 | 13 | 10 | 13 | 11 | 17 | 8 |
| 3 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 3 | 3 |
| 5 | 5 | 5 | 5 | 8 | 5 | 7 | 5 | 5 | 5 | 5 | 5 |
| 8 | 8 | 11 | 8 | 11 | 8 | 11 | 8 | 11 | 8 | 11 | 8 |
| 13 | 11 | 16 | 13 | 19 | 13 | 16 | 11 | 16 | 13 | 16 | 12 |
| 19 | 19 | 27 | 19 | 32 | 21 | 27 | 19 | 24 | 21 | 28 | 19 |
| 27 | 27 | 32 | 27 | 41 | 32 | 38 | 27 | 37 | 33 | 38 | 27 |
| 46 | 56 | 57 | 53 | 76 | 57 | 62 | 57 | 76 | 57 | 76 | 38 |
| 24 | 22 | 28 | 22 | 35 | 24 | 30 | 22 | 24 | 22 | 27 | 24 |
| 20 | 16 | 23 | 14 | 33 | 19 | 23 | 19 | 16 | 17 | 32 | 26 |
| 21 | 24 | 27 | 26 | 31 | 24 | 24 | 21 | 26 | 19 | 29 | 18 |
| 12 | 20 | 16 | 16 | 24 | 18 | 14 | 15 | 26 | 13 | 37 | 9 |
| 21 | 30 | 25 | 22 | 30 | 24 | 28 | 20 | 22 | 24 | 19 | 14 |
| 13 | 18 | 15 | 12 | 18 | 16 | 22 | 14 | 12 | 18 | 11 | 7 |
| 170 | 227 | 227 | 144 | 454 | 144 | 227 | 251 | 170 | 256 | 454 | 170 |
| 255 | 340 | 227 | 178 | 454 | 240 | 624 | 284 | 284 | 304 | 470 | 170 |
| 23 | 23 | 27 | 23 | 32 | 23 | 26 | 21 | 24 | 20 | 23 | 18 |
| 18 | 17 | 21 | 17 | 27 | 18 | 21 | 17 | 18 | 17 | 32 | 18 |
| 8 | 8 | 8 | 8 | 10 | 8 | 8 | 8 | 8 | 8 | 8 | 8 |
| 16 | 16 | 16 | 16 | 16 | 16 | 16 | 15 | 16 | 10 | 16 | 8 |
| 16 | 16 | 24 | 16 | 24 | 16 | 24 | 16 | 16 | 16 | 16 | 16 |
| 24 | 24 | 32 | 24 | 32 | 24 | 32 | 24 | 24 | 24 | 24 | 20 |
| 40 | 40 | 48 | 40 | 57 | 40 | 48 | 32 | 40 | 32 | 38 | 28 |
| 57 | 57 | 57 | 57 | 84 | 57 | 57 | 57 | 57 | 57 | 57 | 36 |
| 112 | 113 | 113 | 85 | 113 | 113 | 113 | 85 | 113 | 85 | 113 | 113 |
| 170 | 113 | 227 | 144 | 454 | 144 | 227 | 227 | 146 | 240 | 454 | 170 |
| 6.3 | 7.2 | 3.9 | 8.1 | 8.8 | 29.6 | 15.4 | 21.7 | .9 | 5.5 | 1.8 | 1.8 |
| 1.0 | 1.3 | .7 | 2.0 | .6 | 1.8 | .6 | .8 | .1 | .7 | .4 | .5 |

TABLE 7.13.--LAMB
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 1.5 | 2.6 | 1.4 | .6 | .5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.4 | 1.7 | 1.4 | .6 | .5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .1 | .9 | 0 | 0 | 0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 1.4 | 1.0 | 1.0 | 1.0 |
| STANDARD DEVIATION.....DO..... | .3 | .5 | .0 | .0 | .0 |
| MAXIMUM.....DO..... | 3.0 | 2.0 | 1.0 | 1.0 | 1.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 54 | 24 | 24 | 30 | 46 |
| STANDARD DEVIATION.....GM..... | 33 | 13 | 15 | 20 | 29 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 15 | | | | |
| 25TH.....GM..... | 36 | | | | |
| 50TH.....GM..... | 46 | | | | |
| 75TH.....GM..... | 73 | | | | |
| 90TH.....GM..... | 93 | | | | |
| 95TH.....GM..... | 112 | | | | |
| 99TH.....GM..... | 149 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 149 | 56 | 72 | 90 | 139 |
| STANDARD DEVIATION.....GM..... | 88 | 34 | 46 | 59 | 86 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 141 | 49 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 63 | 15 | 0 | 0 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 64 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 0 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 864 | 128 | 128 | 218 | 336 |
| MAXIMUM OVER 3 DAYS.....GM..... | 864 | 136 | 128 | 218 | 336 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 146 | 52 | 72 | 90 | 139 |
| STANDARD DEVIATION.....GM..... | 84 | 29 | 46 | 59 | 86 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 43 | | | | |
| 25TH.....GM..... | 88 | | | | |
| 50TH.....GM..... | 123 | | | | |
| 75TH.....GM..... | 184 | | | | |
| 90TH.....GM..... | 227 | | | | |
| 95TH.....GM..... | 280 | | | | |
| 99TH.....GM..... | 448 | | | | |
| MAXIMUM.....GM..... | 864 | 128 | 128 | 218 | 336 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 9.6 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.5 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,156 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| .9 | .7 | .5 | 1.0 | 1.2 | 1.4 | 1.8 | 1.8 | 3.5 | 3.0 | 2.6 | 2.8 |
| .8 | .7 | .5 | .9 | .9 | 1.3 | 1.8 | 1.8 | 3.0 | 2.7 | 2.6 | 1.9 |
| .1 | 0 | .1 | .1 | .3 | .1 | .1 | .1 | .5 | .4 | 0 | .9 |
| 0 | 0 | 0 | 0 | 0 | (2) | 0 | 0 | 0 | 0 | 0 | 0 |
| 1.1 | 1.0 | 1.1 | 1.1 | 1.3 | 1.1 | 1.1 | 1.0 | 1.2 | 1.1 | 1.0 | 1.3 |
| .3 | .0 | .3 | .3 | .5 | .3 | .3 | .2 | .4 | .4 | .0 | .5 |
| 2.0 | 1.0 | 2.0 | 2.0 | 3.0 | 3.0 | 3.0 | 2.0 | 2.0 | 2.0 | 1.0 | 2.0 |
| 61 | 42 | 58 | 41 | 63 | 50 | 64 | 48 | 70 | 61 | 42 | 55 |
| 42 | 23 | 30 | 17 | 36 | 25 | 43 | 27 | 38 | 28 | 14 | 26 |
| | | | | | 19 | 28 | 15 | | 31 | | |
| | | | | | 31 | 38 | 28 | | 36 | | |
| | | | | | 37 | 56 | 46 | | 48 | | |
| | | | | | 73 | 73 | 65 | | 79 | | |
| | | | | | 79 | 97 | 76 | | 112 | | |
| | | | | | 109 | 149 | 93 | | 113 | | |
| | | | | | 112 | 288 | 149 | | 118 | | |
| 154 | 127 | 157 | 116 | 167 | 145 | 186 | 135 | 193 | 160 | 126 | 135 |
| 73 | 68 | 83 | 47 | 91 | 74 | 126 | 65 | 106 | 65 | 43 | 56 |
| 280 | 0 | 154 | 96 | 130 | 112 | 190 | 199 | 157 | 172 | 0 | 113 |
| 0 | 0 | 0 | 0 | 65 | 0 | 47 | 29 | 55 | 4 | 0 | 40 |
| 0 | 0 | 0 | 0 | 0 | 64 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 336 | 280 | 280 | 218 | 504 | 336 | 864 | 280 | 448 | 336 | 218 | 218 |
| 560 | 280 | 308 | 218 | 504 | 336 | 864 | 448 | 448 | 354 | 218 | 356 |
| 171 | 127 | 156 | 112 | 150 | 139 | 182 | 140 | 175 | 158 | 126 | 124 |
| 80 | 68 | 81 | 43 | 87 | 73 | 121 | 66 | 89 | 60 | 43 | 53 |
| | | | | | 40 | 84 | 45 | | 92 | | |
| | | | | | 92 | 112 | 85 | | 109 | | |
| | | | | | 112 | 168 | 138 | | 144 | | |
| | | | | | 168 | 218 | 196 | | 210 | | |
| | | | | | 237 | 280 | 224 | | 236 | | |
| | | | | | 253 | 290 | 276 | | 276 | | |
| | | | | | 336 | 864 | 280 | | 336 | | |
| 336 | 280 | 280 | 218 | 504 | 336 | 864 | 280 | 448 | 336 | 218 | 218 |
| .8 | 0 | 0 | 0 | 0 | 0 | 2.1 | 1.9 | 0 | 0 | 0 | 4.8 |
| 3.7 | 0 | 0 | 0 | 0 | 0 | 2.2 | 1.4 | 0 | 0 | 0 | 13.0 |

TABLE 7.14.--VEAL
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.3 | 3.2 | 1.2 | 1.6 | 2.0 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.1 | 1.9 | 1.2 | 1.6 | 2.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .2 | .5 | 0 | 0 | (2) |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | .8 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 1.1 | 2.0 | 1.0 | 1.0 | 1.0 |
| STANDARD DEVIATION.....DO..... | .4 | 1.2 | .0 | .0 | .2 |
| MAXIMUM.....DO..... | 5.0 | 5.0 | 1.0 | 1.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 49 | 36 | 27 | 25 | 40 |
| STANDARD DEVIATION.....GM..... | 27 | 30 | 9 | 11 | 26 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 15 | | | | |
| 25TH.....GM..... | 28 | | | | |
| 50TH.....GM..... | 38 | | | | |
| 75TH.....GM..... | 59 | | | | |
| 90TH.....GM..... | 90 | | | | |
| 95TH.....GM..... | 95 | | | | |
| 99TH.....GM..... | 144 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 138 | 67 | 80 | 75 | 118 |
| STANDARD DEVIATION.....GM..... | 74 | 44 | 28 | 33 | 78 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 121 | 48 | 0 | 0 | 110 |
| STANDARD DEVIATION.....GM..... | 45 | 19 | 0 | 0 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 72 | 72 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 35 | 35 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 448 | 128 | 128 | 138 | 336 |
| MAXIMUM OVER 3 DAYS.....GM..... | 504 | 340 | 128 | 138 | 336 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 130 | 54 | 80 | 75 | 115 |
| STANDARD DEVIATION.....GM..... | 71 | 37 | 28 | 33 | 72 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 42 | | | | |
| 25TH.....GM..... | 84 | | | | |
| 50TH.....GM..... | 112 | | | | |
| 75TH.....GM..... | 168 | | | | |
| 90TH.....GM..... | 224 | | | | |
| 95TH.....GM..... | 276 | | | | |
| 99TH.....GM..... | 352 | | | | |
| MAXIMUM.....GM..... | 448 | 128 | 128 | 138 | 336 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 26.2 | 0 | 0 | 3.9 | .8 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.7 | 0 | 0 | 13.9 | 2.0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.5 | 1.5 | 1.5 | 2.1 | 2.1 | 1.9 | 2.6 | 2.7 | 2.4 | 4.2 | 3.8 | 3.2 |
| 1.4 | 1.4 | 1.3 | 2.0 | 2.0 | 1.7 | 2.4 | 2.5 | 1.9 | 3.2 | 3.1 | 2.9 |
| .1 | .1 | .2 | .1 | .1 | .1 | .2 | .2 | .5 | 1.0 | .7 | .3 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.3 | 1.4 | 1.1 |
| .3 | .3 | .3 | .3 | .5 | .3 | .4 | .3 | .4 | .6 | .9 | .3 |
| 2.0 | 2.0 | 2.0 | 2.0 | 4.0 | 2.0 | 3.0 | 3.0 | 2.0 | 3.0 | 4.0 | 2.0 |
| 45 | 35 | 64 | 48 | 57 | 50 | 60 | 48 | 61 | 49 | 42 | 44 |
| 28 | 21 | 34 | 25 | 27 | 21 | 32 | 24 | 29 | 24 | 24 | 29 |
| | | | | 19 | 28 | 24 | 19 | | 9 | | |
| | | | | 37 | 37 | 37 | 32 | | 36 | | |
| | | | | 56 | 43 | 56 | 39 | | 44 | | |
| | | | | 75 | 72 | 75 | 59 | | 59 | | |
| | | | | 93 | 77 | 93 | 84 | | 89 | | |
| | | | | 112 | 93 | 117 | 90 | | 90 | | |
| | | | | 135 | 93 | 168 | 114 | | 112 | | |
| 124 | 99 | 175 | 136 | 165 | 142 | 168 | 140 | 171 | 127 | 101 | 113 |
| 81 | 61 | 98 | 67 | 78 | 56 | 83 | 70 | 93 | 56 | 27 | 47 |
| 122 | 112 | 152 | 142 | 164 | 130 | 169 | 105 | 114 | 103 | 114 | 159 |
| 9 | 0 | 2 | 35 | 21 | 12 | 53 | 27 | 22 | 42 | 51 | 68 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 448 | 280 | 405 | 336 | 405 | 280 | 448 | 405 | 360 | 270 | 172 | 224 |
| 448 | 280 | 405 | 337 | 405 | 280 | 504 | 405 | 360 | 336 | 344 | 448 |
| 124 | 96 | 170 | 131 | 156 | 140 | 164 | 131 | 150 | 111 | 93 | 121 |
| 75 | 46 | 87 | 62 | 76 | 53 | 84 | 63 | 86 | 57 | 25 | 54 |
| | | | | 56 | 84 | 71 | 43 | | 28 | | |
| | | | | 88 | 112 | 112 | 84 | | 65 | | |
| | | | | 138 | 116 | 140 | 112 | | 112 | | |
| | | | | 224 | 168 | 220 | 168 | | 140 | | |
| | | | | 276 | 224 | 280 | 224 | | 211 | | |
| | | | | 280 | 232 | 336 | 264 | | 220 | | |
| | | | | 405 | 280 | 432 | 336 | | 270 | | |
| 448 | 224 | 405 | 336 | 405 | 280 | 448 | 342 | 360 | 270 | 168 | 224 |
| 3.2 | 0 | .9 | .9 | 0 | 0 | 3.5 | 4.7 | 0 | 8.3 | 0 | 0 |
| 9.4 | 0 | 3.9 | 2.8 | 0 | 0 | 2.5 | 2.2 | 0 | 8.7 | 0 | 0 |

TABLE 7.15.--POULTRY
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,900 | 524 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 42.8 | 18.2 | 42.2 | 42.6 | 45.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 33.0 | 11.6 | 32.1 | 34.1 | 36.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 8.8 | 5.3 | 8.4 | 8.1 | 7.9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.1 | 1.2 | 1.7 | .4 | 1.0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 1.6 | 1.4 | 1.3 | 1.3 |
| STANDARD DEVIATION.....DO..... | .6 | .8 | .7 | .5 | .6 |
| MAXIMUM.....DO..... | 7.0 | 4.0 | 4.0 | 4.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 57 | 31 | 33 | 38 | 43 |
| STANDARD DEVIATION.....GM..... | 39 | 25 | 23 | 24 | 27 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 17 | 3 | 9 | 12 | 14 |
| 25TH.....GM..... | 32 | 14 | 18 | 20 | 27 |
| 50TH.....GM..... | 46 | 28 | 28 | 33 | 37 |
| 75TH.....GM..... | 73 | 43 | 38 | 47 | 56 |
| 90TH.....GM..... | 109 | 58 | 66 | 69 | 75 |
| 95TH.....GM..... | 132 | 84 | 85 | 82 | 100 |
| 99TH.....GM..... | 191 | 134 | 112 | 114 | 141 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 140 | 64 | 77 | 96 | 110 |
| STANDARD DEVIATION.....GM..... | 84 | 45 | 43 | 52 | 59 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 133 | 65 | 82 | 90 | 102 |
| STANDARD DEVIATION.....GM..... | 68 | 34 | 42 | 44 | 48 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 122 | 75 | 65 | 93 | 97 |
| STANDARD DEVIATION.....GM..... | 55 | 42 | 30 | 27 | 45 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,282 | 227 | 336 | 485 | 679 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,282 | 411 | 418 | 582 | 679 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 128 | 60 | 73 | 90 | 103 |
| STANDARD DEVIATION.....GM..... | 77 | 38 | 44 | 50 | 56 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 42 | 9 | 20 | 28 | 32 |
| 25TH.....GM..... | 82 | 33 | 40 | 56 | 60 |
| 50TH.....GM..... | 112 | 56 | 60 | 85 | 97 |
| 75TH.....GM..... | 168 | 84 | 100 | 112 | 113 |
| 90TH.....GM..... | 224 | 128 | 113 | 150 | 186 |
| 95TH.....GM..... | 280 | 128 | 140 | 194 | 220 |
| 99TH.....GM..... | 388 | 142 | 222 | 234 | 291 |
| MAXIMUM.....GM..... | 990 | 213 | 336 | 448 | 567 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 115.1 | 1.7 | 2.5 | 4.0 | 7.8 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .5 | 1.1 | .4 | .4 | .7 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 44.3 | 44.0 | 43.8 | 43.7 | 39.9 | 43.2 | 43.1 | 44.6 | 42.4 | 45.1 | 36.1 | 40.3 |
| 35.8 | 35.9 | 34.4 | 36.0 | 30.4 | 33.3 | 33.7 | 33.8 | 29.1 | 32.7 | 24.2 | 27.6 |
| 7.9 | 7.6 | 8.4 | 6.9 | 8.2 | 9.0 | 8.4 | 9.7 | 11.5 | 10.8 | 9.9 | 10.4 |
| .7 | .5 | 1.0 | .9 | 1.2 | .9 | 1.1 | 1.0 | 1.8 | 1.7 | 1.9 | 2.3 |
| 1.3 | 1.3 | 1.3 | 1.3 | 1.3 | 1.3 | 1.3 | 1.4 | 1.5 | 1.4 | 1.5 | 1.5 |
| .6 | .6 | .6 | .5 | .6 | .6 | .6 | .6 | .7 | .7 | .7 | .8 |
| 4.0 | 6.0 | 5.0 | 4.0 | 5.0 | 5.0 | 7.0 | 6.0 | 5.0 | 6.0 | 4.0 | 6.0 |
| 56 | 47 | 68 | 51 | 76 | 54 | 71 | 55 | 65 | 55 | 59 | 55 |
| 37 | 29 | 45 | 31 | 49 | 33 | 45 | 35 | 41 | 35 | 36 | 37 |
| 19 | 17 | 19 | 19 | 19 | 17 | 20 | 19 | 20 | 18 | 16 | 18 |
| 32 | 28 | 37 | 30 | 38 | 32 | 38 | 32 | 37 | 31 | 32 | 32 |
| 43 | 37 | 56 | 42 | 65 | 45 | 60 | 44 | 56 | 42 | 50 | 46 |
| 71 | 61 | 93 | 65 | 97 | 66 | 91 | 68 | 85 | 70 | 75 | 71 |
| 109 | 82 | 129 | 93 | 140 | 98 | 129 | 97 | 132 | 97 | 118 | 94 |
| 124 | 104 | 159 | 109 | 170 | 119 | 151 | 118 | 151 | 124 | 131 | 122 |
| 187 | 141 | 208 | 162 | 241 | 173 | 231 | 175 | 189 | 187 | 151 | 253 |
| 141 | 120 | 166 | 133 | 186 | 132 | 173 | 134 | 148 | 126 | 139 | 125 |
| 80 | 61 | 104 | 73 | 109 | 70 | 94 | 77 | 90 | 71 | 82 | 66 |
| 136 | 115 | 161 | 125 | 178 | 127 | 168 | 125 | 148 | 121 | 118 | 121 |
| 67 | 50 | 68 | 52 | 82 | 55 | 78 | 60 | 63 | 54 | 48 | 71 |
| 133 | 122 | 160 | 96 | 132 | 125 | 161 | 116 | 115 | 125 | 119 | 100 |
| 43 | 64 | 64 | 38 | 63 | 44 | 66 | 47 | 31 | 45 | 38 | 34 |
| 681 | 509 | 1,282 | 582 | 990 | 595 | 968 | 968 | 568 | 560 | 454 | 531 |
| 867 | 837 | 1,282 | 634 | 1,029 | 709 | 1,136 | 1,132 | 673 | 691 | 638 | 782 |
| 131 | 112 | 153 | 123 | 169 | 124 | 158 | 121 | 134 | 114 | 115 | 108 |
| 75 | 58 | 85 | 68 | 101 | 66 | 88 | 69 | 70 | 61 | 58 | 60 |
| 51 | 43 | 54 | 43 | 56 | 43 | 56 | 43 | 51 | 43 | 46 | 38 |
| 84 | 70 | 97 | 82 | 98 | 84 | 97 | 84 | 86 | 80 | 84 | 62 |
| 112 | 101 | 130 | 112 | 149 | 112 | 140 | 111 | 113 | 102 | 112 | 97 |
| 170 | 140 | 194 | 152 | 220 | 164 | 200 | 142 | 170 | 138 | 131 | 114 |
| 224 | 194 | 279 | 220 | 291 | 209 | 276 | 209 | 227 | 194 | 173 | 194 |
| 291 | 222 | 320 | 254 | 346 | 246 | 321 | 228 | 280 | 227 | 209 | 227 |
| 385 | 302 | 420 | 388 | 485 | 336 | 440 | 376 | 340 | 340 | 284 | 340 |
| 681 | 485 | 679 | 520 | 990 | 595 | 968 | 968 | 456 | 454 | 454 | 393 |
| 4.5 | 5.7 | 5.5 | 3.3 | 19.1 | 22.1 | 20.0 | 6.5 | .8 | 6.5 | 2.2 | 3.0 |
| .4 | .5 | .7 | .4 | .9 | .7 | .7 | .2 | .1 | .6 | .7 | .5 |

TABLE 7.16.--CHICKEN
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1: | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 38.7 | 15.6 | 38.8 | 39.3 | 41.4 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 31.2 | 10.5 | 30.7 | 32.7 | 34.9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 6.8 | 4.1 | 6.9 | 6.2 | 6.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .6 | 1.1 | 1.2 | .4 | .5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 1.5 | 1.3 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .6 | .8 | .6 | .5 | .5 |
| MAXIMUM.....DO..... | 7.0 | 4.0 | 4.0 | 4.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 56 | 32 | 32 | 37 | 43 |
| STANDARD DEVIATION.....GM..... | 36 | 26 | 22 | 23 | 26 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 17 | 5 | 9 | 12 | 14 |
| 25TH.....GM..... | 32 | 14 | 17 | 21 | 27 |
| 50TH.....GM..... | 45 | 24 | 27 | 32 | 37 |
| 75TH.....GM..... | 70 | 38 | 37 | 43 | 55 |
| 90TH.....GM..... | 102 | 58 | 65 | 65 | 75 |
| 95TH.....GM..... | 129 | 85 | 81 | 78 | 99 |
| 99TH.....GM..... | 185 | 137 | 103 | 114 | 129 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 141 | 65 | 76 | 97 | 111 |
| STANDARD DEVIATION.....GM..... | 83 | 39 | 42 | 52 | 58 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 133 | 66 | 83 | 93 | 105 |
| STANDARD DEVIATION.....GM..... | 66 | 37 | 41 | 46 | 46 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 124 | 81 | 60 | 88 | 121 |
| STANDARD DEVIATION.....GM..... | 55 | 44 | 33 | 25 | 46 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,282 | 213 | 336 | 485 | 390 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,282 | 411 | 418 | 582 | 630 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 131 | 62 | 73 | 92 | 106 |
| STANDARD DEVIATION.....GM..... | 76 | 39 | 43 | 50 | 55 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 43 | 14 | 20 | 28 | 38 |
| 25TH.....GM..... | 84 | 35 | 40 | 56 | 61 |
| 50TH.....GM..... | 112 | 56 | 61 | 89 | 97 |
| 75TH.....GM..... | 170 | 85 | 101 | 112 | 116 |
| 90TH.....GM..... | 224 | 128 | 113 | 150 | 194 |
| 95TH.....GM..... | 280 | 128 | 131 | 194 | 220 |
| 99TH.....GM..... | 388 | 135 | 222 | 234 | 291 |
| MAXIMUM.....GM..... | 990 | 213 | 336 | 448 | 390 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 80.2 | 1.7 | 1.6 | 4.0 | 5.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 1.4 | .3 | .5 | .6 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 39.8 | 39.6 | 38.9 | 39.5 | 34.7 | 39.3 | 38.8 | 40.3 | 38.7 | 41.4 | 33.3 | 37.8 |
| 33.7 | 33.4 | 32.5 | 34.4 | 27.6 | 31.6 | 31.9 | 32.1 | 28.5 | 31.2 | 24.8 | 26.3 |
| 6.0 | 6.0 | 6.1 | 4.9 | 6.5 | 7.2 | 6.2 | 7.6 | 9.1 | 8.9 | 6.9 | 9.4 |
| .2 | .2 | .3 | .2 | .6 | .5 | .7 | .6 | 1.1 | 1.3 | 1.6 | 2.1 |
| 1.2 | 1.2 | 1.2 | 1.2 | 1.3 | 1.3 | 1.3 | 1.3 | 1.4 | 1.4 | 1.5 | 1.5 |
| .5 | .5 | .5 | .4 | .5 | .5 | .6 | .6 | .6 | .7 | .7 | .7 |
| 3.0 | 4.0 | 4.0 | 4.0 | 3.0 | 5.0 | 7.0 | 6.0 | 5.0 | 6.0 | 4.0 | 6.0 |
| 55 | 47 | 66 | 51 | 75 | 53 | 69 | 52 | 63 | 53 | 58 | 55 |
| 35 | 27 | 43 | 29 | 46 | 32 | 42 | 32 | 40 | 35 | 34 | 36 |
| 19 | 18 | 20 | 19 | 27 | 18 | 23 | 19 | 19 | 18 | 16 | 18 |
| 32 | 28 | 37 | 32 | 41 | 32 | 38 | 32 | 34 | 31 | 32 | 32 |
| 42 | 37 | 56 | 41 | 65 | 43 | 60 | 42 | 50 | 42 | 48 | 41 |
| 70 | 59 | 86 | 65 | 97 | 65 | 87 | 65 | 80 | 70 | 75 | 71 |
| 102 | 78 | 129 | 87 | 135 | 95 | 124 | 93 | 129 | 94 | 115 | 93 |
| 122 | 102 | 153 | 102 | 166 | 113 | 150 | 113 | 151 | 122 | 131 | 113 |
| 173 | 138 | 191 | 162 | 228 | 168 | 202 | 161 | 187 | 187 | 151 | 253 |
| 144 | 121 | 170 | 136 | 193 | 135 | 177 | 133 | 148 | 126 | 139 | 127 |
| 79 | 60 | 103 | 73 | 106 | 69 | 91 | 74 | 90 | 72 | 82 | 66 |
| 140 | 118 | 167 | 129 | 175 | 126 | 164 | 123 | 145 | 124 | 135 | 117 |
| 72 | 50 | 72 | 50 | 83 | 54 | 75 | 56 | 62 | 51 | 44 | 70 |
| 117 | 118 | 154 | 98 | 140 | 135 | 169 | 111 | 122 | 127 | 106 | 102 |
| 31 | 17 | 29 | 17 | 57 | 44 | 67 | 49 | 35 | 50 | 25 | 35 |
| 681 | 485 | 1,282 | 582 | 990 | 520 | 907 | 880 | 568 | 560 | 454 | 531 |
| 867 | 539 | 1,282 | 582 | 990 | 709 | 1,136 | 907 | 582 | 691 | 500 | 782 |
| 136 | 115 | 160 | 128 | 177 | 127 | 162 | 122 | 134 | 116 | 120 | 109 |
| 77 | 57 | 87 | 70 | 99 | 65 | 85 | 67 | 70 | 60 | 58 | 59 |
| 52 | 45 | 56 | 51 | 60 | 49 | 57 | 47 | 51 | 45 | 49 | 38 |
| 85 | 78 | 97 | 85 | 111 | 84 | 97 | 84 | 85 | 80 | 85 | 65 |
| 112 | 105 | 140 | 112 | 162 | 112 | 148 | 112 | 112 | 104 | 112 | 97 |
| 170 | 140 | 200 | 164 | 224 | 168 | 209 | 149 | 170 | 140 | 145 | 114 |
| 224 | 194 | 284 | 224 | 304 | 209 | 279 | 209 | 227 | 194 | 182 | 189 |
| 291 | 224 | 328 | 255 | 352 | 244 | 320 | 227 | 280 | 227 | 224 | 226 |
| 388 | 284 | 448 | 388 | 485 | 336 | 440 | 352 | 340 | 336 | 454 | 340 |
| 681 | 485 | 679 | 520 | 990 | 520 | 907 | 680 | 456 | 454 | 454 | 393 |
| 3.2 | 2.8 | 3.7 | 2.5 | 13.4 | 19.3 | 9.7 | 3.5 | .8 | 4.8 | 2.2 | 1.1 |
| .3 | .3 | .6 | .4 | .8 | .7 | .4 | .1 | .1 | .5 | .8 | .2 |

TABLE 7.17.--TURKEY
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.8 | 5.1 | 4.4 | 4.5 | 5.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.9 | 4.6 | 3.9 | 3.8 | 4.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .8 | .2 | .4 | .7 | .7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | .2 | .1 | 0 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 1.3 | 1.2 | 1.2 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .6 | .5 | .5 | .5 | .5 |
| MAXIMUM.....DO..... | 5.0 | 3.0 | 3.0 | 3.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 45 | 21 | 30 | 29 | 30 |
| STANDARD DEVIATION.....GM..... | 35 | 15 | 23 | 17 | 18 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 9 | | | 9 | 9 |
| 25TH.....GM..... | 19 | | | 19 | 19 |
| 50TH.....GM..... | 37 | | | 28 | 28 |
| 75TH.....GM..... | 57 | | | 38 | 38 |
| 90TH.....GM..... | 85 | | | 53 | 54 |
| 95TH.....GM..... | 113 | | | 75 | 60 |
| 99TH.....GM..... | 161 | | | 84 | 106 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 114 | 58 | 79 | 78 | 80 |
| STANDARD DEVIATION.....GM..... | 85 | 42 | 64 | 43 | 46 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 116 | 50 | 84 | 73 | 62 |
| STANDARD DEVIATION.....GM..... | 66 | 0 | 37 | 30 | 29 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 92 | 47 | 54 | 0 | 67 |
| STANDARD DEVIATION.....GM..... | 40 | 0 | 0 | 0 | 33 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 968 | 142 | 318 | 248 | 318 |
| MAXIMUM OVER 3 DAYS.....GM..... | 968 | 142 | 318 | 253 | 350 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 105 | 53 | 73 | 74 | 74 |
| STANDARD DEVIATION.....GM..... | 73 | 34 | 59 | 39 | 44 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 28 | | | 28 | 28 |
| 25TH.....GM..... | 57 | | | 56 | 43 |
| 50TH.....GM..... | 86 | | | 57 | 57 |
| 75TH.....GM..... | 129 | | | 85 | 86 |
| 90TH.....GM..... | 172 | | | 140 | 129 |
| 95TH.....GM..... | 240 | | | 159 | 168 |
| 99TH.....GM..... | 350 | | | 248 | 224 |
| MAXIMUM.....GM..... | 968 | 142 | 318 | 248 | 318 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 26.8 | 0 | .9 | 0 | 1.8 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.0 | 0 | 1.6 | 0 | 1.5 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 6.5 | 6.2 | 7.5 | 6.2 | 6.7 | 5.5 | 6.1 | 6.1 | 4.3 | 4.8 | 3.7 | 3.8 |
| 5.7 | 5.4 | 6.5 | 4.9 | 5.5 | 4.8 | 5.1 | 5.0 | 3.2 | 3.5 | 1.7 | 3.7 |
| .7 | .7 | .8 | 1.0 | .9 | .6 | .9 | .9 | .5 | 1.1 | 2.0 | .2 |
| .1 | (2) | .1 | .3 | .2 | .1 | .1 | .1 | .6 | .2 | 0 | 0 |
| 1.2 | 1.2 | 1.2 | 1.3 | 1.3 | 1.2 | 1.2 | 1.3 | 1.4 | 1.4 | 1.7 | 1.1 |
| .5 | .4 | .5 | .6 | .6 | .5 | .5 | .6 | .7 | .7 | .7 | .3 |
| 3.0 | 3.0 | 4.0 | 4.0 | 4.0 | 4.0 | 5.0 | 4.0 | 3.0 | 4.0 | 3.0 | 2.0 |
| 42 | 36 | 49 | 39 | 52 | 39 | 54 | 49 | 60 | 48 | 48 | 33 |
| 28 | 26 | 39 | 27 | 44 | 26 | 40 | 39 | 31 | 33 | 20 | 20 |
| 14 | 9 | 14 | 11 | 9 | 9 | 14 | 9 | | 14 | | |
| 23 | 19 | 27 | 23 | 23 | 19 | 28 | 28 | | 28 | | |
| 29 | 28 | 38 | 29 | 38 | 35 | 43 | 38 | | 38 | | |
| 55 | 43 | 57 | 43 | 72 | 57 | 75 | 57 | | 56 | | |
| 84 | 57 | 104 | 76 | 103 | 76 | 113 | 95 | | 105 | | |
| 100 | 86 | 140 | 106 | 129 | 86 | 124 | 113 | | 132 | | |
| 123 | 144 | 208 | 151 | 261 | 106 | 173 | 233 | | 152 | | |
| 106 | 97 | 120 | 95 | 125 | 107 | 139 | 125 | 150 | 122 | 119 | 94 |
| 59 | 73 | 80 | 58 | 87 | 66 | 105 | 105 | 72 | 86 | 66 | 56 |
| 129 | 88 | 153 | 102 | 164 | 100 | 136 | 120 | 115 | 82 | 84 | 99 |
| 52 | 37 | 74 | 55 | 102 | 50 | 58 | 57 | 38 | 37 | 20 | 0 |
| 95 | 86 | 152 | 66 | 78 | 66 | 88 | 107 | 105 | 134 | 0 | 0 |
| 28 | 0 | 76 | 8 | 28 | 0 | 57 | 30 | 20 | 17 | 0 | 0 |
| 344 | 509 | 484 | 484 | 562 | 340 | 968 | 968 | 454 | 454 | 284 | 227 |
| 547 | 509 | 680 | 484 | 842 | 480 | 968 | 968 | 454 | 455 | 284 | 227 |
| 103 | 90 | 120 | 89 | 119 | 99 | 129 | 110 | 129 | 102 | 86 | 90 |
| 56 | 54 | 68 | 47 | 82 | 63 | 92 | 80 | 59 | 63 | 46 | 54 |
| 43 | 28 | 42 | 28 | 28 | 28 | 43 | 28 | | 36 | | |
| 57 | 57 | 70 | 57 | 57 | 56 | 84 | 57 | | 57 | | |
| 86 | 85 | 112 | 85 | 113 | 85 | 113 | 86 | | 86 | | |
| 113 | 112 | 142 | 113 | 159 | 113 | 170 | 113 | | 113 | | |
| 170 | 159 | 227 | 159 | 227 | 170 | 227 | 171 | | 140 | | |
| 227 | 172 | 258 | 210 | 258 | 247 | 284 | 240 | | 170 | | |
| 327 | 350 | 340 | 242 | 454 | 318 | 363 | 376 | | 454 | | |
| 327 | 432 | 340 | 248 | 454 | 340 | 968 | 968 | 454 | 454 | 284 | 227 |
| 0 | .9 | 1.8 | .8 | 4.5 | 2.8 | 6.7 | 3.0 | 0 | 1.7 | 0 | 1.9 |
| 0 | .6 | 1.4 | .7 | 1.3 | .8 | 1.8 | .5 | 0 | 1.5 | 0 | 4.6 |

TABLE 7.18.--LIVER
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1: | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.8 | 1.5 | 3.5 | 2.9 | 2.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.6 | 1.4 | 3.3 | 2.8 | 2.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .2 | .2 | .1 | .1 | .1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 1.1 | 1.1 | 1.0 | 1.0 |
| STANDARD DEVIATION.....DO..... | .3 | .3 | .2 | .2 | .2 |
| MAXIMUM.....DO..... | 3.0 | 2.0 | 2.0 | 2.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 43 | 29 | 16 | 24 | 26 |
| STANDARD DEVIATION.....GM..... | 28 | 20 | 10 | 15 | 14 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 9 | | | 6 | 9 |
| 25TH.....GM..... | 25 | | | 14 | 19 |
| 50TH.....GM..... | 37 | | | 19 | 28 |
| 75TH.....GM..... | 56 | | | 35 | 37 |
| 90TH.....GM..... | 75 | | | 37 | 38 |
| 95TH.....GM..... | 90 | | | 56 | 47 |
| 99TH.....GM..... | 149 | | | 75 | 75 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 123 | 67 | 49 | 71 | 76 |
| STANDARD DEVIATION.....GM..... | 80 | 36 | 30 | 44 | 38 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 102 | 112 | 28 | 56 | 72 |
| STANDARD DEVIATION.....GM..... | 49 | 0 | 0 | 0 | 20 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 112 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 0 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 907 | 128 | 126 | 224 | 224 |
| MAXIMUM OVER 3 DAYS.....GM..... | 907 | 224 | 126 | 224 | 224 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 118 | 77 | 46 | 71 | 75 |
| STANDARD DEVIATION.....GM..... | 76 | 37 | 27 | 43 | 37 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 27 | | | 18 | 28 |
| 25TH.....GM..... | 70 | | | 43 | 56 |
| 50TH.....GM..... | 112 | | | 56 | 75 |
| 75TH.....GM..... | 150 | | | 86 | 100 |
| 90TH.....GM..... | 224 | | | 112 | 113 |
| 95TH.....GM..... | 227 | | | 168 | 113 |
| 99TH.....GM..... | 360 | | | 224 | 224 |
| MAXIMUM.....GM..... | 907 | 128 | 113 | 224 | 224 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 42.9 | 0 | 1.1 | 1.7 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.8 | 0 | 2.9 | 3.4 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.
2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 2.8 | 2.3 | 3.6 | 2.1 | 3.3 | 2.9 | 4.6 | 4.8 | 5.8 | 5.8 | 4.2 | 5.2 |
| 2.8 | 2.3 | 3.5 | 2.1 | 3.3 | 2.9 | 4.4 | 4.5 | 5.4 | 4.7 | 4.2 | 4.2 |
| 0 | 0 | .1 | 0 | (2) | (2) | .2 | .4 | .4 | 1.0 | 0 | 1.0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | (2) | 0 | 0 | 0 | 0 |
| 1.1 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.1 | 1.1 | 1.1 | 1.3 | 1.0 | 1.3 |
| .2 | .0 | .2 | .0 | .1 | .1 | .3 | .3 | .3 | .5 | .0 | .4 |
| 2.0 | 1.0 | 2.0 | 1.0 | 2.0 | 2.0 | 2.0 | 3.0 | 2.0 | 3.0 | 1.0 | 2.0 |
| 35 | 30 | 44 | 34 | 56 | 42 | 49 | 43 | 44 | 49 | 44 | 41 |
| 21 | 17 | 27 | 19 | 33 | 37 | 25 | 25 | 23 | 30 | 23 | 27 |
| 8 | 7 | 9 | | 12 | 9 | 18 | 13 | 9 | 9 | | 9 |
| 19 | 19 | 28 | | 33 | 28 | 33 | 28 | 28 | 28 | | 21 |
| 28 | 28 | 37 | | 48 | 37 | 47 | 37 | 37 | 37 | | 37 |
| 47 | 37 | 56 | | 75 | 50 | 57 | 56 | 56 | 75 | | 56 |
| 57 | 56 | 76 | | 93 | 75 | 76 | 75 | 75 | 76 | | 66 |
| 68 | 56 | 93 | | 113 | 75 | 93 | 82 | 93 | 105 | | 75 |
| 112 | 93 | 131 | | 149 | 302 | 149 | 149 | 112 | 149 | | 151 |
| 105 | 91 | 131 | 103 | 167 | 126 | 144 | 121 | 128 | 126 | 131 | 116 |
| 63 | 50 | 83 | 57 | 100 | 111 | 71 | 66 | 66 | 80 | 68 | 88 |
| 0 | 0 | 57 | 0 | 98 | 156 | 116 | 106 | 82 | 123 | 0 | 76 |
| 0 | 0 | 0 | 0 | 0 | 0 | 53 | 54 | 50 | 38 | 0 | 21 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 112 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 336 | 280 | 392 | 336 | 448 | 907 | 492 | 448 | 336 | 448 | 280 | 454 |
| 336 | 280 | 392 | 336 | 448 | 907 | 504 | 504 | 336 | 448 | 280 | 454 |
| 99 | 91 | 128 | 103 | 166 | 125 | 138 | 118 | 120 | 115 | 131 | 98 |
| 61 | 50 | 82 | 57 | 99 | 110 | 70 | 66 | 61 | 57 | 68 | 52 |
| 28 | 22 | 28 | | 35 | 20 | 43 | 25 | 26 | 36 | | 43 |
| 56 | 56 | 57 | | 98 | 84 | 86 | 84 | 84 | 70 | | 57 |
| 84 | 84 | 112 | | 140 | 112 | 125 | 112 | 112 | 112 | | 85 |
| 112 | 112 | 168 | | 224 | 150 | 168 | 150 | 168 | 140 | | 112 |
| 168 | 168 | 227 | | 280 | 198 | 224 | 224 | 196 | 224 | | 168 |
| 204 | 168 | 280 | | 340 | 224 | 280 | 227 | 224 | 224 | | 227 |
| 336 | 280 | 392 | | 448 | 907 | 448 | 360 | 280 | 324 | | 227 |
| 336 | 280 | 392 | 336 | 448 | 907 | 492 | 448 | 280 | 324 | 280 | 227 |
| .7 | 0 | .7 | 1.3 | 2.9 | 6.7 | 9.4 | 14.1 | 2.2 | 0 | 1.0 | 1.0 |
| 1.1 | 0 | 1.4 | 4.3 | 2.2 | 4.2 | 3.9 | 3.8 | 3.1 | 0 | 4.5 | 1.6 |

TABLE 7.19.--FISH AND SHELLFISH
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 24.5 | 2.3 | 17.3 | 20.4 | 22.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 20.5 | 2.2 | 14.2 | 18.2 | 19.9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.6 | .2 | 2.5 | 2.1 | 2.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .4 | 0 | .6 | .2 | .3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER.... | 1.2 | 1.1 | 1.3 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .5 | .3 | .7 | .5 | .4 |
| MAXIMUM.....DO..... | 6.0 | 2.0 | 5.0 | 5.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 48 | 15 | 22 | 27 | 32 |
| STANDARD DEVIATION.....GM..... | 42 | 9 | 18 | 19 | 24 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 8 | | 3 | 4 | 6 |
| 25TH.....GM..... | 20 | | 9 | 13 | 19 |
| 50TH.....GM..... | 37 | | 19 | 25 | 28 |
| 75TH.....GM..... | 57 | | 28 | 37 | 38 |
| 90TH.....GM..... | 94 | | 47 | 49 | 57 |
| 95TH.....GM..... | 128 | | 56 | 57 | 75 |
| 99TH.....GM..... | 215 | | 93 | 96 | 131 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 124 | 48 | 55 | 75 | 86 |
| STANDARD DEVIATION.....GM..... | 106 | 26 | 40 | 53 | 61 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 120 | 7 | 56 | 65 | 83 |
| STANDARD DEVIATION.....GM..... | 83 | 0 | 38 | 37 | 57 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 111 | 0 | 54 | 59 | 56 |
| STANDARD DEVIATION.....GM..... | 53 | 0 | 10 | 12 | 15 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 2,350 | 91 | 228 | 432 | 524 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,350 | 91 | 324 | 432 | 580 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 117 | 42 | 52 | 70 | 81 |
| STANDARD DEVIATION.....GM..... | 98 | 28 | 38 | 51 | 58 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 20 | | 8 | 12 | 19 |
| 25TH.....GM..... | 57 | | 28 | 36 | 40 |
| 50TH.....GM..... | 85 | | 43 | 57 | 72 |
| 75TH.....GM..... | 152 | | 58 | 85 | 112 |
| 90TH.....GM..... | 227 | | 112 | 113 | 160 |
| 95TH.....GM..... | 284 | | 125 | 170 | 170 |
| 99TH.....GM..... | 456 | | 168 | 240 | 288 |
| MAXIMUM.....GM..... | 1,410 | 91 | 228 | 432 | 448 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER.... | 736.0 | .8 | 8.3 | 23.1 | 34.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 6.5 | 6.1 | 3.6 | 5.7 | 7.1 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 22.0 | 22.9 | 21.1 | 24.2 | 25.1 | 25.7 | 26.9 | 28.0 | 27.5 | 24.7 | 20.7 | 19.6 |
| 19.1 | 20.4 | 18.6 | 20.7 | 20.5 | 20.9 | 21.9 | 23.2 | 21.7 | 20.6 | 17.9 | 17.1 |
| 2.5 | 2.1 | 2.3 | 3.2 | 4.3 | 4.2 | 4.4 | 4.4 | 5.0 | 3.8 | 2.0 | 2.1 |
| .3 | .3 | .2 | .4 | .3 | .6 | .6 | .3 | .7 | .3 | .9 | .3 |
| 1.2 | 1.2 | 1.2 | 1.2 | 1.3 | 1.3 | 1.3 | 1.2 | 1.3 | 1.2 | 1.2 | 1.2 |
| .5 | .4 | .5 | .5 | .5 | .6 | .6 | .5 | .5 | .5 | .5 | .4 |
| 4.0 | 4.0 | 4.0 | 3.0 | 4.0 | 6.0 | 6.0 | 4.0 | 4.0 | 4.0 | 3.0 | 3.0 |
| 40 | 33 | 46 | 45 | 62 | 44 | 62 | 49 | 62 | 50 | 51 | 45 |
| 31 | 25 | 45 | 42 | 58 | 35 | 49 | 40 | 47 | 41 | 28 | 28 |
| 9 | 8 | 9 | 9 | 11 | 8 | 13 | 7 | 12 | 9 | 18 | 7 |
| 19 | 19 | 27 | 19 | 27 | 20 | 28 | 23 | 33 | 25 | 28 | 24 |
| 30 | 28 | 37 | 33 | 47 | 37 | 51 | 38 | 49 | 38 | 47 | 38 |
| 47 | 38 | 57 | 57 | 80 | 56 | 76 | 61 | 76 | 61 | 71 | 57 |
| 76 | 61 | 84 | 85 | 128 | 85 | 123 | 93 | 121 | 93 | 89 | 76 |
| 96 | 82 | 113 | 112 | 165 | 112 | 151 | 122 | 151 | 119 | 112 | 97 |
| 154 | 151 | 297 | 217 | 224 | 180 | 245 | 224 | 253 | 215 | 114 | 141 |
| 107 | 90 | 129 | 121 | 157 | 110 | 157 | 127 | 160 | 126 | 144 | 121 |
| 81 | 62 | 117 | 112 | 156 | 84 | 116 | 105 | 106 | 86 | 77 | 74 |
| 100 | 94 | 109 | 105 | 153 | 102 | 148 | 119 | 138 | 130 | 103 | 105 |
| 59 | 63 | 109 | 87 | 96 | 60 | 95 | 72 | 108 | 96 | 49 | 42 |
| 73 | 49 | 80 | 91 | 134 | 122 | 152 | 102 | 110 | 149 | 87 | 101 |
| 29 | 26 | 32 | 48 | 33 | 42 | 66 | 29 | 32 | 50 | 15 | 38 |
| 840 | 567 | 1,344 | 932 | 2,350 | 758 | 996 | 830 | 768 | 624 | 432 | 450 |
| 840 | 582 | 1,429 | 1,158 | 2,350 | 758 | 1,087 | 853 | 1,152 | 900 | 432 | 450 |
| 101 | 86 | 117 | 111 | 149 | 104 | 147 | 119 | 145 | 123 | 124 | 112 |
| 78 | 62 | 115 | 102 | 125 | 74 | 116 | 98 | 109 | 87 | 68 | 69 |
| 28 | 19 | 20 | 24 | 28 | 20 | 28 | 20 | 35 | 24 | 36 | 20 |
| 56 | 45 | 57 | 56 | 64 | 57 | 80 | 57 | 75 | 61 | 80 | 61 |
| 84 | 79 | 85 | 85 | 113 | 85 | 113 | 85 | 113 | 103 | 106 | 112 |
| 113 | 112 | 142 | 130 | 196 | 135 | 180 | 152 | 180 | 168 | 170 | 151 |
| 170 | 168 | 200 | 225 | 284 | 184 | 258 | 227 | 270 | 227 | 227 | 196 |
| 255 | 206 | 252 | 270 | 362 | 227 | 360 | 280 | 392 | 304 | 227 | 225 |
| 425 | 288 | 454 | 568 | 643 | 394 | 577 | 480 | 480 | 448 | 336 | 360 |
| 840 | 567 | 1,344 | 864 | 1,410 | 758 | 996 | 830 | 768 | 624 | 336 | 450 |
| 58.8 | 35.3 | 21.7 | 34.7 | 59.8 | 78.2 | 143.4 | 173.7 | 14.4 | 31.6 | 6.2 | 11.7 |
| 10.8 | 6.2 | 6.2 | 8.0 | 4.8 | 4.5 | 8.5 | 7.1 | 3.7 | 6.0 | 4.5 | 5.1 |

TABLE 7.20.--FINFISH OTHER THAN CANNED, DRIED, AND RAW
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 14.5 | 1.7 | 10.7 | 13.2 | 13.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 13.3 | 1.7 | 9.2 | 12.2 | 13.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.2 | 0 | 1.5 | 1.0 | .7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 1.0 | 1.1 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .0 | .4 | .3 | .3 |
| MAXIMUM.....DO..... | 5.0 | 1.0 | 3.0 | 2.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 54 | 15 | 26 | 31 | 36 |
| STANDARD DEVIATION.....GM..... | 43 | 9 | 17 | 20 | 22 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 17 | | 9 | 9 | 10 |
| 25TH.....GM..... | 28 | | 14 | 19 | 19 |
| 50TH.....GM..... | 38 | | 19 | 28 | 29 |
| 75TH.....GM..... | 67 | | 37 | 38 | 45 |
| 90TH.....GM..... | 96 | | 47 | 56 | 57 |
| 95TH.....GM..... | 132 | | 56 | 70 | 76 |
| 99TH.....GM..... | 221 | | 93 | 142 | 131 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 148 | 44 | 70 | 89 | 105 |
| STANDARD DEVIATION.....GM..... | 113 | 26 | 40 | 55 | 61 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 151 | 0 | 63 | 74 | 94 |
| STANDARD DEVIATION.....GM..... | 91 | 0 | 38 | 45 | 52 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 106 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 48 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 2,350 | 91 | 228 | 432 | 448 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,350 | 91 | 324 | 432 | 560 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 145 | 44 | 68 | 85 | 101 |
| STANDARD DEVIATION.....GM..... | 106 | 26 | 39 | 55 | 59 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 43 | | 21 | 28 | 28 |
| 25TH.....GM..... | 84 | | 38 | 56 | 57 |
| 50TH.....GM..... | 113 | | 57 | 84 | 85 |
| 75TH.....GM..... | 170 | | 85 | 112 | 113 |
| 90TH.....GM..... | 255 | | 113 | 144 | 170 |
| 95TH.....GM..... | 340 | | 144 | 170 | 210 |
| 99TH.....GM..... | 565 | | 168 | 340 | 336 |
| MAXIMUM.....GM..... | 1,410 | 91 | 228 | 432 | 448 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 681.0 | .8 | 8.3 | 23.1 | 32.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 11.1 | 8.7 | 6.5 | 9.3 | 11.8 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 15.1 | 13.8 | 12.5 | 14.7 | 14.0 | 13.0 | 16.5 | 16.2 | 18.4 | 16.6 | 12.5 | 13.1 |
| 13.9 | 13.1 | 12.0 | 13.5 | 12.4 | 12.1 | 14.9 | 15.0 | 15.7 | 14.7 | 10.7 | 11.8 |
| 1.1 | .6 | .5 | 1.1 | 1.5 | 1.0 | 1.5 | 1.2 | 2.3 | 1.9 | 1.3 | 1.2 |
| .1 | 0 | 0 | .1 | .1 | 0 | .1 | (2) | .3 | 0 | .5 | .2 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.1 | 1.3 | 1.1 |
| .3 | .3 | .3 | .4 | .4 | .3 | .4 | .3 | .5 | .4 | .5 | .4 |
| 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 5.0 | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| 43 | 36 | 52 | 52 | 72 | 47 | 68 | 55 | 70 | 57 | 58 | 49 |
| 31 | 24 | 49 | 46 | 61 | 32 | 48 | 39 | 51 | 44 | 27 | 26 |
| 13 | 12 | 19 | 19 | 19 | 17 | 19 | 19 | 19 | 19 | 19 | 19 |
| 27 | 19 | 28 | 28 | 37 | 28 | 37 | 28 | 37 | 28 | 38 | 28 |
| 37 | 28 | 38 | 38 | 57 | 38 | 57 | 47 | 57 | 47 | 57 | 44 |
| 56 | 38 | 57 | 63 | 90 | 57 | 80 | 70 | 76 | 75 | 76 | 57 |
| 85 | 66 | 88 | 85 | 132 | 80 | 131 | 93 | 135 | 101 | 98 | 76 |
| 108 | 84 | 117 | 136 | 169 | 112 | 152 | 128 | 151 | 139 | 113 | 101 |
| 149 | 144 | 299 | 288 | 267 | 180 | 227 | 221 | 284 | 253 | 144 | 150 |
| 121 | 103 | 154 | 145 | 198 | 128 | 189 | 154 | 178 | 148 | 168 | 139 |
| 86 | 59 | 148 | 116 | 171 | 75 | 132 | 106 | 104 | 90 | 77 | 73 |
| 116 | 125 | 111 | 152 | 190 | 154 | 175 | 147 | 200 | 174 | 101 | 90 |
| 48 | 64 | 26 | 130 | 96 | 73 | 76 | 78 | 130 | 114 | 51 | 30 |
| 95 | 0 | 0 | 75 | 57 | 0 | 179 | 85 | 105 | 0 | 86 | 141 |
| 43 | 0 | 0 | 0 | 0 | 0 | 32 | 0 | 18 | 0 | 21 | 0 |
| 840 | 454 | 1,344 | 932 | 2,350 | 565 | 996 | 830 | 768 | 624 | 432 | 450 |
| 840 | 582 | 1,344 | 1,158 | 2,350 | 663 | 996 | 853 | 1,152 | 900 | 432 | 450 |
| 117 | 102 | 145 | 139 | 191 | 129 | 181 | 150 | 175 | 150 | 137 | 130 |
| 82 | 61 | 138 | 114 | 136 | 73 | 125 | 102 | 117 | 94 | 69 | 69 |
| 28 | 28 | 56 | 54 | 57 | 48 | 56 | 55 | 56 | 56 | 56 | 56 |
| 68 | 57 | 85 | 84 | 112 | 84 | 105 | 85 | 100 | 85 | 85 | 84 |
| 90 | 85 | 112 | 112 | 168 | 112 | 168 | 120 | 140 | 113 | 113 | 113 |
| 140 | 113 | 170 | 170 | 227 | 168 | 227 | 170 | 227 | 180 | 170 | 170 |
| 224 | 170 | 227 | 252 | 340 | 224 | 336 | 252 | 340 | 280 | 227 | 224 |
| 284 | 227 | 304 | 304 | 432 | 270 | 432 | 336 | 432 | 340 | 280 | 227 |
| 425 | 324 | 897 | 649 | 649 | 396 | 680 | 589 | 577 | 454 | 336 | 450 |
| 840 | 454 | 1,344 | 864 | 1,410 | 540 | 996 | 830 | 768 | 624 | 336 | 450 |
| 58.8 | 28.8 | 21.7 | 32.9 | 52.9 | 67.7 | 134.3 | 160.7 | 13.0 | 29.9 | 6.2 | 9.5 |
| 16.7 | 9.0 | 11.4 | 13.5 | 8.4 | 8.8 | 14.6 | 12.8 | 5.3 | 9.1 | 7.3 | 6.5 |

TABLE 7.21.--CANNED TUNA
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 7.2 | .6 | 6.2 | 6.2 | 7.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 6.2 | .4 | 5.3 | 5.4 | 6.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .9 | .2 | .6 | .6 | .9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | .3 | .1 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.2 | 1.3 | 1.2 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .4 | .5 | .6 | .4 | .5 |
| MAXIMUM.....DO..... | 4.0 | 2.0 | 4.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 28 | 17 | 13 | 18 | 22 |
| STANDARD DEVIATION.....GM..... | 22 | 8 | 9 | 12 | 18 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 7 | | 2 | 3 | 3 |
| 25TH.....GM..... | 13 | | 7 | 7 | 9 |
| 50TH.....GM..... | 27 | | 13 | 13 | 18 |
| 75TH.....GM..... | 34 | | 13 | 27 | 27 |
| 90TH.....GM..... | 56 | | 27 | 33 | 42 |
| 95TH.....GM..... | 66 | | 28 | 43 | 54 |
| 99TH.....GM..... | 107 | | 43 | 57 | 113 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 73 | 65 | 32 | 48 | 52 |
| STANDARD DEVIATION.....GM..... | 46 | 11 | 19 | 32 | 36 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 76 | 7 | 30 | 53 | 60 |
| STANDARD DEVIATION.....GM..... | 52 | 0 | 15 | 21 | 41 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 80 | 0 | 34 | 28 | 58 |
| STANDARD DEVIATION.....GM..... | 41 | 0 | 7 | 0 | 16 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 480 | 80 | 113 | 170 | 284 |
| MAXIMUM OVER 3 DAYS.....GM..... | 960 | 80 | 128 | 170 | 340 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 73 | 39 | 31 | 47 | 54 |
| STANDARD DEVIATION.....GM..... | 47 | 30 | 18 | 31 | 39 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 19 | | 7 | 10 | 10 |
| 25TH.....GM..... | 40 | | 19 | 20 | 28 |
| 50TH.....GM..... | 61 | | 28 | 40 | 40 |
| 75TH.....GM..... | 85 | | 40 | 61 | 61 |
| 90TH.....GM..... | 120 | | 57 | 80 | 85 |
| 95TH.....GM..... | 169 | | 57 | 99 | 113 |
| 99TH.....GM..... | 213 | | 85 | 170 | 200 |
| MAXIMUM.....GM..... | 480 | 80 | 113 | 170 | 284 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 16.5 | 0 | 0 | 0 | .8 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .5 | 0 | 0 | 0 | .5 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.7 | 6.9 | 6.3 | 7.4 | 8.0 | 9.4 | 6.4 | 8.7 | 4.5 | 5.5 | 5.8 | 3.6 |
| 5.0 | 6.4 | 5.7 | 6.0 | 7.1 | 7.8 | 5.5 | 7.4 | 4.2 | 5.0 | 5.8 | 2.9 |
| .6 | .5 | .6 | 1.2 | .9 | 1.3 | .8 | 1.2 | .4 | .5 | 0 | .7 |
| .1 | .1 | .1 | .1 | (2) | .3 | .1 | .1 | 0 | .1 | 0 | 0 |
| 1.1 | 1.1 | 1.1 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.1 | 1.1 | 1.0 | 1.2 |
| .4 | .4 | .4 | .5 | .4 | .5 | .4 | .4 | .3 | .3 | .0 | .4 |
| 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 2.0 | 3.0 | 1.0 | 2.0 |
| 26 | 23 | 27 | 29 | 35 | 31 | 31 | 28 | 28 | 26 | 36 | 35 |
| 19 | 14 | 16 | 22 | 35 | 23 | 20 | 19 | 15 | 15 | 21 | 28 |
| 9 | 7 | 5 | 7 | 9 | 7 | 10 | 7 | 12 | 7 | | |
| 13 | 13 | 13 | 13 | 18 | 14 | 19 | 13 | 19 | 18 | | |
| 23 | 19 | 27 | 20 | 27 | 27 | 27 | 27 | 27 | 20 | | |
| 31 | 28 | 33 | 38 | 40 | 38 | 38 | 33 | 32 | 33 | | |
| 53 | 47 | 53 | 59 | 66 | 57 | 57 | 53 | 53 | 47 | | |
| 53 | 53 | 60 | 80 | 95 | 71 | 67 | 61 | 57 | 57 | | |
| 154 | 66 | 67 | 113 | 184 | 113 | 92 | 99 | 61 | 66 | | |
| 72 | 64 | 77 | 71 | 90 | 75 | 83 | 71 | 79 | 74 | 107 | 86 |
| 40 | 38 | 44 | 54 | 60 | 46 | 45 | 40 | 42 | 40 | 62 | 58 |
| 65 | 59 | 56 | 77 | 106 | 84 | 73 | 77 | 61 | 53 | 0 | 98 |
| 56 | 27 | 28 | 29 | 112 | 29 | 37 | 38 | 22 | 25 | 0 | 58 |
| 43 | 63 | 60 | 63 | 184 | 92 | 100 | 102 | 0 | 93 | 0 | 0 |
| 27 | 3 | 0 | 25 | 0 | 40 | 28 | 16 | 0 | 0 | 0 | 0 |
| 320 | 170 | 200 | 340 | 480 | 312 | 262 | 320 | 184 | 198 | 198 | 223 |
| 462 | 217 | 200 | 340 | 960 | 553 | 454 | 640 | 184 | 278 | 198 | 397 |
| 69 | 62 | 71 | 70 | 91 | 76 | 80 | 71 | 75 | 71 | 107 | 89 |
| 45 | 37 | 42 | 47 | 70 | 41 | 43 | 39 | 39 | 38 | 62 | 59 |
| 20 | 19 | 15 | 20 | 20 | 19 | 20 | 20 | 35 | 10 | | |
| 40 | 40 | 40 | 40 | 40 | 40 | 57 | 40 | 40 | 40 | | |
| 57 | 57 | 61 | 57 | 80 | 80 | 80 | 61 | 66 | 61 | | |
| 80 | 80 | 80 | 80 | 113 | 96 | 92 | 85 | 92 | 85 | | |
| 120 | 113 | 120 | 113 | 170 | 120 | 160 | 113 | 160 | 113 | | |
| 160 | 160 | 170 | 160 | 198 | 169 | 169 | 142 | 169 | 160 | | |
| 198 | 170 | 200 | 340 | 480 | 184 | 198 | 184 | 184 | 198 | | |
| 320 | 170 | 200 | 340 | 480 | 213 | 262 | 320 | 184 | 198 | 198 | 223 |
| 0 | 1.7 | 0 | 0 | 3.9 | 4.4 | 0 | 4.4 | 0 | 0 | 0 | 1.3 |
| 0 | 1.0 | 0 | 0 | 1.1 | .7 | 0 | .6 | 0 | 0 | 0 | 3.0 |

TABLE 7.22.--SHRIMP
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.5 | 0 | 1.0 | 1.0 | 1.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.4 | 0 | 1.0 | 1.0 | 1.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .1 | 0 | 0 | 0 | 0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 1.1 | 0 | 1.0 | 1.0 | 1.0 |
| STANDARD DEVIATION.....DO..... | .3 | 0 | .0 | .0 | .2 |
| MAXIMUM.....DO..... | 3.0 | 0 | 1.0 | 1.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 24 | 0 | 7 | 10 | 14 |
| STANDARD DEVIATION.....GM..... | 30 | 0 | 10 | 11 | 12 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 3 | | | | |
| 25TH.....GM..... | 8 | | | | |
| 50TH.....GM..... | 14 | | | | |
| 75TH.....GM..... | 27 | | | | |
| 90TH.....GM..... | 47 | | | | |
| 95TH.....GM..... | 76 | | | | |
| 99TH.....GM..... | 151 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 66 | 0 | 22 | 31 | 42 |
| STANDARD DEVIATION.....GM..... | 76 | 0 | 31 | 34 | 37 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 89 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 102 | 0 | 0 | 0 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 30 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 0 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 907 | 0 | 135 | 135 | 135 |
| MAXIMUM OVER 3 DAYS.....GM..... | 971 | 0 | 135 | 135 | 135 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 67 | 0 | 22 | 31 | 41 |
| STANDARD DEVIATION.....GM..... | 82 | 0 | 31 | 34 | 37 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 8 | | | | |
| 25TH.....GM..... | 22 | | | | |
| 50TH.....GM..... | 40 | | | | |
| 75TH.....GM..... | 70 | | | | |
| 90TH.....GM..... | 135 | | | | |
| 95TH.....GM..... | 227 | | | | |
| 99TH.....GM..... | 454 | | | | |
| MAXIMUM.....GM..... | 907 | 0 | 135 | 135 | 135 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 11.9 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.2 | 0 | 0 | 0 | 0 |

^{1/} SEE TABLE NOTES IN APPENDIX.

^{2/} MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY, 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.0 | 2.1 | 2.9 | 2.2 | 2.8 | 3.3 | 3.1 | 3.4 | 2.7 | 1.4 | .7 | 1.5 |
| 1.0 | 2.1 | 2.6 | 2.2 | 2.7 | 3.0 | 2.9 | 3.2 | 2.5 | 1.4 | .7 | 1.5 |
| 0 | (2) | .2 | 0 | .1 | .3 | .2 | .1 | .2 | 0 | 0 | 0 |
| 0 | 0 | .1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1.0 | 1.0 | 1.1 | 1.0 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.0 | 1.0 | 1.1 |
| .0 | .1 | .4 | .0 | .2 | .3 | .3 | .2 | .5 | .0 | .0 | .3 |
| 1.0 | 2.0 | 3.0 | 1.0 | 2.0 | 2.0 | 2.0 | 2.0 | 3.0 | 1.0 | 1.0 | 2.0 |
| 21 | 15 | 32 | 23 | 27 | 22 | 29 | 22 | 36 | 23 | 24 | 14 |
| 16 | 15 | 32 | 24 | 33 | 25 | 40 | 29 | 40 | 16 | 23 | 8 |
| | | | | 2 | 3 | 3 | 3 | | | | |
| | | | | 11 | 8 | 8 | 5 | | | | |
| | | | | 16 | 16 | 19 | 11 | | | | |
| | | | | 32 | 27 | 34 | 23 | | | | |
| | | | | 57 | 45 | 50 | 46 | | | | |
| | | | | 81 | 76 | 76 | 97 | | | | |
| | | | | 218 | 151 | 197 | 151 | | | | |
| 63 | 45 | 96 | 68 | 71 | 63 | 75 | 63 | 105 | 69 | 72 | 42 |
| 49 | 46 | 101 | 71 | 69 | 72 | 79 | 82 | 126 | 47 | 70 | 25 |
| 0 | 30 | 49 | 0 | 178 | 56 | 143 | 72 | 64 | 0 | 0 | 0 |
| 0 | 0 | 14 | 0 | 115 | 44 | 161 | 73 | 0 | 0 | 0 | 0 |
| 0 | 0 | 30 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 227 | 227 | 454 | 340 | 453 | 454 | 907 | 454 | 454 | 135 | 170 | 97 |
| 227 | 227 | 454 | 340 | 654 | 454 | 971 | 486 | 454 | 135 | 170 | 97 |
| 63 | 44 | 87 | 68 | 78 | 62 | 82 | 62 | 93 | 69 | 72 | 38 |
| 49 | 45 | 94 | 71 | 79 | 69 | 111 | 79 | 116 | 47 | 70 | 17 |
| | | | | 6 | 6 | 8 | 8 | | | | |
| | | | | 32 | 24 | 24 | 16 | | | | |
| | | | | 48 | 43 | 57 | 32 | | | | |
| | | | | 101 | 68 | 85 | 68 | | | | |
| | | | | 170 | 135 | 135 | 135 | | | | |
| | | | | 227 | 169 | 227 | 227 | | | | |
| | | | | 453 | 454 | 561 | 340 | | | | |
| 227 | 227 | 454 | 340 | 453 | 454 | 907 | 454 | 454 | 135 | 170 | 68 |
| 0 | 2.7 | 0 | .9 | 0 | 1.2 | 5.2 | 1.0 | 0 | 0 | 0 | .9 |
| 0 | 5.9 | 0 | 2.7 | 0 | .6 | 3.2 | .4 | 0 | 0 | 0 | 5.6 |

TABLE 7.23.--FRANKFURTERS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 17.8 | 2.9 | 28.6 | 29.7 | 27.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 15.4 | 2.1 | 22.5 | 23.7 | 24.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.1 | .3 | 5.2 | 5.2 | 3.1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | .5 | .9 | .9 | .6 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.2 | 1.5 | 1.3 | 1.3 | 1.2 |
| STANDARD DEVIATION.....DO..... | .5 | .8 | .6 | .5 | .5 |
| MAXIMUM.....DO..... | 7.0 | 3.0 | 3.0 | 4.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 29 | 16 | 24 | 24 | 25 |
| STANDARD DEVIATION.....GM..... | 19 | 14 | 17 | 16 | 15 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 15 | | 7 | 9 | 14 |
| 25TH.....GM..... | 15 | | 15 | 15 | 15 |
| 50TH.....GM..... | 29 | | 15 | 15 | 19 |
| 75TH.....GM..... | 29 | | 29 | 29 | 29 |
| 90TH.....GM..... | 52 | | 44 | 44 | 44 |
| 95TH.....GM..... | 59 | | 59 | 48 | 57 |
| 99TH.....GM..... | 103 | | 95 | 89 | 88 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 76 | 28 | 55 | 58 | 63 |
| STANDARD DEVIATION.....GM..... | 41 | 14 | 32 | 31 | 29 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 77 | 35 | 60 | 57 | 69 |
| STANDARD DEVIATION.....GM..... | 37 | 8 | 29 | 25 | 31 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 78 | 40 | 57 | 60 | 63 |
| STANDARD DEVIATION.....GM..... | 45 | 15 | 26 | 37 | 30 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 660 | 57 | 280 | 280 | 308 |
| MAXIMUM OVER 3 DAYS.....GM..... | 836 | 157 | 352 | 567 | 484 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 75 | 33 | 55 | 56 | 62 |
| STANDARD DEVIATION.....GM..... | 40 | 15 | 29 | 29 | 30 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 43 | | 22 | 28 | 35 |
| 25TH.....GM..... | 44 | | 44 | 44 | 44 |
| 50TH.....GM..... | 57 | | 44 | 44 | 44 |
| 75TH.....GM..... | 88 | | 57 | 57 | 88 |
| 90TH.....GM..... | 132 | | 88 | 88 | 88 |
| 95TH.....GM..... | 142 | | 113 | 112 | 113 |
| 99TH.....GM..... | 220 | | 176 | 170 | 170 |
| MAXIMUM.....GM..... | 454 | 57 | 280 | 280 | 280 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 13.3 | 0 | 0 | 0 | 2.4 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .2 | 0 | 0 | 0 | .4 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 25.9 | 24.9 | 20.8 | 18.9 | 15.9 | 18.7 | 13.9 | 13.8 | 13.2 | 9.0 | 9.7 | 11.4 |
| 22.4 | 21.9 | 17.7 | 16.6 | 13.9 | 15.9 | 12.4 | 12.7 | 12.0 | 8.0 | 8.3 | 9.4 |
| 3.1 | 2.6 | 3.1 | 2.0 | 1.8 | 2.6 | 1.2 | 1.0 | .9 | 1.0 | 1.2 | 1.7 |
| .4 | .3 | .1 | .2 | .3 | .2 | .2 | .1 | .3 | .1 | .2 | .3 |
| 1.2 | 1.1 | 1.2 | 1.2 | 1.2 | 1.2 | 1.1 | 1.1 | 1.1 | 1.2 | 1.2 | 1.3 |
| .5 | .4 | .4 | .5 | .4 | .4 | .4 | .3 | .4 | .4 | .6 | .8 |
| 4.0 | 3.0 | 4.0 | 4.0 | 3.0 | 4.0 | 4.0 | 3.0 | 3.0 | 3.0 | 4.0 | 7.0 |
| 29 | 25 | 36 | 26 | 40 | 28 | 36 | 26 | 33 | 28 | 35 | 27 |
| 19 | 16 | 21 | 16 | 23 | 19 | 24 | 15 | 19 | 15 | 35 | 16 |
| 15 | 12 | 15 | 15 | 15 | 15 | 15 | 15 | 15 | 15 | 12 | 15 |
| 15 | 15 | 19 | 15 | 29 | 15 | 22 | 15 | 28 | 15 | 19 | 15 |
| 29 | 19 | 29 | 23 | 29 | 29 | 29 | 29 | 29 | 29 | 29 | 29 |
| 29 | 29 | 44 | 29 | 47 | 29 | 44 | 29 | 37 | 29 | 37 | 29 |
| 48 | 44 | 59 | 44 | 59 | 47 | 59 | 44 | 44 | 56 | 66 | 44 |
| 59 | 59 | 76 | 59 | 88 | 59 | 76 | 57 | 76 | 59 | 76 | 59 |
| 113 | 76 | 113 | 113 | 117 | 103 | 132 | 88 | 113 | 67 | 235 | 103 |
| 77 | 65 | 93 | 68 | 106 | 71 | 95 | 72 | 88 | 75 | 82 | 69 |
| 42 | 36 | 49 | 29 | 56 | 33 | 46 | 34 | 35 | 38 | 44 | 27 |
| 73 | 70 | 94 | 78 | 104 | 82 | 91 | 76 | 95 | 70 | 73 | 59 |
| 35 | 25 | 38 | 38 | 41 | 39 | 39 | 31 | 62 | 25 | 30 | 27 |
| 88 | 65 | 73 | 56 | 77 | 74 | 149 | 73 | 81 | 59 | 235 | 95 |
| 40 | 22 | 0 | 11 | 19 | 31 | 60 | 19 | 30 | 0 | 0 | 7 |
| 454 | 444 | 352 | 227 | 660 | 284 | 454 | 264 | 255 | 179 | 308 | 176 |
| 572 | 444 | 352 | 340 | 660 | 425 | 836 | 340 | 340 | 264 | 704 | 308 |
| 75 | 65 | 90 | 67 | 102 | 73 | 95 | 71 | 88 | 71 | 84 | 65 |
| 41 | 35 | 42 | 30 | 48 | 36 | 48 | 32 | 43 | 31 | 47 | 27 |
| 44 | 43 | 44 | 44 | 44 | 43 | 44 | 44 | 44 | 44 | 43 | 44 |
| 44 | 44 | 44 | 44 | 85 | 44 | 57 | 44 | 44 | 44 | 44 | 44 |
| 57 | 44 | 88 | 56 | 88 | 66 | 88 | 57 | 88 | 66 | 88 | 44 |
| 88 | 88 | 113 | 88 | 132 | 88 | 113 | 88 | 88 | 88 | 88 | 88 |
| 132 | 88 | 140 | 88 | 176 | 113 | 132 | 112 | 132 | 99 | 170 | 88 |
| 132 | 132 | 176 | 132 | 176 | 132 | 176 | 132 | 176 | 132 | 176 | 112 |
| 220 | 176 | 220 | 170 | 227 | 176 | 264 | 176 | 255 | 176 | 227 | 142 |
| 454 | 444 | 264 | 227 | 440 | 284 | 454 | 227 | 255 | 179 | 227 | 142 |
| 3.4 | .8 | 0 | 0 | 0 | 0 | .8 | 3.3 | 0 | 2.6 | 0 | 0 |
| .5 | .1 | 0 | 0 | 0 | 0 | .1 | .3 | 0 | 1.4 | 0 | 0 |

TABLE 7.24.--BOLOGNA
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 15.9 | 2.4 | 20.4 | 23.8 | 22.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 11.7 | 1.7 | 14.2 | 18.3 | 15.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.2 | .5 | 4.4 | 4.1 | 5.3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.0 | .2 | 1.8 | 1.4 | 1.5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.4 | 1.4 | 1.4 | 1.3 | 1.4 |
| STANDARD DEVIATION.....DO..... | .6 | .6 | .7 | .6 | .6 |
| MAXIMUM.....DO..... | 6.0 | 3.0 | 3.0 | 3.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 21 | 19 | 15 | 16 | 18 |
| STANDARD DEVIATION.....GM..... | 16 | 22 | 11 | 11 | 12 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 7 | | 5 | 5 | 9 |
| 25TH.....GM..... | 9 | | 9 | 9 | 9 |
| 50TH.....GM..... | 19 | | 9 | 9 | 19 |
| 75TH.....GM..... | 28 | | 19 | 19 | 19 |
| 90TH.....GM..... | 38 | | 28 | 28 | 37 |
| 95TH.....GM..... | 56 | | 37 | 38 | 38 |
| 99TH.....GM..... | 76 | | 75 | 65 | 57 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 49 | 27 | 32 | 37 | 42 |
| STANDARD DEVIATION.....GM..... | 32 | 18 | 16 | 20 | 23 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 48 | 82 | 37 | 37 | 39 |
| STANDARD DEVIATION.....GM..... | 26 | 40 | 23 | 20 | 19 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 47 | 13 | 27 | 36 | 34 |
| STANDARD DEVIATION.....GM..... | 24 | 0 | 11 | 16 | 12 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 312 | 112 | 140 | 142 | 170 |
| MAXIMUM OVER 3 DAYS.....GM..... | 448 | 224 | 227 | 280 | 227 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 47 | 41 | 32 | 36 | 40 |
| STANDARD DEVIATION.....GM..... | 29 | 37 | 18 | 20 | 21 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 14 | | 14 | 14 | 14 |
| 25TH.....GM..... | 28 | | 28 | 28 | 28 |
| 50TH.....GM..... | 28 | | 28 | 28 | 28 |
| 75TH.....GM..... | 57 | | 28 | 43 | 56 |
| 90TH.....GM..... | 85 | | 57 | 57 | 57 |
| 95TH.....GM..... | 112 | | 57 | 84 | 84 |
| 99TH.....GM..... | 168 | | 112 | 112 | 113 |
| MAXIMUM.....GM..... | 284 | 112 | 113 | 140 | 140 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 14.5 | 0 | .9 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .2 | 0 | .3 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 19.7 | 21.3 | 20.2 | 15.9 | 19.1 | 15.0 | 16.5 | 11.5 | 12.8 | 8.3 | 13.3 | 7.0 |
| 14.8 | 15.9 | 15.3 | 12.5 | 12.7 | 11.7 | 11.3 | 9.2 | 8.9 | 5.9 | 11.1 | 4.8 |
| 4.0 | 4.3 | 4.0 | 2.8 | 4.6 | 2.7 | 3.7 | 1.9 | 2.9 | 2.1 | 1.5 | 1.5 |
| .9 | 1.1 | .9 | .6 | 1.8 | .6 | 1.5 | .4 | .9 | .3 | .7 | .6 |
| 1.3 | 1.3 | 1.3 | 1.3 | 1.5 | 1.3 | 1.4 | 1.3 | 1.4 | 1.4 | 1.2 | 1.4 |
| .6 | .6 | .6 | .6 | .8 | .5 | .7 | .6 | .6 | .6 | .5 | .7 |
| 4.0 | 5.0 | 6.0 | 4.0 | 5.0 | 3.0 | 4.0 | 4.0 | 3.0 | 3.0 | 3.0 | 4.0 |
| 20 | 18 | 26 | 19 | 29 | 19 | 27 | 18 | 22 | 19 | 25 | 20 |
| 14 | 13 | 19 | 14 | 21 | 14 | 19 | 13 | 13 | 11 | 19 | 13 |
| 7 | 7 | 9 | 9 | 9 | 5 | 9 | 5 | 9 | 9 | 9 | 9 |
| 9 | 9 | 19 | 9 | 19 | 9 | 19 | 9 | 9 | 9 | 9 | 9 |
| 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 |
| 28 | 19 | 37 | 19 | 37 | 19 | 37 | 19 | 28 | 19 | 38 | 28 |
| 38 | 37 | 56 | 37 | 56 | 38 | 56 | 37 | 38 | 37 | 38 | 38 |
| 47 | 47 | 69 | 47 | 75 | 47 | 65 | 42 | 47 | 38 | 47 | 47 |
| 76 | 75 | 113 | 75 | 104 | 66 | 76 | 66 | 75 | 56 | 104 | 57 |
| 49 | 42 | 62 | 46 | 66 | 44 | 62 | 43 | 50 | 45 | 66 | 49 |
| 32 | 24 | 37 | 28 | 40 | 27 | 39 | 25 | 21 | 23 | 56 | 34 |
| 47 | 41 | 59 | 44 | 56 | 47 | 58 | 46 | 50 | 42 | 62 | 38 |
| 23 | 19 | 34 | 21 | 29 | 25 | 29 | 25 | 26 | 19 | 28 | 17 |
| 38 | 43 | 60 | 56 | 63 | 43 | 51 | 46 | 36 | 31 | 39 | 35 |
| 15 | 21 | 34 | 27 | 26 | 17 | 26 | 15 | 9 | 11 | 9 | 10 |
| 284 | 227 | 258 | 255 | 282 | 170 | 255 | 227 | 141 | 113 | 312 | 170 |
| 284 | 255 | 448 | 336 | 392 | 340 | 420 | 340 | 224 | 170 | 312 | 170 |
| 46 | 41 | 60 | 45 | 59 | 44 | 57 | 43 | 48 | 42 | 60 | 42 |
| 28 | 23 | 36 | 27 | 34 | 26 | 34 | 24 | 23 | 22 | 40 | 27 |
| 14 | 14 | 28 | 21 | 28 | 14 | 28 | 14 | 21 | 14 | 28 | 14 |
| 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 |
| 28 | 28 | 56 | 28 | 56 | 28 | 56 | 28 | 56 | 28 | 56 | 28 |
| 57 | 56 | 57 | 56 | 57 | 56 | 57 | 57 | 57 | 56 | 85 | 56 |
| 85 | 64 | 112 | 84 | 112 | 84 | 112 | 57 | 84 | 70 | 113 | 84 |
| 113 | 85 | 113 | 112 | 113 | 112 | 113 | 85 | 85 | 84 | 113 | 85 |
| 113 | 113 | 170 | 113 | 170 | 140 | 170 | 113 | 113 | 113 | 227 | 170 |
| 284 | 227 | 227 | 255 | 280 | 170 | 255 | 227 | 113 | 113 | 227 | 170 |
| 3.3 | 1.1 | 0 | 3.0 | .8 | 0 | 1.2 | 2.3 | 0 | 1.9 | 0 | 0 |
| .6 | .2 | 0 | 1.0 | .1 | 0 | .1 | .2 | 0 | 1.0 | 0 | 0 |

TABLE 7.25.--SALAMI
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.4 | .2 | 1.9 | 3.0 | 4.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.4 | .2 | 1.5 | 2.4 | 3.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .7 | 0 | .3 | .5 | .9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | 0 | .1 | .1 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.3 | 1.0 | 1.3 | 1.3 | 1.3 |
| STANDARD DEVIATION.....DO..... | .6 | .0 | .6 | .5 | .7 |
| MAXIMUM.....DO..... | 5.0 | 1.0 | 3.0 | 3.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 25 | 9 | 15 | 18 | 21 |
| STANDARD DEVIATION.....GM..... | 20 | 0 | 9 | 14 | 18 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 7 | | | 3 | 9 |
| 25TH.....GM..... | 9 | | | 9 | 9 |
| 50TH.....GM..... | 19 | | | 9 | 15 |
| 75TH.....GM..... | 37 | | | 28 | 28 |
| 90TH.....GM..... | 47 | | | 38 | 38 |
| 95TH.....GM..... | 65 | | | 38 | 47 |
| 99TH.....GM..... | 95 | | | 65 | 93 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 60 | 28 | 40 | 43 | 46 |
| STANDARD DEVIATION.....GM..... | 42 | 0 | 22 | 31 | 26 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 60 | 0 | 22 | 39 | 53 |
| STANDARD DEVIATION.....GM..... | 35 | 0 | 6 | 16 | 33 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 54 | 0 | 38 | 56 | 62 |
| STANDARD DEVIATION.....GM..... | 31 | 0 | 0 | 9 | 32 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 284 | 28 | 112 | 113 | 140 |
| MAXIMUM OVER 3 DAYS.....GM..... | 420 | 28 | 113 | 196 | 280 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 58 | 28 | 35 | 43 | 47 |
| STANDARD DEVIATION.....GM..... | 39 | 0 | 20 | 28 | 22 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 20 | | | 10 | 25 |
| 25TH.....GM..... | 28 | | | 28 | 28 |
| 50TH.....GM..... | 56 | | | 28 | 42 |
| 75TH.....GM..... | 57 | | | 57 | 57 |
| 90TH.....GM..... | 113 | | | 84 | 84 |
| 95TH.....GM..... | 113 | | | 113 | 85 |
| 99TH.....GM..... | 224 | | | 113 | 113 |
| MAXIMUM.....GM..... | 284 | 28 | 112 | 113 | 113 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 4.8 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .2 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.6 | 4.3 | 5.7 | 4.5 | 6.8 | 3.8 | 7.7 | 3.3 | 3.5 | 1.9 | 2.5 | 2.4 |
| 3.0 | 3.1 | 4.8 | 4.1 | 5.0 | 3.3 | 5.5 | 2.8 | 2.4 | 1.5 | 2.1 | 1.8 |
| .5 | 1.0 | .7 | .2 | 1.6 | .4 | 1.5 | .4 | .7 | .3 | .2 | .4 |
| .2 | .2 | .2 | .2 | .2 | (2) | .7 | .1 | .3 | .1 | .2 | .3 |
| 1.3 | 1.3 | 1.3 | 1.2 | 1.3 | 1.1 | 1.4 | 1.2 | 1.5 | 1.3 | 1.3 | 1.4 |
| .6 | .6 | .6 | .5 | .6 | .4 | .8 | .4 | .8 | .6 | .6 | .7 |
| 3.0 | 3.0 | 3.0 | 3.0 | 4.0 | 3.0 | 5.0 | 3.0 | 4.0 | 3.0 | 3.0 | 3.0 |
| 26 | 23 | 30 | 22 | 29 | 20 | 31 | 18 | 31 | 16 | 23 | 16 |
| 19 | 16 | 21 | 14 | 19 | 14 | 26 | 13 | 25 | 10 | 21 | 11 |
| 9 | 5 | 7 | 7 | 9 | 5 | 9 | 5 | | | | |
| 13 | 9 | 19 | 9 | 19 | 9 | 14 | 9 | | | | |
| 19 | 19 | 19 | 19 | 19 | 19 | 25 | 14 | | | | |
| 28 | 28 | 38 | 28 | 38 | 22 | 38 | 19 | | | | |
| 56 | 38 | 56 | 38 | 56 | 38 | 75 | 37 | | | | |
| 76 | 47 | 66 | 47 | 75 | 56 | 76 | 43 | | | | |
| 93 | 75 | 112 | 75 | 84 | 76 | 132 | 57 | | | | |
| 69 | 55 | 77 | 58 | 74 | 54 | 68 | 45 | 77 | 34 | 51 | 37 |
| 53 | 35 | 47 | 32 | 46 | 36 | 49 | 28 | 59 | 15 | 38 | 31 |
| 54 | 56 | 58 | 55 | 58 | 51 | 77 | 53 | 76 | 52 | 28 | 38 |
| 36 | 31 | 36 | 17 | 26 | 29 | 44 | 23 | 52 | 10 | 0 | 7 |
| 48 | 28 | 88 | 51 | 65 | 47 | 59 | 41 | 32 | 28 | 76 | 28 |
| 11 | 0 | 19 | 20 | 29 | 0 | 36 | 16 | 7 | 0 | 0 | 0 |
| 280 | 227 | 170 | 142 | 252 | 227 | 284 | 170 | 243 | 84 | 113 | 112 |
| 280 | 340 | 336 | 225 | 336 | 227 | 420 | 227 | 340 | 113 | 227 | 112 |
| 59 | 52 | 71 | 57 | 67 | 53 | 65 | 46 | 65 | 38 | 54 | 35 |
| 44 | 33 | 44 | 32 | 41 | 35 | 43 | 27 | 54 | 15 | 34 | 24 |
| 28 | 25 | 14 | 25 | 28 | 14 | 20 | 14 | | | | |
| 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | | | | |
| 56 | 43 | 57 | 56 | 56 | 56 | 57 | 40 | | | | |
| 57 | 57 | 112 | 57 | 85 | 57 | 85 | 57 | | | | |
| 112 | 85 | 142 | 112 | 113 | 85 | 113 | 84 | | | | |
| 113 | 140 | 168 | 140 | 140 | 112 | 140 | 112 | | | | |
| 280 | 142 | 170 | 142 | 227 | 227 | 224 | 168 | | | | |
| 280 | 170 | 170 | 142 | 252 | 227 | 284 | 170 | 243 | 84 | 113 | 112 |
| 0 | 0 | 0 | 0 | 1.7 | 0 | 1.2 | 0 | 0 | 1.8 | 0 | 0 |
| 0 | 0 | 0 | 0 | .5 | 0 | .2 | 0 | 0 | 4.2 | 0 | 0 |

TABLE 7.26.--PORK BREAKFAST SAUSAGE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 13.5 | 1.4 | 11.9 | 13.4 | 12.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 10.6 | 1.1 | 8.8 | 10.4 | 10.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.3 | .3 | 2.8 | 2.8 | 2.1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .6 | 0 | .3 | .2 | .3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.3 | 1.2 | 1.3 | 1.3 | 1.2 |
| STANDARD DEVIATION.....DO..... | .6 | .4 | .5 | .5 | .5 |
| MAXIMUM.....DO..... | 5.0 | 2.0 | 3.0 | 4.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 24 | 19 | 15 | 16 | 18 |
| STANDARD DEVIATION.....GM..... | 21 | 16 | 14 | 14 | 15 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 4 | | 4 | 4 | 4 |
| 25TH.....GM..... | 9 | | 7 | 9 | 9 |
| 50TH.....GM..... | 18 | | 9 | 9 | 13 |
| 75TH.....GM..... | 28 | | 18 | 19 | 22 |
| 90TH.....GM..... | 47 | | 28 | 37 | 38 |
| 95TH.....GM..... | 64 | | 38 | 47 | 57 |
| 99TH.....GM..... | 113 | | 72 | 76 | 75 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 59 | 50 | 33 | 42 | 47 |
| STANDARD DEVIATION.....GM..... | 49 | 52 | 20 | 35 | 37 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 55 | 40 | 37 | 32 | 40 |
| STANDARD DEVIATION.....GM..... | 42 | 15 | 32 | 19 | 24 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 46 | 0 | 33 | 66 | 50 |
| STANDARD DEVIATION.....GM..... | 27 | 0 | 13 | 18 | 20 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 452 | 160 | 162 | 227 | 227 |
| MAXIMUM OVER 3 DAYS.....GM..... | 510 | 160 | 324 | 255 | 255 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 55 | 46 | 34 | 39 | 44 |
| STANDARD DEVIATION.....GM..... | 45 | 42 | 26 | 31 | 34 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 13 | | 10 | 13 | 13 |
| 25TH.....GM..... | 26 | | 14 | 22 | 26 |
| 50TH.....GM..... | 52 | | 27 | 27 | 28 |
| 75TH.....GM..... | 57 | | 43 | 54 | 57 |
| 90TH.....GM..... | 113 | | 57 | 85 | 85 |
| 95TH.....GM..... | 142 | | 85 | 85 | 113 |
| 99TH.....GM..... | 227 | | 113 | 170 | 170 |
| MAXIMUM.....GM..... | 452 | 160 | 162 | 227 | 227 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 37.0 | 0 | 0 | 0 | 3.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | 0 | 0 | 0 | 1.1 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 16.3 | 13.2 | 14.3 | 13.4 | 16.1 | 11.0 | 17.4 | 12.0 | 16.8 | 10.1 | 15.4 | 11.4 |
| 13.7 | 11.0 | 11.6 | 11.4 | 12.1 | 8.9 | 13.3 | 9.4 | 12.5 | 8.8 | 12.5 | 7.6 |
| 2.0 | 1.7 | 1.9 | 1.4 | 3.3 | 1.7 | 3.1 | 2.1 | 3.0 | 1.0 | 2.3 | 2.2 |
| .5 | .5 | .8 | .5 | .8 | .3 | 1.0 | .5 | 1.2 | .3 | .6 | 1.5 |
| 1.2 | 1.2 | 1.3 | 1.2 | 1.3 | 1.2 | 1.3 | 1.3 | 1.4 | 1.2 | 1.3 | 1.5 |
| .5 | .5 | .6 | .5 | .6 | .5 | .6 | .6 | .6 | .4 | .6 | .8 |
| 3.0 | 3.0 | 4.0 | 3.0 | 4.0 | 5.0 | 5.0 | 5.0 | 4.0 | 3.0 | 4.0 | 4.0 |
| 20 | 19 | 27 | 22 | 30 | 22 | 30 | 22 | 26 | 17 | 27 | 22 |
| 17 | 16 | 25 | 16 | 26 | 18 | 25 | 19 | 22 | 13 | 20 | 22 |
| 4 | 4 | 4 | 4 | 4 | 4 | 9 | 4 | 9 | 4 | 4 | 4 |
| 9 | 9 | 9 | 9 | 13 | 9 | 13 | 9 | 9 | 9 | 9 | 9 |
| 17 | 17 | 19 | 18 | 19 | 18 | 19 | 18 | 19 | 13 | 20 | 18 |
| 27 | 22 | 38 | 30 | 38 | 28 | 38 | 28 | 36 | 19 | 38 | 28 |
| 38 | 38 | 57 | 45 | 57 | 42 | 66 | 43 | 57 | 38 | 57 | 38 |
| 47 | 52 | 80 | 57 | 76 | 54 | 76 | 57 | 76 | 53 | 66 | 57 |
| 76 | 85 | 113 | 76 | 142 | 85 | 124 | 95 | 113 | 57 | 91 | 170 |
| 51 | 48 | 67 | 59 | 75 | 58 | 71 | 57 | 67 | 44 | 68 | 48 |
| 43 | 33 | 62 | 47 | 64 | 45 | 55 | 43 | 59 | 33 | 48 | 36 |
| 49 | 51 | 60 | 52 | 69 | 46 | 75 | 49 | 57 | 48 | 70 | 37 |
| 37 | 39 | 49 | 20 | 49 | 36 | 48 | 38 | 29 | 23 | 44 | 15 |
| 45 | 39 | 56 | 40 | 46 | 46 | 52 | 41 | 41 | 42 | 29 | 49 |
| 9 | 19 | 34 | 16 | 17 | 15 | 33 | 26 | 29 | 15 | 1 | 41 |
| 340 | 255 | 340 | 227 | 452 | 364 | 373 | 284 | 340 | 170 | 264 | 227 |
| 454 | 369 | 454 | 227 | 454 | 454 | 509 | 454 | 340 | 197 | 274 | 510 |
| 50 | 46 | 63 | 55 | 68 | 53 | 67 | 52 | 59 | 44 | 61 | 44 |
| 41 | 33 | 57 | 41 | 56 | 41 | 50 | 40 | 44 | 32 | 43 | 30 |
| 13 | 13 | 13 | 13 | 13 | 13 | 14 | 13 | 13 | 13 | 13 | 13 |
| 26 | 26 | 26 | 26 | 27 | 26 | 27 | 26 | 27 | 26 | 27 | 27 |
| 39 | 39 | 54 | 52 | 54 | 45 | 54 | 43 | 54 | 28 | 54 | 27 |
| 57 | 56 | 81 | 57 | 85 | 57 | 85 | 57 | 81 | 56 | 85 | 57 |
| 85 | 85 | 113 | 113 | 135 | 113 | 113 | 113 | 113 | 85 | 113 | 85 |
| 113 | 113 | 170 | 113 | 170 | 126 | 170 | 113 | 142 | 113 | 150 | 113 |
| 227 | 170 | 312 | 227 | 227 | 227 | 227 | 227 | 227 | 170 | 170 | 170 |
| 340 | 227 | 340 | 227 | 452 | 364 | 373 | 284 | 227 | 170 | 170 | 170 |
| 2.6 | .9 | 0 | 0 | 3.7 | 1.4 | 13.1 | 4.9 | 0 | 0 | .9 | 6.4 |
| .6 | .2 | 0 | 0 | .4 | .2 | 1.1 | .5 | 0 | 0 | .8 | 3.8 |

TABLE 7.27.--SAUSAGE OTHER THAN BOLOGNA, FRANKFURTERS, SALAMI, AND PORK BREAKFAST SAUSAGE CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.2 | .8 | 4.8 | 4.0 | 3.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.4 | .8 | 4.2 | 3.6 | 3.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .6 | 0 | .5 | .3 | .3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | 0 | .1 | .1 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 1.0 | 1.2 | 1.1 | 1.2 |
| STANDARD DEVIATION.....DO..... | .6 | .0 | .4 | .4 | .5 |
| MAXIMUM.....DO..... | 5.0 | 1.0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 32 | 12 | 17 | 20 | 22 |
| STANDARD DEVIATION.....GM..... | 30 | 5 | 11 | 13 | 18 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 5 | | 2 | 5 | 4 |
| 25TH.....GM..... | 14 | | 9 | 9 | 9 |
| 50TH.....GM..... | 25 | | 16 | 18 | 19 |
| 75TH.....GM..... | 38 | | 24 | 25 | 28 |
| 90TH.....GM..... | 68 | | 33 | 38 | 38 |
| 95TH.....GM..... | 76 | | 38 | 47 | 66 |
| 99TH.....GM..... | 151 | | 50 | 50 | 93 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 87 | 35 | 49 | 59 | 63 |
| STANDARD DEVIATION.....GM..... | 76 | 15 | 33 | 40 | 52 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 71 | 0 | 38 | 30 | 46 |
| STANDARD DEVIATION.....GM..... | 47 | 0 | 15 | 10 | 31 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 73 | 0 | 16 | 18 | 15 |
| STANDARD DEVIATION.....GM..... | 65 | 0 | 0 | 3 | (2) |
| MAXIMUM IN ANY 1 DAY.....GM..... | 907 | 57 | 151 | 150 | 280 |
| MAXIMUM OVER 3 DAYS.....GM..... | 907 | 57 | 151 | 150 | 280 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 79 | 35 | 44 | 52 | 55 |
| STANDARD DEVIATION.....GM..... | 70 | 15 | 31 | 38 | 49 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 14 | | 8 | 14 | 11 |
| 25TH.....GM..... | 28 | | 19 | 25 | 22 |
| 50TH.....GM..... | 57 | | 34 | 35 | 34 |
| 75TH.....GM..... | 113 | | 64 | 71 | 71 |
| 90TH.....GM..... | 168 | | 85 | 114 | 113 |
| 95TH.....GM..... | 227 | | 113 | 140 | 140 |
| 99TH.....GM..... | 340 | | 151 | 150 | 280 |
| MAXIMUM.....GM..... | 907 | 57 | 151 | 150 | 280 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 60.0 | 0 | 1.0 | 6.6 | 5.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.5 | 0 | 1.7 | 8.5 | 6.0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.5 | 3.9 | 4.6 | 4.0 | 5.9 | 4.6 | 7.7 | 5.6 | 6.0 | 3.8 | 3.5 | 3.4 |
| 4.6 | 3.5 | 3.7 | 3.6 | 5.0 | 4.3 | 6.3 | 4.8 | 5.2 | 3.0 | 2.5 | 2.9 |
| .8 | .3 | .9 | .3 | .7 | .2 | 1.0 | .6 | .6 | .7 | .9 | .5 |
| .1 | .1 | .1 | 0 | .2 | .1 | .4 | .2 | .3 | .1 | .2 | 0 |
| 1.3 | 1.1 | 1.3 | 1.2 | 1.2 | 1.1 | 1.3 | 1.2 | 1.2 | 1.2 | 1.4 | 1.2 |
| .6 | .4 | .6 | .4 | .6 | .4 | .7 | .6 | .5 | .5 | .6 | .5 |
| 4.0 | 3.0 | 3.0 | 2.0 | 4.0 | 3.0 | 5.0 | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| 28 | 24 | 30 | 27 | 49 | 31 | 38 | 30 | 26 | 26 | 50 | 23 |
| 21 | 18 | 23 | 25 | 42 | 38 | 31 | 22 | 19 | 16 | 52 | 15 |
| 4 | 5 | 5 | 5 | 5 | 5 | 7 | 5 | 5 | 5 | | |
| 11 | 9 | 17 | 9 | 19 | 14 | 19 | 15 | 11 | 16 | | |
| 19 | 19 | 25 | 19 | 38 | 23 | 28 | 25 | 19 | 19 | | |
| 38 | 31 | 38 | 32 | 67 | 38 | 56 | 38 | 38 | 38 | | |
| 56 | 47 | 73 | 57 | 95 | 60 | 76 | 57 | 57 | 47 | | |
| 75 | 75 | 85 | 101 | 140 | 76 | 101 | 76 | 66 | 57 | | |
| 113 | 76 | 95 | 127 | 175 | 274 | 151 | 113 | 76 | 76 | | |
| 75 | 67 | 79 | 74 | 134 | 78 | 100 | 84 | 66 | 72 | 137 | 68 |
| 60 | 50 | 59 | 65 | 125 | 59 | 81 | 58 | 46 | 49 | 165 | 49 |
| 70 | 69 | 68 | 82 | 104 | 62 | 89 | 54 | 90 | 49 | 64 | 37 |
| 31 | 38 | 45 | 65 | 50 | 59 | 54 | 30 | 20 | 15 | 23 | 7 |
| 15 | 28 | 57 | 0 | 91 | 204 | 71 | 59 | 34 | 57 | 150 | 0 |
| 0 | 0 | 0 | 0 | 44 | 107 | 34 | 41 | 11 | 0 | 0 | 0 |
| 340 | 227 | 284 | 340 | 907 | 454 | 454 | 340 | 170 | 227 | 454 | 168 |
| 340 | 227 | 284 | 380 | 907 | 822 | 454 | 425 | 227 | 227 | 454 | 168 |
| 67 | 64 | 71 | 69 | 119 | 84 | 87 | 74 | 66 | 62 | 108 | 58 |
| 53 | 46 | 52 | 57 | 107 | 76 | 68 | 54 | 43 | 42 | 127 | 45 |
| 11 | 11 | 17 | 11 | 15 | 14 | 16 | 11 | 17 | 16 | | |
| 28 | 28 | 32 | 28 | 57 | 28 | 42 | 28 | 32 | 28 | | |
| 57 | 56 | 57 | 57 | 85 | 57 | 64 | 57 | 56 | 57 | | |
| 87 | 85 | 85 | 76 | 152 | 113 | 113 | 113 | 85 | 76 | | |
| 142 | 113 | 142 | 114 | 227 | 180 | 170 | 152 | 114 | 113 | | |
| 168 | 142 | 198 | 227 | 284 | 227 | 227 | 170 | 168 | 170 | | |
| 227 | 227 | 284 | 304 | 454 | 454 | 340 | 227 | 170 | 227 | | |
| 340 | 227 | 284 | 304 | 907 | 454 | 454 | 340 | 170 | 227 | 454 | 168 |
| 11.0 | 0 | 0 | 1.0 | 10.6 | 8.3 | 7.2 | 9.1 | 0 | 0 | 0 | 0 |
| 7.6 | 0 | 0 | 1.5 | 3.7 | 3.1 | 1.4 | 1.9 | 0 | 0 | 0 | 0 |

TABLE 7.28.--BEEF STEW WITH VEGETABLES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.2 | .7 | 5.1 | 4.6 | 5.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.7 | .3 | 5.1 | 4.2 | 5.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | .2 | 0 | .4 | .4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | .2 | 0 | 0 | (2) |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 1.9 | 1.0 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .9 | .1 | .3 | .3 |
| MAXIMUM.....DO..... | 4.0 | 3.0 | 2.0 | 2.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 133 | 109 | 60 | 75 | 90 |
| STANDARD DEVIATION.....GM..... | 92 | 80 | 45 | 43 | 49 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 41 | | 20 | 20 | 41 |
| 25TH.....GM..... | 82 | | 41 | 41 | 50 |
| 50TH.....GM..... | 123 | | 41 | 82 | 82 |
| 75TH.....GM..... | 163 | | 82 | 82 | 102 |
| 90TH.....GM..... | 245 | | 82 | 163 | 163 |
| 95TH.....GM..... | 327 | | 163 | 163 | 204 |
| 99TH.....GM..... | 490 | 225 | 327 | 190 | 245 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 357 | 122 | 181 | 210 | 247 |
| STANDARD DEVIATION.....GM..... | 199 | 0 | 136 | 115 | 125 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 382 | 122 | 0 | 191 | 247 |
| STANDARD DEVIATION.....GM..... | 200 | 0 | 0 | 85 | 66 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 451 | 225 | 0 | 0 | 223 |
| STANDARD DEVIATION.....GM..... | 221 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,960 | 245 | 980 | 613 | 735 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,940 | 674 | 980 | 613 | 735 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 355 | 173 | 178 | 207 | 247 |
| STANDARD DEVIATION.....GM..... | 189 | 55 | 112 | 111 | 119 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 123 | | 61 | 61 | 123 |
| 25TH.....GM..... | 245 | | 123 | 123 | 151 |
| 50TH.....GM..... | 306 | | 123 | 245 | 245 |
| 75TH.....GM..... | 490 | | 245 | 245 | 245 |
| 90TH.....GM..... | 490 | | 245 | 326 | 368 |
| 95TH.....GM..... | 735 | | 490 | 490 | 490 |
| 99TH.....GM..... | 981 | 490 | 490 | 490 | 613 |
| MAXIMUM.....GM..... | 1,470 | 245 | 490 | 613 | 735 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 28.0 | 0 | 0 | 0 | 1.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.3 | 0 | 0 | 0 | 1.4 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.0 | 6.7 | 5.5 | 5.4 | 4.9 | 4.4 | 5.7 | 5.0 | 6.9 | 6.0 | 6.5 | 4.0 |
| 4.7 | 6.4 | 5.0 | 5.1 | 4.5 | 4.2 | 5.0 | 4.7 | 5.8 | 4.9 | 5.4 | 2.9 |
| .3 | .3 | .5 | .2 | .4 | .3 | .6 | .3 | .8 | .9 | .6 | 1.1 |
| 0 | 0 | 0 | .1 | (2) | 0 | .1 | (2) | .3 | .2 | .5 | .1 |
| 1.1 | 1.0 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.1 | 1.2 | 1.3 | 1.4 | 1.3 |
| .2 | .2 | .3 | .3 | .4 | .3 | .4 | .4 | .5 | .6 | .7 | .6 |
| 2.0 | 2.0 | 2.0 | 3.0 | 3.0 | 3.0 | 3.0 | 4.0 | 3.0 | 4.0 | 3.0 | 3.0 |
| 110 | 104 | 162 | 114 | 179 | 121 | 168 | 128 | 163 | 140 | 194 | 141 |
| 69 | 50 | 89 | 65 | 126 | 64 | 107 | 75 | 94 | 111 | 143 | 82 |
| 31 | 41 | 41 | 41 | 51 | 41 | 50 | 41 | 82 | 41 | | |
| 82 | 82 | 82 | 82 | 82 | 82 | 82 | 82 | 102 | 82 | | |
| 82 | 82 | 163 | 82 | 163 | 123 | 163 | 123 | 163 | 113 | | |
| 123 | 123 | 204 | 163 | 245 | 163 | 163 | 163 | 163 | 163 | | |
| 163 | 163 | 327 | 163 | 290 | 163 | 327 | 204 | 327 | 245 | | |
| 245 | 163 | 327 | 245 | 368 | 245 | 327 | 245 | 408 | 327 | | |
| 327 | 245 | 490 | 408 | 898 | 327 | 490 | 327 | 490 | 612 | | |
| 312 | 297 | 454 | 325 | 471 | 342 | 444 | 355 | 397 | 324 | 443 | 320 |
| 176 | 135 | 252 | 177 | 243 | 162 | 233 | 157 | 127 | 171 | 311 | 103 |
| 338 | 299 | 398 | 339 | 580 | 353 | 450 | 321 | 434 | 361 | 532 | 314 |
| 172 | 93 | 100 | 119 | 366 | 153 | 134 | 94 | 158 | 204 | 91 | 103 |
| 0 | 0 | 0 | 245 | 490 | 0 | 466 | 523 | 450 | 490 | 490 | 490 |
| 0 | 0 | 0 | 0 | 0 | 0 | 312 | 346 | 41 | 141 | 0 | 0 |
| 980 | 735 | 1,470 | 1,225 | 1,960 | 980 | 1,470 | 1,470 | 735 | 980 | 1,470 | 490 |
| 1,225 | 980 | 1,470 | 1,225 | 2,695 | 1,470 | 2,940 | 2,695 | 1,470 | 1,838 | 1,470 | 1,470 |
| 314 | 297 | 440 | 315 | 471 | 336 | 439 | 344 | 405 | 327 | 426 | 316 |
| 179 | 132 | 234 | 163 | 219 | 158 | 216 | 151 | 137 | 168 | 248 | 114 |
| 113 | 123 | 123 | 123 | 123 | 123 | 151 | 123 | 245 | 93 | | |
| 245 | 245 | 245 | 245 | 245 | 245 | 245 | 245 | 245 | 245 | | |
| 245 | 245 | 490 | 245 | 490 | 245 | 490 | 368 | 490 | 245 | | |
| 368 | 368 | 490 | 368 | 490 | 490 | 490 | 490 | 490 | 490 | | |
| 490 | 490 | 736 | 490 | 736 | 490 | 735 | 490 | 490 | 551 | | |
| 735 | 490 | 980 | 490 | 980 | 490 | 736 | 490 | 490 | 735 | | |
| 736 | 735 | 1,470 | 1,224 | 980 | 736 | 1,225 | 736 | 735 | 735 | | |
| 980 | 735 | 1,470 | 1,225 | 980 | 980 | 1,470 | 980 | 735 | 735 | 1,470 | 490 |
| .8 | .9 | 1.0 | 0 | 4.5 | 1.4 | 7.7 | 6.6 | 0 | 3.6 | 0 | 0 |
| .7 | .6 | 1.1 | 0 | 2.1 | .6 | 2.4 | 1.7 | 0 | 2.6 | 0 | 0 |

TABLE 7.29.--CHILI CON CARNE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.5 | .7 | 4.2 | 4.6 | 6.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.0 | .7 | 3.6 | 4.3 | 6.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | 0 | .5 | .3 | .3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | .1 | (2) | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 1.0 | 1.2 | 1.1 | 1.1 |
| STANDARD DEVIATION.....CO..... | .4 | .0 | .4 | .4 | .2 |
| MAXIMUM.....DO..... | 5.0 | 1.0 | 3.0 | 4.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 116 | 30 | 63 | 71 | 81 |
| STANDARD DEVIATION.....GM..... | 90 | 10 | 39 | 48 | 54 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 11 | | | 5 | 10 |
| 25TH.....GM..... | 64 | | | 43 | 43 |
| 50TH.....GM..... | 85 | | | 64 | 85 |
| 75TH.....GM..... | 170 | | | 85 | 85 |
| 90TH.....GM..... | 213 | | | 170 | 170 |
| 95TH.....GM..... | 255 | | | 170 | 170 |
| 99TH.....GM..... | 510 | | | 170 | 340 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 308 | 91 | 156 | 197 | 228 |
| STANDARD DEVIATION.....GM..... | 200 | 31 | 82 | 132 | 135 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 362 | 0 | 189 | 223 | 303 |
| STANDARD DEVIATION.....GM..... | 222 | 0 | 38 | 32 | 134 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 426 | 0 | 175 | 4 | 0 |
| STANDARD DEVIATION.....GM..... | 228 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,530 | 128 | 383 | 510 | 510 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,550 | 128 | 526 | 510 | 1,020 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 308 | 91 | 164 | 187 | 232 |
| STANDARD DEVIATION.....GM..... | 198 | 31 | 82 | 130 | 139 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 32 | | | 5 | 24 |
| 25TH.....GM..... | 171 | | | 84 | 128 |
| 50TH.....GM..... | 255 | | | 191 | 255 |
| 75TH.....GM..... | 383 | | | 255 | 255 |
| 90TH.....GM..... | 510 | | | 383 | 510 |
| 95TH.....GM..... | 765 | | | 510 | 510 |
| 99TH.....GM..... | 1,020 | | | 510 | 510 |
| MAXIMUM.....GM..... | 1,530 | 128 | 383 | 510 | 510 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 42.0 | 0 | 1.0 | 0 | 1.0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.8 | 0 | 2.0 | 0 | .8 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 7.0 | 6.4 | 7.4 | 5.2 | 6.6 | 5.5 | 6.3 | 5.3 | 5.8 | 1.8 | 4.1 | 1.3 |
| 6.7 | 5.9 | 7.0 | 4.9 | 5.8 | 5.0 | 5.6 | 4.9 | 5.1 | 1.8 | 4.1 | 1.1 |
| .3 | .5 | .4 | .2 | .8 | .4 | .6 | .4 | .7 | 0 | 0 | .2 |
| 0 | 0 | 0 | .1 | .1 | .1 | .1 | (2) | 0 | 0 | 0 | 0 |
| 1.0 | 1.1 | 1.1 | 1.1 | 1.2 | 1.1 | 1.2 | 1.1 | 1.1 | 1.0 | 1.0 | 1.4 |
| .2 | .3 | .3 | .5 | .5 | .4 | .5 | .4 | .3 | .2 | .0 | .5 |
| 2.0 | 3.0 | 3.0 | 5.0 | 4.0 | 3.0 | 4.0 | 4.0 | 2.0 | 2.0 | 1.0 | 2.0 |
| 106 | 99 | 129 | 92 | 156 | 112 | 140 | 110 | 123 | 94 | 98 | 126 |
| 72 | 60 | 98 | 64 | 133 | 94 | 103 | 65 | 78 | 61 | 48 | 67 |
| 11 | 16 | 11 | 5 | 11 | 11 | 21 | 11 | 11 | | | |
| 59 | 64 | 64 | 43 | 85 | 53 | 85 | 85 | 85 | | | |
| 85 | 85 | 106 | 85 | 128 | 85 | 128 | 85 | 85 | | | |
| 128 | 128 | 170 | 128 | 170 | 170 | 170 | 133 | 170 | | | |
| 213 | 170 | 255 | 170 | 301 | 170 | 255 | 170 | 213 | | | |
| 255 | 170 | 340 | 170 | 425 | 255 | 340 | 234 | 340 | | | |
| 340 | 298 | 510 | 340 | 596 | 510 | 552 | 340 | 340 | | | |
| 303 | 279 | 362 | 255 | 388 | 284 | 366 | 305 | 324 | 283 | 295 | 310 |
| 197 | 155 | 256 | 169 | 254 | 177 | 217 | 165 | 193 | 184 | 143 | 130 |
| 338 | 246 | 427 | 310 | 460 | 413 | 390 | 299 | 366 | 0 | 0 | 382 |
| 155 | 148 | 271 | 124 | 282 | 261 | 227 | 138 | 100 | 0 | 0 | 0 |
| 0 | 0 | 0 | 170 | 850 | 433 | 521 | 267 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 10 | 106 | 37 | 0 | 0 | 0 | 0 |
| 1,020 | 765 | 1,530 | 765 | 1,275 | 1,275 | 1,275 | 1,020 | 1,020 | 893 | 510 | 510 |
| 1,275 | 1,020 | 1,785 | 1,020 | 2,550 | 2,040 | 2,040 | 1,020 | 1,020 | 893 | 510 | 765 |
| 305 | 270 | 365 | 244 | 395 | 294 | 363 | 294 | 332 | 271 | 295 | 272 |
| 196 | 157 | 250 | 168 | 245 | 199 | 208 | 154 | 180 | 141 | 143 | 130 |
| 32 | 32 | 32 | 16 | 32 | 16 | 64 | 32 | 32 | | | |
| 158 | 128 | 191 | 128 | 255 | 158 | 255 | 224 | 255 | | | |
| 255 | 255 | 319 | 255 | 383 | 255 | 383 | 255 | 255 | | | |
| 383 | 383 | 510 | 319 | 510 | 383 | 510 | 383 | 383 | | | |
| 510 | 510 | 510 | 510 | 766 | 510 | 637 | 510 | 510 | | | |
| 765 | 510 | 766 | 510 | 766 | 510 | 765 | 510 | 637 | | | |
| 766 | 637 | 1,021 | 765 | 1,020 | 1,021 | 1,020 | 765 | 1,020 | | | |
| 1,020 | 765 | 1,530 | 765 | 1,020 | 1,275 | 1,020 | 1,020 | 1,020 | 510 | 510 | 510 |
| 1.9 | 1.0 | 1.3 | 2.7 | 8.6 | 4.7 | 9.8 | 5.9 | 0 | 4.2 | 0 | 0 |
| 1.2 | .6 | 1.2 | 3.1 | 2.8 | 1.4 | 2.7 | 1.4 | 0 | 12.5 | 0 | 0 |

TABLE 7.30.--MEAT LOAF
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 8.8 | 2.7 | 9.3 | 8.5 | 10.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 7.8 | 2.2 | 8.7 | 7.8 | 10.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .9 | .5 | .5 | .5 | .6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | 0 | .2 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 1.2 | 1.2 | 1.1 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .4 | .3 | .4 | .3 |
| MAXIMUM.....DO..... | 4.0 | 2.0 | 2.0 | 3.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 48 | 18 | 26 | 40 | 35 |
| STANDARD DEVIATION.....GM..... | 38 | 13 | 24 | 48 | 27 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 14 | | 7 | 9 | 9 |
| 25TH.....GM..... | 28 | | 14 | 19 | 19 |
| 50TH.....GM..... | 38 | | 19 | 28 | 28 |
| 75TH.....GM..... | 57 | | 29 | 38 | 38 |
| 90TH.....GM..... | 88 | | 38 | 66 | 57 |
| 95TH.....GM..... | 117 | | 56 | 119 | 88 |
| 99TH.....GM..... | 188 | | 150 | 300 | 150 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 133 | 45 | 72 | 98 | 98 |
| STANDARD DEVIATION.....GM..... | 101 | 34 | 67 | 84 | 76 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 111 | 44 | 89 | 85 | 97 |
| STANDARD DEVIATION.....GM..... | 62 | 15 | 35 | 50 | 54 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 142 | 0 | 0 | 300 | 0 |
| STANDARD DEVIATION.....GM..... | 104 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,056 | 113 | 528 | 675 | 528 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,406 | 113 | 528 | 900 | 528 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 125 | 45 | 72 | 105 | 96 |
| STANDARD DEVIATION.....GM..... | 93 | 30 | 66 | 92 | 75 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 42 | | 22 | 28 | 28 |
| 25TH.....GM..... | 57 | | 43 | 56 | 56 |
| 50TH.....GM..... | 113 | | 57 | 85 | 85 |
| 75TH.....GM..... | 169 | | 85 | 113 | 113 |
| 90TH.....GM..... | 225 | | 113 | 225 | 170 |
| 95TH.....GM..... | 284 | | 113 | 338 | 225 |
| 99TH.....GM..... | 450 | | 450 | 450 | 450 |
| MAXIMUM.....GM..... | 1,056 | 113 | 528 | 675 | 528 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 137.6 | 0 | 2.4 | 4.0 | 9.0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 3.6 | 0 | 2.3 | 2.4 | 4.3 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 11.0 | 9.2 | 10.3 | 8.9 | 8.6 | 8.2 | 8.8 | 8.9 | 8.9 | 6.8 | 7.0 | 7.8 |
| 10.4 | 8.7 | 9.3 | 8.0 | 7.8 | 7.3 | 7.5 | 7.7 | 6.9 | 5.7 | 6.2 | 6.3 |
| .5 | .5 | 1.0 | .8 | .8 | .8 | 1.2 | 1.2 | 1.8 | 1.0 | .8 | 1.5 |
| (2) | 0 | .1 | .1 | 0 | .1 | .1 | .1 | .2 | .1 | 0 | 0 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.2 | 1.2 | 1.2 | 1.3 | 1.2 | 1.1 | 1.3 |
| .3 | .3 | .4 | .4 | .4 | .4 | .4 | .4 | .5 | .5 | .4 | .5 |
| 3.0 | 2.0 | 3.0 | 3.0 | 3.0 | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 | 2.0 | 3.0 |
| 44 | 41 | 59 | 48 | 63 | 47 | 59 | 47 | 54 | 40 | 49 | 45 |
| 33 | 34 | 37 | 31 | 44 | 31 | 38 | 39 | 33 | 25 | 49 | 46 |
| 14 | 14 | 19 | 14 | 15 | 14 | 17 | 14 | 19 | 7 | | 7 |
| 27 | 19 | 38 | 28 | 38 | 28 | 29 | 28 | 29 | 19 | | 21 |
| 38 | 29 | 47 | 38 | 57 | 38 | 56 | 38 | 44 | 38 | | 28 |
| 57 | 44 | 75 | 66 | 76 | 57 | 76 | 57 | 71 | 57 | | 57 |
| 76 | 75 | 113 | 76 | 113 | 85 | 112 | 76 | 113 | 76 | | 103 |
| 85 | 110 | 150 | 101 | 138 | 113 | 132 | 110 | 132 | 76 | | 150 |
| 175 | 225 | 176 | 176 | 227 | 149 | 191 | 192 | 150 | 113 | | 235 |
| 127 | 118 | 164 | 139 | 180 | 127 | 162 | 129 | 133 | 102 | 147 | 113 |
| 99 | 97 | 101 | 93 | 117 | 84 | 106 | 102 | 70 | 61 | 150 | 128 |
| 86 | 105 | 135 | 91 | 135 | 113 | 124 | 98 | 126 | 115 | 80 | 115 |
| 30 | 76 | 61 | 37 | 108 | 46 | 62 | 47 | 58 | 23 | 66 | 71 |
| 131 | 0 | 189 | 57 | 0 | 120 | 132 | 137 | 132 | 57 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 23 | 27 | 156 | 0 | 0 | 0 | 0 |
| 792 | 792 | 528 | 528 | 1,056 | 450 | 675 | 1,056 | 352 | 281 | 675 | 704 |
| 792 | 792 | 567 | 528 | 1,276 | 450 | 732 | 1,406 | 450 | 338 | 675 | 704 |
| 120 | 113 | 157 | 125 | 162 | 121 | 145 | 119 | 127 | 98 | 129 | 106 |
| 93 | 81 | 93 | 86 | 115 | 75 | 98 | 96 | 60 | 51 | 123 | 86 |
| 43 | 43 | 43 | 43 | 43 | 43 | 43 | 43 | 43 | 28 | | 22 |
| 71 | 57 | 112 | 82 | 86 | 74 | 85 | 66 | 85 | 57 | | 57 |
| 86 | 86 | 129 | 113 | 114 | 113 | 113 | 88 | 113 | 113 | | 85 |
| 149 | 113 | 176 | 169 | 225 | 169 | 176 | 132 | 170 | 113 | | 113 |
| 225 | 225 | 264 | 225 | 264 | 225 | 264 | 225 | 224 | 170 | | 220 |
| 227 | 330 | 340 | 227 | 352 | 227 | 338 | 227 | 225 | 170 | | 352 |
| 524 | 450 | 495 | 528 | 676 | 397 | 495 | 454 | 225 | 281 | | 450 |
| 792 | 450 | 528 | 528 | 1,056 | 450 | 675 | 1,056 | 352 | 281 | 675 | 450 |
| 13.8 | 9.8 | 1.2 | 7.8 | 13.4 | 19.5 | 19.8 | 25.9 | 3.7 | 7.3 | 0 | 0 |
| 5.5 | 4.5 | .7 | 5.2 | 3.4 | 3.9 | 3.8 | 3.4 | 2.9 | 5.0 | 0 | 0 |

TABLE 7.31.--TUNA SALAD
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|------------|------------|------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE | MALE | MALE | MALE |
| | | AND FEMALE | AND FEMALE | AND FEMALE | AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.4 | .4 | 2.6 | 4.1 | 3.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.9 | .4 | 2.4 | 3.6 | 3.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | 0 | .2 | .5 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | .1 | .1 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.2 | 1.0 | 1.1 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .4 | .0 | .4 | .4 | .5 |
| MAXIMUM.....DO..... | 4.0 | 1.0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 39 | 11 | 20 | 26 | 27 |
| STANDARD DEVIATION.....GM..... | 30 | 6 | 18 | 20 | 17 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 9 | | | 4 | 8 |
| 25TH.....GM..... | 17 | | | 9 | 17 |
| 50TH.....GM..... | 34 | | | 17 | 26 |
| 75TH.....GM..... | 51 | | | 34 | 34 |
| 90TH.....GM..... | 68 | | | 51 | 51 |
| 95TH.....GM..... | 103 | | | 68 | 68 |
| 99TH.....GM..... | 137 | | | 85 | 85 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 106 | 33 | 56 | 75 | 78 |
| STANDARD DEVIATION.....GM..... | 82 | 19 | 53 | 60 | 47 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 87 | 0 | 62 | 54 | 55 |
| STANDARD DEVIATION.....GM..... | 53 | 0 | 13 | 25 | 27 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 79 | 0 | 20 | 18 | 41 |
| STANDARD DEVIATION.....GM..... | 45 | 0 | 0 | 0 | 30 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 666 | 51 | 205 | 256 | 205 |
| MAXIMUM OVER 3 DAYS.....GM..... | 666 | 51 | 205 | 256 | 256 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 99 | 33 | 54 | 68 | 68 |
| STANDARD DEVIATION.....GM..... | 75 | 19 | 49 | 56 | 42 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 25 | | | 12 | 13 |
| 25TH.....GM..... | 51 | | | 26 | 51 |
| 50TH.....GM..... | 85 | | | 51 | 51 |
| 75TH.....GM..... | 103 | | | 103 | 103 |
| 90TH.....GM..... | 205 | | | 154 | 103 |
| 95TH.....GM..... | 205 | | | 205 | 154 |
| 99TH.....GM..... | 410 | | | 256 | 205 |
| MAXIMUM.....GM..... | 666 | 51 | 205 | 256 | 205 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 22.2 | 0 | 1.5 | 0 | 1.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.1 | 0 | 4.7 | 0 | 1.8 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.8 | 4.0 | 4.2 | 5.3 | 4.4 | 5.5 | 3.6 | 6.0 | 3.4 | 4.2 | 1.9 | 1.8 |
| 3.2 | 3.6 | 3.9 | 4.6 | 3.9 | 4.7 | 3.2 | 5.3 | 2.9 | 3.6 | 1.9 | 1.8 |
| .5 | .4 | .2 | .5 | .4 | .6 | .4 | .5 | .5 | .6 | 0 | 0 |
| .1 | 0 | .2 | .2 | .1 | .2 | 0 | .2 | 0 | .1 | 0 | 0 |
| 1.2 | 1.1 | 1.2 | 1.2 | 1.2 | 1.2 | 1.1 | 1.2 | 1.1 | 1.2 | 1.0 | 1.0 |
| .5 | .3 | .5 | .5 | .4 | .5 | .3 | .5 | .4 | .4 | .0 | .0 |
| 3.0 | 2.0 | 3.0 | 4.0 | 3.0 | 3.0 | 2.0 | 3.0 | 2.0 | 3.0 | 1.0 | 1.0 |
| 42 | 35 | 43 | 38 | 40 | 37 | 48 | 40 | 34 | 39 | 41 | 27 |
| 33 | 27 | 37 | 34 | 26 | 28 | 39 | 31 | 19 | 27 | 36 | 21 |
| 9 | 9 | 9 | 9 | 13 | 9 | 13 | 9 | | 8 | | |
| 17 | 17 | 17 | 17 | 23 | 17 | 23 | 17 | | 17 | | |
| 28 | 34 | 34 | 28 | 34 | 34 | 34 | 34 | | 34 | | |
| 68 | 42 | 68 | 51 | 51 | 51 | 61 | 51 | | 51 | | |
| 103 | 68 | 92 | 68 | 68 | 68 | 137 | 71 | | 91 | | |
| 111 | 85 | 137 | 103 | 91 | 92 | 137 | 111 | | 103 | | |
| 137 | 137 | 171 | 137 | 137 | 137 | 137 | 137 | | 103 | | |
| 109 | 93 | 121 | 98 | 112 | 102 | 135 | 110 | 91 | 110 | 124 | 80 |
| 87 | 67 | 101 | 76 | 72 | 78 | 109 | 83 | 52 | 76 | 109 | 63 |
| 107 | 98 | 61 | 90 | 92 | 78 | 116 | 89 | 82 | 76 | 0 | 0 |
| 57 | 59 | 57 | 63 | 47 | 45 | 71 | 51 | 25 | 39 | 0 | 0 |
| 64 | 0 | 103 | 128 | 85 | 60 | 0 | 101 | 0 | 102 | 0 | 0 |
| 38 | 0 | 57 | 60 | 0 | 40 | 0 | 25 | 0 | 0 | 0 | 0 |
| 410 | 308 | 513 | 410 | 410 | 410 | 666 | 410 | 205 | 308 | 308 | 205 |
| 410 | 410 | 513 | 615 | 410 | 410 | 666 | 410 | 205 | 308 | 308 | 205 |
| 103 | 93 | 111 | 97 | 102 | 93 | 130 | 104 | 89 | 102 | 124 | 80 |
| 77 | 65 | 92 | 74 | 65 | 71 | 101 | 73 | 47 | 70 | 109 | 63 |
| 26 | 26 | 26 | 25 | 26 | 26 | 26 | 25 | | 25 | | |
| 51 | 51 | 51 | 51 | 51 | 51 | 57 | 51 | | 51 | | |
| 85 | 103 | 85 | 78 | 103 | 68 | 103 | 103 | | 103 | | |
| 103 | 103 | 137 | 103 | 103 | 103 | 154 | 115 | | 113 | | |
| 205 | 205 | 205 | 205 | 205 | 205 | 205 | 205 | | 205 | | |
| 205 | 205 | 308 | 205 | 205 | 205 | 410 | 205 | | 284 | | |
| 410 | 308 | 512 | 410 | 308 | 410 | 410 | 410 | | 308 | | |
| 410 | 308 | 513 | 410 | 410 | 410 | 666 | 410 | 205 | 308 | 308 | 205 |
| 0 | 0 | 1.3 | 1.9 | 0 | 2.8 | 5.0 | 6.8 | 1.5 | 0 | 0 | 0 |
| 0 | 0 | 1.9 | 2.0 | 0 | .8 | 2.5 | 1.4 | 3.6 | 0 | 0 | 0 |

TABLE 7.32.--POT PIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.1 | .4 | 2.3 | 2.0 | 2.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.0 | .4 | 2.1 | 1.9 | 2.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .1 | 0 | .1 | .1 | (2) |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | .2 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.0 | 1.0 | 1.3 | 1.1 | 1.0 |
| STANDARD DEVIATION.....DO..... | .2 | .0 | .7 | .2 | .1 |
| MAXIMUM.....DO..... | 4.0 | 1.0 | 4.0 | 2.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 91 | 38 | 75 | 68 | 67 |
| STANDARD DEVIATION.....GM..... | 49 | 22 | 57 | 31 | 23 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 38 | | | | |
| 25TH.....GM..... | 76 | | | | |
| 50TH.....GM..... | 76 | | | | |
| 75TH.....GM..... | 95 | | | | |
| 90TH.....GM..... | 160 | | | | |
| 95TH.....GM..... | 167 | | | | |
| 99TH.....GM..... | 250 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 260 | 115 | 178 | 186 | 197 |
| STANDARD DEVIATION.....GM..... | 126 | 65 | 96 | 64 | 59 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 296 | 0 | 227 | 234 | 227 |
| STANDARD DEVIATION.....GM..... | 110 | 0 | 0 | 6 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 232 | 0 | 232 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 42 | 0 | 42 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 960 | 188 | 454 | 360 | 360 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,200 | 188 | 850 | 480 | 454 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 261 | 115 | 180 | 192 | 198 |
| STANDARD DEVIATION.....GM..... | 127 | 65 | 79 | 62 | 58 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 113 | | | | |
| 25TH.....GM..... | 227 | | | | |
| 50TH.....GM..... | 227 | | | | |
| 75TH.....GM..... | 250 | | | | |
| 90TH.....GM..... | 454 | | | | |
| 95TH.....GM..... | 480 | | | | |
| 99TH.....GM..... | 750 | | | | |
| MAXIMUM.....GM..... | 960 | 188 | 454 | 360 | 360 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 5.6 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .7 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY, 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.1 | 2.7 | 2.4 | 1.8 | 2.6 | 2.1 | 1.6 | 1.7 | 2.2 | 3.0 | 2.0 | 1.6 |
| 2.9 | 2.6 | 2.2 | 1.8 | 2.5 | 2.1 | 1.5 | 1.7 | 2.0 | 2.9 | 1.7 | 1.6 |
| .2 | .1 | .2 | .1 | .1 | (2) | .1 | (2) | .2 | (2) | .2 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1.1 | 1.0 | 1.1 | 1.0 | 1.0 | 1.0 | 1.1 | 1.0 | 1.1 | 1.0 | 1.1 | 1.0 |
| .3 | .2 | .3 | .2 | .2 | .1 | .2 | .2 | .3 | .1 | .3 | .0 |
| 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 1.0 |
| 90 | 85 | 119 | 81 | 107 | 88 | 114 | 86 | 94 | 75 | 116 | 88 |
| 39 | 40 | 49 | 31 | 53 | 34 | 82 | 43 | 37 | 30 | 61 | 28 |
| 40 | 38 | | | 47 | 53 | 38 | 33 | | 38 | | |
| 76 | 72 | | | 76 | 76 | 76 | 76 | | 76 | | |
| 76 | 76 | | | 76 | 76 | 76 | 76 | | 76 | | |
| 95 | 83 | | | 151 | 83 | 151 | 83 | | 76 | | |
| 151 | 151 | | | 167 | 151 | 200 | 151 | | 100 | | |
| 160 | 160 | | | 227 | 160 | 303 | 160 | | 100 | | |
| 240 | 240 | | | 240 | 160 | 400 | 240 | | 250 | | |
| 259 | 240 | 346 | 231 | 312 | 259 | 303 | 251 | 266 | 217 | 334 | 263 |
| 110 | 97 | 148 | 81 | 150 | 100 | 177 | 120 | 104 | 60 | 191 | 83 |
| 234 | 288 | 233 | 240 | 340 | 227 | 544 | 288 | 227 | 375 | 227 | 0 |
| 6 | 66 | 6 | 0 | 0 | 0 | 71 | 47 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 720 | 625 | 720 | 500 | 750 | 750 | 960 | 720 | 480 | 454 | 750 | 500 |
| 720 | 720 | 720 | 500 | 750 | 750 | 1,200 | 720 | 480 | 750 | 750 | 500 |
| 252 | 244 | 327 | 231 | 310 | 258 | 326 | 253 | 260 | 222 | 311 | 263 |
| 100 | 98 | 142 | 77 | 147 | 98 | 204 | 117 | 96 | 66 | 175 | 83 |
| 120 | 114 | | | 142 | 158 | 113 | 99 | | 114 | | |
| 227 | 227 | | | 227 | 227 | 227 | 227 | | 227 | | |
| 227 | 227 | | | 227 | 227 | 227 | 227 | | 227 | | |
| 240 | 240 | | | 454 | 250 | 454 | 284 | | 227 | | |
| 360 | 375 | | | 480 | 454 | 500 | 375 | | 300 | | |
| 454 | 480 | | | 568 | 480 | 901 | 480 | | 300 | | |
| 709 | 625 | | | 721 | 480 | 960 | 720 | | 454 | | |
| 720 | 625 | 720 | 500 | 750 | 750 | 960 | 720 | 480 | 454 | 750 | 500 |
| 0 | 1.0 | 0 | .9 | 0 | 0 | 1.2 | .8 | 0 | 1.7 | 0 | 0 |
| 0 | 1.6 | 0 | 3.2 | 0 | 0 | 1.5 | .6 | 0 | 3.2 | 0 | 0 |

TABLE 7.33.--FROZEN PLATE MEALS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 1.4 | .4 | 1.6 | 1.2 | 1.0 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.2 | .4 | 1.6 | 1.0 | .9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .1 | 0 | .1 | .1 | 0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | 0 | .1 | (2) |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 1.0 | 1.0 | 1.2 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .0 | .2 | .5 | .4 |
| MAXIMUM.....DO..... | 3.0 | 1.0 | 2.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 122 | 19 | 96 | 117 | 111 |
| STANDARD DEVIATION.....GM..... | 57 | (2) | 34 | 72 | 52 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 57 | | | | |
| 25TH.....GM..... | 93 | | | | |
| 50TH.....GM..... | 104 | | | | |
| 75TH.....GM..... | 142 | | | | |
| 90TH.....GM..... | 196 | | | | |
| 95TH.....GM..... | 238 | | | | |
| 99TH.....GM..... | 335 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 324 | 56 | 271 | 295 | 308 |
| STANDARD DEVIATION.....GM..... | 110 | (2) | 70 | 67 | 105 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 307 | 0 | 312 | 142 | 0 |
| STANDARD DEVIATION.....GM..... | 60 | 0 | 0 | 0 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 300 | 0 | 0 | 369 | 301 |
| STANDARD DEVIATION.....GM..... | 50 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 907 | 57 | 340 | 539 | 510 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,106 | 57 | 624 | 1,106 | 903 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 319 | 56 | 274 | 296 | 307 |
| STANDARD DEVIATION.....GM..... | 103 | (2) | 68 | 80 | 99 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 142 | | | | |
| 25TH.....GM..... | 279 | | | | |
| 50TH.....GM..... | 312 | | | | |
| 75TH.....GM..... | 340 | | | | |
| 90TH.....GM..... | 454 | | | | |
| 95TH.....GM..... | 510 | | | | |
| 99TH.....GM..... | 680 | | | | |
| MAXIMUM.....GM..... | 907 | 57 | 340 | 539 | 510 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 2.7 | 0 | 0 | 0 | .7 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | 0 | 0 | 3.6 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.3 | 1.1 | 1.1 | 1.5 | 1.5 | 1.5 | 1.2 | 1.4 | 2.5 | 1.7 | 1.5 | 2.1 |
| 1.3 | 1.0 | .9 | 1.4 | 1.3 | 1.3 | 1.0 | 1.3 | 2.0 | 1.7 | 1.4 | 1.7 |
| (2) | .1 | .2 | .2 | .1 | .2 | .1 | .1 | .3 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | .1 | 0 | .1 | (2) | .3 | 0 | .1 | .4 |
| 1.0 | 1.1 | 1.1 | 1.1 | 1.2 | 1.1 | 1.2 | 1.1 | 1.3 | 1.0 | 1.2 | 1.4 |
| .2 | .3 | .3 | .3 | .5 | .3 | .6 | .4 | .7 | .0 | .6 | .8 |
| 2.0 | 2.0 | 2.0 | 2.0 | 3.0 | 2.0 | 3.0 | 3.0 | 3.0 | 1.0 | 3.0 | 3.0 |
| 106 | 102 | 134 | 107 | 134 | 119 | 151 | 121 | 135 | 105 | 110 | 133 |
| 37 | 47 | 71 | 38 | 64 | 46 | 72 | 50 | 60 | 16 | 46 | 82 |
| | | | | 80 | 57 | 78 | 47 | | | | |
| | | | | 93 | 95 | 102 | 95 | | | | |
| | | | | 104 | 104 | 113 | 109 | | | | |
| | | | | 151 | 149 | 196 | 149 | | | | |
| | | | | 217 | 186 | 265 | 180 | | | | |
| | | | | 301 | 210 | 302 | 217 | | | | |
| | | | | 340 | 248 | 343 | 333 | | | | |
| 308 | 280 | 340 | 298 | 340 | 321 | 386 | 339 | 328 | 315 | 286 | 282 |
| 97 | 108 | 146 | 97 | 83 | 97 | 163 | 115 | 92 | 49 | 8 | 56 |
| 312 | 276 | 399 | 248 | 343 | 309 | 325 | 300 | 287 | 0 | 0 | 0 |
| 0 | 88 | 60 | 42 | 21 | 43 | 46 | 40 | 8 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 328 | 0 | 318 | 256 | 266 | 0 | 256 | 299 |
| 0 | 0 | 0 | 0 | 16 | 0 | 19 | 83 | 45 | 0 | 0 | 16 |
| 539 | 539 | 680 | 482 | 539 | 539 | 907 | 680 | 510 | 397 | 312 | 340 |
| 624 | 680 | 922 | 567 | 1,021 | 744 | 1,028 | 999 | 1,020 | 397 | 769 | 936 |
| 309 | 279 | 354 | 288 | 338 | 318 | 364 | 330 | 306 | 315 | 279 | 289 |
| 94 | 105 | 144 | 91 | 72 | 91 | 138 | 113 | 100 | 49 | 31 | 45 |
| | | | | 241 | 142 | 248 | 113 | | | | |
| | | | | 312 | 284 | 284 | 280 | | | | |
| | | | | 312 | 312 | 312 | 312 | | | | |
| | | | | 347 | 340 | 397 | 369 | | | | |
| | | | | 454 | 447 | 532 | 454 | | | | |
| | | | | 482 | 538 | 680 | 540 | | | | |
| | | | | 538 | 539 | 907 | 680 | | | | |
| 539 | 539 | 680 | 482 | 539 | 539 | 907 | 680 | 510 | 397 | 312 | 340 |
| 0 | .7 | 0 | 0 | 0 | 0 | 1.3 | 0 | 0 | 0 | 0 | 0 |
| 0 | 2.6 | 0 | 0 | 0 | 0 | 1.9 | 0 | 0 | 0 | 0 | 0 |

TABLE 7.34.--HAMBURGER AND CHEESEBURGER SANDWICHES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 10.0 | 0 | 7.1 | 9.4 | 10.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 8.5 | 0 | 6.3 | 8.5 | 9.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.2 | 0 | .7 | .8 | .9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | 0 | 0 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 0 | 1.1 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .5 | 0 | .4 | .3 | .4 |
| MAXIMUM.....DO..... | 6.0 | 0 | 3.0 | 2.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 65 | 0 | 38 | 43 | 51 |
| STANDARD DEVIATION.....GM..... | 44 | 0 | 23 | 27 | 36 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 32 | | 9 | 16 | 28 |
| 25TH.....GM..... | 37 | | 19 | 32 | 32 |
| 50TH.....GM..... | 52 | | 32 | 32 | 37 |
| 75TH.....GM..... | 69 | | 51 | 52 | 56 |
| 90TH.....GM..... | 113 | | 65 | 65 | 77 |
| 95TH.....GM..... | 154 | | 95 | 103 | 113 |
| 99TH.....GM..... | 244 | | 120 | 162 | 248 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 160 | 0 | 98 | 118 | 133 |
| STANDARD DEVIATION.....GM..... | 75 | 0 | 53 | 64 | 60 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 174 | 0 | 120 | 130 | 157 |
| STANDARD DEVIATION.....GM..... | 74 | 0 | 36 | 57 | 73 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 195 | 0 | 0 | 0 | 310 |
| STANDARD DEVIATION.....GM..... | 77 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 770 | 0 | 336 | 485 | 558 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,386 | 0 | 359 | 558 | 930 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 161 | 0 | 99 | 116 | 138 |
| STANDARD DEVIATION.....GM..... | 74 | 0 | 49 | 58 | 65 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 97 | | 42 | 49 | 84 |
| 25TH.....GM..... | 97 | | 73 | 97 | 97 |
| 50TH.....GM..... | 157 | | 97 | 97 | 111 |
| 75TH.....GM..... | 186 | | 111 | 113 | 168 |
| 90TH.....GM..... | 266 | | 157 | 170 | 211 |
| 95TH.....GM..... | 308 | | 186 | 222 | 266 |
| 99TH.....GM..... | 444 | | 336 | 336 | 372 |
| MAXIMUM.....GM..... | 616 | 0 | 336 | 485 | 558 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 2.3 | 0 | 1.0 | 1.2 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | (2) | 0 | 1.3 | .7 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 13.1 | 12.9 | 19.0 | 15.1 | 18.0 | 11.6 | 10.4 | 5.2 | 3.6 | 3.2 | 2.1 | 1.2 |
| 11.7 | 12.1 | 14.3 | 12.7 | 13.6 | 9.6 | 8.9 | 4.8 | 3.2 | 3.1 | 2.1 | 1.2 |
| 1.2 | .6 | 3.7 | 1.6 | 3.4 | 1.9 | 1.0 | .3 | .4 | (2) | 0 | 0 |
| .2 | .2 | 1.0 | .7 | 1.0 | .1 | .5 | .1 | 0 | 0 | 0 | 0 |
| 1.1 | 1.1 | 1.4 | 1.3 | 1.3 | 1.2 | 1.2 | 1.1 | 1.2 | 1.0 | 1.0 | 1.0 |
| .4 | .4 | .7 | .7 | .6 | .4 | .5 | .4 | .4 | .1 | .0 | .0 |
| 4.0 | 4.0 | 5.0 | 6.0 | 4.0 | 3.0 | 5.0 | 3.0 | 2.0 | 2.0 | 1.0 | 1.0 |
| 58 | 50 | 82 | 60 | 84 | 65 | 71 | 53 | 54 | 46 | 53 | 45 |
| 36 | 24 | 59 | 40 | 56 | 35 | 43 | 31 | 22 | 18 | 12 | 10 |
| 32 | 30 | 32 | 32 | 32 | 32 | 32 | 28 | | 28 | | |
| 32 | 32 | 51 | 32 | 52 | 37 | 51 | 32 | | 32 | | |
| 52 | 38 | 61 | 52 | 62 | 56 | 61 | 51 | | 32 | | |
| 65 | 61 | 99 | 65 | 105 | 69 | 89 | 61 | | 61 | | |
| 105 | 76 | 157 | 97 | 154 | 111 | 124 | 89 | | 74 | | |
| 124 | 89 | 198 | 126 | 210 | 150 | 177 | 103 | | 89 | | |
| 174 | 154 | 333 | 207 | 266 | 186 | 238 | 177 | | 89 | | |
| 155 | 138 | 180 | 147 | 188 | 165 | 181 | 143 | 142 | 137 | 160 | 136 |
| 73 | 50 | 86 | 62 | 89 | 64 | 88 | 55 | 41 | 53 | 35 | 31 |
| 169 | 136 | 195 | 162 | 199 | 161 | 158 | 178 | 152 | 111 | 0 | 0 |
| 94 | 51 | 87 | 65 | 70 | 60 | 63 | 94 | 20 | 0 | 0 | 0 |
| 143 | 134 | 218 | 146 | 220 | 191 | 189 | 127 | 0 | 0 | 0 | 0 |
| 21 | 45 | 96 | 83 | 75 | 52 | 55 | 28 | 0 | 0 | 0 | 0 |
| 732 | 340 | 616 | 532 | 770 | 497 | 532 | 454 | 266 | 266 | 200 | 186 |
| 1,098 | 652 | 1,386 | 1,029 | 1,330 | 798 | 873 | 871 | 340 | 266 | 200 | 186 |
| 154 | 136 | 182 | 143 | 191 | 162 | 175 | 145 | 138 | 136 | 160 | 136 |
| 73 | 49 | 90 | 53 | 80 | 62 | 82 | 61 | 38 | 53 | 35 | 31 |
| 97 | 85 | 97 | 97 | 97 | 97 | 97 | 85 | | 97 | | |
| 97 | 97 | 111 | 97 | 154 | 111 | 111 | 97 | | 97 | | |
| 154 | 111 | 157 | 142 | 183 | 157 | 157 | 154 | | 97 | | |
| 183 | 157 | 194 | 183 | 222 | 186 | 186 | 170 | | 183 | | |
| 266 | 194 | 308 | 194 | 266 | 266 | 266 | 222 | | 186 | | |
| 314 | 227 | 366 | 266 | 340 | 266 | 366 | 266 | | 266 | | |
| 399 | 266 | 532 | 308 | 532 | 336 | 532 | 340 | | 266 | | |
| 523 | 340 | 608 | 334 | 616 | 485 | 532 | 454 | 266 | 266 | 200 | 186 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 7.35.--SOUPS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 31.3 | 12.4 | 47.0 | 39.7 | 31.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 22.1 | 6.5 | 29.6 | 27.5 | 24.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 7.1 | 4.8 | 13.1 | 9.6 | 5.5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.1 | 1.1 | 4.3 | 2.7 | 1.5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.4 | 1.7 | 1.5 | 1.4 | 1.3 |
| STANDARD DEVIATION.....DO..... | .7 | .8 | .8 | .7 | .6 |
| MAXIMUM.....DO..... | 10.0 | 4.0 | 7.0 | 5.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 139 | 80 | 96 | 101 | 110 |
| STANDARD DEVIATION.....GM..... | 100 | 69 | 80 | 66 | 70 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 40 | 10 | 20 | 31 | 31 |
| 25TH.....GM..... | 80 | 20 | 41 | 60 | 76 |
| 50TH.....GM..... | 107 | 63 | 80 | 82 | 82 |
| 75TH.....GM..... | 163 | 122 | 121 | 123 | 160 |
| 90TH.....GM..... | 245 | 160 | 163 | 189 | 204 |
| 95TH.....GM..... | 327 | 188 | 242 | 242 | 243 |
| 99TH.....GM..... | 509 | 395 | 327 | 320 | 345 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 306 | 113 | 188 | 224 | 259 |
| STANDARD DEVIATION.....GM..... | 178 | 90 | 102 | 117 | 136 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 307 | 173 | 196 | 219 | 274 |
| STANDARD DEVIATION.....GM..... | 150 | 80 | 110 | 89 | 121 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 310 | 177 | 216 | 215 | 224 |
| STANDARD DEVIATION.....GM..... | 145 | 105 | 142 | 95 | 68 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 2,450 | 735 | 1,800 | 827 | 1,080 |
| MAXIMUM OVER 3 DAYS.....GM..... | 4,595 | 1,185 | 3,180 | 2,040 | 1,455 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 290 | 140 | 186 | 211 | 249 |
| STANDARD DEVIATION.....GM..... | 155 | 97 | 103 | 100 | 125 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 114 | 40 | 60 | 81 | 92 |
| 25TH.....GM..... | 240 | 61 | 120 | 123 | 180 |
| 50TH.....GM..... | 245 | 123 | 180 | 240 | 240 |
| 75TH.....GM..... | 360 | 188 | 245 | 245 | 245 |
| 90TH.....GM..... | 490 | 245 | 245 | 250 | 480 |
| 95TH.....GM..... | 600 | 250 | 368 | 375 | 490 |
| 99TH.....GM..... | 736 | 368 | 601 | 601 | 721 |
| MAXIMUM.....GM..... | 1,960 | 735 | 735 | 735 | 858 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 191.4 | 0 | 8.8 | 6.9 | 2.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.1 | 0 | 1.2 | .7 | .3 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 27.8 | 29.4 | 24.1 | 24.0 | 27.0 | 29.9 | 32.2 | 31.4 | 37.5 | 36.9 | 34.6 | 42.1 |
| 22.0 | 22.7 | 18.3 | 19.0 | 20.8 | 22.5 | 21.6 | 21.4 | 23.6 | 23.7 | 19.7 | 25.7 |
| 4.9 | 5.5 | 4.5 | 3.9 | 5.2 | 5.8 | 7.8 | 7.8 | 10.4 | 9.4 | 9.9 | 11.7 |
| .9 | 1.3 | .8 | 1.2 | 1.0 | 1.6 | 2.8 | 2.1 | 3.6 | 3.8 | 5.0 | 4.8 |
| 1.3 | 1.3 | 1.3 | 1.3 | 1.3 | 1.4 | 1.5 | 1.5 | 1.5 | 1.6 | 1.7 | 1.6 |
| .6 | .6 | .6 | .6 | .6 | .7 | .8 | .8 | .8 | .8 | 1.0 | .8 |
| 10.0 | 5.0 | 4.0 | 5.0 | 4.0 | 6.0 | 6.0 | 9.0 | 5.0 | 5.0 | 6.0 | 5.0 |
| 126 | 124 | 156 | 127 | 152 | 131 | 166 | 141 | 172 | 150 | 194 | 145 |
| 99 | 79 | 120 | 85 | 104 | 96 | 112 | 95 | 117 | 98 | 162 | 97 |
| 41 | 41 | 40 | 27 | 41 | 40 | 50 | 40 | 60 | 41 | 41 | 41 |
| 80 | 80 | 80 | 80 | 82 | 80 | 82 | 80 | 82 | 82 | 82 | 80 |
| 82 | 90 | 120 | 83 | 123 | 85 | 151 | 120 | 160 | 123 | 160 | 123 |
| 162 | 160 | 179 | 163 | 203 | 163 | 208 | 163 | 213 | 194 | 242 | 163 |
| 240 | 204 | 323 | 240 | 280 | 243 | 323 | 245 | 327 | 250 | 395 | 250 |
| 250 | 245 | 404 | 306 | 327 | 323 | 400 | 323 | 405 | 322 | 512 | 320 |
| 483 | 485 | 613 | 487 | 571 | 551 | 561 | 492 | 605 | 478 | 817 | 490 |
| 298 | 297 | 336 | 310 | 357 | 289 | 361 | 306 | 359 | 316 | 355 | 286 |
| 148 | 150 | 181 | 183 | 207 | 156 | 215 | 159 | 229 | 164 | 305 | 133 |
| 316 | 295 | 405 | 276 | 359 | 311 | 355 | 295 | 356 | 312 | 353 | 288 |
| 160 | 143 | 199 | 115 | 149 | 150 | 168 | 130 | 148 | 149 | 151 | 126 |
| 305 | 260 | 442 | 329 | 384 | 337 | 329 | 323 | 326 | 288 | 411 | 301 |
| 293 | 96 | 180 | 94 | 151 | 138 | 117 | 134 | 150 | 107 | 222 | 109 |
| 1,940 | 1,470 | 1,470 | 1,225 | 2,000 | 1,470 | 1,960 | 1,470 | 1,816 | 1,715 | 2,450 | 980 |
| 4,595 | 1,573 | 2,373 | 1,508 | 2,002 | 1,960 | 2,573 | 2,293 | 2,695 | 2,817 | 3,341 | 2,200 |
| 294 | 282 | 358 | 290 | 347 | 282 | 334 | 287 | 333 | 284 | 334 | 270 |
| 158 | 131 | 198 | 153 | 181 | 139 | 183 | 142 | 160 | 129 | 175 | 109 |
| 120 | 120 | 120 | 61 | 120 | 91 | 123 | 120 | 123 | 120 | 123 | 120 |
| 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 |
| 245 | 245 | 305 | 245 | 300 | 245 | 245 | 245 | 250 | 245 | 245 | 245 |
| 360 | 360 | 480 | 368 | 485 | 360 | 480 | 360 | 454 | 360 | 454 | 300 |
| 490 | 490 | 612 | 490 | 612 | 490 | 500 | 490 | 500 | 490 | 562 | 480 |
| 500 | 490 | 720 | 540 | 720 | 500 | 659 | 490 | 613 | 495 | 626 | 490 |
| 751 | 735 | 1,001 | 740 | 980 | 731 | 981 | 736 | 930 | 637 | 961 | 613 |
| 1,470 | 735 | 1,470 | 980 | 1,225 | 1,200 | 1,960 | 1,470 | 980 | 1,102 | 1,470 | 735 |
| 8.3 | 6.8 | 10.0 | 9.0 | 19.9 | 21.5 | 28.3 | 40.3 | 4.0 | 12.8 | 5.1 | 7.0 |
| 1.1 | .8 | 2.3 | 1.9 | 1.4 | 1.0 | 1.2 | 1.2 | .6 | 1.3 | 1.6 | 1.0 |

TABLE 8.01.--FLUID MILK, MILK BEVERAGES, AND MILK-BASED INFANT FORMULAS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 82.5 | 89.0 | 96.9 | 97.0 | 98.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 11.5 | 1.8 | 4.1 | 4.2 | 3.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 15.8 | 3.0 | 9.6 | 13.3 | 9.9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 55.2 | 84.3 | 83.2 | 79.5 | 84.7 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 4.9 | 11.8 | 7.1 | 5.9 | 6.2 |
| STANDARD DEVIATION.....DO..... | 3.1 | 4.8 | 3.5 | 2.7 | 2.7 |
| MAXIMUM.....DO..... | 39.0 | 34.0 | 24.0 | 17.0 | 17.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 328 | 667 | 424 | 392 | 468 |
| STANDARD DEVIATION.....GM..... | 270 | 308 | 266 | 215 | 240 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 31 | 117 | 73 | 83 | 122 |
| 25TH.....GM..... | 122 | 488 | 240 | 225 | 286 |
| 50TH.....GM..... | 248 | 683 | 368 | 366 | 447 |
| 75TH.....GM..... | 470 | 857 | 572 | 528 | 611 |
| 90TH.....GM..... | 692 | 1,030 | 773 | 691 | 793 |
| 95TH.....GM..... | 833 | 1,140 | 904 | 793 | 899 |
| 99TH.....GM..... | 1,221 | 1,443 | 1,189 | 977 | 1,139 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 246 | 195 | 181 | 269 | 280 |
| STANDARD DEVIATION.....GM..... | 171 | 155 | 183 | 204 | 186 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 289 | 331 | 275 | 286 | 333 |
| STANDARD DEVIATION.....GM..... | 189 | 329 | 161 | 149 | 161 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 418 | 695 | 470 | 441 | 514 |
| STANDARD DEVIATION.....GM..... | 279 | 289 | 256 | 201 | 223 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 4,410 | 2,083 | 1,960 | 1,960 | 2,318 |
| MAXIMUM OVER 3 DAYS.....GM..... | 10,053 | 5,856 | 5,856 | 4,087 | 4,911 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 203 | 170 | 179 | 198 | 227 |
| STANDARD DEVIATION.....GM..... | 134 | 71 | 80 | 83 | 89 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 15 | 61 | 61 | 81 | 122 |
| 25TH.....GM..... | 122 | 123 | 122 | 123 | 183 |
| 50TH.....GM..... | 244 | 171 | 183 | 184 | 244 |
| 75TH.....GM..... | 245 | 244 | 244 | 244 | 244 |
| 90TH.....GM..... | 366 | 248 | 245 | 245 | 281 |
| 95TH.....GM..... | 488 | 248 | 245 | 306 | 375 |
| 99TH.....GM..... | 552 | 310 | 488 | 488 | 490 |
| MAXIMUM.....GM..... | 1,960 | 496 | 976 | 1,069 | 1,220 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 466.1 | 4.5 | 21.8 | 31.2 | 40.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | .1 | .3 | .3 | .4 |

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|--------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 97.4 | 95.1 | 93.2 | 88.0 | 78.8 | 77.4 | 75.2 | 72.4 | 81.5 | 79.9 | 83.1 | 83.6 |
| 5.5 | 5.6 | 8.1 | 12.1 | 14.5 | 16.6 | 13.7 | 15.0 | 10.3 | 10.9 | 11.2 | 9.0 |
| 12.3 | 15.7 | 15.1 | 20.1 | 18.6 | 19.4 | 16.6 | 16.3 | 14.0 | 15.0 | 13.5 | 13.3 |
| 79.6 | 73.9 | 70.1 | 55.7 | 45.7 | 41.3 | 44.9 | 41.1 | 57.2 | 54.1 | 58.4 | 61.3 |
| 5.9 | 5.4 | 5.3 | 4.4 | 4.2 | 3.9 | 4.4 | 4.1 | 4.8 | 4.3 | 4.8 | 4.5 |
| 2.7 | 2.5 | 2.7 | 2.5 | 2.8 | 2.8 | 3.1 | 2.9 | 3.3 | 2.7 | 3.0 | 2.7 |
| 15.0 | 15.0 | 23.0 | 17.0 | 26.0 | 29.0 | 21.0 | 26.0 | 29.0 | 23.0 | 14.0 | 39.0 |
| 517 | 433 | 558 | 359 | 355 | 242 | 258 | 195 | 263 | 207 | 269 | 237 |
| 301 | 238 | 375 | 246 | 306 | 205 | 243 | 174 | 225 | 166 | 249 | 180 |
| 122 | 82 | 81 | 81 | 30 | 20 | 18 | 15 | 20 | 20 | 29 | 21 |
| 305 | 245 | 287 | 165 | 126 | 82 | 82 | 81 | 92 | 81 | 95 | 107 |
| 488 | 407 | 488 | 325 | 259 | 184 | 183 | 158 | 224 | 163 | 183 | 204 |
| 672 | 573 | 737 | 488 | 488 | 327 | 351 | 264 | 359 | 285 | 376 | 326 |
| 895 | 734 | 1,051 | 693 | 772 | 500 | 565 | 429 | 553 | 434 | 576 | 474 |
| 1,079 | 857 | 1,224 | 832 | 955 | 650 | 732 | 531 | 733 | 542 | 755 | 570 |
| 1,466 | 1,140 | 1,879 | 1,151 | 1,423 | 952 | 1,189 | 746 | 977 | 732 | 1,222 | 819 |
| 347 | 287 | 343 | 303 | 298 | 234 | 243 | 210 | 228 | 201 | 209 | 186 |
| 214 | 183 | 220 | 188 | 196 | 147 | 173 | 151 | 133 | 127 | 108 | 137 |
| 408 | 349 | 470 | 335 | 347 | 269 | 269 | 218 | 230 | 226 | 251 | 255 |
| 205 | 180 | 304 | 158 | 232 | 158 | 179 | 143 | 153 | 125 | 195 | 136 |
| 583 | 501 | 661 | 464 | 486 | 337 | 341 | 260 | 324 | 250 | 331 | 277 |
| 288 | 219 | 363 | 242 | 327 | 225 | 272 | 196 | 237 | 179 | 265 | 187 |
| 3,172 | 2,982 | 4,157 | 2,187 | 4,410 | 2,235 | 2,928 | 2,713 | 2,806 | 1,490 | 2,196 | 1,496 |
| 6,710 | 5,420 | 10,053 | 5,359 | 7,381 | 4,715 | 6,019 | 6,159 | 7,015 | 3,484 | 4,392 | 4,452 |
| 265 | 242 | 314 | 244 | 254 | 185 | 178 | 142 | 165 | 143 | 167 | 157 |
| 125 | 103 | 164 | 113 | 181 | 131 | 153 | 120 | 127 | 111 | 125 | 109 |
| 122 | 122 | 122 | 61 | 15 | 11 | 10 | 5 | 5 | 5 | 5 | 10 |
| 244 | 206 | 244 | 244 | 122 | 61 | 31 | 31 | 31 | 31 | 61 | 61 |
| 244 | 244 | 245 | 244 | 244 | 244 | 183 | 122 | 176 | 123 | 183 | 123 |
| 250 | 245 | 368 | 245 | 366 | 245 | 245 | 244 | 244 | 244 | 244 | 244 |
| 488 | 366 | 490 | 366 | 488 | 306 | 366 | 245 | 305 | 245 | 305 | 245 |
| 488 | 488 | 500 | 488 | 490 | 469 | 488 | 306 | 368 | 305 | 366 | 305 |
| 733 | 500 | 977 | 521 | 976 | 490 | 611 | 490 | 488 | 488 | 490 | 488 |
| 1,464 | 1,776 | 1,960 | 980 | 1,960 | 1,500 | 1,464 | 980 | 1,000 | 976 | 976 | 624 |
| 46.7 | 27.7 | 41.4 | 23.9 | 61.5 | 32.6 | 51.3 | 43.0 | 7.3 | 16.5 | 3.9 | 11.8 |
| .4 | .3 | .6 | .4 | .5 | .2 | .3 | .2 | .2 | .3 | .2 | .3 |

TABLE 8.02.--FLUID MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 78.9 | 50.8 | 95.3 | 95.2 | 96.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 12.4 | 3.3 | 4.7 | 5.7 | 5.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 16.1 | 2.8 | 11.3 | 14.0 | 12.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 50.3 | 44.7 | 79.4 | 75.4 | 79.5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 4.6 | 9.5 | 6.9 | 5.7 | 5.8 |
| STANDARD DEVIATION.....DO..... | 3.0 | 4.5 | 3.5 | 2.7 | 2.7 |
| MAXIMUM.....DO..... | 36.0 | 26.0 | 24.0 | 17.0 | 16.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 309 | 558 | 409 | 372 | 434 |
| STANDARD DEVIATION.....GM..... | 259 | 338 | 265 | 212 | 236 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 31 | 61 | 61 | 81 | 82 |
| 25TH.....GM..... | 122 | 306 | 214 | 204 | 245 |
| 50TH.....GM..... | 244 | 539 | 366 | 346 | 407 |
| 75TH.....GM..... | 434 | 777 | 561 | 498 | 570 |
| 90TH.....GM..... | 651 | 976 | 763 | 651 | 733 |
| 95TH.....GM..... | 794 | 1,061 | 885 | 753 | 876 |
| 99TH.....GM..... | 1,160 | 1,466 | 1,184 | 940 | 1,058 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 243 | 247 | 175 | 271 | 284 |
| STANDARD DEVIATION.....GM..... | 172 | 280 | 174 | 201 | 181 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 284 | 219 | 269 | 277 | 323 |
| STANDARD DEVIATION.....GM..... | 186 | 231 | 157 | 149 | 155 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 403 | 618 | 463 | 428 | 490 |
| STANDARD DEVIATION.....GM..... | 270 | 311 | 254 | 197 | 219 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 4,410 | 2,083 | 1,960 | 1,960 | 2,318 |
| MAXIMUM OVER 3 DAYS.....GM..... | 9,053 | 5,856 | 5,856 | 4,087 | 4,911 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 201 | 176 | 179 | 195 | 224 |
| STANDARD DEVIATION.....GM..... | 133 | 78 | 79 | 82 | 89 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 15 | 61 | 61 | 61 | 122 |
| 25TH.....GM..... | 122 | 122 | 122 | 122 | 183 |
| 50TH.....GM..... | 244 | 183 | 183 | 184 | 244 |
| 75TH.....GM..... | 244 | 244 | 244 | 244 | 244 |
| 90TH.....GM..... | 366 | 245 | 245 | 245 | 245 |
| 95TH.....GM..... | 488 | 245 | 245 | 305 | 368 |
| 99TH.....GM..... | 550 | 337 | 429 | 488 | 488 |
| MAXIMUM.....GM..... | 1,960 | 488 | 976 | 1,069 | 1,220 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 406.1 | 3.4 | 20.9 | 27.9 | 39.3 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | .1 | .3 | .3 | .4 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 96.2 | 93.0 | 91.0 | 85.2 | 76.6 | 74.1 | 71.5 | 68.5 | 76.6 | 74.1 | 77.5 | 79.0 |
| 7.6 | 7.6 | 9.3 | 14.7 | 15.2 | 16.5 | 14.2 | 15.7 | 11.1 | 12.0 | 11.1 | 10.0 |
| 14.7 | 18.1 | 15.0 | 20.5 | 18.6 | 19.1 | 16.3 | 15.7 | 14.7 | 14.7 | 14.1 | 14.0 |
| 73.9 | 67.4 | 66.7 | 50.0 | 42.7 | 38.4 | 40.9 | 37.1 | 50.8 | 47.3 | 52.3 | 55.1 |
| 5.5 | 5.0 | 5.1 | 4.2 | 4.0 | 3.8 | 4.2 | 4.0 | 4.4 | 4.1 | 4.6 | 4.3 |
| 2.6 | 2.5 | 2.7 | 2.5 | 2.8 | 2.7 | 3.0 | 2.9 | 2.8 | 2.6 | 2.9 | 2.6 |
| 14.0 | 14.0 | 23.0 | 17.0 | 26.0 | 29.0 | 21.0 | 26.0 | 17.0 | 23.0 | 14.0 | 36.0 |
| 472 | 396 | 523 | 333 | 340 | 231 | 249 | 186 | 250 | 198 | 262 | 223 |
| 293 | 232 | 365 | 243 | 301 | 199 | 235 | 171 | 213 | 159 | 246 | 164 |
| 81 | 81 | 81 | 51 | 29 | 20 | 20 | 15 | 25 | 20 | 24 | 20 |
| 245 | 224 | 245 | 163 | 122 | 81 | 82 | 66 | 86 | 81 | 92 | 92 |
| 427 | 366 | 449 | 285 | 244 | 166 | 171 | 142 | 203 | 163 | 166 | 183 |
| 630 | 530 | 711 | 447 | 488 | 325 | 327 | 245 | 336 | 265 | 369 | 325 |
| 854 | 711 | 978 | 651 | 735 | 489 | 534 | 407 | 510 | 417 | 570 | 452 |
| 1,017 | 814 | 1,198 | 814 | 916 | 592 | 693 | 528 | 691 | 528 | 753 | 564 |
| 1,423 | 1,098 | 1,872 | 1,139 | 1,423 | 976 | 1,060 | 740 | 936 | 732 | 1,222 | 733 |
| 329 | 284 | 338 | 288 | 292 | 230 | 250 | 203 | 234 | 188 | 197 | 169 |
| 206 | 173 | 233 | 185 | 189 | 148 | 182 | 144 | 153 | 123 | 108 | 125 |
| 390 | 341 | 459 | 321 | 342 | 263 | 269 | 212 | 232 | 220 | 260 | 231 |
| 199 | 173 | 301 | 155 | 226 | 156 | 182 | 140 | 163 | 123 | 193 | 131 |
| 551 | 476 | 629 | 452 | 475 | 325 | 335 | 254 | 315 | 249 | 327 | 271 |
| 281 | 215 | 356 | 242 | 326 | 221 | 265 | 194 | 227 | 169 | 265 | 167 |
| 3,172 | 1,952 | 4,157 | 1,960 | 4,410 | 1,952 | 2,928 | 2,713 | 2,806 | 1,220 | 2,196 | 996 |
| 6,710 | 4,880 | 9,053 | 5,124 | 7,381 | 4,404 | 6,019 | 6,159 | 7,015 | 2,724 | 4,392 | 2,952 |
| 260 | 238 | 310 | 241 | 252 | 181 | 177 | 140 | 170 | 144 | 172 | 154 |
| 122 | 102 | 161 | 114 | 180 | 130 | 151 | 118 | 125 | 110 | 123 | 106 |
| 122 | 122 | 122 | 61 | 15 | 10 | 10 | 5 | 10 | 5 | 10 | 10 |
| 244 | 183 | 244 | 244 | 122 | 61 | 31 | 31 | 61 | 31 | 61 | 61 |
| 244 | 244 | 244 | 244 | 244 | 244 | 183 | 122 | 183 | 123 | 183 | 123 |
| 245 | 244 | 368 | 245 | 366 | 244 | 245 | 244 | 244 | 244 | 244 | 244 |
| 488 | 366 | 488 | 366 | 488 | 306 | 366 | 245 | 305 | 245 | 305 | 245 |
| 488 | 488 | 490 | 488 | 490 | 427 | 488 | 305 | 366 | 305 | 366 | 275 |
| 732 | 490 | 976 | 549 | 976 | 490 | 610 | 490 | 488 | 488 | 488 | 488 |
| 1,464 | 1,776 | 1,960 | 980 | 1,960 | 1,220 | 1,464 | 980 | 976 | 976 | 976 | 490 |
| 38.7 | 26.9 | 36.5 | 20.7 | 53.7 | 26.2 | 43.5 | 35.2 | 5.3 | 12.1 | 3.9 | 11.8 |
| .4 | .3 | .6 | .4 | .4 | .2 | .3 | .2 | .1 | .2 | .2 | .3 |

TABLE 8.03.--FLUID WHOLE MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 64.6 | 37.4 | 76.0 | 79.1 | 86.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 13.6 | 3.1 | 9.6 | 9.5 | 9.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 13.9 | 3.2 | 10.1 | 13.2 | 13.9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 37.1 | 31.1 | 56.3 | 56.4 | 63.2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 4.3 | 9.1 | 6.2 | 5.2 | 5.2 |
| STANDARD DEVIATION.....DO..... | 3.0 | 4.7 | 3.7 | 2.8 | 2.8 |
| MAXIMUM.....DO..... | 36.0 | 24.0 | 24.0 | 16.0 | 16.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 282 | 542 | 375 | 341 | 391 |
| STANDARD DEVIATION.....GM..... | 250 | 350 | 266 | 213 | 239 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 20 | 51 | 41 | 44 | 81 |
| 25TH.....GM..... | 81 | 244 | 164 | 163 | 203 |
| 50TH.....GM..... | 214 | 529 | 325 | 316 | 366 |
| 75TH.....GM..... | 407 | 777 | 529 | 488 | 529 |
| 90TH.....GM..... | 611 | 956 | 733 | 650 | 732 |
| 95TH.....GM..... | 752 | 1,138 | 864 | 733 | 854 |
| 99TH.....GM..... | 1,100 | 1,465 | 1,058 | 915 | 1,056 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 243 | 225 | 211 | 288 | 289 |
| STANDARD DEVIATION.....GM..... | 187 | 214 | 184 | 241 | 198 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 284 | 232 | 290 | 272 | 318 |
| STANDARD DEVIATION.....GM..... | 196 | 217 | 169 | 147 | 156 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 390 | 629 | 460 | 421 | 473 |
| STANDARD DEVIATION.....GM..... | 267 | 316 | 252 | 195 | 220 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 3,294 | 1,952 | 1,952 | 1,830 | 2,318 |
| MAXIMUM OVER 3 DAYS.....GM..... | 7,930 | 5,856 | 5,856 | 4,087 | 4,911 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 198 | 178 | 180 | 197 | 226 |
| STANDARD DEVIATION.....GM..... | 135 | 79 | 80 | 83 | 90 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 15 | 46 | 61 | 61 | 122 |
| 25TH.....GM..... | 122 | 122 | 122 | 122 | 183 |
| 50TH.....GM..... | 244 | 183 | 183 | 183 | 244 |
| 75TH.....GM..... | 244 | 244 | 244 | 244 | 244 |
| 90TH.....GM..... | 366 | 244 | 244 | 244 | 244 |
| 95TH.....GM..... | 488 | 244 | 244 | 305 | 427 |
| 99TH.....GM..... | 529 | 366 | 488 | 488 | 488 |
| MAXIMUM.....GM..... | 1,952 | 488 | 976 | 1,069 | 1,220 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 313.9 | 2.6 | 17.9 | 21.6 | 32.7 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | .2 | .4 | .3 | .4 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 87.9 | 82.7 | 80.1 | 71.7 | 65.0 | 57.8 | 57.7 | 51.9 | 59.2 | 54.6 | 66.9 | 63.1 |
| 10.7 | 10.9 | 12.7 | 16.7 | 16.9 | 16.1 | 14.3 | 15.3 | 12.0 | 11.7 | 10.7 | 10.4 |
| 17.0 | 18.1 | 16.8 | 16.9 | 16.5 | 15.0 | 13.4 | 11.8 | 10.6 | 10.7 | 12.9 | 12.3 |
| 60.1 | 53.7 | 50.6 | 38.1 | 31.6 | 26.7 | 30.0 | 24.8 | 36.7 | 32.3 | 43.3 | 40.3 |
| 4.8 | 4.5 | 4.5 | 3.8 | 3.7 | 3.7 | 4.0 | 3.7 | 4.3 | 3.9 | 4.6 | 4.2 |
| 2.7 | 2.5 | 2.7 | 2.5 | 2.8 | 2.8 | 3.1 | 2.9 | 2.9 | 2.7 | 3.0 | 2.8 |
| 14.0 | 14.0 | 23.0 | 14.0 | 26.0 | 29.0 | 21.0 | 26.0 | 17.0 | 23.0 | 14.0 | 36.0 |
| 416 | 355 | 450 | 303 | 305 | 208 | 221 | 162 | 235 | 173 | 257 | 212 |
| 287 | 228 | 340 | 238 | 288 | 192 | 215 | 162 | 225 | 153 | 254 | 169 |
| 81 | 81 | 81 | 41 | 20 | 10 | 12 | 10 | 15 | 12 | 20 | 20 |
| 203 | 163 | 203 | 122 | 97 | 81 | 81 | 46 | 81 | 70 | 81 | 81 |
| 356 | 325 | 402 | 244 | 224 | 163 | 163 | 117 | 163 | 122 | 168 | 164 |
| 569 | 488 | 611 | 407 | 407 | 285 | 300 | 225 | 325 | 244 | 325 | 305 |
| 773 | 651 | 855 | 611 | 671 | 454 | 488 | 366 | 528 | 407 | 570 | 452 |
| 936 | 773 | 1,057 | 744 | 874 | 570 | 650 | 488 | 692 | 488 | 772 | 569 |
| 1,343 | 1,017 | 1,709 | 1,138 | 1,423 | 976 | 976 | 712 | 977 | 718 | 1,464 | 733 |
| 322 | 290 | 354 | 272 | 288 | 228 | 251 | 190 | 204 | 176 | 187 | 159 |
| 189 | 183 | 229 | 173 | 209 | 157 | 217 | 146 | 148 | 125 | 111 | 122 |
| 363 | 329 | 442 | 311 | 344 | 251 | 263 | 203 | 218 | 219 | 269 | 222 |
| 193 | 169 | 336 | 157 | 230 | 158 | 196 | 151 | 179 | 128 | 201 | 132 |
| 521 | 454 | 584 | 438 | 455 | 311 | 306 | 235 | 315 | 224 | 329 | 273 |
| 280 | 214 | 333 | 242 | 327 | 224 | 249 | 192 | 243 | 171 | 278 | 174 |
| 3,172 | 1,952 | 3,294 | 1,952 | 3,172 | 1,952 | 2,928 | 2,713 | 2,806 | 1,220 | 2,196 | 996 |
| 6,710 | 4,880 | 7,930 | 5,124 | 7,381 | 4,209 | 6,019 | 6,159 | 7,015 | 2,684 | 4,392 | 2,952 |
| 258 | 238 | 303 | 239 | 244 | 171 | 166 | 129 | 165 | 133 | 169 | 153 |
| 120 | 100 | 155 | 115 | 181 | 130 | 152 | 119 | 131 | 114 | 128 | 109 |
| 122 | 122 | 122 | 31 | 15 | 10 | 8 | 5 | 8 | 5 | 5 | 10 |
| 244 | 183 | 244 | 198 | 122 | 31 | 31 | 29 | 31 | 31 | 61 | 61 |
| 244 | 244 | 244 | 244 | 244 | 183 | 122 | 93 | 153 | 122 | 153 | 122 |
| 244 | 244 | 366 | 244 | 305 | 244 | 244 | 244 | 244 | 244 | 244 | 244 |
| 488 | 366 | 488 | 366 | 488 | 305 | 366 | 244 | 305 | 244 | 305 | 244 |
| 488 | 488 | 488 | 488 | 488 | 366 | 488 | 305 | 366 | 305 | 366 | 305 |
| 732 | 488 | 976 | 549 | 976 | 488 | 610 | 488 | 488 | 488 | 488 | 488 |
| 1,464 | 1,769 | 1,464 | 976 | 1,952 | 1,220 | 1,464 | 976 | 976 | 976 | 976 | 488 |
| 34.0 | 22.6 | 20.5 | 12.3 | 47.5 | 19.9 | 33.0 | 27.9 | .9 | 12.1 | 3.9 | 4.6 |
| .4 | .3 | .4 | .3 | .5 | .2 | .3 | .2 | (2) | .3 | .2 | .2 |

TABLE 8.04.--FLUID LOWFAT MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 17.3 | 14.0 | 24.2 | 21.7 | 21.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.2 | 1.8 | 3.4 | 2.4 | 2.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 4.5 | 1.1 | 3.1 | 5.3 | 5.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 8.6 | 11.0 | 17.7 | 14.1 | 13.6 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 3.9 | 8.8 | 6.3 | 5.2 | 4.6 |
| STANDARD DEVIATION.....DO..... | 2.6 | 4.3 | 3.7 | 2.7 | 2.4 |
| MAXIMUM.....DO..... | 26.0 | 26.0 | 21.0 | 13.0 | 12.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 279 | 513 | 365 | 321 | 328 |
| STANDARD DEVIATION.....GM..... | 237 | 332 | 274 | 203 | 200 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 41 | 31 | 41 | 61 | 62 |
| 25TH.....GM..... | 91 | 245 | 163 | 163 | 163 |
| 50TH.....GM..... | 223 | 489 | 327 | 286 | 306 |
| 75TH.....GM..... | 388 | 736 | 490 | 449 | 459 |
| 90TH.....GM..... | 581 | 980 | 674 | 613 | 572 |
| 95TH.....GM..... | 735 | 1,030 | 858 | 695 | 673 |
| 99TH.....GM..... | 1,062 | 1,836 | 1,472 | 858 | 959 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 263 | 338 | 215 | 253 | 320 |
| STANDARD DEVIATION.....GM..... | 187 | 329 | 176 | 142 | 204 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 311 | 478 | 307 | 322 | 333 |
| STANDARD DEVIATION.....GM..... | 211 | 163 | 223 | 220 | 188 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 409 | 600 | 448 | 400 | 413 |
| STANDARD DEVIATION.....GM..... | 251 | 313 | 265 | 188 | 187 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 4,410 | 2,083 | 1,960 | 1,960 | 1,348 |
| MAXIMUM OVER 3 DAYS.....GM..... | 6,125 | 5,513 | 5,145 | 3,063 | 3,430 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 217 | 175 | 173 | 185 | 213 |
| STANDARD DEVIATION.....GM..... | 130 | 76 | 78 | 79 | 85 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 25 | 61 | 61 | 61 | 92 |
| 25TH.....GM..... | 123 | 123 | 123 | 123 | 123 |
| 50TH.....GM..... | 245 | 184 | 153 | 184 | 245 |
| 75TH.....GM..... | 245 | 245 | 245 | 245 | 245 |
| 90TH.....GM..... | 368 | 245 | 245 | 245 | 245 |
| 95TH.....GM..... | 490 | 245 | 245 | 306 | 368 |
| 99TH.....GM..... | 613 | 276 | 368 | 490 | 490 |
| MAXIMUM.....GM..... | 1,960 | 368 | 735 | 613 | 551 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 68.5 | .8 | 3.0 | 6.3 | 6.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | .1 | .2 | .3 | .4 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 19.4 | 19.6 | 19.7 | 17.6 | 16.1 | 17.3 | 15.9 | 15.3 | 16.9 | 17.5 | 10.6 | 14.2 |
| 3.7 | 4.1 | 2.9 | 4.4 | 4.2 | 5.7 | 4.4 | 5.0 | 3.5 | 4.3 | 1.9 | 3.3 |
| 3.9 | 5.0 | 4.7 | 4.7 | 4.7 | 4.9 | 4.8 | 4.3 | 4.2 | 4.2 | 2.2 | 2.2 |
| 11.8 | 10.6 | 12.1 | 8.5 | 7.1 | 6.8 | 6.6 | 5.9 | 9.2 | 8.9 | 6.5 | 8.6 |
| 4.4 | 4.0 | 4.5 | 3.7 | 3.4 | 3.1 | 3.4 | 3.2 | 3.7 | 3.6 | 3.7 | 3.8 |
| 2.3 | 2.2 | 2.3 | 2.3 | 2.3 | 2.2 | 2.5 | 2.4 | 2.4 | 2.4 | 2.4 | 2.1 |
| 11.0 | 11.0 | 13.0 | 16.0 | 13.0 | 13.0 | 17.0 | 17.0 | 12.0 | 14.0 | 9.0 | 10.0 |
| 384 | 311 | 502 | 305 | 322 | 223 | 249 | 181 | 229 | 207 | 230 | 199 |
| 262 | 204 | 354 | 209 | 274 | 186 | 250 | 166 | 161 | 162 | 194 | 153 |
| 82 | 54 | 82 | 61 | 41 | 31 | 31 | 20 | 41 | 27 | 41 | 30 |
| 174 | 163 | 245 | 163 | 123 | 82 | 82 | 61 | 102 | 82 | 82 | 82 |
| 327 | 265 | 408 | 245 | 245 | 163 | 163 | 123 | 201 | 163 | 136 | 163 |
| 510 | 449 | 613 | 408 | 470 | 327 | 327 | 245 | 277 | 306 | 368 | 286 |
| 735 | 593 | 1,041 | 571 | 695 | 490 | 541 | 408 | 449 | 439 | 530 | 412 |
| 878 | 695 | 1,225 | 654 | 857 | 602 | 736 | 530 | 573 | 490 | 673 | 490 |
| 1,144 | 939 | 1,878 | 1,061 | 1,224 | 899 | 1,328 | 762 | 775 | 826 | 736 | 660 |
| 375 | 339 | 465 | 354 | 330 | 238 | 245 | 205 | 240 | 184 | 196 | 202 |
| 245 | 241 | 367 | 217 | 215 | 146 | 153 | 116 | 101 | 110 | 59 | 120 |
| 414 | 336 | 529 | 359 | 408 | 295 | 281 | 225 | 247 | 235 | 141 | 184 |
| 226 | 199 | 279 | 144 | 291 | 168 | 194 | 142 | 93 | 167 | 77 | 96 |
| 500 | 428 | 641 | 437 | 480 | 363 | 408 | 299 | 315 | 300 | 322 | 270 |
| 255 | 184 | 360 | 210 | 296 | 199 | 299 | 195 | 166 | 157 | 194 | 153 |
| 2,573 | 1,776 | 2,940 | 1,960 | 4,410 | 1,409 | 2,029 | 1,470 | 1,103 | 1,118 | 919 | 980 |
| 5,513 | 3,185 | 6,125 | 4,043 | 6,003 | 2,940 | 5,189 | 3,430 | 2,328 | 2,618 | 2,205 | 2,328 |
| 264 | 236 | 336 | 247 | 284 | 214 | 218 | 169 | 186 | 174 | 187 | 159 |
| 120 | 109 | 183 | 108 | 179 | 125 | 145 | 111 | 103 | 97 | 83 | 101 |
| 123 | 92 | 123 | 77 | 31 | 15 | 15 | 10 | 15 | 15 | 31 | 10 |
| 245 | 184 | 245 | 245 | 245 | 123 | 123 | 61 | 123 | 123 | 123 | 92 |
| 245 | 245 | 245 | 245 | 245 | 245 | 245 | 184 | 245 | 184 | 184 | 123 |
| 245 | 245 | 490 | 245 | 368 | 245 | 245 | 245 | 245 | 245 | 245 | 245 |
| 490 | 368 | 490 | 368 | 490 | 368 | 368 | 245 | 277 | 245 | 245 | 245 |
| 490 | 490 | 735 | 490 | 490 | 490 | 490 | 368 | 368 | 306 | 260 | 245 |
| 735 | 490 | 980 | 520 | 980 | 552 | 735 | 490 | 490 | 368 | 490 | 490 |
| 980 | 1,776 | 1,960 | 980 | 1,960 | 919 | 980 | 613 | 490 | 490 | 490 | 490 |
| 4.7 | 4.3 | 8.5 | 4.2 | 4.3 | 5.2 | 9.6 | 3.8 | 4.4 | 0 | 0 | 2.6 |
| .3 | .3 | .7 | .4 | .2 | .2 | .4 | .1 | .6 | 0 | 0 | .5 |

TABLE 8.05.--FLUID SKIM MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.6 | 2.3 | 4.2 | 4.0 | 3.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.7 | 0 | .6 | .7 | .7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.4 | .9 | .8 | .8 | .8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.6 | 1.4 | 2.8 | 2.5 | 2.4 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 3.6 | 8.0 | 6.3 | 5.4 | 4.4 |
| STANDARD DEVIATION.....DO..... | 2.6 | 5.1 | 3.3 | 2.9 | 2.4 |
| MAXIMUM.....DO..... | 21.0 | 21.0 | 13.0 | 13.0 | 9.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 226 | 380 | 374 | 336 | 309 |
| STANDARD DEVIATION.....GM..... | 201 | 230 | 245 | 206 | 189 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 20 | | | 41 | 41 |
| 25TH.....GM..... | 82 | | | 163 | 163 |
| 50TH.....GM..... | 163 | | | 327 | 286 |
| 75TH.....GM..... | 327 | | | 464 | 449 |
| 90TH.....GM..... | 490 | | | 592 | 531 |
| 95TH.....GM..... | 623 | | | 653 | 694 |
| 99TH.....GM..... | 899 | | | 1,061 | 817 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 250 | 0 | 213 | 241 | 220 |
| STANDARD DEVIATION.....GM..... | 194 | 0 | 108 | 102 | 90 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 278 | 260 | 297 | 399 | 378 |
| STANDARD DEVIATION.....GM..... | 169 | 222 | 150 | 153 | 152 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 341 | 516 | 496 | 431 | 393 |
| STANDARD DEVIATION.....GM..... | 228 | 164 | 208 | 189 | 174 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 2,940 | 827 | 1,103 | 1,225 | 980 |
| MAXIMUM OVER 3 DAYS.....GM..... | 4,410 | 2,435 | 2,818 | 3,185 | 2,450 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 190 | 143 | 179 | 186 | 209 |
| STANDARD DEVIATION.....GM..... | 123 | 57 | 75 | 76 | 82 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 15 | | | 61 | 61 |
| 25TH.....GM..... | 123 | | | 123 | 123 |
| 50TH.....GM..... | 245 | | | 184 | 245 |
| 75TH.....GM..... | 245 | | | 245 | 245 |
| 90TH.....GM..... | 306 | | | 245 | 245 |
| 95TH.....GM..... | 368 | | | 245 | 368 |
| 99TH.....GM..... | 490 | | | 490 | 490 |
| MAXIMUM.....GM..... | 980 | 260 | 735 | 490 | 613 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 20.7 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.5 | 4.3 | 4.3 | 5.3 | 3.4 | 6.2 | 5.5 | 8.2 | 6.3 | 9.1 | 2.6 | 7.4 |
| .8 | 1.2 | 1.4 | 1.7 | 1.2 | 2.2 | 2.2 | 2.3 | 1.3 | 2.6 | 0 | 1.8 |
| .8 | 1.1 | .9 | 1.5 | .8 | 1.5 | 1.3 | 2.3 | 1.5 | 2.0 | .6 | 1.6 |
| 1.9 | 2.0 | 2.1 | 2.1 | 1.4 | 2.4 | 2.1 | 3.6 | 3.5 | 4.5 | 2.0 | 4.0 |
| 4.1 | 3.9 | 3.7 | 3.1 | 3.5 | 2.9 | 3.1 | 3.5 | 3.8 | 3.4 | 4.3 | 3.7 |
| 2.4 | 2.3 | 2.2 | 2.1 | 2.3 | 2.2 | 2.5 | 2.6 | 2.6 | 2.2 | 1.8 | 2.2 |
| 9.0 | 9.0 | 10.0 | 10.0 | 9.0 | 12.0 | 13.0 | 14.0 | 9.0 | 10.0 | 8.0 | 9.0 |
| 391 | 304 | 374 | 244 | 304 | 188 | 190 | 179 | 213 | 174 | 225 | 191 |
| 312 | 207 | 235 | 209 | 293 | 165 | 181 | 154 | 182 | 137 | 144 | 144 |
| 61 | 41 | 44 | 41 | 20 | 20 | 41 | 20 | 10 | 20 | | 20 |
| 123 | 123 | 123 | 82 | 85 | 82 | 82 | 71 | 61 | 82 | | 82 |
| 368 | 286 | 408 | 184 | 204 | 163 | 123 | 123 | 163 | 134 | | 143 |
| 552 | 449 | 572 | 327 | 408 | 245 | 245 | 245 | 286 | 245 | | 255 |
| 837 | 531 | 694 | 510 | 736 | 398 | 408 | 408 | 490 | 378 | | 408 |
| 981 | 694 | 775 | 735 | 899 | 530 | 530 | 490 | 612 | 408 | | 449 |
| 1,448 | 919 | 873 | 981 | 1,186 | 715 | 837 | 735 | 877 | 572 | | 653 |
| 246 | 311 | 476 | 262 | 330 | 240 | 256 | 225 | 227 | 188 | 0 | 163 |
| 70 | 222 | 371 | 149 | 307 | 135 | 187 | 174 | 255 | 123 | 0 | 72 |
| 473 | 389 | 519 | 274 | 305 | 259 | 261 | 238 | 250 | 218 | 236 | 216 |
| 255 | 198 | 391 | 120 | 132 | 109 | 150 | 144 | 139 | 110 | 14 | 111 |
| 565 | 448 | 533 | 409 | 530 | 297 | 313 | 259 | 286 | 251 | 244 | 273 |
| 308 | 183 | 141 | 233 | 328 | 203 | 225 | 180 | 201 | 145 | 158 | 141 |
| 1,715 | 1,348 | 1,470 | 1,470 | 2,940 | 1,470 | 1,470 | 1,235 | 980 | 746 | 735 | 735 |
| 4,349 | 3,185 | 2,622 | 2,940 | 4,410 | 4,165 | 3,001 | 2,450 | 2,634 | 2,205 | 1,470 | 1,960 |
| 288 | 234 | 300 | 238 | 263 | 193 | 186 | 152 | 169 | 155 | 156 | 154 |
| 156 | 108 | 159 | 108 | 155 | 119 | 122 | 115 | 110 | 92 | 105 | 86 |
| 123 | 123 | 123 | 61 | 31 | 15 | 15 | 10 | 10 | 15 | | 15 |
| 245 | 184 | 245 | 214 | 184 | 123 | 123 | 31 | 61 | 61 | | 93 |
| 245 | 245 | 245 | 245 | 245 | 245 | 184 | 123 | 184 | 123 | | 123 |
| 368 | 245 | 368 | 245 | 368 | 245 | 245 | 245 | 245 | 245 | | 245 |
| 490 | 368 | 490 | 368 | 490 | 306 | 306 | 245 | 276 | 245 | | 245 |
| 490 | 490 | 490 | 490 | 490 | 368 | 368 | 306 | 306 | 245 | | 245 |
| 980 | 490 | 980 | 490 | 736 | 490 | 490 | 490 | 490 | 368 | | 274 |
| 980 | 980 | 980 | 980 | 980 | 919 | 980 | 980 | 551 | 490 | 490 | 306 |
| 0 | 0 | 7.6 | 4.3 | 1.8 | 0 | .9 | 1.6 | 0 | 0 | 0 | 4.5 |
| 0 | 0 | 3.4 | 1.8 | .4 | 0 | .1 | .1 | 0 | 0 | 0 | 1.7 |

TABLE 8.06.--RECONSTITUTED DRY MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 1.9 | 1.9 | 2.0 | 2.3 | 2.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | .6 | 0 | .4 | .4 | .4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .5 | .4 | .7 | .9 | .5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .8 | 1.5 | 1.0 | .9 | 1.2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 3.4 | 9.7 | 5.0 | 4.0 | 3.8 |
| STANDARD DEVIATION.....DO..... | 2.5 | 5.0 | 3.2 | 2.6 | 2.1 |
| MAXIMUM.....DO..... | 22.0 | 22.0 | 15.0 | 9.0 | 9.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 206 | 515 | 262 | 234 | 223 |
| STANDARD DEVIATION.....GM..... | 198 | 302 | 247 | 185 | 149 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 14 | | | | |
| 25TH.....GM..... | 68 | | | | |
| 50TH.....GM..... | 163 | | | | |
| 75TH.....GM..... | 265 | | | | |
| 90TH.....GM..... | 449 | | | | |
| 95TH.....GM..... | 600 | | | | |
| 99TH.....GM..... | 981 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 191 | 0 | 213 | 197 | 166 |
| STANDARD DEVIATION.....GM..... | 162 | 0 | 205 | 114 | 62 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 258 | 364 | 269 | 271 | 348 |
| STANDARD DEVIATION.....GM..... | 147 | 30 | 181 | 143 | 183 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 322 | 583 | 389 | 363 | 283 |
| STANDARD DEVIATION.....GM..... | 233 | 301 | 282 | 203 | 138 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,960 | 1,164 | 1,348 | 1,225 | 919 |
| MAXIMUM OVER 3 DAYS.....GM..... | 3,675 | 3,063 | 3,308 | 2,205 | 2,083 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 183 | 160 | 158 | 177 | 179 |
| STANDARD DEVIATION.....GM..... | 128 | 70 | 84 | 94 | 75 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 15 | | | | |
| 25TH.....GM..... | 123 | | | | |
| 50TH.....GM..... | 184 | | | | |
| 75TH.....GM..... | 245 | | | | |
| 90TH.....GM..... | 306 | | | | |
| 95TH.....GM..... | 368 | | | | |
| 99TH.....GM..... | 735 | | | | |
| MAXIMUM.....GM..... | 980 | 245 | 368 | 490 | 490 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 4.7 | 0 | 1.0 | 1.4 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .2 | 0 | 1.0 | .9 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.3 | 1.4 | 1.2 | 1.2 | 1.5 | 1.6 | 1.7 | 2.3 | 3.5 | 3.5 | 2.3 | 3.3 |
| .4 | .5 | .2 | .3 | .5 | .5 | .5 | .7 | .8 | 1.1 | .4 | 1.3 |
| .6 | .5 | .3 | .3 | .3 | .6 | .4 | .6 | .5 | .8 | .6 | .4 |
| .3 | .4 | .7 | .6 | .6 | .5 | .7 | 1.0 | 2.1 | 1.6 | 1.2 | 1.5 |
| 2.9 | 3.2 | 4.1 | 3.1 | 2.9 | 2.9 | 3.3 | 3.1 | 4.2 | 3.2 | 2.8 | 2.9 |
| 2.0 | 2.4 | 1.8 | 1.6 | 1.8 | 2.4 | 2.3 | 2.1 | 3.2 | 1.9 | 1.3 | 1.8 |
| 9.0 | 10.0 | 6.0 | 7.0 | 8.0 | 12.0 | 9.0 | 10.0 | 12.0 | 7.0 | 6.0 | 8.0 |
| 192 | 206 | 358 | 261 | 277 | 154 | 220 | 163 | 263 | 169 | 165 | 128 |
| 159 | 156 | 255 | 244 | 307 | 129 | 245 | 136 | 200 | 150 | 57 | 114 |
| | | | | 3 | 5 | 3 | 10 | | 10 | | |
| | | | | 61 | 61 | 56 | 41 | | 41 | | |
| | | | | 184 | 113 | 163 | 123 | | 143 | | |
| | | | | 408 | 245 | 245 | 245 | | 245 | | |
| | | | | 898 | 327 | 490 | 368 | | 432 | | |
| | | | | 980 | 416 | 817 | 425 | | 459 | | |
| | | | | 1,225 | 571 | 1,199 | 602 | | 613 | | |
| 310 | 243 | 329 | 295 | 192 | 161 | 221 | 134 | 121 | 226 | 261 | 219 |
| 274 | 229 | 116 | 99 | 179 | 97 | 165 | 111 | 113 | 188 | 121 | 167 |
| 243 | 292 | 360 | 281 | 320 | 219 | 264 | 229 | 469 | 183 | 206 | 151 |
| 128 | 142 | 114 | 127 | 146 | 124 | 126 | 114 | 282 | 86 | 57 | 92 |
| 388 | 359 | 466 | 389 | 497 | 274 | 344 | 256 | 338 | 257 | 205 | 182 |
| 193 | 143 | 268 | 296 | 355 | 145 | 315 | 140 | 174 | 174 | 29 | 134 |
| 858 | 854 | 1,103 | 1,715 | 1,960 | 735 | 1,960 | 827 | 980 | 980 | 368 | 551 |
| 2,083 | 1,899 | 3,001 | 2,940 | 3,675 | 1,715 | 3,597 | 1,807 | 2,205 | 1,838 | 735 | 1,409 |
| 201 | 192 | 263 | 251 | 289 | 162 | 198 | 157 | 187 | 159 | 177 | 132 |
| 140 | 91 | 150 | 170 | 222 | 105 | 160 | 101 | 106 | 112 | 56 | 82 |
| | | | | 31 | 8 | 10 | 15 | | 5 | | |
| | | | | 123 | 93 | 61 | 61 | | 61 | | |
| | | | | 245 | 152 | 245 | 164 | | 123 | | |
| | | | | 337 | 245 | 245 | 245 | | 245 | | |
| | | | | 490 | 274 | 306 | 245 | | 245 | | |
| | | | | 736 | 306 | 490 | 306 | | 306 | | |
| | | | | 980 | 490 | 980 | 490 | | 490 | | |
| 735 | 732 | 735 | 980 | 980 | 490 | 980 | 490 | 490 | 490 | 306 | 245 |
| 0 | 0 | 0 | 0 | 0 | 0 | 2.3 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | .8 | 0 | 0 | 0 | 0 | 0 |

TABLE 8.07.--CHOCOLATE MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.5 | .7 | 4.6 | 9.5 | 15.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.4 | .7 | 3.5 | 6.2 | 8.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.4 | 0 | .7 | 2.3 | 4.4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .7 | 0 | .4 | .9 | 2.4 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.6 | 1.0 | 1.6 | 1.6 | 1.7 |
| STANDARD DEVIATION.....DO..... | 1.0 | .0 | 1.3 | 1.0 | 1.0 |
| MAXIMUM.....DO..... | 10.0 | 1.0 | 9.0 | 6.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 159 | 53 | 98 | 119 | 152 |
| STANDARD DEVIATION.....GM..... | 128 | 18 | 64 | 83 | 99 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 63 | | | 42 | 63 |
| 25TH.....GM..... | 83 | | | 63 | 83 |
| 50TH.....GM..... | 125 | | | 83 | 104 |
| 75TH.....GM..... | 167 | | | 146 | 167 |
| 90TH.....GM..... | 333 | | | 250 | 250 |
| 95TH.....GM..... | 365 | | | 333 | 333 |
| 99TH.....GM..... | 583 | | | 417 | 500 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 305 | 158 | 226 | 236 | 267 |
| STANDARD DEVIATION.....GM..... | 160 | 55 | 121 | 104 | 86 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 310 | 0 | 216 | 248 | 293 |
| STANDARD DEVIATION.....GM..... | 146 | 0 | 50 | 124 | 116 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 346 | 0 | 230 | 266 | 296 |
| STANDARD DEVIATION.....GM..... | 209 | 0 | 80 | 85 | 104 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 2,500 | 250 | 563 | 1,000 | 1,000 |
| MAXIMUM OVER 3 DAYS.....GM..... | 5,500 | 250 | 1,000 | 1,500 | 1,750 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 290 | 158 | 186 | 222 | 260 |
| STANDARD DEVIATION.....GM..... | 139 | 55 | 83 | 76 | 84 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 125 | | | 125 | 125 |
| 25TH.....GM..... | 250 | | | 188 | 250 |
| 50TH.....GM..... | 250 | | | 250 | 250 |
| 75TH.....GM..... | 250 | | | 250 | 250 |
| 90TH.....GM..... | 500 | | | 250 | 375 |
| 95TH.....GM..... | 500 | | | 313 | 500 |
| 99TH.....GM..... | 1,000 | | | 500 | 500 |
| MAXIMUM.....GM..... | 1,500 | 250 | 500 | 625 | 750 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 16.8 | 0 | 0 | .8 | 1.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .5 | 0 | 0 | .3 | .3 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 18.2 | 16.9 | 12.6 | 11.7 | 4.0 | 3.1 | 1.6 | .9 | .9 | .3 | .5 | .4 |
| 10.2 | 8.7 | 7.5 | 6.9 | 3.0 | 2.5 | 1.0 | .7 | .4 | .2 | .2 | .4 |
| 5.3 | 6.0 | 3.6 | 3.6 | .6 | .4 | .4 | .1 | .3 | .1 | 0 | 0 |
| 2.7 | 2.3 | 1.4 | 1.2 | .4 | .2 | .1 | .1 | .2 | 0 | .3 | 0 |
| 1.8 | 1.8 | 1.6 | 1.6 | 1.5 | 1.4 | 1.6 | 1.3 | 1.9 | 1.3 | 2.2 | 1.0 |
| 1.2 | 1.1 | .9 | .8 | .9 | .9 | 1.0 | .6 | .9 | .4 | 1.0 | .0 |
| 9.0 | 10.0 | 5.0 | 6.0 | 5.0 | 6.0 | 7.0 | 3.0 | 3.0 | 2.0 | 3.0 | 1.0 |
| 182 | 165 | 191 | 152 | 168 | 134 | 188 | 114 | 217 | 105 | 185 | 83 |
| 154 | 122 | 139 | 96 | 137 | 97 | 236 | 53 | 142 | 37 | 81 | 0 |
| 83 | 83 | 83 | 83 | 63 | 63 | 63 | 56 | | | | |
| 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | | | | |
| 167 | 167 | 167 | 125 | 156 | 83 | 125 | 83 | | | | |
| 250 | 171 | 250 | 167 | 167 | 167 | 167 | 167 | | | | |
| 333 | 333 | 333 | 250 | 333 | 250 | 333 | 167 | | | | |
| 500 | 333 | 417 | 333 | 333 | 333 | 500 | 250 | | | | |
| 958 | 500 | 646 | 458 | 833 | 500 | 1,833 | 250 | | | | |
| 321 | 291 | 390 | 295 | 369 | 314 | 320 | 290 | 379 | 250 | 250 | 250 |
| 165 | 135 | 219 | 109 | 209 | 184 | 161 | 103 | 311 | 0 | 0 | 0 |
| 329 | 296 | 389 | 308 | 338 | 314 | 346 | 216 | 299 | 250 | 0 | 0 |
| 155 | 109 | 253 | 117 | 108 | 134 | 135 | 49 | 61 | 0 | 0 | 0 |
| 393 | 338 | 333 | 301 | 425 | 335 | 656 | 250 | 437 | 0 | 250 | 0 |
| 242 | 198 | 137 | 128 | 234 | 123 | 577 | 0 | 0 | 0 | 0 | 0 |
| 1,750 | 2,250 | 2,000 | 750 | 1,500 | 1,500 | 2,500 | 500 | 1,000 | 250 | 250 | 250 |
| 3,750 | 4,688 | 4,000 | 2,250 | 2,500 | 1,875 | 5,500 | 750 | 1,313 | 500 | 750 | 250 |
| 306 | 281 | 352 | 282 | 343 | 290 | 356 | 268 | 348 | 250 | 250 | 250 |
| 156 | 114 | 185 | 97 | 166 | 150 | 202 | 90 | 176 | 0 | 0 | 0 |
| 125 | 188 | 250 | 125 | 188 | 188 | 188 | 125 | | | | |
| 250 | 250 | 250 | 250 | 250 | 250 | 250 | 250 | | | | |
| 250 | 250 | 250 | 250 | 250 | 250 | 250 | 250 | | | | |
| 250 | 250 | 500 | 250 | 500 | 250 | 500 | 250 | | | | |
| 500 | 500 | 500 | 500 | 500 | 500 | 500 | 375 | | | | |
| 500 | 500 | 626 | 500 | 500 | 500 | 1,000 | 500 | | | | |
| 1,000 | 750 | 1,000 | 500 | 1,000 | 751 | 1,000 | 500 | | | | |
| 1,125 | 1,000 | 1,250 | 500 | 1,000 | 1,500 | 1,000 | 500 | 1,000 | 250 | 250 | 250 |
| 5.5 | 0 | 0 | 3.1 | 0 | 2.2 | 0 | 1.6 | 2.0 | 0 | 0 | 0 |
| .8 | 0 | 0 | 1.1 | 0 | 1.0 | 0 | 1.9 | 10.2 | 0 | 0 | 0 |

TABLE 8.08.--COCOA AND HOT CHOCOLATE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.8 | 0 | 4.1 | 5.8 | 8.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.0 | 0 | 3.0 | 4.1 | 5.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.1 | 0 | 1.0 | .9 | 2.1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .7 | 0 | .2 | .7 | 1.2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.7 | 0 | 1.5 | 1.6 | 1.6 |
| STANDARD DEVIATION.....DO..... | 1.0 | 0 | .9 | 1.1 | .9 |
| MAXIMUM.....DO..... | 9.0 | 0 | 5.0 | 8.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 142 | 0 | 98 | 121 | 129 |
| STANDARD DEVIATION.....GM..... | 104 | 0 | 88 | 98 | 85 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 53 | | | 42 | 42 |
| 25TH.....GM..... | 83 | | | 63 | 83 |
| 50TH.....GM..... | 85 | | | 83 | 83 |
| 75TH.....GM..... | 169 | | | 167 | 167 |
| 90TH.....GM..... | 256 | | | 250 | 250 |
| 95TH.....GM..... | 333 | | | 256 | 292 |
| 99TH.....GM..... | 562 | | | 666 | 363 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 262 | 0 | 186 | 237 | 236 |
| STANDARD DEVIATION.....GM..... | 105 | 0 | 85 | 116 | 145 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 282 | 0 | 223 | 236 | 265 |
| STANDARD DEVIATION.....GM..... | 107 | 0 | 87 | 76 | 57 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 305 | 0 | 432 | 306 | 266 |
| STANDARD DEVIATION.....GM..... | 139 | 0 | 62 | 135 | 63 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,563 | 0 | 750 | 1,250 | 1,563 |
| MAXIMUM OVER 3 DAYS.....GM..... | 3,250 | 0 | 1,500 | 2,000 | 1,563 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 255 | 0 | 196 | 228 | 236 |
| STANDARD DEVIATION.....GM..... | 90 | 0 | 78 | 67 | 61 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 128 | | | 125 | 125 |
| 25TH.....GM..... | 250 | | | 188 | 219 |
| 50TH.....GM..... | 250 | | | 250 | 250 |
| 75TH.....GM..... | 256 | | | 250 | 250 |
| 90TH.....GM..... | 313 | | | 256 | 256 |
| 95TH.....GM..... | 500 | | | 275 | 256 |
| 99TH.....GM..... | 513 | | | 500 | 500 |
| MAXIMUM.....GM..... | 1,250 | 0 | 500 | 500 | 576 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 6.9 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .2 | 0 | 0 | 0 | 0 |

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 7.8 | 8.0 | 7.5 | 7.1 | 4.0 | 5.5 | 2.8 | 3.5 | 2.3 | 2.9 | 2.2 | 4.8 |
| 4.3 | 4.9 | 5.1 | 4.5 | 2.6 | 3.2 | 1.7 | 2.4 | .9 | 1.7 | 1.2 | 2.8 |
| 2.1 | 2.0 | 1.6 | 2.2 | .8 | 1.4 | .6 | .8 | .9 | .4 | .6 | 1.2 |
| 1.4 | 1.1 | .8 | .5 | .6 | .9 | .5 | .4 | .6 | .7 | .4 | .9 |
| 1.8 | 1.6 | 1.6 | 1.6 | 1.7 | 1.7 | 1.9 | 1.6 | 2.3 | 1.7 | 1.6 | 1.7 |
| 1.0 | 1.0 | .8 | 1.2 | 1.1 | .9 | 1.4 | 1.0 | 1.4 | 1.0 | .8 | .8 |
| 7.0 | 8.0 | 5.0 | 9.0 | 5.0 | 6.0 | 8.0 | 6.0 | 5.0 | 4.0 | 3.0 | 4.0 |
| 153 | 139 | 149 | 132 | 157 | 144 | 162 | 133 | 182 | 142 | 147 | 147 |
| 103 | 95 | 96 | 106 | 146 | 93 | 137 | 84 | 117 | 90 | 64 | 90 |
| 63 | 63 | 63 | 63 | 63 | 63 | 63 | 63 | | 64 | | |
| 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | | 83 | | |
| 128 | 85 | 104 | 83 | 92 | 92 | 92 | 85 | | 85 | | |
| 169 | 167 | 171 | 167 | 167 | 171 | 206 | 169 | | 167 | | |
| 256 | 250 | 333 | 171 | 254 | 254 | 275 | 256 | | 333 | | |
| 339 | 271 | 333 | 252 | 363 | 313 | 513 | 333 | | 335 | | |
| 445 | 513 | 421 | 667 | 1,082 | 563 | 708 | 412 | | 341 | | |
| 275 | 267 | 301 | 252 | 272 | 267 | 251 | 271 | 246 | 243 | 312 | 275 |
| 97 | 88 | 152 | 55 | 100 | 93 | 77 | 107 | 17 | 33 | 146 | 98 |
| 269 | 266 | 360 | 249 | 317 | 284 | 298 | 302 | 284 | 295 | 250 | 252 |
| 86 | 118 | 125 | 60 | 140 | 121 | 93 | 105 | 116 | 114 | 0 | 3 |
| 302 | 295 | 276 | 442 | 382 | 272 | 370 | 268 | 320 | 257 | 250 | 293 |
| 128 | 109 | 91 | 196 | 242 | 99 | 171 | 96 | 108 | 72 | 0 | 94 |
| 1,000 | 1,024 | 1,000 | 1,000 | 1,500 | 1,500 | 1,125 | 768 | 549 | 512 | 500 | 576 |
| 2,500 | 2,500 | 1,375 | 2,250 | 3,250 | 1,706 | 2,125 | 1,250 | 1,250 | 1,024 | 750 | 1,500 |
| 259 | 253 | 282 | 247 | 283 | 259 | 260 | 250 | 240 | 245 | 271 | 266 |
| 83 | 73 | 128 | 55 | 125 | 105 | 85 | 86 | 26 | 65 | 89 | 75 |
| 160 | 188 | 125 | 172 | 188 | 160 | 156 | 125 | | 128 | | |
| 250 | 250 | 250 | 250 | 250 | 250 | 250 | 250 | | 212 | | |
| 250 | 250 | 250 | 250 | 250 | 250 | 250 | 250 | | 250 | | |
| 256 | 250 | 256 | 250 | 256 | 256 | 256 | 256 | | 256 | | |
| 375 | 275 | 500 | 256 | 375 | 375 | 375 | 275 | | 256 | | |
| 500 | 375 | 500 | 275 | 500 | 469 | 500 | 500 | | 384 | | |
| 500 | 500 | 1,000 | 500 | 751 | 550 | 512 | 512 | | 500 | | |
| 512 | 768 | 1,000 | 500 | 1,125 | 1,250 | 512 | 549 | 275 | 500 | 500 | 576 |
| 1.8 | .8 | .9 | 0 | 1.8 | 1.7 | 0 | 0 | 0 | 0 | 0 | 0 |
| .6 | .3 | .6 | 0 | .7 | .3 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 8.09.--NATURAL AND PROCESSED CHEESE
 CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 40.0 | 6.1 | 35.9 | 37.0 | 35.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 25.0 | 4.9 | 23.3 | 23.7 | 23.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 10.8 | .8 | 9.1 | 10.5 | 9.5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 4.2 | .4 | 3.4 | 2.8 | 2.7 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.6 | 1.3 | 1.6 | 1.6 | 1.6 |
| STANDARD DEVIATION.....DO..... | .9 | .6 | .9 | .8 | .9 |
| MAXIMUM.....DO..... | 8.0 | 3.0 | 8.0 | 5.0 | 7.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 22 | 11 | 17 | 16 | 18 |
| STANDARD DEVIATION.....GM..... | 19 | 11 | 13 | 11 | 15 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 5 | | 4 | 3 | 4 |
| 25TH.....GM..... | 9 | | 9 | 9 | 9 |
| 50TH.....GM..... | 19 | | 9 | 13 | 13 |
| 75TH.....GM..... | 28 | | 19 | 19 | 23 |
| 90TH.....GM..... | 46 | | 35 | 29 | 37 |
| 95TH.....GM..... | 57 | | 45 | 37 | 47 |
| 99TH.....GM..... | 95 | | 65 | 49 | 66 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 42 | 22 | 31 | 34 | 36 |
| STANDARD DEVIATION.....GM..... | 29 | 22 | 19 | 19 | 24 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 46 | 26 | 38 | 34 | 38 |
| STANDARD DEVIATION.....GM..... | 29 | 3 | 20 | 17 | 20 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 49 | 40 | 36 | 34 | 42 |
| STANDARD DEVIATION.....GM..... | 29 | 9 | 16 | 16 | 26 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 700 | 113 | 142 | 198 | 227 |
| MAXIMUM OVER 3 DAYS.....GM..... | 765 | 142 | 282 | 312 | 453 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 41 | 25 | 31 | 31 | 35 |
| STANDARD DEVIATION.....GM..... | 28 | 21 | 19 | 17 | 23 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 14 | | 7 | 7 | 7 |
| 25TH.....GM..... | 28 | | 19 | 28 | 28 |
| 50TH.....GM..... | 28 | | 28 | 28 | 28 |
| 75TH.....GM..... | 56 | | 28 | 28 | 38 |
| 90TH.....GM..... | 58 | | 57 | 57 | 57 |
| 95TH.....GM..... | 85 | | 61 | 57 | 71 |
| 99TH.....GM..... | 140 | | 85 | 85 | 140 |
| MAXIMUM.....GM..... | 700 | 113 | 113 | 140 | 227 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 756.5 | 1.3 | 5.1 | 19.3 | 40.3 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 3.0 | 3.3 | .8 | 2.0 | 4.0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 31.2 | 34.9 | 39.0 | 39.8 | 47.0 | 45.5 | 43.1 | 42.6 | 34.5 | 35.8 | 32.8 | 31.9 |
| 22.8 | 24.5 | 25.4 | 26.4 | 27.5 | 27.5 | 25.2 | 27.4 | 21.2 | 22.1 | 17.9 | 19.0 |
| 6.6 | 8.7 | 9.4 | 10.0 | 13.3 | 12.8 | 12.0 | 11.5 | 8.8 | 10.0 | 9.0 | 9.3 |
| 1.8 | 1.8 | 4.2 | 3.4 | 6.2 | 5.2 | 6.0 | 3.7 | 4.6 | 3.7 | 5.9 | 3.5 |
| 1.4 | 1.5 | 1.6 | 1.6 | 1.8 | 1.7 | 1.7 | 1.6 | 1.6 | 1.6 | 1.7 | 1.6 |
| .8 | .8 | 1.0 | .9 | 1.1 | 1.0 | 1.0 | .9 | .9 | .8 | .9 | .8 |
| 5.0 | 6.0 | 6.0 | 6.0 | 8.0 | 8.0 | 7.0 | 7.0 | 6.0 | 6.0 | 5.0 | 4.0 |
| 18 | 18 | 25 | 19 | 29 | 23 | 26 | 20 | 24 | 20 | 25 | 19 |
| 14 | 14 | 22 | 16 | 25 | 20 | 24 | 15 | 21 | 15 | 21 | 13 |
| 5 | 5 | 7 | 5 | 7 | 5 | 6 | 5 | 5 | 5 | 5 | 5 |
| 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 |
| 15 | 13 | 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 |
| 21 | 20 | 28 | 26 | 37 | 28 | 33 | 28 | 29 | 28 | 34 | 23 |
| 37 | 37 | 47 | 38 | 57 | 47 | 56 | 38 | 47 | 38 | 56 | 38 |
| 47 | 42 | 66 | 52 | 76 | 66 | 66 | 52 | 57 | 47 | 61 | 52 |
| 73 | 77 | 112 | 84 | 117 | 95 | 112 | 75 | 122 | 76 | 136 | 57 |
| 40 | 38 | 49 | 37 | 53 | 41 | 48 | 40 | 44 | 40 | 42 | 39 |
| 21 | 25 | 34 | 24 | 37 | 29 | 33 | 26 | 28 | 30 | 30 | 25 |
| 45 | 40 | 51 | 43 | 56 | 47 | 53 | 42 | 50 | 38 | 61 | 35 |
| 23 | 22 | 29 | 21 | 39 | 25 | 37 | 22 | 29 | 17 | 43 | 15 |
| 44 | 42 | 57 | 47 | 60 | 52 | 53 | 43 | 49 | 43 | 33 | 38 |
| 21 | 23 | 36 | 23 | 32 | 28 | 34 | 21 | 33 | 18 | 12 | 15 |
| 227 | 284 | 395 | 227 | 339 | 310 | 700 | 252 | 255 | 228 | 306 | 170 |
| 280 | 312 | 529 | 349 | 765 | 423 | 765 | 414 | 510 | 255 | 408 | 210 |
| 39 | 35 | 46 | 37 | 49 | 40 | 46 | 37 | 44 | 37 | 43 | 36 |
| 22 | 23 | 30 | 23 | 33 | 27 | 37 | 23 | 28 | 22 | 36 | 21 |
| 14 | 7 | 14 | 10 | 14 | 10 | 14 | 14 | 14 | 14 | 5 | 14 |
| 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 |
| 28 | 28 | 28 | 28 | 43 | 28 | 28 | 28 | 28 | 28 | 28 | 28 |
| 56 | 56 | 57 | 56 | 57 | 57 | 57 | 56 | 57 | 56 | 57 | 56 |
| 57 | 57 | 84 | 57 | 85 | 61 | 84 | 57 | 84 | 57 | 85 | 57 |
| 84 | 81 | 112 | 84 | 113 | 85 | 112 | 84 | 91 | 84 | 113 | 58 |
| 113 | 113 | 170 | 113 | 170 | 128 | 168 | 116 | 170 | 113 | 170 | 113 |
| 168 | 284 | 227 | 227 | 315 | 280 | 700 | 244 | 170 | 170 | 306 | 170 |
| 33.7 | 50.7 | 29.4 | 46.4 | 94.4 | 115.7 | 108.3 | 148.8 | 19.9 | 22.8 | 11.0 | 9.5 |
| 3.6 | 4.5 | 3.3 | 5.0 | 2.9 | 2.8 | 3.0 | 3.1 | 3.1 | 2.3 | 3.6 | 1.9 |

TABLE 8.10.--CHEDDAR CHEESE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.4 | .7 | 4.6 | 3.5 | 2.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.1 | .7 | 3.0 | 2.8 | 2.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.0 | 0 | 1.4 | .3 | .5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | .2 | .3 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.4 | 1.0 | 1.5 | 1.3 | 1.2 |
| STANDARD DEVIATION.....DO..... | .7 | .0 | .7 | .8 | .4 |
| MAXIMUM.....DO..... | 6.0 | 1.0 | 3.0 | 5.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 20 | 4 | 16 | 15 | 15 |
| STANDARD DEVIATION.....GM..... | 17 | 1 | 12 | 9 | 11 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 5 | | | 5 | 5 |
| 25TH.....GM..... | 9 | | | 9 | 9 |
| 50TH.....GM..... | 19 | | | 9 | 9 |
| 75TH.....GM..... | 28 | | | 19 | 19 |
| 90TH.....GM..... | 38 | | | 28 | 28 |
| 95TH.....GM..... | 57 | | | 33 | 38 |
| 99TH.....GM..... | 85 | | | 56 | 56 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 46 | 12 | 34 | 36 | 36 |
| STANDARD DEVIATION.....GM..... | 34 | 3 | 23 | 21 | 22 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 48 | 0 | 38 | 33 | 47 |
| STANDARD DEVIATION.....GM..... | 28 | 0 | 21 | 9 | 20 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 49 | 0 | 28 | 32 | 0 |
| STANDARD DEVIATION.....GM..... | 25 | 0 | (2) | 10 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 306 | 14 | 113 | 113 | 113 |
| MAXIMUM OVER 3 DAYS.....GM..... | 510 | 14 | 142 | 168 | 169 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 45 | 12 | 33 | 34 | 38 |
| STANDARD DEVIATION.....GM..... | 31 | 3 | 23 | 18 | 22 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 14 | | | 14 | 14 |
| 25TH.....GM..... | 28 | | | 28 | 28 |
| 50TH.....GM..... | 28 | | | 28 | 28 |
| 75TH.....GM..... | 57 | | | 28 | 56 |
| 90TH.....GM..... | 85 | | | 57 | 57 |
| 95TH.....GM..... | 112 | | | 77 | 77 |
| 99TH.....GM..... | 170 | | | 85 | 113 |
| MAXIMUM.....GM..... | 306 | 14 | 113 | 113 | 113 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 52.2 | 0 | 0 | 0 | 2.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.9 | 0 | 0 | 0 | 3.8 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.1 | 3.6 | 3.9 | 4.5 | 5.6 | 7.0 | 6.7 | 5.9 | 7.6 | 5.5 | 7.6 | 4.8 |
| 2.8 | 3.0 | 3.2 | 3.6 | 4.6 | 5.1 | 5.3 | 4.4 | 5.8 | 3.6 | 3.8 | 3.6 |
| .3 | .5 | .7 | .6 | .7 | 1.5 | 1.1 | 1.2 | 1.1 | 1.3 | 1.5 | 1.1 |
| 0 | 0 | 0 | .2 | .2 | .4 | .3 | .4 | .7 | .6 | 2.2 | .1 |
| 1.1 | 1.2 | 1.2 | 1.3 | 1.3 | 1.5 | 1.3 | 1.4 | 1.5 | 1.5 | 1.8 | 1.3 |
| .4 | .5 | .5 | .7 | .6 | .8 | .8 | .7 | 1.1 | .7 | .9 | .5 |
| 3.0 | 3.0 | 3.0 | 5.0 | 4.0 | 6.0 | 6.0 | 5.0 | 6.0 | 3.0 | 3.0 | 3.0 |
| 14 | 16 | 23 | 17 | 21 | 22 | 23 | 19 | 27 | 23 | 28 | 18 |
| 8 | 12 | 17 | 15 | 15 | 18 | 18 | 15 | 25 | 17 | 29 | 13 |
| 5 | 2 | 9 | 2 | 5 | 5 | 5 | 5 | 9 | 9 | | |
| 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 17 | 9 | | |
| 9 | 9 | 19 | 9 | 19 | 19 | 19 | 12 | 19 | 19 | | |
| 19 | 19 | 28 | 23 | 28 | 28 | 28 | 21 | 28 | 28 | | |
| 24 | 38 | 47 | 38 | 38 | 47 | 47 | 38 | 61 | 45 | | |
| 28 | 38 | 66 | 47 | 47 | 57 | 61 | 47 | 76 | 64 | | |
| 47 | 50 | 76 | 84 | 76 | 85 | 91 | 91 | 170 | 85 | | |
| 40 | 42 | 65 | 40 | 54 | 45 | 52 | 40 | 56 | 53 | 53 | 48 |
| 24 | 33 | 50 | 31 | 39 | 32 | 41 | 28 | 24 | 32 | 44 | 33 |
| 30 | 37 | 41 | 45 | 49 | 56 | 58 | 43 | 55 | 33 | 70 | 32 |
| 10 | 22 | 27 | 15 | 17 | 28 | 26 | 23 | 33 | 12 | 80 | 17 |
| 0 | 0 | 0 | 51 | 48 | 48 | 54 | 48 | 79 | 59 | 32 | 64 |
| 0 | 0 | 0 | 24 | 21 | 22 | 25 | 19 | 46 | 19 | 6 | 0 |
| 142 | 150 | 227 | 140 | 227 | 254 | 272 | 204 | 170 | 142 | 306 | 136 |
| 142 | 168 | 227 | 253 | 255 | 310 | 272 | 272 | 510 | 255 | 408 | 192 |
| 37 | 38 | 56 | 39 | 50 | 45 | 51 | 40 | 54 | 46 | 47 | 42 |
| 22 | 27 | 46 | 26 | 33 | 31 | 37 | 26 | 26 | 27 | 54 | 31 |
| 14 | 7 | 28 | 13 | 14 | 14 | 14 | 14 | 21 | 26 | | |
| 28 | 26 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | | |
| 28 | 28 | 28 | 28 | 56 | 28 | 43 | 28 | 56 | 28 | | |
| 56 | 57 | 57 | 56 | 57 | 57 | 57 | 57 | 57 | 57 | | |
| 57 | 57 | 113 | 57 | 85 | 84 | 85 | 68 | 85 | 85 | | |
| 71 | 112 | 170 | 113 | 112 | 112 | 113 | 85 | 113 | 113 | | |
| 113 | 136 | 227 | 140 | 227 | 170 | 227 | 113 | 113 | 136 | | |
| 113 | 136 | 227 | 140 | 227 | 227 | 272 | 204 | 113 | 136 | 306 | 136 |
| .9 | 3.7 | 2.1 | 3.5 | 1.8 | 7.3 | 14.6 | 9.7 | 1.0 | 2.1 | 0 | 3.0 |
| 1.3 | 3.9 | 3.2 | 3.9 | .7 | 1.3 | 3.3 | 1.7 | .8 | 1.5 | 0 | 4.8 |

TABLE 8.11.--SWISS CHEESE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.2 | 0 | 1.0 | 1.1 | 1.4 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.4 | 0 | .4 | .9 | 1.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .7 | 0 | .5 | .2 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | 0 | 0 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.4 | 0 | 1.6 | 1.2 | 1.7 |
| STANDARD DEVIATION.....DO..... | .7 | 0 | .5 | .4 | 1.0 |
| MAXIMUM.....DO..... | 5.0 | 0 | 2.0 | 2.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 20 | 0 | 18 | 13 | 30 |
| STANDARD DEVIATION.....GM..... | 17 | 0 | 8 | 6 | 39 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 5 | | | | |
| 25TH.....GM..... | 9 | | | | |
| 50TH.....GM..... | 19 | | | | |
| 75TH.....GM..... | 23 | | | | |
| 90TH.....GM..... | 38 | | | | |
| 95TH.....GM..... | 56 | | | | |
| 99TH.....GM..... | 94 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 45 | 0 | 42 | 31 | 43 |
| STANDARD DEVIATION.....GM..... | 29 | 0 | 28 | 14 | 34 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 49 | 0 | 32 | 32 | 30 |
| STANDARD DEVIATION.....GM..... | 31 | 0 | 8 | 7 | 14 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 59 | 0 | 0 | 0 | 97 |
| STANDARD DEVIATION.....GM..... | 30 | 0 | 0 | 0 | 50 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 227 | 0 | 84 | 57 | 227 |
| MAXIMUM OVER 3 DAYS.....GM..... | 453 | 0 | 85 | 85 | 453 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 45 | 0 | 35 | 32 | 53 |
| STANDARD DEVIATION.....GM..... | 27 | 0 | 19 | 13 | 51 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 14 | | | | |
| 25TH.....GM..... | 28 | | | | |
| 50TH.....GM..... | 28 | | | | |
| 75TH.....GM..... | 57 | | | | |
| 90TH.....GM..... | 85 | | | | |
| 95TH.....GM..... | 112 | | | | |
| 99TH.....GM..... | 140 | | | | |
| MAXIMUM.....GM..... | 227 | 0 | 84 | 57 | 227 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 42.6 | 0 | 0 | 0 | 3.4 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.6 | 0 | 0 | 0 | 7.9 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.4 | 1.6 | 2.4 | 2.6 | 5.1 | 3.8 | 4.4 | 4.0 | 3.3 | 1.9 | 2.1 | 2.5 |
| 1.2 | 1.4 | 2.1 | 2.1 | 3.8 | 2.9 | 3.1 | 3.1 | 2.1 | 1.4 | 1.5 | 1.9 |
| .1 | .2 | .2 | .4 | 1.1 | .8 | 1.0 | .8 | 1.0 | .5 | .6 | .6 |
| (2) | .1 | .1 | .1 | .1 | .1 | .2 | .1 | .1 | 0 | 0 | 0 |
| 1.2 | 1.2 | 1.2 | 1.3 | 1.4 | 1.3 | 1.5 | 1.3 | 1.4 | 1.3 | 1.3 | 1.2 |
| .6 | .4 | .6 | .7 | .6 | .5 | .9 | .6 | .6 | .4 | .4 | .4 |
| 4.0 | 3.0 | 4.0 | 4.0 | 4.0 | 3.0 | 5.0 | 4.0 | 3.0 | 2.0 | 2.0 | 2.0 |
| 18 | 14 | 23 | 17 | 25 | 18 | 23 | 18 | 26 | 18 | 30 | 15 |
| 13 | 10 | 19 | 14 | 19 | 14 | 21 | 14 | 16 | 10 | 13 | 6 |
| | | | | 9 | 5 | 5 | 5 | | | | |
| | | | | 9 | 9 | 9 | 9 | | | | |
| | | | | 19 | 11 | 19 | 14 | | | | |
| | | | | 37 | 19 | 28 | 19 | | | | |
| | | | | 56 | 38 | 38 | 38 | | | | |
| | | | | 66 | 45 | 57 | 49 | | | | |
| | | | | 94 | 85 | 113 | 85 | | | | |
| 43 | 40 | 59 | 35 | 56 | 40 | 44 | 42 | 57 | 40 | 71 | 42 |
| 25 | 29 | 42 | 14 | 37 | 24 | 25 | 27 | 20 | 15 | 26 | 20 |
| 53 | 28 | 58 | 48 | 61 | 40 | 67 | 38 | 47 | 48 | 71 | 28 |
| 25 | 10 | 35 | 29 | 34 | 20 | 44 | 16 | 9 | 13 | 0 | (2) |
| 54 | 21 | 85 | 48 | 59 | 67 | 37 | 62 | 85 | 0 | 0 | 0 |
| 0 | 0 | 0 | 10 | 33 | 20 | 13 | 17 | 0 | 0 | 0 | 0 |
| 113 | 170 | 168 | 113 | 224 | 168 | 227 | 170 | 113 | 84 | 113 | 85 |
| 170 | 170 | 255 | 170 | 339 | 255 | 340 | 255 | 255 | 113 | 142 | 85 |
| 43 | 36 | 57 | 38 | 53 | 41 | 46 | 41 | 55 | 43 | 71 | 37 |
| 26 | 27 | 40 | 15 | 28 | 25 | 24 | 24 | 20 | 17 | 22 | 17 |
| | | | | 21 | 14 | 14 | 14 | | | | |
| | | | | 28 | 28 | 28 | 28 | | | | |
| | | | | 56 | 28 | 56 | 28 | | | | |
| | | | | 57 | 57 | 57 | 57 | | | | |
| | | | | 88 | 60 | 70 | 57 | | | | |
| | | | | 113 | 112 | 105 | 85 | | | | |
| | | | | 140 | 113 | 113 | 140 | | | | |
| 113 | 170 | 168 | 57 | 142 | 113 | 168 | 170 | 113 | 84 | 113 | 85 |
| 0 | 1.3 | 0 | 1.4 | 13.7 | 6.7 | 1.5 | 12.7 | 0 | 0 | 0 | 2.1 |
| 0 | 3.1 | 0 | 2.7 | 4.9 | 2.6 | .5 | 3.4 | 0 | 0 | 0 | 6.9 |

TABLE 8.12.--NATURAL CHEESE OTHER THAN CHEDDAR, SWISS, AND PARMESAN
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.3 | .5 | 1.8 | 2.1 | 1.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.4 | .5 | 1.6 | 1.7 | 1.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .6 | 0 | .2 | .2 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | 0 | .2 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.4 | 1.0 | 1.2 | 1.3 | 1.2 |
| STANDARD DEVIATION.....DO..... | .8 | .0 | .4 | .6 | .5 |
| MAXIMUM.....DO..... | 6.0 | 1.0 | 2.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 22 | 5 | 14 | 14 | 15 |
| STANDARD DEVIATION.....GM..... | 20 | 4 | 8 | 7 | 10 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 5 | | | | |
| 25TH.....GM..... | 9 | | | | |
| 50TH.....GM..... | 19 | | | | |
| 75TH.....GM..... | 28 | | | | |
| 90TH.....GM..... | 38 | | | | |
| 95TH.....GM..... | 57 | | | | |
| 99TH.....GM..... | 92 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 49 | 16 | 35 | 36 | 36 |
| STANDARD DEVIATION.....GM..... | 40 | 11 | 17 | 18 | 25 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 48 | 0 | 49 | 27 | 40 |
| STANDARD DEVIATION.....GM..... | 31 | 0 | 7 | 13 | 18 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 48 | 0 | 0 | 24 | 28 |
| STANDARD DEVIATION.....GM..... | 30 | 0 | 0 | 6 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 680 | 28 | 57 | 57 | 113 |
| MAXIMUM OVER 3 DAYS.....GM..... | 737 | 28 | 113 | 85 | 113 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 46 | 16 | 35 | 32 | 36 |
| STANDARD DEVIATION.....GM..... | 35 | 11 | 18 | 17 | 22 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 14 | | | | |
| 25TH.....GM..... | 28 | | | | |
| 50TH.....GM..... | 28 | | | | |
| 75TH.....GM..... | 57 | | | | |
| 90TH.....GM..... | 85 | | | | |
| 95TH.....GM..... | 113 | | | | |
| 99TH.....GM..... | 140 | | | | |
| MAXIMUM.....GM..... | 680 | 28 | 57 | 57 | 113 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 38.9 | 0 | 0 | .7 | 6.4 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.2 | 0 | 0 | 1.6 | 16.2 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.3 | 1.8 | 2.5 | 1.9 | 4.0 | 5.6 | 3.6 | 3.8 | 3.0 | 3.4 | 3.3 | 2.6 |
| 1.1 | 1.6 | 1.9 | 1.4 | 3.0 | 4.3 | 2.5 | 2.6 | 1.9 | 2.3 | 2.0 | 1.9 |
| .1 | .2 | .5 | .5 | .6 | .7 | .8 | .9 | .8 | .9 | .8 | .3 |
| (2) | (2) | .1 | .1 | .4 | .6 | .3 | .3 | .4 | .2 | .5 | .5 |
| 1.2 | 1.2 | 1.4 | 1.3 | 1.5 | 1.4 | 1.5 | 1.4 | 1.5 | 1.5 | 1.7 | 1.5 |
| .5 | .5 | 1.0 | .5 | .9 | .8 | .7 | .7 | .7 | .7 | 1.1 | .9 |
| 3.0 | 4.0 | 6.0 | 3.0 | 6.0 | 5.0 | 4.0 | 3.0 | 3.0 | 3.0 | 4.0 | 4.0 |
| 20 | 19 | 22 | 15 | 27 | 22 | 28 | 19 | 18 | 22 | 28 | 23 |
| 13 | 16 | 13 | 10 | 20 | 18 | 34 | 12 | 10 | 17 | 19 | 21 |
| | | | | 6 | 5 | 9 | 5 | | 9 | | |
| | | | | 14 | 9 | 9 | 9 | | 9 | | |
| | | | | 19 | 19 | 19 | 19 | | 19 | | |
| | | | | 38 | 28 | 28 | 26 | | 28 | | |
| | | | | 53 | 43 | 56 | 37 | | 56 | | |
| | | | | 57 | 57 | 75 | 42 | | 62 | | |
| | | | | 113 | 76 | 227 | 62 | | 66 | | |
| 50 | 48 | 55 | 39 | 60 | 46 | 62 | 45 | 42 | 43 | 45 | 31 |
| 24 | 46 | 31 | 31 | 33 | 27 | 77 | 29 | 18 | 30 | 23 | 18 |
| 62 | 49 | 42 | 29 | 66 | 54 | 55 | 37 | 30 | 53 | 75 | 70 |
| 29 | 17 | 18 | 10 | 46 | 30 | 37 | 20 | 12 | 24 | 22 | 20 |
| 33 | 38 | 46 | 28 | 50 | 53 | 63 | 34 | 34 | 43 | 47 | 56 |
| 0 | 0 | 12 | 0 | 16 | 19 | 62 | 12 | 9 | 18 | 0 | 8 |
| 112 | 284 | 113 | 112 | 226 | 170 | 680 | 170 | 57 | 140 | 103 | 98 |
| 196 | 284 | 170 | 112 | 339 | 340 | 737 | 224 | 140 | 198 | 181 | 210 |
| 51 | 47 | 45 | 34 | 55 | 47 | 58 | 39 | 36 | 44 | 49 | 45 |
| 25 | 40 | 26 | 26 | 31 | 26 | 63 | 23 | 18 | 25 | 25 | 21 |
| | | | | 14 | 14 | 14 | 14 | | 15 | | |
| | | | | 28 | 28 | 28 | 28 | | 28 | | |
| | | | | 57 | 56 | 38 | 28 | | 28 | | |
| | | | | 57 | 57 | 57 | 57 | | 57 | | |
| | | | | 113 | 85 | 113 | 62 | | 84 | | |
| | | | | 113 | 112 | 140 | 85 | | 85 | | |
| | | | | 128 | 128 | 284 | 113 | | 113 | | |
| 112 | 284 | 113 | 112 | 142 | 170 | 680 | 113 | 57 | 113 | 103 | 85 |
| 1.6 | 2.2 | 0 | 0 | 2.3 | 9.3 | 2.3 | 14.1 | 0 | 0 | 0 | 0 |
| 4.9 | 4.8 | 0 | 0 | 1.0 | 2.2 | .9 | 3.7 | 0 | 0 | 0 | 0 |

TABLE 8.13.--PROCESSED CHEESE, CHEESE FOOD, AND CHEESE SPREAD
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 22.4 | 3.9 | 23.5 | 24.8 | 22.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 16.2 | 2.9 | 16.3 | 17.1 | 16.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 4.8 | .6 | 5.1 | 6.4 | 5.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.4 | .4 | 2.1 | 1.3 | 1.1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1</u> /NUMBER.... | 1.4 | 1.4 | 1.5 | 1.4 | 1.4 |
| STANDARD DEVIATION.....DO..... | .7 | .7 | .9 | .7 | .7 |
| MAXIMUM.....DO..... | 7.0 | 3.0 | 6.0 | 4.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1</u> /GM..... | 19 | 14 | 16 | 16 | 17 |
| STANDARD DEVIATION.....GM..... | 15 | 13 | 13 | 10 | 12 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> / | | | | | |
| 5TH.....GM..... | 5 | | 5 | 5 | 5 |
| 25TH.....GM..... | 9 | | 9 | 9 | 9 |
| 50TH.....GM..... | 19 | | 9 | 9 | 12 |
| 75TH.....GM..... | 22 | | 19 | 19 | 19 |
| 90TH.....GM..... | 37 | | 30 | 29 | 28 |
| 95TH.....GM..... | 47 | | 38 | 37 | 38 |
| 99TH.....GM..... | 76 | | 65 | 47 | 56 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 42 | 28 | 32 | 35 | 40 |
| STANDARD DEVIATION.....GM..... | 28 | 27 | 20 | 20 | 27 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 44 | 26 | 34 | 34 | 35 |
| STANDARD DEVIATION.....GM..... | 24 | 3 | 16 | 16 | 16 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 44 | 40 | 39 | 32 | 37 |
| STANDARD DEVIATION.....GM..... | 25 | 9 | 19 | 14 | 15 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 700 | 113 | 140 | 142 | 168 |
| MAXIMUM OVER 3 DAYS.....GM..... | 765 | 142 | 282 | 227 | 280 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1</u> /GM..... | 40 | 29 | 32 | 33 | 36 |
| STANDARD DEVIATION.....GM..... | 26 | 23 | 19 | 17 | 22 |
| PERCENTILES: <u>1</u> / | | | | | |
| 5TH.....GM..... | 14 | | 9 | 14 | 14 |
| 25TH.....GM..... | 28 | | 21 | 28 | 28 |
| 50TH.....GM..... | 28 | | 28 | 28 | 28 |
| 75TH.....GM..... | 56 | | 28 | 31 | 43 |
| 90TH.....GM..... | 57 | | 57 | 57 | 57 |
| 95TH.....GM..... | 85 | | 78 | 57 | 71 |
| 99TH.....GM..... | 132 | | 85 | 85 | 140 |
| MAXIMUM.....GM..... | 700 | 113 | 112 | 140 | 168 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> / | | | | | |
| EATING OCCASIONS.....NUMBER.... | 126.1 | 0 | 0 | 3.7 | 6.8 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.0 | 0 | 0 | .6 | 1.2 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 17.9 | 20.6 | 23.5 | 21.8 | 25.7 | 24.8 | 23.1 | 23.7 | 17.0 | 18.5 | 16.4 | 17.3 |
| 14.1 | 16.8 | 17.5 | 15.7 | 17.0 | 17.7 | 16.2 | 17.8 | 13.2 | 13.3 | 11.6 | 12.0 |
| 2.9 | 2.9 | 4.4 | 4.8 | 6.4 | 5.6 | 5.2 | 5.0 | 2.8 | 3.7 | 4.0 | 3.8 |
| .9 | .9 | 1.5 | 1.3 | 2.2 | 1.5 | 1.7 | 1.0 | 1.0 | 1.5 | .9 | 1.5 |
| 1.3 | 1.3 | 1.4 | 1.4 | 1.5 | 1.5 | 1.4 | 1.4 | 1.3 | 1.4 | 1.4 | 1.4 |
| .7 | .7 | .8 | .8 | .8 | .8 | .7 | .7 | .6 | .7 | .6 | .7 |
| 4.0 | 5.0 | 5.0 | 5.0 | 6.0 | 7.0 | 5.0 | 6.0 | 4.0 | 3.0 | 4.0 | 4.0 |
| 18 | 16 | 21 | 18 | 25 | 19 | 22 | 17 | 18 | 16 | 18 | 17 |
| 13 | 12 | 17 | 14 | 19 | 15 | 20 | 12 | 11 | 11 | 13 | 10 |
| 7 | 5 | 7 | 5 | 9 | 5 | 7 | 5 | 5 | 5 | 2 | 5 |
| 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 |
| 14 | 9 | 19 | 15 | 19 | 19 | 19 | 13 | 19 | 9 | 15 | 19 |
| 19 | 19 | 28 | 19 | 28 | 23 | 28 | 19 | 19 | 19 | 20 | 19 |
| 37 | 28 | 38 | 37 | 47 | 38 | 38 | 31 | 37 | 29 | 37 | 28 |
| 43 | 38 | 57 | 47 | 65 | 50 | 56 | 39 | 42 | 38 | 38 | 38 |
| 71 | 75 | 85 | 76 | 103 | 75 | 85 | 65 | 57 | 56 | 61 | 57 |
| 41 | 40 | 47 | 39 | 52 | 40 | 49 | 39 | 44 | 35 | 39 | 39 |
| 19 | 24 | 30 | 22 | 33 | 25 | 40 | 22 | 28 | 23 | 25 | 23 |
| 47 | 39 | 51 | 41 | 51 | 45 | 47 | 43 | 43 | 39 | 42 | 33 |
| 23 | 16 | 28 | 19 | 31 | 22 | 26 | 23 | 16 | 13 | 23 | 10 |
| 42 | 44 | 48 | 49 | 53 | 47 | 50 | 37 | 33 | 33 | 37 | 33 |
| 19 | 22 | 24 | 17 | 27 | 23 | 37 | 15 | 8 | 10 | 12 | 12 |
| 140 | 227 | 196 | 210 | 315 | 258 | 700 | 170 | 170 | 228 | 122 | 170 |
| 245 | 280 | 298 | 241 | 504 | 308 | 765 | 308 | 170 | 228 | 183 | 170 |
| 40 | 37 | 45 | 38 | 48 | 40 | 46 | 37 | 41 | 36 | 40 | 35 |
| 21 | 20 | 27 | 21 | 32 | 25 | 36 | 22 | 25 | 19 | 24 | 19 |
| 14 | 14 | 14 | 14 | 18 | 14 | 15 | 14 | 14 | 14 | 9 | 14 |
| 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 |
| 28 | 28 | 28 | 28 | 40 | 28 | 28 | 28 | 28 | 28 | 28 | 28 |
| 56 | 56 | 56 | 56 | 57 | 56 | 57 | 56 | 56 | 45 | 56 | 56 |
| 57 | 57 | 84 | 57 | 84 | 57 | 76 | 57 | 57 | 57 | 61 | 57 |
| 84 | 70 | 112 | 61 | 112 | 85 | 85 | 84 | 84 | 57 | 85 | 57 |
| 113 | 113 | 168 | 113 | 170 | 122 | 142 | 116 | 142 | 113 | 122 | 70 |
| 140 | 140 | 170 | 210 | 315 | 258 | 700 | 170 | 170 | 158 | 122 | 170 |
| 7.7 | 12.3 | 8.1 | 10.5 | 11.6 | 11.6 | 14.6 | 26.7 | 8.0 | .9 | 3.5 | 0 |
| 1.5 | 2.1 | 1.7 | 2.3 | .7 | .6 | .9 | 1.2 | 3.2 | .2 | 2.9 | 0 |

TABLE 8.14---COTTAGE CHEESE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 8.6 | 2.7 | 4.9 | 5.0 | 4.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.9 | 1.6 | 3.3 | 4.1 | 3.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.1 | .5 | 1.5 | .9 | .4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .6 | .6 | .1 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.5 | 1.7 | 1.5 | 1.3 | 1.1 |
| STANDARD DEVIATION.....DO..... | .8 | 1.0 | .6 | .5 | .3 |
| MAXIMUM.....DO..... | 6.0 | 4.0 | 4.0 | 3.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 55 | 44 | 37 | 40 | 40 |
| STANDARD DEVIATION.....GM..... | 45 | 45 | 29 | 28 | 26 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 9 | | 5 | 9 | 9 |
| 25TH.....GM..... | 28 | | 19 | 19 | 19 |
| 50TH.....GM..... | 38 | | 38 | 38 | 38 |
| 75TH.....GM..... | 75 | | 42 | 57 | 38 |
| 90TH.....GM..... | 113 | | 75 | 75 | 75 |
| 95TH.....GM..... | 151 | | 94 | 80 | 75 |
| 99TH.....GM..... | 226 | | 177 | 151 | 151 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 117 | 73 | 85 | 100 | 113 |
| STANDARD DEVIATION.....GM..... | 75 | 37 | 65 | 64 | 76 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 123 | 67 | 78 | 101 | 84 |
| STANDARD DEVIATION.....GM..... | 74 | 14 | 34 | 51 | 42 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 119 | 98 | 177 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 75 | 71 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 678 | 199 | 339 | 226 | 452 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,130 | 597 | 532 | 452 | 452 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 113 | 77 | 77 | 93 | 105 |
| STANDARD DEVIATION.....GM..... | 71 | 54 | 49 | 56 | 61 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 19 | | 14 | 28 | 27 |
| 25TH.....GM..... | 57 | | 42 | 57 | 57 |
| 50TH.....GM..... | 113 | | 57 | 85 | 113 |
| 75TH.....GM..... | 113 | | 113 | 113 | 113 |
| 90TH.....GM..... | 226 | | 133 | 226 | 226 |
| 95TH.....GM..... | 226 | | 170 | 226 | 226 |
| 99TH.....GM..... | 339 | | 226 | 226 | 226 |
| MAXIMUM.....GM..... | 452 | 199 | 226 | 226 | 226 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 29.7 | 0 | .9 | 0 | 4.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | 0 | 1.2 | 0 | 4.8 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.2 | 4.5 | 3.9 | 4.6 | 5.3 | 8.4 | 9.9 | 13.8 | 11.1 | 17.3 | 7.8 | 16.7 |
| 2.5 | 3.8 | 3.2 | 4.0 | 3.9 | 5.8 | 7.0 | 9.1 | 6.4 | 10.7 | 6.2 | 8.5 |
| .7 | .6 | .6 | .5 | 1.3 | 2.2 | 2.4 | 3.5 | 2.9 | 5.3 | .9 | 6.1 |
| (2) | .1 | .1 | .1 | .1 | .4 | .5 | 1.2 | 1.8 | 1.3 | .8 | 2.1 |
| 1.2 | 1.2 | 1.2 | 1.2 | 1.3 | 1.5 | 1.4 | 1.5 | 1.8 | 1.6 | 1.3 | 1.7 |
| .5 | .5 | .5 | .6 | .5 | .8 | .7 | .8 | 1.2 | .8 | .6 | .9 |
| 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 5.0 | 6.0 | 5.0 | 6.0 | 6.0 | 3.0 | 5.0 |
| 47 | 43 | 48 | 49 | 63 | 59 | 58 | 53 | 66 | 53 | 69 | 59 |
| 34 | 38 | 34 | 36 | 44 | 49 | 45 | 41 | 63 | 46 | 74 | 50 |
| 5 | 5 | 9 | 9 | 19 | 14 | 9 | 9 | 9 | 7 | | 5 |
| 29 | 19 | 28 | 25 | 38 | 29 | 38 | 29 | 24 | 19 | | 28 |
| 38 | 38 | 38 | 38 | 47 | 38 | 38 | 38 | 38 | 38 | | 38 |
| 57 | 50 | 75 | 75 | 75 | 75 | 75 | 75 | 75 | 75 | | 75 |
| 94 | 75 | 75 | 94 | 114 | 113 | 113 | 94 | 151 | 132 | | 113 |
| 113 | 113 | 151 | 113 | 151 | 151 | 151 | 132 | 226 | 151 | | 151 |
| 151 | 226 | 151 | 188 | 226 | 226 | 226 | 207 | 264 | 207 | | 283 |
| 106 | 102 | 113 | 131 | 155 | 123 | 125 | 114 | 119 | 96 | 135 | 102 |
| 59 | 63 | 59 | 102 | 94 | 80 | 74 | 65 | 96 | 66 | 77 | 63 |
| 135 | 113 | 147 | 119 | 138 | 143 | 139 | 114 | 112 | 126 | 92 | 108 |
| 62 | 78 | 66 | 42 | 79 | 86 | 80 | 67 | 81 | 73 | 46 | 63 |
| 76 | 167 | 20 | 104 | 171 | 120 | 120 | 105 | 144 | 105 | 277 | 123 |
| 0 | 56 | 0 | 9 | 19 | 81 | 68 | 66 | 87 | 64 | 47 | 74 |
| 339 | 283 | 339 | 565 | 678 | 678 | 678 | 678 | 452 | 509 | 339 | 452 |
| 452 | 678 | 452 | 565 | 904 | 1,130 | 904 | 1,017 | 791 | 735 | 904 | 848 |
| 114 | 106 | 117 | 119 | 145 | 122 | 126 | 106 | 111 | 103 | 160 | 106 |
| 64 | 67 | 68 | 79 | 85 | 73 | 75 | 63 | 80 | 65 | 98 | 67 |
| 28 | 14 | 28 | 28 | 42 | 28 | 27 | 27 | 14 | 14 | | 14 |
| 57 | 57 | 85 | 57 | 113 | 75 | 85 | 57 | 57 | 57 | | 57 |
| 113 | 113 | 113 | 113 | 113 | 113 | 113 | 113 | 113 | 113 | | 113 |
| 113 | 113 | 113 | 113 | 226 | 113 | 170 | 113 | 113 | 113 | | 113 |
| 226 | 226 | 226 | 226 | 226 | 226 | 226 | 226 | 226 | 226 | | 226 |
| 226 | 226 | 226 | 226 | 226 | 226 | 226 | 226 | 226 | 226 | | 226 |
| 339 | 226 | 339 | 452 | 452 | 452 | 339 | 226 | 452 | 228 | | 226 |
| 339 | 283 | 339 | 452 | 452 | 452 | 452 | 452 | 452 | 339 | 339 | 339 |
| .8 | 2.8 | 0 | 0 | .8 | 0 | 0 | 9.7 | 0 | 8.0 | 0 | 2.6 |
| 1.0 | 2.4 | 0 | 0 | .3 | 0 | 0 | .7 | 0 | 1.7 | 0 | .9 |

TABLE 8.15.--FLUID CREAM
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 7.0 | .1 | .7 | .6 | .5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.3 | 0 | .6 | .5 | .3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.6 | 0 | .1 | (2) | (2) |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.1 | .1 | 0 | .1 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 2.7 | 18.0 | 1.2 | 1.5 | 1.5 |
| STANDARD DEVIATION.....DO..... | 2.2 | .0 | .4 | 1.2 | .8 |
| MAXIMUM.....DO..... | 19.0 | 18.0 | 2.0 | 5.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 23 | 685 | 25 | 24 | 32 |
| STANDARD DEVIATION.....GM..... | 35 | 0 | 15 | 23 | 30 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 2 | | | | |
| 25TH.....GM..... | 5 | | | | |
| 50TH.....GM..... | 12 | | | | |
| 75TH.....GM..... | 27 | | | | |
| 90TH.....GM..... | 50 | | | | |
| 95TH.....GM..... | 75 | | | | |
| 99TH.....GM..... | 140 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 31 | 0 | 65 | 55 | 110 |
| STANDARD DEVIATION.....GM..... | 36 | 0 | 44 | 42 | 99 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 32 | 0 | 60 | 60 | 60 |
| STANDARD DEVIATION.....GM..... | 29 | 0 | 0 | 0 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 43 | 685 | 0 | 48 | 12 |
| STANDARD DEVIATION.....GM..... | 54 | 0 | 0 | 39 | 10 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 726 | 726 | 121 | 121 | 240 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,057 | 2,057 | 121 | 273 | 240 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 26 | 114 | 63 | 48 | 65 |
| STANDARD DEVIATION.....GM..... | 32 | 19 | 37 | 41 | 84 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 5 | | | | |
| 25TH.....GM..... | 10 | | | | |
| 50TH.....GM..... | 15 | | | | |
| 75TH.....GM..... | 30 | | | | |
| 90TH.....GM..... | 60 | | | | |
| 95TH.....GM..... | 76 | | | | |
| 99TH.....GM..... | 161 | | | | |
| MAXIMUM.....GM..... | 484 | 121 | 121 | 121 | 240 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 108.2 | 0 | 1.0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.5 | 0 | 11.5 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACF.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.1 | .9 | 2.9 | 2.7 | 9.0 | 6.5 | 11.8 | 9.9 | 11.3 | 10.3 | 11.5 | 11.9 |
| .7 | .7 | 1.8 | 1.6 | 4.2 | 3.9 | 5.6 | 4.4 | 5.3 | 4.2 | 2.7 | 4.9 |
| .2 | .2 | .6 | .8 | 2.2 | 1.5 | 2.6 | 2.1 | 2.3 | 2.6 | 4.1 | 2.3 |
| .2 | .1 | .6 | .3 | 2.5 | 1.5 | 3.7 | 3.4 | 3.6 | 3.6 | 4.6 | 4.7 |
| 1.8 | 1.4 | 1.7 | 2.0 | 2.5 | 2.3 | 3.0 | 2.8 | 2.4 | 2.8 | 3.4 | 2.7 |
| 1.3 | .7 | 1.0 | 1.6 | 2.0 | 2.1 | 2.6 | 2.3 | 1.6 | 2.1 | 2.6 | 1.9 |
| 6.0 | 3.0 | 5.0 | 8.0 | 16.0 | 17.0 | 15.0 | 19.0 | 8.0 | 9.0 | 10.0 | 10.0 |
| 18 | 16 | 21 | 14 | 21 | 15 | 23 | 21 | 34 | 23 | 60 | 27 |
| 16 | 16 | 26 | 15 | 35 | 16 | 33 | 26 | 63 | 26 | 76 | 31 |
| | | | | 2 | 2 | 2 | 2 | 3 | 2 | 3 | 2 |
| | | | | 5 | 5 | 5 | 5 | 7 | 5 | 20 | 5 |
| | | | | 12 | 10 | 10 | 13 | 10 | 15 | 40 | 20 |
| | | | | 25 | 20 | 30 | 25 | 30 | 30 | 81 | 40 |
| | | | | 40 | 35 | 55 | 50 | 82 | 55 | 131 | 55 |
| | | | | 60 | 45 | 71 | 70 | 121 | 81 | 161 | 68 |
| | | | | 140 | 76 | 130 | 140 | 242 | 101 | 364 | 176 |
| 41 | 36 | 47 | 29 | 29 | 25 | 34 | 27 | 33 | 32 | 87 | 33 |
| 34 | 29 | 65 | 39 | 31 | 23 | 41 | 27 | 28 | 42 | 70 | 27 |
| 48 | 29 | 22 | 26 | 31 | 23 | 34 | 30 | 28 | 33 | 60 | 39 |
| 37 | 17 | 17 | 21 | 32 | 21 | 27 | 28 | 21 | 28 | 52 | 30 |
| 14 | 37 | 43 | 29 | 39 | 30 | 42 | 38 | 77 | 38 | 97 | 44 |
| 4 | 31 | 35 | 12 | 56 | 23 | 48 | 34 | 96 | 33 | 102 | 40 |
| 121 | 121 | 242 | 180 | 484 | 165 | 484 | 240 | 726 | 242 | 545 | 218 |
| 240 | 240 | 360 | 180 | 1,119 | 334 | 1,452 | 625 | 1,634 | 489 | 1,091 | 529 |
| 30 | 33 | 36 | 21 | 25 | 19 | 24 | 23 | 42 | 25 | 53 | 30 |
| 33 | 29 | 46 | 25 | 30 | 16 | 31 | 24 | 57 | 28 | 55 | 34 |
| | | | | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| | | | | 10 | 10 | 10 | 10 | 15 | 10 | 15 | 15 |
| | | | | 15 | 15 | 15 | 15 | 15 | 15 | 30 | 15 |
| | | | | 30 | 30 | 30 | 30 | 30 | 30 | 66 | 30 |
| | | | | 45 | 30 | 45 | 53 | 121 | 60 | 121 | 61 |
| | | | | 61 | 60 | 61 | 61 | 182 | 66 | 182 | 121 |
| | | | | 151 | 75 | 121 | 121 | 242 | 126 | 187 | 169 |
| 121 | 121 | 242 | 180 | 242 | 121 | 484 | 182 | 303 | 242 | 242 | 197 |
| 2.0 | 5.2 | 1.7 | 2.9 | 12.6 | 9.6 | 17.1 | 41.5 | 2.5 | 3.1 | 0 | 9.2 |
| 4.8 | 18.4 | 2.3 | 3.6 | 1.4 | 1.1 | 1.0 | 2.1 | .8 | .6 | 0 | 2.9 |

TABLE 8.16.--LIQUID AND FROZEN CREAM SUBSTITUTES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FCOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 1.7 | 0 | 0 | .2 | .2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | .6 | 0 | 0 | .1 | .1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | 0 | 0 | .1 | 0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .7 | 0 | 0 | 0 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 3.1 | 0 | 0 | 1.6 | 1.8 |
| STANDARD DEVIATION.....DO..... | 2.7 | 0 | 0 | .5 | 1.0 |
| MAXIMUM.....DO..... | 16.0 | 0 | 0 | 2.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 22 | 0 | 0 | 52 | 58 |
| STANDARD DEVIATION.....GM..... | 28 | 0 | 0 | 17 | 53 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 2 | | | | |
| 25TH.....GM..... | 5 | | | | |
| 50TH.....GM..... | 10 | | | | |
| 75TH.....GM..... | 27 | | | | |
| 90TH.....GM..... | 47 | | | | |
| 95TH.....GM..... | 77 | | | | |
| 99TH.....GM..... | 150 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 34 | 0 | 0 | 91 | 40 |
| STANDARD DEVIATION.....GM..... | 52 | 0 | 0 | 0 | 22 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 27 | 0 | 0 | 96 | 0 |
| STANDARD DEVIATION.....GM..... | 29 | 0 | 0 | 10 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 33 | 0 | 0 | 0 | 120 |
| STANDARD DEVIATION.....GM..... | 36 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 600 | 0 | 0 | 120 | 120 |
| MAXIMUM OVER 3 DAYS.....GM..... | 600 | 0 | 0 | 211 | 360 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 21 | 0 | 0 | 95 | 95 |
| STANDARD DEVIATION.....GM..... | 29 | 0 | 0 | 14 | 39 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 5 | | | | |
| 25TH.....GM..... | 5 | | | | |
| 50TH.....GM..... | 10 | | | | |
| 75TH.....GM..... | 29 | | | | |
| 90TH.....GM..... | 60 | | | | |
| 95TH.....GM..... | 68 | | | | |
| 99TH.....GM..... | 120 | | | | |
| MAXIMUM.....GM..... | 240 | 0 | 0 | 120 | 120 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 43.1 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.2 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| (2) | .1 | .3 | .5 | 1.3 | 1.4 | 2.9 | 2.8 | 3.7 | 3.3 | 3.9 | 3.5 |
| (2) | (2) | .2 | .3 | .8 | .7 | 1.0 | .9 | 1.3 | 1.2 | .5 | .7 |
| 0 | 0 | .1 | .1 | .2 | .2 | .7 | .7 | 1.2 | .7 | 1.2 | 1.1 |
| 0 | .1 | 0 | .1 | .3 | .5 | 1.2 | 1.3 | 1.2 | 1.4 | 2.2 | 1.8 |
| 1.0 | 2.2 | 1.3 | 3.0 | 2.3 | 3.4 | 3.2 | 3.3 | 2.8 | 2.5 | 4.2 | 2.8 |
| .0 | 1.0 | .5 | 3.7 | 2.6 | 3.4 | 2.3 | 2.9 | 2.1 | 1.4 | 3.1 | 1.4 |
| 1.0 | 3.0 | 2.0 | 11.0 | 14.0 | 16.0 | 11.0 | 16.0 | 9.0 | 7.0 | 12.0 | 6.0 |
| 2 | 71 | 47 | 11 | 10 | 17 | 21 | 21 | 29 | 20 | 29 | 35 |
| 0 | 57 | 25 | 6 | 6 | 29 | 29 | 26 | 31 | 18 | 28 | 45 |
| | | | | 2 | 2 | 2 | 2 | | 2 | | |
| | | | | 5 | 5 | 5 | 5 | | 7 | | |
| | | | | 10 | 8 | 11 | 10 | | 17 | | |
| | | | | 15 | 18 | 25 | 29 | | 25 | | |
| | | | | 20 | 43 | 40 | 45 | | 40 | | |
| | | | | 20 | 60 | 75 | 68 | | 55 | | |
| | | | | 23 | 200 | 200 | 150 | | 120 | | |
| 5 | 15 | 97 | 27 | 24 | 45 | 29 | 30 | 36 | 47 | 11 | 13 |
| 0 | 0 | 39 | 19 | 16 | 106 | 36 | 33 | 19 | 44 | 6 | 8 |
| 0 | 0 | 120 | 15 | 17 | 32 | 24 | 17 | 27 | 27 | 39 | 58 |
| 0 | 0 | 0 | 0 | 9 | 45 | 20 | 15 | 22 | 11 | 37 | 37 |
| 0 | 120 | 0 | 18 | 15 | 19 | 33 | 34 | 58 | 25 | 37 | 44 |
| 0 | 0 | 0 | 0 | 3 | 16 | 40 | 33 | 36 | 23 | 30 | 58 |
| 5 | 120 | 120 | 60 | 60 | 600 | 240 | 240 | 180 | 120 | 90 | 240 |
| 5 | 360 | 240 | 60 | 68 | 600 | 600 | 450 | 375 | 360 | 269 | 600 |
| 5 | 99 | 108 | 11 | 13 | 15 | 20 | 19 | 32 | 24 | 21 | 37 |
| 0 | 42 | 31 | 13 | 10 | 22 | 31 | 24 | 31 | 28 | 20 | 50 |
| | | | | 3 | 3 | 5 | 5 | | 5 | | |
| | | | | 5 | 5 | 5 | 5 | | 10 | | |
| | | | | 10 | 5 | 11 | 10 | | 15 | | |
| | | | | 15 | 15 | 15 | 15 | | 30 | | |
| | | | | 30 | 43 | 45 | 60 | | 60 | | |
| | | | | 30 | 60 | 60 | 60 | | 120 | | |
| | | | | 60 | 120 | 180 | 120 | | 120 | | |
| 5 | 120 | 120 | 60 | 60 | 120 | 240 | 166 | 180 | 120 | 80 | 240 |
| 0 | .9 | 0 | 0 | 0 | .9 | 11.3 | 25.5 | 2.2 | 0 | 2.2 | 0 |
| 0 | 19.7 | 0 | 0 | 0 | .3 | 2.5 | 3.8 | 1.9 | 0 | 2.5 | 0 |

TABLE 8.17.--POWDERED CREAM SUBSTITUTES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|------------|------------|------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE | MALE | MALE | MALE |
| | | AND FEMALE | AND FEMALE | AND FEMALE | AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.3 | 0 | 0 | .2 | .4 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.2 | 0 | 0 | 0 | .4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.0 | 0 | 0 | .2 | 0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.1 | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 3.4 | 0 | 0 | 2.0 | 1.0 |
| STANDARD DEVIATION.....DO..... | 2.6 | 0 | 0 | .0 | .0 |
| MAXIMUM.....DO..... | 20.0 | 0 | 0 | 2.0 | 1.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 4 | 0 | 0 | 4 | 3 |
| STANDARD DEVIATION.....GM..... | 7 | 0 | 0 | 5 | 3 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 1 | | | | |
| 25TH.....GM..... | 1 | | | | |
| 50TH.....GM..... | 3 | | | | |
| 75TH.....GM..... | 5 | | | | |
| 90TH.....GM..... | 9 | | | | |
| 95TH.....GM..... | 12 | | | | |
| 99TH.....GM..... | 27 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 5 | 0 | 0 | 0 | 8 |
| STANDARD DEVIATION.....GM..... | 6 | 0 | 0 | 0 | 10 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 5 | 0 | 0 | 6 | 0 |
| STANDARD DEVIATION.....GM..... | 6 | 0 | 0 | 7 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 6 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 9 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 106 | 0 | 0 | 24 | 28 |
| MAXIMUM OVER 3 DAYS.....GM..... | 317 | 0 | 0 | 36 | 28 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 4 | 0 | 0 | 6 | 8 |
| STANDARD DEVIATION.....GM..... | 5 | 0 | 0 | 8 | 10 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 1 | | | | |
| 25TH.....GM..... | 2 | | | | |
| 50TH.....GM..... | 2 | | | | |
| 75TH.....GM..... | 4 | | | | |
| 90TH.....GM..... | 6 | | | | |
| 95TH.....GM..... | 12 | | | | |
| 99TH.....GM..... | 28 | | | | |
| MAXIMUM.....GM..... | 94 | 0 | 0 | 24 | 28 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 67.4 | 0 | 0 | 2.1 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.2 | 0 | 0 | 28.1 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| .2 | .2 | .6 | 1.7 | 4.6 | 5.0 | 6.6 | 7.4 | 4.8 | 8.3 | 5.6 | 6.6 |
| .2 | .1 | .4 | .6 | 1.5 | 1.9 | 1.8 | 1.9 | .8 | 1.5 | .8 | .9 |
| (2) | (2) | 0 | .4 | 1.3 | 1.1 | 1.6 | 1.6 | 1.0 | 1.9 | 1.1 | 1.2 |
| (2) | .1 | .1 | .6 | 1.8 | 2.0 | 3.3 | 3.9 | 3.1 | 4.9 | 3.6 | 4.5 |
| 1.5 | 2.2 | 1.7 | 2.6 | 3.1 | 3.1 | 3.4 | 3.6 | 3.7 | 3.6 | 3.8 | 4.3 |
| .8 | .9 | .8 | 1.8 | 2.7 | 2.6 | 2.4 | 2.7 | 2.1 | 2.9 | 2.7 | 2.6 |
| 3.0 | 3.0 | 3.0 | 7.0 | 20.0 | 12.0 | 16.0 | 20.0 | 9.0 | 17.0 | 12.0 | 10.0 |
| 2 | 2 | 2 | 2 | 4 | 3 | 4 | 4 | 4 | 4 | 4 | 11 |
| 1 | 1 | 1 | 1 | 5 | 4 | 4 | 5 | 3 | 6 | 3 | 24 |
| | | | | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| | | | | 1 | 1 | 1 | 2 | 1 | 1 | 1 | 2 |
| | | | | 2 | 2 | 3 | 3 | 3 | 2 | 2 | 4 |
| | | | | 5 | 4 | 6 | 5 | 5 | 5 | 5 | 9 |
| | | | | 10 | 8 | 10 | 9 | 8 | 9 | 9 | 14 |
| | | | | 12 | 10 | 14 | 12 | 9 | 12 | 12 | 57 |
| | | | | 28 | 20 | 22 | 24 | 12 | 36 | | 106 |
| 4 | 2 | 5 | 4 | 4 | 4 | 5 | 6 | 3 | 3 | 3 | 4 |
| 3 | 0 | 1 | 2 | 4 | 3 | 8 | 6 | 1 | 1 | 2 | 2 |
| 2 | 2 | 0 | 4 | 7 | 5 | 5 | 5 | 4 | 4 | 3 | 7 |
| 0 | 0 | 0 | 2 | 10 | 5 | 4 | 5 | 4 | 3 | 1 | 6 |
| 3 | 3 | 5 | 3 | 6 | 6 | 6 | 6 | 5 | 6 | 5 | 15 |
| 0 | 1 | (2) | 1 | 6 | 5 | 5 | 5 | 3 | 7 | 3 | 28 |
| 10 | 4 | 6 | 12 | 80 | 72 | 57 | 71 | 20 | 98 | 12 | 106 |
| 10 | 12 | 15 | 16 | 133 | 116 | 113 | 135 | 37 | 108 | 29 | 317 |
| 3 | 2 | 4 | 3 | 4 | 3 | 4 | 4 | 3 | 4 | 3 | 8 |
| 2 | 1 | 2 | 1 | 4 | 3 | 4 | 4 | 2 | 6 | 2 | 11 |
| | | | | 2 | 1 | 1 | 1 | 2 | 1 | | 2 |
| | | | | 2 | 2 | 2 | 2 | 2 | 2 | | 2 |
| | | | | 2 | 2 | 2 | 2 | 2 | 2 | | 3 |
| | | | | 6 | 4 | 4 | 4 | 4 | 4 | | 6 |
| | | | | 6 | 6 | 8 | 6 | 6 | 6 | | 35 |
| | | | | 12 | 8 | 12 | 9 | 6 | 8 | | 35 |
| | | | | 18 | 18 | 18 | 24 | 12 | 12 | | 35 |
| 10 | 4 | 6 | 6 | 28 | 47 | 28 | 53 | 18 | 94 | 8 | 35 |
| 0 | 0 | 0 | 0 | 5.0 | 27.1 | 19.3 | 8.3 | 0 | 4.7 | 0 | .8 |
| 0 | 0 | 0 | 0 | .9 | 3.2 | 1.7 | .4 | 0 | .9 | 0 | .3 |

TABLE 8.18.--ICE CREAM AND ICE MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 23.6 | 3.7 | 24.0 | 29.8 | 30.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 15.9 | 3.3 | 16.9 | 19.1 | 21.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 5.6 | .3 | 5.8 | 8.5 | 7.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.0 | .1 | 1.3 | 2.1 | 2.5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.5 | 1.1 | 1.4 | 1.5 | 1.5 |
| STANDARD DEVIATION.....DO..... | .8 | .4 | .7 | .7 | .7 |
| MAXIMUM.....DO..... | 7.0 | 3.0 | 4.0 | 5.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 57 | 17 | 32 | 43 | 50 |
| STANDARD DEVIATION.....GM..... | 45 | 14 | 23 | 28 | 34 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 15 | | 11 | 11 | 17 |
| 25TH.....GM..... | 22 | | 22 | 22 | 22 |
| 50TH.....GM..... | 44 | | 22 | 44 | 44 |
| 75TH.....GM..... | 67 | | 44 | 44 | 66 |
| 90TH.....GM..... | 111 | | 64 | 89 | 89 |
| 95TH.....GM..... | 133 | | 89 | 89 | 122 |
| 99TH.....GM..... | 222 | | 133 | 133 | 177 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 118 | 47 | 72 | 94 | 107 |
| STANDARD DEVIATION.....GM..... | 70 | 42 | 35 | 50 | 54 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 122 | 47 | 69 | 87 | 102 |
| STANDARD DEVIATION.....GM..... | 68 | 0 | 35 | 41 | 44 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 129 | 33 | 79 | 85 | 114 |
| STANDARD DEVIATION.....GM..... | 68 | 0 | 44 | 33 | 47 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,064 | 133 | 233 | 360 | 399 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,596 | 133 | 466 | 532 | 798 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 116 | | 69 | 86 | 101 |
| STANDARD DEVIATION.....GM..... | 66 | | 36 | 43 | 47 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 33 | | 17 | 33 | 33 |
| 25TH.....GM..... | 67 | | 44 | 67 | 67 |
| 50TH.....GM..... | 133 | | 67 | 67 | 100 |
| 75TH.....GM..... | 133 | | 67 | 133 | 133 |
| 90TH.....GM..... | 200 | | 133 | 133 | 133 |
| 95TH.....GM..... | 266 | | 133 | 133 | 175 |
| 99TH.....GM..... | 333 | | 133 | 266 | 266 |
| MAXIMUM.....GM..... | 532 | 133 | 200 | 266 | 267 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 93.2 | 0 | 2.4 | 1.7 | 8.4 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .7 | 0 | .7 | .2 | 1.0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 31.5 | 30.1 | 27.1 | 23.6 | 19.9 | 18.8 | 26.4 | 19.4 | 27.6 | 21.9 | 21.3 | 27.7 |
| 20.1 | 20.2 | 16.6 | 16.8 | 14.7 | 14.3 | 17.1 | 13.7 | 15.9 | 14.0 | 9.7 | 15.3 |
| 8.2 | 7.7 | 8.4 | 5.8 | 3.9 | 3.5 | 6.4 | 4.3 | 7.1 | 5.2 | 7.8 | 8.4 |
| 3.2 | 2.2 | 2.0 | 1.0 | 1.3 | 1.1 | 2.9 | 1.4 | 4.6 | 2.6 | 3.9 | 4.0 |
| 1.6 | 1.5 | 1.5 | 1.4 | 1.4 | 1.4 | 1.5 | 1.4 | 1.7 | 1.5 | 1.9 | 1.7 |
| .8 | .8 | .8 | .7 | .8 | .7 | .8 | .7 | .9 | .8 | 1.0 | 1.0 |
| 7.0 | 5.0 | 5.0 | 4.0 | 7.0 | 5.0 | 5.0 | 5.0 | 6.0 | 5.0 | 6.0 | 6.0 |
| 68 | 58 | 82 | 57 | 67 | 50 | 67 | 49 | 64 | 52 | 66 | 53 |
| 49 | 43 | 70 | 47 | 50 | 35 | 53 | 35 | 49 | 41 | 40 | 42 |
| 18 | 17 | 22 | 17 | 19 | 15 | 17 | 11 | 15 | 11 | 22 | 11 |
| 37 | 28 | 44 | 22 | 33 | 22 | 33 | 22 | 23 | 22 | 44 | 22 |
| 44 | 44 | 67 | 44 | 44 | 44 | 44 | 44 | 44 | 44 | 55 | 44 |
| 89 | 67 | 89 | 67 | 89 | 67 | 89 | 67 | 89 | 67 | 89 | 67 |
| 133 | 100 | 155 | 100 | 133 | 89 | 133 | 89 | 133 | 100 | 133 | 111 |
| 177 | 133 | 200 | 133 | 156 | 122 | 155 | 124 | 155 | 133 | 133 | 133 |
| 244 | 222 | 355 | 222 | 266 | 177 | 266 | 177 | 266 | 222 | 222 | 222 |
| 139 | 121 | 149 | 121 | 145 | 112 | 132 | 109 | 114 | 102 | 121 | 89 |
| 82 | 69 | 77 | 66 | 81 | 58 | 77 | 67 | 66 | 51 | 64 | 46 |
| 142 | 129 | 168 | 133 | 160 | 119 | 140 | 104 | 109 | 98 | 126 | 99 |
| 71 | 73 | 93 | 64 | 86 | 53 | 76 | 45 | 49 | 48 | 52 | 48 |
| 134 | 116 | 216 | 155 | 153 | 116 | 144 | 114 | 140 | 119 | 99 | 114 |
| 66 | 58 | 120 | 111 | 62 | 52 | 72 | 47 | 55 | 61 | 45 | 54 |
| 665 | 665 | 1,064 | 532 | 532 | 532 | 532 | 798 | 399 | 333 | 266 | 532 |
| 1,131 | 1,032 | 1,596 | 1,264 | 1,064 | 931 | 1,197 | 798 | 798 | 732 | 665 | 798 |
| 130 | 116 | 159 | 123 | 142 | 110 | 131 | 104 | 115 | 100 | 103 | 92 |
| 68 | 60 | 92 | 75 | 75 | 57 | 73 | 57 | 60 | 52 | 54 | 47 |
| 50 | 34 | 67 | 44 | 50 | 33 | 50 | 33 | 33 | 28 | 33 | 33 |
| 67 | 67 | 100 | 67 | 82 | 67 | 67 | 67 | 67 | 67 | 67 | 67 |
| 133 | 133 | 133 | 133 | 133 | 133 | 133 | 100 | 131 | 67 | 131 | 67 |
| 133 | 133 | 200 | 133 | 141 | 133 | 133 | 133 | 133 | 133 | 133 | 133 |
| 262 | 200 | 266 | 200 | 266 | 134 | 266 | 133 | 200 | 133 | 133 | 133 |
| 266 | 266 | 357 | 266 | 266 | 227 | 266 | 200 | 266 | 200 | 200 | 133 |
| 393 | 266 | 532 | 402 | 399 | 266 | 399 | 266 | 266 | 266 | 266 | 266 |
| 399 | 399 | 532 | 532 | 532 | 532 | 532 | 532 | 399 | 266 | 266 | 266 |
| 5.5 | 10.8 | 3.0 | 1.8 | 9.0 | 14.0 | 9.1 | 8.2 | 7.1 | 7.9 | 0 | 4.3 |
| .5 | 1.1 | .5 | .4 | .8 | 1.0 | .5 | .4 | 1.4 | 1.4 | 0 | .9 |

TABLE 8.19.--NONDAIRY WHIPPED TOPPING
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.4 | .2 | 1.3 | 1.8 | 2.0 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.0 | .2 | 1.3 | 1.7 | 1.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .3 | 0 | 0 | .1 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 1.2 | 1.0 | 1.0 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .5 | .0 | .0 | .4 | .4 |
| MAXIMUM.....DO..... | 4.0 | 1.0 | 1.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 5 | 3 | 3 | 4 | 4 |
| STANDARD DEVIATION.....GM..... | 7 | 0 | 2 | 3 | 4 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 1 | | | | |
| 25TH.....GM..... | 2 | | | | |
| 50TH.....GM..... | 3 | | | | |
| 75TH.....GM..... | 6 | | | | |
| 90TH.....GM..... | 10 | | | | |
| 95TH.....GM..... | 13 | | | | |
| 99TH.....GM..... | 31 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 12 | 9 | 9 | 11 | 11 |
| STANDARD DEVIATION.....GM..... | 12 | 0 | 7 | 8 | 12 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 15 | 0 | 0 | 9 | 7 |
| STANDARD DEVIATION.....GM..... | 22 | 0 | 0 | 7 | 5 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 17 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 9 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 160 | 9 | 25 | 38 | 57 |
| MAXIMUM OVER 3 DAYS.....GM..... | 320 | 9 | 25 | 38 | 57 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 12 | 9 | 9 | 11 | 10 |
| STANDARD DEVIATION.....GM..... | 15 | 0 | 7 | 8 | 11 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 2 | | | | |
| 25TH.....GM..... | 5 | | | | |
| 50TH.....GM..... | 9 | | | | |
| 75TH.....GM..... | 14 | | | | |
| 90TH.....GM..... | 20 | | | | |
| 95TH.....GM..... | 38 | | | | |
| 99TH.....GM..... | 75 | | | | |
| MAXIMUM.....GM..... | 160 | | | | |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 168.3 | 0 | 1.8 | 2.5 | 4.2 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 15.6 | 0 | 12.6 | 7.5 | 10.2 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 2.1 | 2.6 | 1.9 | 2.1 | 2.0 | 1.7 | 2.2 | 2.9 | 3.6 | 4.0 | 3.7 | 4.5 |
| 1.9 | 2.5 | 1.4 | 1.8 | 1.8 | 1.5 | 1.8 | 2.5 | 3.0 | 2.8 | 1.6 | 3.2 |
| .2 | .1 | .5 | .2 | .2 | .1 | .3 | .3 | .6 | 1.1 | 1.7 | 1.2 |
| 0 | 0 | 0 | .1 | 0 | (2) | .1 | (2) | 0 | .1 | .5 | 0 |
| 1.1 | 1.0 | 1.3 | 1.2 | 1.1 | 1.1 | 1.3 | 1.1 | 1.3 | 1.4 | 1.8 | 1.3 |
| .3 | .2 | .6 | .5 | .3 | .3 | .7 | .4 | .5 | .6 | .8 | .7 |
| 2.0 | 2.0 | 3.0 | 3.0 | 2.0 | 3.0 | 4.0 | 3.0 | 3.0 | 4.0 | 4.0 | 4.0 |
| 3 | 4 | 7 | 5 | 7 | 3 | 7 | 5 | 6 | 5 | 8 | 3 |
| 2 | 3 | 7 | 5 | 6 | 3 | 12 | 9 | 8 | 6 | 5 | 2 |
| | 1 | | | 1 | 1 | 1 | 1 | | 1 | | |
| | 2 | | | 3 | 2 | 3 | 3 | | 2 | | |
| | 3 | | | 6 | 3 | 3 | 3 | | 3 | | |
| | 5 | | | 7 | 3 | 6 | 5 | | 6 | | |
| | 6 | | | 13 | 7 | 14 | 9 | | 10 | | |
| | 7 | | | 18 | 11 | 17 | 16 | | 13 | | |
| | 13 | | | 35 | 13 | 38 | 41 | | 47 | | |
| 8 | 11 | 16 | 11 | 18 | 10 | 13 | 12 | 11 | 11 | 17 | 9 |
| 6 | 8 | 13 | 7 | 17 | 8 | 17 | 13 | 7 | 8 | 12 | 5 |
| 7 | 7 | 16 | 10 | 18 | 9 | 25 | 21 | 26 | 9 | 12 | 7 |
| 3 | 2 | 15 | 2 | 9 | 3 | 38 | 34 | 21 | 6 | 5 | 5 |
| 0 | 0 | 0 | 30 | 0 | 11 | 14 | 19 | 0 | 21 | 15 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 5 | 5 | 0 | 18 | 1 | 0 |
| 30 | 38 | 56 | 38 | 106 | 40 | 160 | 160 | 75 | 75 | 38 | 23 |
| 30 | 38 | 94 | 89 | 106 | 40 | 320 | 320 | 131 | 141 | 50 | 38 |
| 8 | 10 | 15 | 12 | 17 | 9 | 15 | 14 | 14 | 11 | 13 | 8 |
| 6 | 8 | 12 | 9 | 13 | 8 | 22 | 19 | 15 | 9 | 8 | 6 |
| | 3 | | | 4 | 2 | 4 | 2 | | 2 | | |
| | 5 | | | 9 | 5 | 8 | 5 | | 5 | | |
| | 9 | | | 16 | 9 | 9 | 9 | | 9 | | |
| | 10 | | | 20 | 9 | 14 | 14 | | 15 | | |
| | 19 | | | 38 | 20 | 20 | 25 | | 19 | | |
| | 21 | | | 38 | 28 | 38 | 40 | | 26 | | |
| | 38 | | | 80 | 40 | 160 | 75 | | 38 | | |
| | 38 | 42 | 38 | 80 | 40 | 160 | 160 | 75 | 75 | 38 | 23 |
| 8.7 | 8.3 | 3.2 | 7.1 | 11.9 | 13.0 | 28.1 | 42.6 | 14.3 | 7.3 | 10.4 | 4.9 |
| 18.0 | 14.0 | 9.4 | 19.7 | 13.2 | 13.4 | 19.3 | 18.2 | 27.7 | 7.5 | 29.2 | 8.3 |

TABLE 8.20.--SOUR CREAM
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.9 | .2 | 1.1 | 1.2 | 1.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.6 | .2 | 1.1 | 1.2 | 1.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .2 | 0 | 0 | .1 | 0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.1 | 1.0 | 1.0 | 1.0 | 1.0 |
| STANDARD DEVIATION.....CO..... | .4 | .0 | .0 | .2 | .0 |
| MAXIMUM.....DO..... | 4.0 | 1.0 | 1.0 | 2.0 | 1.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 13 | 2 | 3 | 8 | 7 |
| STANDARD DEVIATION.....GM..... | 16 | 0 | 2 | 9 | 8 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 2 | | | | |
| 25TH.....GM..... | 4 | | | | |
| 50TH.....GM..... | 8 | | | | |
| 75TH.....GM..... | 16 | | | | |
| 90TH.....GM..... | 29 | | | | |
| 95TH.....GM..... | 38 | | | | |
| 99TH.....GM..... | 77 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 33 | 5 | 10 | 24 | 20 |
| STANDARD DEVIATION.....GM..... | 35 | 0 | 6 | 27 | 24 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 41 | 0 | 0 | 24 | 0 |
| STANDARD DEVIATION.....GM..... | 36 | 0 | 0 | 0 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 72 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 67 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 345 | 5 | 24 | 115 | 115 |
| MAXIMUM OVER 3 DAYS.....GM..... | 690 | 5 | 24 | 115 | 115 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 35 | 5 | 10 | 24 | 20 |
| STANDARD DEVIATION.....GM..... | 38 | 0 | 6 | 26 | 24 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 6 | | | | |
| 25TH.....GM..... | 12 | | | | |
| 50TH.....GM..... | 24 | | | | |
| 75TH.....GM..... | 36 | | | | |
| 90TH.....GM..... | 72 | | | | |
| 95TH.....GM..... | 115 | | | | |
| 99TH.....GM..... | 230 | | | | |
| MAXIMUM.....GM..... | 345 | 5 | 24 | 115 | 115 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 58.6 | 0 | 1.2 | 3.3 | 2.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 4.8 | 0 | 9.7 | 14.8 | 8.6 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.5 | 1.3 | 2.1 | 1.7 | 3.1 | 3.9 | 3.8 | 4.0 | 2.7 | 4.0 | 1.5 | 1.9 |
| 1.3 | 1.2 | 2.1 | 1.7 | 2.8 | 3.6 | 3.5 | 3.6 | 2.5 | 3.3 | 1.5 | 1.7 |
| .1 | .1 | 0 | 0 | .3 | .3 | .2 | .3 | .2 | .5 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | (2) | .1 | 0 | .2 | 0 | .2 |
| 1.1 | 1.1 | 1.0 | 1.0 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.3 | 1.0 | 1.2 |
| .3 | .3 | .0 | .0 | .3 | .3 | .4 | .4 | .3 | .7 | .0 | .6 |
| 2.0 | 2.0 | 1.0 | 1.0 | 2.0 | 2.0 | 4.0 | 3.0 | 2.0 | 4.0 | 1.0 | 3.0 |
| 18 | 12 | 15 | 13 | 14 | 12 | 15 | 11 | 11 | 18 | 7 | 15 |
| 15 | 16 | 11 | 9 | 15 | 12 | 18 | 19 | 8 | 22 | 2 | 18 |
| | | | | 3 | 3 | 4 | 2 | | 3 | | |
| | | | | 4 | 4 | 4 | 4 | | 4 | | |
| | | | | 8 | 8 | 8 | 8 | | 8 | | |
| | | | | 16 | 14 | 16 | 12 | | 19 | | |
| | | | | 38 | 27 | 38 | 20 | | 54 | | |
| | | | | 38 | 38 | 40 | 29 | | 57 | | |
| | | | | 85 | 77 | 115 | 66 | | 115 | | |
| 50 | 31 | 44 | 38 | 35 | 35 | 41 | 26 | 30 | 35 | 21 | 29 |
| 42 | 45 | 34 | 26 | 31 | 37 | 51 | 23 | 21 | 30 | 7 | 30 |
| 50 | 49 | 0 | 0 | 49 | 27 | 48 | 30 | 41 | 74 | 0 | 0 |
| 20 | 20 | 0 | 0 | 45 | 15 | 35 | 23 | 0 | 56 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 40 | 113 | 0 | 48 | 0 | 57 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 97 | 0 | 15 | 0 | 0 |
| 154 | 230 | 115 | 115 | 230 | 230 | 345 | 230 | 115 | 230 | 24 | 115 |
| 154 | 230 | 115 | 115 | 254 | 230 | 345 | 690 | 115 | 345 | 24 | 170 |
| 50 | 32 | 44 | 38 | 38 | 33 | 41 | 30 | 31 | 42 | 21 | 37 |
| 39 | 33 | 34 | 26 | 41 | 34 | 47 | 35 | 21 | 42 | 7 | 29 |
| | | | | 10 | 7 | 12 | 6 | | 10 | | |
| | | | | 12 | 12 | 12 | 12 | | 12 | | |
| | | | | 24 | 24 | 24 | 24 | | 24 | | |
| | | | | 36 | 36 | 58 | 28 | | 58 | | |
| | | | | 87 | 58 | 87 | 58 | | 115 | | |
| | | | | 115 | 115 | 115 | 76 | | 115 | | |
| | | | | 230 | 151 | 230 | 230 | | 230 | | |
| 154 | 115 | 115 | 115 | 230 | 230 | 345 | 230 | 115 | 230 | 24 | 115 |
| 3.7 | 2.3 | 4.0 | 1.0 | 7.0 | 12.5 | 4.8 | 9.5 | 0 | 5.4 | 0 | 1.8 |
| 11.1 | 7.5 | 13.7 | 4.3 | 5.1 | 5.3 | 2.4 | 3.0 | 0 | 5.9 | 0 | 8.0 |

TABLE 8.21.--YOGURT
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.2 | 2.9 | 3.8 | 2.9 | 2.4 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 1.9 | .6 | 2.7 | 1.9 | 1.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .9 | 1.3 | .8 | .7 | .6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .4 | 1.1 | .3 | .3 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.6 | 2.4 | 1.4 | 1.6 | 1.3 |
| STANDARD DEVIATION.....DO..... | .9 | 1.1 | .7 | .7 | .5 |
| MAXIMUM.....DO..... | 6.0 | 5.0 | 3.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 99 | 62 | 55 | 71 | 74 |
| STANDARD DEVIATION.....GM..... | 76 | 57 | 46 | 54 | 54 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 16 | | | 20 | |
| 25TH.....GM..... | 41 | | | 38 | |
| 50TH.....GM..... | 82 | | | 57 | |
| 75TH.....GM..... | 143 | | | 82 | |
| 90TH.....GM..... | 204 | | | 143 | |
| 95TH.....GM..... | 245 | | | 163 | |
| 99TH.....GM..... | 378 | | | 306 | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 187 | 139 | 122 | 168 | 150 |
| STANDARD DEVIATION.....GM..... | 102 | 64 | 82 | 105 | 78 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 198 | 48 | 149 | 146 | 221 |
| STANDARD DEVIATION.....GM..... | 94 | 25 | 103 | 59 | 74 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 198 | 106 | 70 | 101 | 0 |
| STANDARD DEVIATION.....GM..... | 115 | 71 | 20 | 109 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 735 | 245 | 490 | 490 | 613 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,593 | 681 | 735 | 918 | 735 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 182 | 78 | 117 | 137 | 173 |
| STANDARD DEVIATION.....GM..... | 94 | 66 | 91 | 84 | 82 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 31 | | | 31 | |
| 25TH.....GM..... | 123 | | | 61 | |
| 50TH.....GM..... | 227 | | | 123 | |
| 75TH.....GM..... | 245 | | | 184 | |
| 90TH.....GM..... | 245 | | | 245 | |
| 95TH.....GM..... | 245 | | | 245 | |
| 99TH.....GM..... | 490 | | | 490 | |
| MAXIMUM.....GM..... | 613 | 245 | 490 | 490 | 368 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 2.4 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .1 | 0 | 0 | 0 | 0 |

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.4 | 1.6 | 2.0 | 2.5 | 3.5 | 5.0 | 2.0 | 4.3 | 1.6 | 4.3 | 0 | 2.5 |
| 1.2 | 1.3 | 1.5 | 1.6 | 2.1 | 2.9 | 1.0 | 2.5 | 1.0 | 2.4 | 0 | 1.2 |
| .2 | .3 | .3 | .8 | .9 | 1.4 | .6 | 1.3 | .1 | 1.4 | 0 | .7 |
| 0 | 0 | .2 | .1 | .5 | .7 | .4 | .6 | .5 | .5 | 0 | .6 |
| 1.3 | 1.3 | 1.4 | 1.4 | 1.6 | 1.7 | 1.8 | 1.6 | 1.9 | 1.6 | 0 | 1.8 |
| .5 | .5 | .7 | .6 | .8 | 1.0 | 1.0 | .8 | 1.0 | .8 | 0 | .9 |
| 3.0 | 3.0 | 3.0 | 3.0 | 6.0 | 6.0 | 5.0 | 4.0 | 4.0 | 6.0 | 0 | 3.0 |
| 81 | 88 | 109 | 98 | 112 | 107 | 118 | 94 | 141 | 106 | 0 | 82 |
| 38 | 59 | 55 | 69 | 83 | 78 | 77 | 73 | 84 | 107 | 0 | 64 |
| | | | | 27 | 20 | 27 | 15 | | 7 | | |
| | | | | 61 | 57 | 76 | 41 | | 41 | | |
| | | | | 82 | 82 | 82 | 82 | | 76 | | |
| | | | | 157 | 163 | 163 | 123 | | 151 | | |
| | | | | 245 | 206 | 227 | 163 | | 245 | | |
| | | | | 265 | 245 | 245 | 245 | | 327 | | |
| | | | | 327 | 378 | 490 | 327 | | 490 | | |
| 204 | 196 | 257 | 188 | 190 | 197 | 214 | 173 | 362 | 174 | 0 | 161 |
| 58 | 110 | 107 | 79 | 72 | 93 | 109 | 92 | 246 | 123 | 0 | 81 |
| 231 | 259 | 215 | 224 | 230 | 211 | 205 | 187 | 64 | 188 | 0 | 183 |
| 44 | 77 | 20 | 108 | 90 | 70 | 80 | 97 | 0 | 136 | 0 | 56 |
| 0 | 0 | 223 | 245 | 240 | 216 | 192 | 189 | 199 | 292 | 0 | 94 |
| 0 | 0 | 17 | 0 | 93 | 123 | 98 | 104 | 55 | 156 | 0 | 102 |
| 454 | 490 | 454 | 490 | 613 | 735 | 554 | 490 | 735 | 735 | 0 | 306 |
| 624 | 735 | 717 | 980 | 1,593 | 1,516 | 1,470 | 1,470 | 735 | 1,470 | 0 | 674 |
| 191 | 209 | 234 | 206 | 209 | 189 | 192 | 173 | 219 | 193 | 0 | 136 |
| 61 | 94 | 74 | 91 | 87 | 83 | 93 | 88 | 124 | 130 | 0 | 91 |
| | | | | 61 | 46 | 46 | 29 | | 29 | | |
| | | | | 123 | 123 | 123 | 114 | | 114 | | |
| | | | | 227 | 227 | 227 | 184 | | 184 | | |
| | | | | 245 | 245 | 245 | 245 | | 245 | | |
| | | | | 245 | 245 | 245 | 245 | | 490 | | |
| | | | | 368 | 245 | 245 | 245 | | 490 | | |
| | | | | 490 | 490 | 554 | 490 | | 490 | | |
| 245 | 490 | 454 | 490 | 490 | 613 | 554 | 490 | 490 | 490 | 0 | 245 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2.4 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | .5 | 0 | 0 | 0 | 0 |

TABLE 9.01.--WHITE SUGAR
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 43.6 | 8.8 | 37.1 | 42.3 | 43.0 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 14.3 | 5.3 | 16.7 | 18.9 | 18.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 10.7 | 1.7 | 12.0 | 13.7 | 13.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 18.7 | 1.8 | 8.4 | 9.8 | 11.4 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 3.3 | 1.8 | 2.1 | 2.2 | 2.1 |
| STANDARD DEVIATION.....DO..... | 2.7 | 1.1 | 1.4 | 1.7 | 1.3 |
| MAXIMUM.....DO..... | 26.0 | 6.0 | 11.0 | 18.0 | 10.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 10 | 3 | 5 | 6 | 6 |
| STANDARD DEVIATION.....GM..... | 12 | 3 | 5 | 7 | 6 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 1 | | 1 | 1 | 1 |
| 25TH.....GM..... | 3 | | 1 | 3 | 3 |
| 50TH.....GM..... | 6 | | 3 | 4 | 4 |
| 75TH.....GM..... | 13 | | 6 | 7 | 8 |
| 90TH.....GM..... | 21 | | 10 | 13 | 14 |
| 95TH.....GM..... | 30 | | 13 | 17 | 19 |
| 99TH.....GM..... | 55 | | 29 | 38 | 33 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 11 | 6 | 7 | 10 | 9 |
| STANDARD DEVIATION.....GM..... | 10 | 6 | 7 | 12 | 7 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 12 | 8 | 8 | 9 | 11 |
| STANDARD DEVIATION.....GM..... | 11 | 6 | 7 | 6 | 9 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 16 | 3 | 8 | 12 | 11 |
| STANDARD DEVIATION.....GM..... | 15 | 2 | 7 | 11 | 8 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 405 | 38 | 75 | 150 | 102 |
| MAXIMUM OVER 3 DAYS.....GM..... | 980 | 42 | 125 | 213 | 150 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 9 | 5 | 7 | 8 | 9 |
| STANDARD DEVIATION.....GM..... | 9 | 5 | 6 | 8 | 8 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 2 | | 2 | 2 | 4 |
| 25TH.....GM..... | 4 | | 4 | 4 | 4 |
| 50TH.....GM..... | 8 | | 4 | 6 | 8 |
| 75TH.....GM..... | 8 | | 8 | 8 | 8 |
| 90TH.....GM..... | 17 | | 13 | 13 | 13 |
| 95TH.....GM..... | 25 | | 13 | 25 | 25 |
| 99TH.....GM..... | 50 | | 28 | 38 | 38 |
| MAXIMUM.....GM..... | 350 | 38 | 50 | 150 | 100 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 4284.7 | 6.9 | 71.6 | 193.2 | 145.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 8.0 | 8.6 | 8.8 | 12.1 | 8.7 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 44.6 | 41.4 | 43.4 | 39.0 | 45.8 | 44.4 | 49.7 | 41.1 | 50.8 | 43.5 | 56.0 | 41.0 |
| 19.6 | 19.1 | 19.2 | 17.8 | 14.5 | 14.8 | 11.1 | 11.6 | 10.5 | 11.2 | 10.9 | 8.9 |
| 13.9 | 13.4 | 11.9 | 11.8 | 10.7 | 10.9 | 10.3 | 8.2 | 11.8 | 9.0 | 11.8 | 8.0 |
| 11.1 | 9.0 | 12.3 | 9.4 | 20.6 | 18.8 | 28.3 | 21.3 | 28.6 | 23.2 | 33.3 | 24.0 |
| 2.1 | 2.1 | 2.3 | 2.3 | 3.4 | 3.3 | 4.3 | 3.8 | 3.9 | 3.7 | 4.1 | 3.7 |
| 1.4 | 1.3 | 1.6 | 1.7 | 2.8 | 2.8 | 3.2 | 2.9 | 2.8 | 2.9 | 2.7 | 2.4 |
| 9.0 | 12.0 | 16.0 | 12.0 | 24.0 | 26.0 | 21.0 | 23.0 | 17.0 | 23.0 | 13.0 | 14.0 |
| 8 | 7 | 10 | 8 | 12 | 9 | 13 | 10 | 13 | 9 | 12 | 9 |
| 9 | 7 | 11 | 9 | 15 | 11 | 14 | 15 | 13 | 9 | 11 | 9 |
| 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 3 | 3 | 3 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 |
| 6 | 4 | 6 | 5 | 8 | 6 | 8 | 7 | 8 | 6 | 8 | 7 |
| 9 | 8 | 13 | 8 | 15 | 11 | 17 | 13 | 16 | 11 | 15 | 11 |
| 17 | 15 | 21 | 17 | 25 | 19 | 28 | 22 | 31 | 18 | 25 | 18 |
| 24 | 19 | 29 | 26 | 36 | 28 | 41 | 29 | 39 | 25 | 35 | 25 |
| 42 | 34 | 56 | 38 | 75 | 53 | 67 | 54 | 67 | 44 | 47 | 50 |
| 12 | 10 | 14 | 11 | 14 | 11 | 11 | 9 | 12 | 9 | 10 | 9 |
| 10 | 9 | 15 | 14 | 13 | 9 | 8 | 8 | 14 | 9 | 10 | 9 |
| 12 | 12 | 16 | 13 | 14 | 12 | 12 | 10 | 11 | 8 | 11 | 10 |
| 10 | 11 | 16 | 11 | 14 | 15 | 11 | 9 | 7 | 6 | 9 | 7 |
| 14 | 12 | 17 | 15 | 19 | 15 | 19 | 16 | 18 | 13 | 16 | 12 |
| 15 | 8 | 14 | 12 | 19 | 12 | 15 | 19 | 15 | 10 | 11 | 10 |
| 225 | 150 | 150 | 188 | 350 | 228 | 240 | 405 | 125 | 100 | 106 | 75 |
| 425 | 179 | 350 | 304 | 700 | 400 | 363 | 980 | 271 | 171 | 190 | 156 |
| 11 | 10 | 13 | 10 | 11 | 9 | 9 | 8 | 10 | 7 | 8 | 7 |
| 10 | 8 | 13 | 8 | 13 | 8 | 8 | 8 | 9 | 6 | 6 | 6 |
| 4 | 4 | 4 | 4 | 4 | 3 | 3 | 2 | 4 | 2 | 2 | 2 |
| 4 | 4 | 8 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 4 | 8 | 4 |
| 13 | 13 | 13 | 13 | 13 | 8 | 8 | 8 | 12 | 8 | 8 | 8 |
| 25 | 17 | 25 | 21 | 21 | 16 | 17 | 15 | 17 | 13 | 17 | 13 |
| 25 | 25 | 38 | 25 | 25 | 24 | 25 | 25 | 25 | 17 | 25 | 25 |
| 50 | 50 | 66 | 50 | 50 | 38 | 38 | 38 | 50 | 38 | 32 | 32 |
| 100 | 100 | 113 | 100 | 350 | 170 | 200 | 170 | 100 | 69 | 50 | 50 |
| 186.4 | 148.4 | 161.0 | 127.2 | 583.8 | 643.7 | 857.6 | 761.0 | 115.1 | 150.9 | 19.3 | 113.7 |
| 9.3 | 8.0 | 11.7 | 9.5 | 9.7 | 8.1 | 8.2 | 6.9 | 5.2 | 5.4 | 1.6 | 7.6 |

TABLE 9.02.--SIRUP
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 10.8 | 4.0 | 14.5 | 18.2 | 17.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 9.1 | 2.8 | 12.9 | 15.4 | 14.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.3 | .3 | 1.3 | 2.2 | 2.4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .4 | .9 | .3 | .5 | .8 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.2 | 3.2 | 1.1 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .6 | 4.3 | .4 | .5 | .5 |
| MAXIMUM.....DO..... | 17.0 | 17.0 | 4.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 28 | 21 | 15 | 22 | 25 |
| STANDARD DEVIATION.....GM..... | 22 | 22 | 13 | 17 | 19 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 7 | | 2 | 4 | 7 |
| 25TH.....GM..... | 13 | | 7 | 13 | 13 |
| 50TH.....GM..... | 26 | | 13 | 17 | 26 |
| 75TH.....GM..... | 31 | | 26 | 27 | 27 |
| 90TH.....GM..... | 53 | | 27 | 40 | 53 |
| 95TH.....GM..... | 62 | | 40 | 53 | 62 |
| 99TH.....GM..... | 113 | | 55 | 91 | 105 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 71 | 28 | 41 | 55 | 66 |
| STANDARD DEVIATION.....GM..... | 49 | 30 | 33 | 34 | 47 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 67 | 83 | 36 | 47 | 52 |
| STANDARD DEVIATION.....GM..... | 45 | 37 | 22 | 24 | 31 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 68 | 46 | 40 | 83 | 61 |
| STANDARD DEVIATION.....GM..... | 43 | 15 | 34 | 32 | 23 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 630 | 178 | 160 | 164 | 320 |
| MAXIMUM OVER 3 DAYS.....GM..... | 788 | 232 | 261 | 473 | 320 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 69 | 20 | 40 | 55 | 62 |
| STANDARD DEVIATION.....GM..... | 48 | 18 | 32 | 32 | 42 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 13 | | 7 | 13 | 20 |
| 25TH.....GM..... | 39 | | 20 | 38 | 39 |
| 50TH.....GM..... | 62 | | 38 | 40 | 59 |
| 75TH.....GM..... | 80 | | 59 | 79 | 79 |
| 90TH.....GM..... | 158 | | 80 | 82 | 104 |
| 95TH.....GM..... | 160 | | 82 | 104 | 158 |
| 99TH.....GM..... | 236 | | 158 | 160 | 197 |
| MAXIMUM.....GM..... | 630 | 82 | 160 | 160 | 320 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 282.2 | 8.8 | 6.6 | 23.3 | 33.0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 5.7 | 13.5 | 3.8 | 6.2 | 8.1 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 18.4 | 14.6 | 14.6 | 12.4 | 10.5 | 9.5 | 10.1 | 6.7 | 7.8 | 6.4 | 7.8 | 5.9 |
| 14.6 | 12.4 | 11.9 | 10.6 | 8.8 | 8.6 | 8.3 | 5.9 | 5.6 | 5.0 | 6.0 | 5.1 |
| 2.9 | 1.7 | 2.0 | 1.5 | 1.3 | .8 | 1.4 | .6 | 1.5 | 1.1 | .7 | .7 |
| .8 | .4 | .7 | .2 | .4 | .1 | .4 | .1 | .7 | .4 | 1.1 | .1 |
| 1.3 | 1.2 | 1.2 | 1.2 | 1.2 | 1.1 | 1.2 | 1.1 | 1.4 | 1.3 | 1.4 | 1.1 |
| .6 | .5 | .5 | .4 | .5 | .3 | .5 | .4 | .8 | .6 | .7 | .4 |
| 4.0 | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 | 5.0 | 3.0 | 5.0 | 3.0 | 3.0 | 3.0 |
| 34 | 26 | 39 | 28 | 36 | 23 | 32 | 24 | 30 | 22 | 24 | 23 |
| 31 | 21 | 32 | 21 | 26 | 16 | 23 | 16 | 21 | 16 | 24 | 12 |
| 7 | 5 | 7 | 7 | 7 | 7 | 9 | 7 | 7 | 5 | | 7 |
| 13 | 13 | 14 | 13 | 20 | 13 | 20 | 13 | 14 | 10 | | 13 |
| 26 | 26 | 27 | 26 | 27 | 20 | 26 | 21 | 27 | 20 | | 26 |
| 40 | 27 | 53 | 27 | 53 | 27 | 35 | 27 | 35 | 27 | | 27 |
| 66 | 53 | 82 | 53 | 55 | 53 | 55 | 53 | 53 | 53 | | 40 |
| 105 | 55 | 106 | 59 | 89 | 53 | 66 | 53 | 60 | 55 | | 53 |
| 160 | 108 | 158 | 107 | 133 | 70 | 133 | 79 | 107 | 66 | | 53 |
| 83 | 66 | 89 | 68 | 91 | 67 | 81 | 67 | 74 | 59 | 58 | 63 |
| 64 | 43 | 61 | 40 | 59 | 43 | 45 | 43 | 44 | 38 | 64 | 33 |
| 78 | 68 | 111 | 85 | 83 | 53 | 70 | 62 | 72 | 44 | 42 | 59 |
| 58 | 39 | 61 | 51 | 42 | 36 | 41 | 33 | 43 | 32 | 22 | 13 |
| 83 | 71 | 102 | 50 | 93 | 26 | 78 | 36 | 38 | 35 | 45 | 35 |
| 58 | 51 | 22 | 9 | 39 | 16 | 48 | 25 | 13 | 19 | 33 | 0 |
| 630 | 315 | 320 | 315 | 473 | 315 | 320 | 223 | 164 | 164 | 320 | 158 |
| 788 | 513 | 480 | 473 | 492 | 320 | 492 | 320 | 320 | 197 | 349 | 158 |
| 80 | 67 | 95 | 70 | 89 | 64 | 78 | 65 | 64 | 52 | 52 | 61 |
| 61 | 43 | 61 | 44 | 55 | 42 | 46 | 41 | 45 | 38 | 55 | 31 |
| 20 | 14 | 20 | 20 | 20 | 14 | 20 | 13 | 13 | 7 | | 13 |
| 39 | 39 | 59 | 39 | 41 | 39 | 40 | 39 | 38 | 20 | | 39 |
| 79 | 60 | 80 | 62 | 79 | 59 | 79 | 60 | 41 | 40 | | 62 |
| 82 | 80 | 122 | 80 | 118 | 80 | 82 | 80 | 82 | 80 | | 80 |
| 160 | 140 | 164 | 158 | 160 | 120 | 160 | 158 | 158 | 80 | | 82 |
| 164 | 160 | 205 | 160 | 164 | 158 | 164 | 160 | 160 | 158 | | 82 |
| 315 | 164 | 315 | 164 | 320 | 164 | 236 | 164 | 164 | 164 | | 158 |
| 630 | 315 | 320 | 315 | 473 | 315 | 320 | 164 | 164 | 164 | 320 | 158 |
| 18.4 | 25.3 | 17.0 | 7.3 | 31.3 | 38.2 | 30.5 | 33.6 | 1.0 | 6.0 | 0 | 1.9 |
| 3.8 | 6.8 | 6.7 | 3.4 | 6.3 | 6.8 | 5.0 | 6.3 | .8 | 4.2 | 0 | 2.8 |

TABLE 9.03.--HONEY
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.2 | 3.7 | 5.1 | 4.6 | 3.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.2 | 1.3 | 2.8 | 2.9 | 2.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .9 | 1.0 | 1.3 | 1.3 | .3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 1.0 | 1.4 | 1.0 | .4 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1</u> /NUMBER..... | 2.0 | 3.1 | 1.8 | 1.6 | 1.4 |
| STANDARD DEVIATION.....DO..... | 1.3 | 2.3 | 1.0 | .9 | .8 |
| MAXIMUM.....DO..... | 14.0 | 9.0 | 6.0 | 6.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1</u> /GM..... | 14 | 32 | 9 | 12 | 11 |
| STANDARD DEVIATION.....GM..... | 19 | 96 | 11 | 17 | 11 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> / | | | | | |
| 5TH.....GM..... | 2 | | 1 | 2 | 2 |
| 25TH.....GM..... | 5 | | 4 | 5 | 5 |
| 50TH.....GM..... | 7 | | 7 | 7 | 7 |
| 75TH.....GM..... | 16 | | 11 | 14 | 18 |
| 90TH.....GM..... | 28 | | 18 | 21 | 28 |
| 95TH.....GM..... | 42 | | 21 | 59 | 35 |
| 99TH.....GM..... | 85 | | 78 | 85 | 57 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 27 | 12 | 17 | 25 | 30 |
| STANDARD DEVIATION.....GM..... | 31 | 7 | 13 | 39 | 30 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 23 | 20 | 12 | 19 | 18 |
| STANDARD DEVIATION.....GM..... | 22 | 8 | 8 | 19 | 16 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1</u> /GM..... | 25 | 70 | 18 | 32 | 22 |
| STANDARD DEVIATION.....GM..... | 30 | 147 | 20 | 32 | 13 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 678 | 678 | 170 | 254 | 170 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,314 | 1,314 | 233 | 254 | 170 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1</u> /GM..... | 22 | 30 | 15 | 22 | 24 |
| STANDARD DEVIATION.....GM..... | 25 | 51 | 18 | 31 | 24 |
| PERCENTILES: <u>1</u> / | | | | | |
| 5TH.....GM..... | 5 | | 4 | 7 | 4 |
| 25TH.....GM..... | 7 | | 7 | 7 | 7 |
| 50TH.....GM..... | 14 | | 11 | 14 | 21 |
| 75TH.....GM..... | 21 | | 21 | 21 | 21 |
| 90TH.....GM..... | 42 | | 21 | 42 | 63 |
| 95TH.....GM..... | 64 | | 32 | 85 | 64 |
| 99TH.....GM..... | 127 | | 170 | 170 | 170 |
| MAXIMUM.....GM..... | 339 | 254 | 170 | 254 | 170 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> / | | | | | |
| EATING OCCASIONS.....NUMBER..... | 171.1 | 4.5 | 5.2 | 7.4 | 13.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 5.5 | 7.7 | 5.4 | 5.7 | 16.0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.7 | 2.1 | 2.9 | 2.3 | 4.2 | 4.5 | 4.6 | 4.1 | 8.2 | 6.2 | 5.5 | 6.6 |
| 1.0 | 1.5 | 1.9 | 1.6 | 2.2 | 2.6 | 2.2 | 2.3 | 2.9 | 2.8 | 2.1 | 2.2 |
| .5 | .5 | .6 | .3 | 1.1 | .9 | .9 | .9 | 2.3 | 1.7 | .5 | 1.7 |
| .1 | .1 | .4 | .5 | .9 | .9 | 1.6 | .9 | 3.0 | 1.8 | 2.9 | 2.7 |
| 1.6 | 1.5 | 1.8 | 1.7 | 1.9 | 1.9 | 2.3 | 1.9 | 2.3 | 2.1 | 2.5 | 2.2 |
| 1.0 | .7 | 1.2 | 1.2 | 1.0 | 1.4 | 1.8 | 1.3 | 1.4 | 1.2 | 1.2 | 1.0 |
| 5.0 | 3.0 | 5.0 | 6.0 | 5.0 | 9.0 | 14.0 | 14.0 | 9.0 | 7.0 | 5.0 | 5.0 |
| 22 | 16 | 23 | 13 | 19 | 11 | 18 | 11 | 17 | 10 | 17 | 11 |
| 29 | 16 | 27 | 11 | 19 | 10 | 20 | 10 | 19 | 8 | 23 | 17 |
| | | | | 2 | 2 | 2 | 2 | 2 | 2 | | 2 |
| | | | | 7 | 5 | 7 | 5 | 7 | 5 | | 5 |
| | | | | 12 | 7 | 12 | 7 | 12 | 7 | | 7 |
| | | | | 28 | 14 | 21 | 14 | 21 | 12 | | 14 |
| | | | | 42 | 24 | 42 | 21 | 42 | 21 | | 16 |
| | | | | 57 | 32 | 49 | 28 | 71 | 28 | | 21 |
| | | | | 89 | 49 | 127 | 42 | 85 | 33 | | 127 |
| 63 | 36 | 41 | 25 | 37 | 22 | 32 | 21 | 17 | 18 | 15 | 14 |
| 101 | 31 | 33 | 22 | 40 | 19 | 27 | 18 | 12 | 11 | 12 | 7 |
| 32 | 39 | 36 | 25 | 33 | 19 | 23 | 24 | 23 | 14 | 7 | 26 |
| 26 | 37 | 18 | 19 | 28 | 13 | 19 | 15 | 17 | 9 | 5 | 46 |
| 37 | 21 | 68 | 26 | 31 | 21 | 30 | 18 | 30 | 16 | 28 | 12 |
| 25 | 0 | 45 | 10 | 22 | 14 | 27 | 11 | 24 | 10 | 27 | 6 |
| 339 | 254 | 191 | 85 | 205 | 85 | 254 | 148 | 127 | 42 | 127 | 191 |
| 339 | 283 | 424 | 125 | 268 | 205 | 381 | 205 | 254 | 99 | 254 | 381 |
| 41 | 32 | 38 | 23 | 30 | 18 | 24 | 18 | 23 | 14 | 21 | 15 |
| 67 | 34 | 32 | 16 | 31 | 15 | 25 | 15 | 21 | 10 | 20 | 24 |
| | | | | 7 | 7 | 7 | 4 | 7 | 4 | | 4 |
| | | | | 11 | 7 | 7 | 7 | 7 | 7 | | 7 |
| | | | | 21 | 14 | 21 | 14 | 14 | 7 | | 7 |
| | | | | 42 | 21 | 32 | 21 | 21 | 21 | | 21 |
| | | | | 64 | 42 | 42 | 42 | 42 | 21 | | 21 |
| | | | | 85 | 42 | 85 | 42 | 64 | 42 | | 21 |
| | | | | 148 | 85 | 129 | 85 | 85 | 42 | | 191 |
| 339 | 254 | 191 | 85 | 170 | 85 | 191 | 148 | 127 | 42 | 85 | 191 |
| 2.8 | 2.5 | 5.7 | 5.7 | 16.8 | 13.1 | 41.4 | 28.3 | 2.8 | 10.8 | 1.7 | 9.0 |
| 5.0 | 3.7 | 7.6 | 9.8 | 5.5 | 2.9 | 8.0 | 5.2 | 1.3 | 4.8 | 2.4 | 6.1 |

TABLE 9.04.--JELLIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 17.0 | .7 | 20.7 | 28.7 | 27.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 10.8 | .7 | 12.9 | 18.2 | 18.9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 4.0 | 0 | 5.6 | 6.9 | 6.8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.1 | 0 | 2.3 | 3.5 | 2.2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.6 | 1.0 | 1.6 | 1.6 | 1.5 |
| STANDARD DEVIATION.....DO..... | .9 | .0 | .9 | .9 | .9 |
| MAXIMUM.....DO..... | 8.0 | 1.0 | 5.0 | 6.0 | 8.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 10 | 5 | 7 | 9 | 9 |
| STANDARD DEVIATION.....GM..... | 11 | 2 | 7 | 8 | 10 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 2 | | 1 | 2 | 2 |
| 25TH.....GM..... | 4 | | 2 | 3 | 4 |
| 50TH.....GM..... | 6 | | 4 | 6 | 6 |
| 75TH.....GM..... | 13 | | 8 | 13 | 13 |
| 90TH.....GM..... | 25 | | 15 | 19 | 19 |
| 95TH.....GM..... | 31 | | 19 | 25 | 25 |
| 99TH.....GM..... | 50 | | 38 | 38 | 50 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 21 | 16 | 15 | 18 | 22 |
| STANDARD DEVIATION.....GM..... | 22 | 5 | 17 | 17 | 20 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 20 | 0 | 13 | 17 | 18 |
| STANDARD DEVIATION.....GM..... | 16 | 0 | 9 | 13 | 13 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 20 | 0 | 13 | 17 | 21 |
| STANDARD DEVIATION.....GM..... | 16 | 0 | 8 | 12 | 19 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 300 | 19 | 114 | 156 | 150 |
| MAXIMUM OVER 3 DAYS.....GM..... | 375 | 19 | 114 | 250 | 227 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 20 | 16 | 13 | 16 | 19 |
| STANDARD DEVIATION.....GM..... | 18 | 5 | 13 | 14 | 17 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 6 | | 3 | 3 | 6 |
| 25TH.....GM..... | 6 | | 6 | 6 | 6 |
| 50TH.....GM..... | 19 | | 6 | 13 | 19 |
| 75TH.....GM..... | 19 | | 19 | 19 | 19 |
| 90TH.....GM..... | 38 | | 19 | 38 | 38 |
| 95TH.....GM..... | 56 | | 38 | 38 | 38 |
| 99TH.....GM..... | 75 | | 75 | 75 | 75 |
| MAXIMUM.....GM..... | 300 | 19 | 114 | 150 | 150 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 358.3 | 0 | 11.5 | 30.7 | 31.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 3.6 | 0 | 3.3 | 3.9 | 4.2 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 26.1 | 21.3 | 19.6 | 15.1 | 13.8 | 13.6 | 15.2 | 13.3 | 19.8 | 15.1 | 15.5 | 17.5 |
| 18.3 | 14.6 | 12.2 | 10.7 | 8.5 | 9.9 | 9.1 | 8.4 | 10.8 | 8.1 | 6.3 | 8.0 |
| 5.2 | 4.7 | 4.5 | 3.1 | 3.0 | 2.8 | 4.0 | 3.3 | 5.1 | 4.6 | 4.1 | 4.5 |
| 2.5 | 2.0 | 2.9 | 1.3 | 2.2 | .9 | 2.1 | 1.7 | 3.9 | 2.4 | 5.1 | 5.0 |
| 1.5 | 1.5 | 1.6 | 1.5 | 1.6 | 1.4 | 1.6 | 1.6 | 1.7 | 1.7 | 2.1 | 1.9 |
| .8 | .7 | .8 | .8 | .9 | .7 | .9 | .8 | .9 | 1.0 | 1.2 | 1.0 |
| 6.0 | 5.0 | 5.0 | 5.0 | 7.0 | 5.0 | 6.0 | 5.0 | 5.0 | 6.0 | 7.0 | 5.0 |
| 12 | 10 | 13 | 11 | 14 | 9 | 12 | 8 | 11 | 9 | 16 | 10 |
| 11 | 10 | 13 | 12 | 14 | 9 | 12 | 8 | 9 | 10 | 17 | 11 |
| 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 4 | 4 | 6 | 4 | 4 | 3 | 4 | 2 | 4 | 2 | 4 | 4 |
| 8 | 6 | 10 | 6 | 8 | 6 | 8 | 6 | 7 | 6 | 13 | 6 |
| 15 | 13 | 19 | 13 | 19 | 13 | 15 | 12 | 13 | 13 | 20 | 10 |
| 25 | 25 | 25 | 25 | 29 | 19 | 25 | 17 | 25 | 19 | 44 | 25 |
| 38 | 25 | 38 | 38 | 38 | 25 | 35 | 25 | 31 | 25 | 50 | 38 |
| 50 | 56 | 57 | 50 | 75 | 43 | 50 | 40 | 41 | 46 | 75 | 52 |
| 27 | 22 | 27 | 23 | 27 | 19 | 25 | 16 | 22 | 13 | 33 | 17 |
| 24 | 20 | 27 | 27 | 27 | 18 | 22 | 17 | 20 | 8 | 55 | 10 |
| 23 | 20 | 27 | 26 | 27 | 20 | 24 | 17 | 18 | 16 | 18 | 15 |
| 15 | 13 | 17 | 18 | 21 | 15 | 22 | 13 | 12 | 10 | 16 | 17 |
| 26 | 22 | 25 | 24 | 25 | 20 | 18 | 16 | 19 | 20 | 25 | 16 |
| 15 | 19 | 18 | 15 | 20 | 14 | 14 | 11 | 10 | 18 | 16 | 14 |
| 169 | 169 | 300 | 300 | 300 | 150 | 263 | 188 | 113 | 113 | 206 | 99 |
| 188 | 244 | 305 | 300 | 300 | 225 | 375 | 188 | 150 | 225 | 225 | 206 |
| 24 | 21 | 25 | 23 | 25 | 19 | 22 | 16 | 19 | 15 | 22 | 16 |
| 20 | 18 | 23 | 23 | 24 | 16 | 20 | 14 | 16 | 12 | 25 | 14 |
| 6 | 6 | 6 | 6 | 6 | 6 | 6 | 3 | 6 | 6 | 6 | 6 |
| 9 | 6 | 13 | 9 | 13 | 6 | 9 | 6 | 13 | 6 | 6 | 6 |
| 19 | 19 | 19 | 19 | 19 | 18 | 19 | 13 | 14 | 9 | 19 | 13 |
| 38 | 19 | 38 | 28 | 38 | 19 | 36 | 19 | 19 | 19 | 19 | 19 |
| 38 | 38 | 38 | 38 | 56 | 38 | 38 | 38 | 38 | 38 | 38 | 38 |
| 57 | 56 | 56 | 75 | 75 | 38 | 56 | 38 | 38 | 38 | 75 | 38 |
| 84 | 75 | 114 | 75 | 114 | 75 | 75 | 75 | 99 | 56 | 150 | 75 |
| 169 | 169 | 300 | 300 | 300 | 150 | 225 | 150 | 113 | 75 | 150 | 99 |
| 31.0 | 29.7 | 19.0 | 15.7 | 27.6 | 35.1 | 45.5 | 37.1 | 12.0 | 14.9 | 3.3 | 13.3 |
| 3.8 | 4.4 | 4.3 | 4.8 | 3.2 | 3.5 | 3.7 | 2.5 | 3.2 | 3.3 | 1.9 | 4.1 |

TABLE 9.05.--JAMS, PRESERVES, AND MARMALADES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 7.1 | 1.4 | 6.7 | 6.8 | 6.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.5 | 1.2 | 4.9 | 4.4 | 4.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.6 | .2 | 1.1 | 1.7 | 1.4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .9 | 0 | .7 | .7 | .7 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER.... | 1.6 | 1.3 | 1.5 | 1.6 | 1.6 |
| STANDARD DEVIATION.....DO..... | .9 | .4 | .8 | .8 | .9 |
| MAXIMUM.....DO..... | 6.0 | 2.0 | 4.0 | 5.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 12 | 5 | 6 | 9 | 10 |
| STANDARD DEVIATION.....GM..... | 14 | 4 | 5 | 6 | 10 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 2 | | 1 | 2 | 2 |
| 25TH.....GM..... | 4 | | 2 | 4 | 4 |
| 50TH.....GM..... | 7 | | 7 | 7 | 7 |
| 75TH.....GM..... | 13 | | 7 | 13 | 13 |
| 90TH.....GM..... | 27 | | 13 | 20 | 20 |
| 95TH.....GM..... | 33 | | 13 | 20 | 33 |
| 99TH.....GM..... | 67 | | 27 | 27 | 53 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 25 | 13 | 16 | 20 | 23 |
| STANDARD DEVIATION.....GM..... | 27 | 13 | 15 | 14 | 21 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 22 | 13 | 14 | 18 | 21 |
| STANDARD DEVIATION.....GM..... | 18 | 0 | 9 | 11 | 15 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 24 | 0 | 9 | 14 | 20 |
| STANDARD DEVIATION.....GM..... | 26 | 0 | 3 | 6 | 14 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 320 | 40 | 80 | 60 | 160 |
| MAXIMUM OVER 3 DAYS.....GM..... | 800 | 40 | 80 | 80 | 160 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 23 | 12 | 13 | 17 | 20 |
| STANDARD DEVIATION.....GM..... | 23 | 8 | 11 | 12 | 17 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 7 | | 3 | 7 | 7 |
| 25TH.....GM..... | 7 | | 7 | 7 | 7 |
| 50TH.....GM..... | 20 | | 7 | 13 | 20 |
| 75TH.....GM..... | 20 | | 20 | 20 | 20 |
| 90TH.....GM..... | 40 | | 20 | 40 | 40 |
| 95TH.....GM..... | 60 | | 40 | 40 | 40 |
| 99TH.....GM..... | 122 | | 40 | 60 | 60 |
| MAXIMUM.....GM..... | 320 | 20 | 80 | 60 | 160 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER.... | 140.5 | 0 | 3.8 | 7.2 | 14.3 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 3.3 | 0 | 3.7 | 3.9 | 7.6 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.8 | 5.4 | 6.5 | 4.8 | 6.4 | 4.7 | 8.3 | 7.2 | 11.7 | 12.6 | 11.6 | 12.4 |
| 4.2 | 4.0 | 3.8 | 3.7 | 4.3 | 3.5 | 5.1 | 4.9 | 4.9 | 6.9 | 4.6 | 5.2 |
| 1.3 | 1.1 | 2.1 | 1.0 | 1.4 | .9 | 1.9 | 1.4 | 3.4 | 3.2 | 4.2 | 3.7 |
| .4 | .4 | .5 | .1 | .6 | .3 | 1.2 | .9 | 3.4 | 2.5 | 2.8 | 3.5 |
| 1.5 | 1.5 | 1.6 | 1.3 | 1.4 | 1.4 | 1.6 | 1.5 | 2.1 | 1.7 | 1.9 | 2.0 |
| .8 | .8 | .8 | .5 | .7 | .7 | 1.0 | .8 | 1.2 | .9 | .9 | 1.0 |
| 4.0 | 4.0 | 5.0 | 3.0 | 3.0 | 5.0 | 6.0 | 6.0 | 6.0 | 4.0 | 5.0 | 5.0 |
| 14 | 11 | 17 | 9 | 14 | 9 | 14 | 11 | 20 | 10 | 19 | 9 |
| 13 | 11 | 13 | 9 | 13 | 8 | 14 | 13 | 35 | 8 | 22 | 7 |
| 2 | 2 | 4 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 1 | 2 |
| 7 | 4 | 7 | 3 | 5 | 4 | 5 | 4 | 7 | 4 | 7 | 4 |
| 11 | 7 | 13 | 7 | 13 | 7 | 8 | 7 | 13 | 7 | 11 | 7 |
| 17 | 13 | 27 | 11 | 19 | 13 | 19 | 13 | 20 | 13 | 20 | 13 |
| 27 | 22 | 33 | 16 | 27 | 13 | 27 | 20 | 40 | 20 | 53 | 20 |
| 40 | 40 | 53 | 20 | 33 | 20 | 39 | 33 | 53 | 27 | 67 | 22 |
| 67 | 53 | 67 | 53 | 80 | 47 | 67 | 67 | 267 | 47 | 93 | 31 |
| 30 | 25 | 40 | 22 | 30 | 23 | 29 | 25 | 21 | 20 | 32 | 14 |
| 25 | 20 | 33 | 29 | 22 | 20 | 30 | 37 | 15 | 15 | 43 | 9 |
| 27 | 25 | 35 | 18 | 25 | 17 | 23 | 19 | 23 | 16 | 40 | 14 |
| 12 | 20 | 23 | 8 | 15 | 10 | 20 | 20 | 14 | 9 | 39 | 7 |
| 45 | 27 | 23 | 20 | 36 | 18 | 27 | 19 | 42 | 17 | 21 | 16 |
| 22 | 19 | 9 | 0 | 24 | 13 | 22 | 12 | 58 | 13 | 20 | 7 |
| 142 | 120 | 160 | 160 | 120 | 160 | 200 | 320 | 320 | 80 | 240 | 40 |
| 240 | 180 | 200 | 160 | 240 | 160 | 346 | 320 | 800 | 142 | 280 | 93 |
| 28 | 23 | 32 | 20 | 29 | 19 | 25 | 21 | 29 | 17 | 30 | 14 |
| 20 | 16 | 26 | 24 | 21 | 17 | 24 | 29 | 30 | 13 | 39 | 9 |
| 7 | 7 | 7 | 7 | 7 | 7 | 7 | 3 | 7 | 7 | 7 | 7 |
| 13 | 10 | 20 | 7 | 13 | 7 | 13 | 7 | 10 | 7 | 7 | 7 |
| 20 | 20 | 20 | 20 | 20 | 20 | 20 | 13 | 20 | 13 | 20 | 13 |
| 40 | 40 | 40 | 20 | 40 | 20 | 30 | 20 | 40 | 20 | 40 | 20 |
| 40 | 40 | 80 | 40 | 60 | 40 | 40 | 40 | 60 | 40 | 40 | 20 |
| 80 | 60 | 80 | 40 | 80 | 40 | 80 | 40 | 80 | 40 | 80 | 40 |
| 120 | 80 | 160 | 160 | 80 | 80 | 160 | 160 | 160 | 60 | 240 | 40 |
| 122 | 80 | 160 | 160 | 120 | 160 | 160 | 320 | 160 | 80 | 240 | 40 |
| 1.0 | 6.3 | 4.1 | 7.5 | 13.5 | 5.9 | 37.3 | 15.0 | 12.3 | 5.6 | .7 | 6.0 |
| .6 | 3.7 | 2.8 | 8.4 | 3.7 | 1.7 | 5.6 | 2.0 | 4.6 | 1.5 | .6 | 2.4 |

TABLE 9.06.--PLAIN GELATIN DESSERTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 6.6 | 3.8 | 7.3 | 7.9 | 11.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.4 | 2.8 | 5.6 | 6.4 | 9.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.0 | .9 | 1.2 | 1.2 | 1.6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | .2 | .5 | .4 | .3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 1.7 | 1.4 | 1.3 | 1.2 |
| STANDARD DEVIATION.....DO..... | .6 | .9 | .8 | .6 | .5 |
| MAXIMUM.....DO..... | 6.0 | 3.0 | 5.0 | 3.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 67 | 50 | 55 | 61 | 62 |
| STANDARD DEVIATION.....GM..... | 48 | 44 | 47 | 40 | 39 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 20 | | 15 | 19 | 20 |
| 25TH.....GM..... | 40 | | 30 | 40 | 40 |
| 50TH.....GM..... | 51 | | 40 | 40 | 40 |
| 75TH.....GM..... | 80 | | 66 | 80 | 80 |
| 90TH.....GM..... | 120 | | 80 | 120 | 100 |
| 95TH.....GM..... | 160 | | 160 | 120 | 160 |
| 99TH.....GM..... | 240 | | 260 | 200 | 200 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 164 | 137 | 122 | 146 | 157 |
| STANDARD DEVIATION.....GM..... | 93 | 143 | 72 | 84 | 83 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 171 | 110 | 132 | 153 | 165 |
| STANDARD DEVIATION.....GM..... | 82 | 16 | 74 | 50 | 83 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 170 | 3 | 132 | 146 | 118 |
| STANDARD DEVIATION.....GM..... | 96 | 0 | 89 | 44 | 31 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 960 | 540 | 480 | 600 | 816 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,440 | 540 | 780 | 720 | 952 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 159 | 89 | 118 | 144 | 151 |
| STANDARD DEVIATION.....GM..... | 85 | 62 | 73 | 74 | 78 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 60 | | 29 | 60 | 60 |
| 25TH.....GM..... | 120 | | 60 | 120 | 120 |
| 50TH.....GM..... | 120 | | 120 | 120 | 120 |
| 75TH.....GM..... | 240 | | 120 | 180 | 240 |
| 90TH.....GM..... | 240 | | 240 | 240 | 240 |
| 95TH.....GM..... | 240 | | 240 | 240 | 240 |
| 99TH.....GM..... | 480 | | 360 | 360 | 480 |
| MAXIMUM.....GM..... | 960 | 240 | 480 | 480 | 480 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 29.5 | 0 | .7 | .7 | 3.3 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .9 | 0 | .6 | .4 | 1.3 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 10.5 | 9.9 | 7.7 | 5.5 | 3.6 | 4.0 | 5.7 | 5.4 | 8.6 | 10.8 | 7.9 | 9.0 |
| 9.1 | 8.0 | 6.5 | 4.7 | 3.1 | 3.4 | 4.5 | 4.7 | 5.8 | 8.5 | 4.9 | 6.7 |
| 1.4 | 1.9 | 1.0 | .7 | .5 | .5 | .9 | .6 | 1.9 | 1.9 | 1.0 | 1.9 |
| (2) | .1 | .2 | .1 | 0 | .1 | .2 | .1 | .9 | .4 | 1.9 | .5 |
| 1.2 | 1.3 | 1.3 | 1.3 | 1.2 | 1.2 | 1.3 | 1.2 | 1.5 | 1.3 | 1.7 | 1.4 |
| .4 | .6 | .5 | .7 | .4 | .5 | .8 | .5 | .9 | .5 | .8 | .7 |
| 3.0 | 4.0 | 3.0 | 6.0 | 3.0 | 4.0 | 6.0 | 6.0 | 6.0 | 3.0 | 3.0 | 4.0 |
| 65 | 66 | 78 | 76 | 70 | 62 | 80 | 64 | 74 | 61 | 91 | 76 |
| 46 | 47 | 56 | 63 | 41 | 40 | 61 | 45 | 41 | 41 | 69 | 55 |
| 20 | 20 | 20 | 20 | 20 | 15 | 20 | 20 | 23 | 10 | | 23 |
| 40 | 40 | 40 | 40 | 40 | 40 | 40 | 40 | 40 | 40 | | 40 |
| 40 | 51 | 80 | 60 | 80 | 40 | 80 | 40 | 77 | 40 | | 60 |
| 80 | 80 | 80 | 80 | 80 | 80 | 80 | 80 | 80 | 80 | | 80 |
| 100 | 120 | 160 | 160 | 120 | 100 | 160 | 110 | 160 | 120 | | 160 |
| 160 | 160 | 183 | 160 | 160 | 160 | 160 | 160 | 160 | 160 | | 200 |
| 306 | 240 | 306 | 440 | 240 | 200 | 360 | 240 | 160 | 240 | | 320 |
| 170 | 162 | 197 | 186 | 180 | 159 | 180 | 161 | 174 | 141 | 154 | 167 |
| 117 | 92 | 126 | 108 | 96 | 79 | 92 | 84 | 87 | 72 | 113 | 77 |
| 172 | 163 | 188 | 181 | 194 | 168 | 210 | 153 | 167 | 158 | 183 | 193 |
| 86 | 95 | 109 | 80 | 64 | 71 | 87 | 67 | 66 | 64 | 49 | 108 |
| 120 | 240 | 224 | 340 | 0 | 156 | 221 | 244 | 96 | 141 | 174 | 150 |
| 0 | 0 | 26 | 100 | 0 | 98 | 127 | 81 | 38 | 59 | 60 | 80 |
| 960 | 480 | 720 | 600 | 720 | 480 | 960 | 600 | 480 | 240 | 480 | 480 |
| 960 | 960 | 918 | 1,320 | 720 | 960 | 1,440 | 1,200 | 480 | 720 | 720 | 960 |
| 165 | 155 | 186 | 181 | 180 | 157 | 182 | 157 | 144 | 144 | 164 | 166 |
| 109 | 79 | 101 | 87 | 90 | 77 | 94 | 74 | 71 | 72 | 84 | 83 |
| 46 | 60 | 60 | 60 | 60 | 45 | 60 | 60 | 51 | 30 | | 60 |
| 120 | 120 | 120 | 120 | 120 | 120 | 120 | 120 | 91 | 91 | | 120 |
| 120 | 120 | 153 | 180 | 180 | 120 | 161 | 120 | 120 | 120 | | 120 |
| 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | | 240 |
| 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | 240 | | 240 |
| 240 | 240 | 459 | 360 | 240 | 240 | 240 | 240 | 240 | 240 | | 240 |
| 720 | 360 | 480 | 480 | 720 | 480 | 480 | 240 | 240 | 240 | | 480 |
| 960 | 480 | 480 | 480 | 720 | 480 | 960 | 480 | 240 | 240 | 480 | 480 |
| 2.8 | 1.9 | 1.8 | 0 | 0 | 8.0 | 1.5 | 4.3 | 0 | 0 | .9 | 3.6 |
| 1.1 | .7 | 1.3 | 0 | 0 | 3.2 | .4 | .9 | 0 | 0 | 1.3 | 2.9 |

TABLE 9.07.--GELATIN DESSERTS WITH FRUIT
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.3 | .4 | 3.4 | 2.4 | 4.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.9 | .2 | 3.2 | 2.1 | 4.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | .2 | .1 | .2 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | .1 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 1.5 | 1.1 | 1.2 | 1.1 |
| STANDARD DEVIATION.....DO..... | .5 | .5 | .4 | .4 | .2 |
| MAXIMUM.....DO..... | 5.0 | 2.0 | 3.0 | 3.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 67 | 20 | 49 | 52 | 55 |
| STANDARD DEVIATION.....GM..... | 49 | 20 | 42 | 33 | 31 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 20 | | | | 20 |
| 25TH.....GM..... | 40 | | | | 40 |
| 50TH.....GM..... | 40 | | | | 40 |
| 75TH.....GM..... | 80 | | | | 80 |
| 90TH.....GM..... | 133 | | | | 90 |
| 95TH.....GM..... | 160 | | | | 120 |
| 99TH.....GM..... | 240 | | | | 160 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 170 | 2 | 128 | 142 | 158 |
| STANDARD DEVIATION.....GM..... | 101 | 0 | 66 | 89 | 88 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 191 | 60 | 37 | 136 | 172 |
| STANDARD DEVIATION.....GM..... | 85 | 0 | 0 | 54 | 37 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 235 | 0 | 240 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 83 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 720 | 60 | 360 | 480 | 480 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,200 | 120 | 720 | 480 | 480 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 167 | 40 | 128 | 135 | 156 |
| STANDARD DEVIATION.....GM..... | 90 | 27 | 81 | 84 | 82 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 60 | | | | 60 |
| 25TH.....GM..... | 120 | | | | 120 |
| 50TH.....GM..... | 120 | | | | 120 |
| 75TH.....GM..... | 240 | | | | 240 |
| 90TH.....GM..... | 240 | | | | 240 |
| 95TH.....GM..... | 306 | | | | 240 |
| 99TH.....GM..... | 480 | | | | 480 |
| MAXIMUM.....GM..... | 612 | 60 | 360 | 480 | 480 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 22.1 | 0 | .8 | 1.2 | 3.0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.5 | 0 | 2.1 | 2.6 | 3.6 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.4 | 3.5 | 4.2 | 3.0 | 2.0 | 1.7 | 3.0 | 3.8 | 4.5 | 5.0 | 4.4 | 6.0 |
| 5.2 | 3.2 | 3.8 | 2.9 | 1.7 | 1.5 | 2.6 | 3.1 | 3.7 | 3.7 | 3.7 | 5.1 |
| .2 | .2 | .4 | .1 | .3 | .1 | .3 | .6 | .7 | 1.3 | .4 | .9 |
| 0 | .1 | .1 | 0 | 0 | 0 | .1 | .1 | .1 | .1 | .2 | 0 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.1 | 1.2 | 1.3 | 1.3 | 1.4 | 1.4 | 1.3 |
| .3 | .4 | .4 | .3 | .5 | .3 | .5 | .6 | .5 | .6 | .9 | .6 |
| 2.0 | 3.0 | 3.0 | 2.0 | 3.0 | 3.0 | 4.0 | 5.0 | 3.0 | 3.0 | 5.0 | 3.0 |
| 59 | 65 | 69 | 63 | 72 | 57 | 75 | 69 | 78 | 86 | 78 | 65 |
| 34 | 46 | 45 | 37 | 38 | 32 | 52 | 55 | 46 | 70 | 74 | 47 |
| 20 | 20 | 20 | | 30 | 20 | 23 | 15 | 26 | 20 | | 11 |
| 40 | 40 | 40 | | 40 | 40 | 40 | 40 | 40 | 40 | | 40 |
| 40 | 40 | 53 | | 80 | 40 | 60 | 40 | 80 | 80 | | 40 |
| 80 | 80 | 80 | | 80 | 80 | 80 | 80 | 94 | 120 | | 80 |
| 91 | 160 | 160 | | 140 | 120 | 160 | 160 | 160 | 160 | | 120 |
| 140 | 160 | 160 | | 160 | 120 | 160 | 180 | 160 | 180 | | 200 |
| 160 | 240 | 240 | | 160 | 160 | 240 | 240 | 160 | 360 | | 204 |
| 170 | 177 | 182 | 185 | 204 | 165 | 185 | 159 | 187 | 169 | 179 | 157 |
| 93 | 119 | 104 | 109 | 110 | 97 | 104 | 92 | 102 | 122 | 83 | 106 |
| 189 | 156 | 198 | 180 | 143 | 132 | 211 | 184 | 236 | 253 | 150 | 198 |
| 46 | 51 | 79 | 0 | 58 | 21 | 60 | 90 | 10 | 106 | 22 | 73 |
| 0 | 190 | 220 | 0 | 0 | 0 | 266 | 242 | 120 | 160 | 400 | 0 |
| 0 | 53 | 0 | 0 | 0 | 0 | 43 | 86 | 0 | 0 | 0 | 0 |
| 480 | 720 | 480 | 544 | 480 | 480 | 600 | 720 | 480 | 720 | 480 | 612 |
| 480 | 720 | 720 | 544 | 480 | 480 | 960 | 1,200 | 480 | 1,080 | 1,200 | 612 |
| 166 | 172 | 186 | 177 | 175 | 157 | 185 | 161 | 182 | 183 | 168 | 149 |
| 80 | 103 | 101 | 100 | 90 | 88 | 86 | 84 | 80 | 101 | 73 | 91 |
| 60 | 60 | 60 | | 60 | 60 | 68 | 60 | 79 | 60 | | 17 |
| 120 | 120 | 120 | | 120 | 120 | 120 | 120 | 120 | 120 | | 120 |
| 120 | 120 | 153 | | 120 | 120 | 180 | 120 | 180 | 120 | | 120 |
| 240 | 240 | 240 | | 240 | 240 | 240 | 240 | 240 | 240 | | 180 |
| 240 | 240 | 272 | | 240 | 240 | 240 | 240 | 240 | 360 | | 240 |
| 240 | 480 | 480 | | 360 | 360 | 272 | 240 | 240 | 360 | | 240 |
| 480 | 480 | 480 | | 480 | 480 | 480 | 480 | 480 | 360 | | 612 |
| 480 | 480 | 480 | 544 | 480 | 480 | 480 | 480 | 480 | 612 | 240 | 612 |
| 3.0 | 2.3 | 0 | .8 | 1.3 | 2.1 | 4.0 | 2.3 | 0 | 1.3 | 0 | 0 |
| 2.4 | 2.7 | 0 | 1.8 | 1.3 | 2.1 | 2.2 | .7 | 0 | 1.1 | 0 | 0 |

TABLE 9.08.--CANDY
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 13.0 | .8 | 17.4 | 20.9 | 19.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 9.4 | .8 | 11.3 | 15.8 | 14.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.8 | 0 | 4.6 | 4.1 | 3.9 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .9 | 0 | 1.5 | 1.0 | 1.3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.5 | 1.0 | 1.6 | 1.4 | 1.5 |
| STANDARD DEVIATION.....DO..... | 1.0 | .0 | .9 | .9 | .9 |
| MAXIMUM.....DO..... | 15.0 | 1.0 | 6.0 | 9.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 22 | 6 | 15 | 16 | 19 |
| STANDARD DEVIATION.....GM..... | 24 | 4 | 18 | 15 | 20 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 3 | | 2 | 2 | 2 |
| 25TH.....GM..... | 9 | | 5 | 6 | 7 |
| 50TH.....GM..... | 14 | | 9 | 11 | 13 |
| 75TH.....GM..... | 28 | | 19 | 20 | 24 |
| 90TH.....GM..... | 47 | | 35 | 35 | 43 |
| 95TH.....GM..... | 66 | | 40 | 47 | 58 |
| 99TH.....GM..... | 116 | | 122 | 73 | 102 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 47 | 18 | 29 | 37 | 41 |
| STANDARD DEVIATION.....GM..... | 46 | 13 | 27 | 31 | 40 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 51 | 0 | 33 | 38 | 50 |
| STANDARD DEVIATION.....GM..... | 45 | 0 | 33 | 27 | 41 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 50 | 0 | 33 | 36 | 42 |
| STANDARD DEVIATION.....GM..... | 38 | 0 | 30 | 20 | 26 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 907 | 37 | 240 | 227 | 340 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,105 | 37 | 421 | 340 | 495 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 43 | 18 | 29 | 33 | 38 |
| STANDARD DEVIATION.....GM..... | 42 | 13 | 31 | 29 | 34 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 7 | | 5 | 6 | 7 |
| 25TH.....GM..... | 18 | | 12 | 14 | 15 |
| 50TH.....GM..... | 30 | | 17 | 28 | 30 |
| 75TH.....GM..... | 51 | | 30 | 43 | 48 |
| 90TH.....GM..... | 85 | | 57 | 60 | 74 |
| 95TH.....GM..... | 113 | | 85 | 85 | 113 |
| 99TH.....GM..... | 220 | | 156 | 170 | 170 |
| MAXIMUM.....GM..... | 907 | 37 | 240 | 227 | 227 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 311.6 | 0 | 8.9 | 22.9 | 25.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 4.1 | 0 | 3.1 | 4.4 | 4.5 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 22.0 | 23.2 | 16.9 | 20.0 | 11.5 | 11.7 | 8.2 | 9.8 | 8.1 | 8.7 | 6.5 | 8.1 |
| 16.1 | 15.7 | 11.8 | 14.0 | 7.9 | 9.0 | 5.6 | 7.4 | 5.9 | 5.7 | 4.5 | 5.9 |
| 4.4 | 5.8 | 3.5 | 5.0 | 2.7 | 2.2 | 1.9 | 1.8 | 1.3 | 2.3 | 1.4 | 1.7 |
| 1.5 | 1.7 | 1.6 | .9 | .9 | .5 | .7 | .6 | .9 | .7 | .6 | .5 |
| 1.5 | 1.6 | 1.6 | 1.5 | 1.6 | 1.4 | 1.5 | 1.4 | 1.9 | 1.6 | 1.8 | 1.4 |
| .9 | 1.0 | 1.0 | .9 | 1.3 | .9 | .8 | 1.1 | 2.7 | .8 | 1.8 | .8 |
| 7.0 | 9.0 | 6.0 | 6.0 | 12.0 | 7.0 | 6.0 | 13.0 | 15.0 | 5.0 | 9.0 | 5.0 |
| 26 | 24 | 28 | 23 | 29 | 20 | 26 | 19 | 23 | 17 | 15 | 15 |
| 26 | 28 | 32 | 21 | 28 | 18 | 30 | 22 | 23 | 16 | 9 | 13 |
| 3 | 3 | 5 | 3 | 5 | 2 | 4 | 3 | 5 | 3 | | 3 |
| 9 | 9 | 10 | 10 | 11 | 9 | 9 | 7 | 9 | 7 | | 6 |
| 18 | 16 | 19 | 15 | 19 | 14 | 18 | 13 | 13 | 11 | | 11 |
| 32 | 28 | 33 | 30 | 38 | 26 | 33 | 21 | 27 | 24 | | 19 |
| 57 | 47 | 57 | 48 | 64 | 40 | 57 | 38 | 53 | 38 | | 32 |
| 84 | 68 | 87 | 71 | 89 | 57 | 74 | 57 | 66 | 43 | | 36 |
| 123 | 132 | 159 | 103 | 141 | 87 | 189 | 95 | 110 | 75 | | 80 |
| 54 | 48 | 58 | 57 | 59 | 47 | 52 | 40 | 40 | 42 | 36 | 34 |
| 52 | 49 | 55 | 55 | 57 | 43 | 51 | 33 | 31 | 48 | 19 | 27 |
| 65 | 55 | 66 | 49 | 65 | 45 | 61 | 45 | 47 | 34 | 33 | 37 |
| 47 | 61 | 72 | 31 | 48 | 30 | 57 | 35 | 22 | 21 | 9 | 29 |
| 59 | 49 | 62 | 43 | 70 | 42 | 56 | 51 | 71 | 23 | 25 | 23 |
| 34 | 33 | 37 | 34 | 29 | 28 | 49 | 54 | 25 | 14 | 13 | 7 |
| 425 | 907 | 737 | 340 | 567 | 340 | 340 | 397 | 168 | 298 | 85 | 213 |
| 510 | 1,105 | 907 | 410 | 567 | 405 | 983 | 808 | 330 | 298 | 108 | 241 |
| 51 | 44 | 53 | 47 | 53 | 41 | 53 | 39 | 35 | 33 | 25 | 31 |
| 44 | 51 | 49 | 45 | 52 | 37 | 53 | 33 | 26 | 34 | 18 | 30 |
| 7 | 6 | 11 | 7 | 8 | 7 | 11 | 6 | 7 | 6 | | 7 |
| 24 | 15 | 28 | 21 | 28 | 18 | 25 | 16 | 20 | 14 | | 14 |
| 36 | 30 | 37 | 34 | 39 | 32 | 38 | 30 | 28 | 28 | | 25 |
| 57 | 55 | 63 | 56 | 57 | 48 | 57 | 50 | 48 | 40 | | 34 |
| 113 | 85 | 105 | 104 | 113 | 85 | 102 | 84 | 60 | 60 | | 57 |
| 151 | 113 | 142 | 113 | 142 | 113 | 144 | 113 | 85 | 85 | | 97 |
| 227 | 227 | 284 | 227 | 255 | 170 | 328 | 170 | 168 | 224 | | 213 |
| 330 | 907 | 454 | 340 | 567 | 340 | 340 | 227 | 168 | 298 | 85 | 213 |
| 31.4 | 25.0 | 15.3 | 19.4 | 23.4 | 37.3 | 28.9 | 49.1 | 6.7 | 8.9 | 1.8 | 7.3 |
| 4.5 | 3.1 | 4.1 | 4.4 | 3.1 | 4.1 | 4.9 | 4.9 | 3.9 | 3.8 | 2.8 | 6.5 |

TABLE 10.01.--WHITE POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 74.4 | 18.1 | 74.5 | 76.3 | 80.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 34.2 | 8.7 | 34.7 | 37.2 | 37.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 28.4 | 4.9 | 30.4 | 29.4 | 31.8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 11.8 | 4.5 | 9.4 | 9.6 | 11.6 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.9 | 2.0 | 1.8 | 1.8 | 1.9 |
| STANDARD DEVIATION.....DO..... | 1.0 | 1.2 | .9 | .8 | .9 |
| MAXIMUM.....DO..... | 10.0 | 6.0 | 6.0 | 5.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 78 | 47 | 43 | 51 | 62 |
| STANDARD DEVIATION.....GM..... | 65 | 51 | 42 | 41 | 47 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 14 | 4 | 6 | 9 | 10 |
| 25TH.....GM..... | 33 | 14 | 18 | 21 | 28 |
| 50TH.....GM..... | 62 | 32 | 35 | 39 | 50 |
| 75TH.....GM..... | 103 | 58 | 55 | 70 | 85 |
| 90TH.....GM..... | 158 | 132 | 92 | 101 | 125 |
| 95TH.....GM..... | 202 | 176 | 112 | 130 | 152 |
| 99TH.....GM..... | 309 | 253 | 168 | 200 | 216 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 130 | 55 | 69 | 90 | 105 |
| STANDARD DEVIATION.....GM..... | 98 | 49 | 54 | 69 | 79 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 135 | 83 | 77 | 93 | 113 |
| STANDARD DEVIATION.....GM..... | 81 | 61 | 46 | 57 | 66 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 151 | 95 | 90 | 96 | 112 |
| STANDARD DEVIATION.....GM..... | 86 | 63 | 78 | 51 | 49 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,560 | 470 | 1,164 | 586 | 702 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,680 | 758 | 2,152 | 1,260 | 1,179 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 125 | 72 | 70 | 86 | 100 |
| STANDARD DEVIATION.....GM..... | 90 | 58 | 56 | 62 | 69 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 29 | 12 | 14 | 16 | 26 |
| 25TH.....GM..... | 63 | 28 | 34 | 46 | 57 |
| 50TH.....GM..... | 105 | 53 | 57 | 70 | 90 |
| 75TH.....GM..... | 170 | 105 | 93 | 105 | 122 |
| 90TH.....GM..... | 235 | 122 | 123 | 179 | 210 |
| 95TH.....GM..... | 280 | 210 | 170 | 210 | 210 |
| 99TH.....GM..... | 426 | 235 | 250 | 276 | 340 |
| MAXIMUM.....GM..... | 1,260 | 368 | 630 | 508 | 588 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 468.8 | 0 | 7.6 | 17.2 | 29.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .9 | 0 | .5 | .7 | 1.0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 81.8 | 77.0 | 81.2 | 77.2 | 79.6 | 70.5 | 78.1 | 71.3 | 76.4 | 69.4 | 70.9 | 66.3 |
| 35.9 | 35.2 | 34.0 | 35.9 | 34.7 | 36.5 | 32.1 | 34.3 | 32.3 | 33.2 | 31.1 | 29.1 |
| 32.5 | 30.2 | 30.9 | 30.5 | 30.2 | 25.3 | 30.5 | 27.2 | 28.5 | 26.7 | 25.4 | 23.9 |
| 13.4 | 11.6 | 16.2 | 10.9 | 14.7 | 8.7 | 15.5 | 9.7 | 15.6 | 9.5 | 14.4 | 13.3 |
| 2.0 | 1.9 | 2.0 | 1.9 | 2.0 | 1.7 | 2.0 | 1.8 | 2.0 | 1.8 | 2.0 | 1.9 |
| 1.0 | .9 | 1.1 | 1.0 | 1.0 | .9 | 1.0 | .9 | 1.0 | .9 | 1.1 | 1.0 |
| 6.0 | 6.0 | 7.0 | 6.0 | 6.0 | 7.0 | 7.0 | 6.0 | 6.0 | 6.0 | 6.0 | 10.0 |
| 81 | 71 | 101 | 73 | 96 | 69 | 99 | 70 | 91 | 73 | 100 | 76 |
| 63 | 54 | 82 | 60 | 80 | 54 | 78 | 54 | 74 | 52 | 79 | 55 |
| 15 | 11 | 19 | 10 | 18 | 13 | 19 | 15 | 18 | 16 | 20 | 18 |
| 35 | 31 | 42 | 31 | 39 | 31 | 45 | 31 | 41 | 35 | 45 | 35 |
| 66 | 57 | 81 | 60 | 76 | 57 | 77 | 58 | 70 | 62 | 78 | 62 |
| 111 | 93 | 140 | 96 | 129 | 89 | 132 | 90 | 117 | 94 | 125 | 101 |
| 159 | 148 | 193 | 140 | 190 | 138 | 202 | 140 | 172 | 140 | 206 | 146 |
| 198 | 182 | 259 | 188 | 242 | 174 | 250 | 173 | 225 | 162 | 260 | 179 |
| 297 | 260 | 407 | 288 | 358 | 266 | 340 | 266 | 393 | 267 | 365 | 274 |
| 135 | 121 | 167 | 123 | 158 | 124 | 154 | 124 | 145 | 133 | 159 | 123 |
| 102 | 90 | 124 | 93 | 123 | 86 | 109 | 89 | 103 | 84 | 114 | 73 |
| 139 | 126 | 168 | 127 | 162 | 126 | 163 | 126 | 141 | 130 | 160 | 126 |
| 84 | 72 | 100 | 69 | 99 | 64 | 91 | 71 | 73 | 64 | 87 | 71 |
| 148 | 130 | 177 | 147 | 174 | 144 | 179 | 135 | 175 | 137 | 191 | 135 |
| 70 | 63 | 105 | 86 | 103 | 71 | 98 | 67 | 96 | 63 | 91 | 60 |
| 985 | 980 | 1,560 | 945 | 1,260 | 816 | 1,134 | 1,074 | 850 | 926 | 736 | 854 |
| 1,701 | 1,260 | 2,143 | 2,019 | 2,680 | 1,294 | 2,652 | 1,709 | 1,865 | 1,242 | 1,510 | 1,129 |
| 124 | 112 | 149 | 116 | 146 | 119 | 149 | 119 | 140 | 124 | 154 | 119 |
| 87 | 80 | 112 | 86 | 110 | 80 | 103 | 78 | 91 | 70 | 94 | 68 |
| 29 | 26 | 29 | 26 | 29 | 29 | 32 | 29 | 39 | 32 | 53 | 29 |
| 57 | 57 | 68 | 57 | 70 | 60 | 85 | 70 | 85 | 90 | 93 | 83 |
| 105 | 93 | 114 | 93 | 114 | 98 | 122 | 98 | 118 | 105 | 123 | 105 |
| 170 | 140 | 210 | 150 | 210 | 158 | 210 | 145 | 210 | 245 | 210 | 210 |
| 210 | 210 | 276 | 210 | 269 | 210 | 263 | 210 | 184 | 158 | 188 | 140 |
| 276 | 250 | 368 | 254 | 368 | 250 | 366 | 250 | 310 | 245 | 340 | 244 |
| 420 | 420 | 508 | 420 | 531 | 420 | 500 | 420 | 490 | 388 | 488 | 368 |
| 716 | 980 | 1,260 | 945 | 1,260 | 816 | 1,134 | 840 | 732 | 630 | 736 | 490 |
| 42.5 | 35.4 | 25.3 | 26.3 | 67.0 | 44.7 | 69.9 | 61.4 | 12.3 | 20.7 | 4.0 | 5.4 |
| 1.3 | 1.1 | 1.1 | 1.2 | 1.1 | .7 | .9 | .7 | .7 | 1.0 | .5 | .4 |

TABLE 10.02.--BAKED POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 18.3 | 2.1 | 14.2 | 13.7 | 14.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 15.6 | 1.9 | 12.4 | 12.6 | 12.9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.4 | .3 | 1.7 | 1.0 | 1.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | .1 | .1 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 1.1 | 1.1 | 1.1 | 1.1 |
| STANDARD DEVIATION.....CO..... | .4 | .3 | .4 | .3 | .3 |
| MAXIMUM.....DO..... | 5.0 | 2.0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 40 | 22 | 24 | 27 | 31 |
| STANDARD DEVIATION.....GM..... | 22 | 18 | 15 | 15 | 16 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 16 | | 6 | 12 | 16 |
| 25TH.....GM..... | 31 | | 16 | 16 | 23 |
| 50TH.....GM..... | 31 | | 23 | 23 | 31 |
| 75TH.....GM..... | 47 | | 31 | 31 | 31 |
| 90TH.....GM..... | 62 | | 43 | 47 | 47 |
| 95TH.....GM..... | 93 | | 62 | 62 | 61 |
| 99TH.....GM..... | 124 | | 78 | 92 | 113 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 103 | 49 | 62 | 76 | 84 |
| STANDARD DEVIATION.....GM..... | 43 | 30 | 36 | 37 | 28 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 105 | 93 | 67 | 72 | 87 |
| STANDARD DEVIATION.....GM..... | 36 | 0 | 27 | 27 | 34 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 104 | 0 | 35 | 66 | 100 |
| STANDARD DEVIATION.....GM..... | 29 | 0 | 0 | 15 | 38 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 784 | 98 | 255 | 276 | 280 |
| MAXIMUM OVER 3 DAYS.....GM..... | 882 | 186 | 255 | 279 | 420 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 103 | 59 | 62 | 75 | 85 |
| STANDARD DEVIATION.....GM..... | 42 | 32 | 31 | 36 | 30 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 47 | | 18 | 35 | 47 |
| 25TH.....GM..... | 93 | | 47 | 47 | 70 |
| 50TH.....GM..... | 93 | | 53 | 70 | 93 |
| 75TH.....GM..... | 105 | | 70 | 93 | 93 |
| 90TH.....GM..... | 140 | | 93 | 98 | 140 |
| 95TH.....GM..... | 184 | | 113 | 140 | 140 |
| 99TH.....GM..... | 280 | | 186 | 210 | 184 |
| MAXIMUM.....GM..... | 784 | 98 | 227 | 276 | 280 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 10.2 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .1 | 0 | 0 | 0 | 0 |

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 15.6 | 15.4 | 14.1 | 16.3 | 19.1 | 17.8 | 23.4 | 21.1 | 21.4 | 19.1 | 16.0 | 16.0 |
| 14.4 | 14.0 | 12.6 | 14.5 | 15.9 | 15.8 | 19.5 | 17.8 | 16.9 | 15.3 | 12.8 | 11.8 |
| 1.1 | 1.3 | 1.4 | 1.6 | 3.0 | 1.7 | 3.4 | 3.0 | 3.8 | 3.3 | 3.0 | 3.6 |
| .1 | .1 | .1 | .2 | .3 | .3 | .5 | .3 | .7 | .6 | .2 | .6 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.1 | 1.2 | 1.2 | 1.3 | 1.2 | 1.2 | 1.3 |
| .3 | .3 | .3 | .4 | .4 | .4 | .5 | .4 | .6 | .5 | .5 | .5 |
| 3.0 | 4.0 | 3.0 | 3.0 | 4.0 | 4.0 | 4.0 | 5.0 | 4.0 | 5.0 | 3.0 | 3.0 |
| 38 | 36 | 44 | 36 | 48 | 38 | 46 | 39 | 43 | 41 | 44 | 39 |
| 17 | 22 | 23 | 16 | 27 | 18 | 25 | 20 | 21 | 21 | 22 | 20 |
| 23 | 16 | 23 | 23 | 23 | 18 | 23 | 23 | 16 | 18 | 23 | 16 |
| 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 |
| 31 | 31 | 31 | 31 | 35 | 31 | 31 | 31 | 31 | 31 | 31 | 31 |
| 47 | 35 | 47 | 38 | 62 | 47 | 48 | 47 | 48 | 47 | 47 | 50 |
| 62 | 54 | 70 | 62 | 93 | 62 | 78 | 62 | 70 | 62 | 93 | 62 |
| 65 | 62 | 93 | 64 | 97 | 72 | 93 | 78 | 93 | 82 | 93 | 70 |
| 98 | 98 | 155 | 93 | 140 | 109 | 140 | 117 | 109 | 117 | 109 | 109 |
| 104 | 97 | 120 | 97 | 120 | 101 | 115 | 99 | 108 | 99 | 103 | 89 |
| 38 | 38 | 56 | 34 | 53 | 36 | 50 | 36 | 43 | 36 | 23 | 22 |
| 101 | 103 | 112 | 97 | 124 | 103 | 116 | 101 | 96 | 99 | 121 | 90 |
| 28 | 71 | 51 | 16 | 40 | 22 | 36 | 32 | 26 | 23 | 26 | 14 |
| 112 | 105 | 109 | 93 | 133 | 110 | 103 | 98 | 98 | 103 | 93 | 110 |
| 16 | 4 | 0 | 0 | 46 | 25 | 29 | 21 | 9 | 23 | 0 | 17 |
| 350 | 784 | 465 | 280 | 454 | 284 | 560 | 465 | 284 | 350 | 186 | 151 |
| 377 | 882 | 558 | 297 | 746 | 558 | 658 | 558 | 336 | 465 | 326 | 420 |
| 104 | 97 | 117 | 96 | 121 | 101 | 114 | 99 | 102 | 99 | 107 | 91 |
| 38 | 50 | 55 | 30 | 51 | 33 | 48 | 36 | 39 | 33 | 24 | 23 |
| 70 | 47 | 70 | 70 | 70 | 53 | 70 | 53 | 47 | 53 | 93 | 49 |
| 93 | 75 | 93 | 93 | 93 | 93 | 93 | 93 | 93 | 93 | 93 | 70 |
| 93 | 93 | 93 | 93 | 93 | 93 | 93 | 93 | 93 | 93 | 93 | 93 |
| 98 | 93 | 140 | 93 | 140 | 98 | 140 | 98 | 140 | 93 | 140 | 93 |
| 140 | 140 | 186 | 140 | 186 | 140 | 145 | 140 | 140 | 140 | 140 | 140 |
| 186 | 140 | 209 | 140 | 210 | 147 | 186 | 140 | 145 | 145 | 140 | 140 |
| 279 | 280 | 290 | 227 | 280 | 210 | 280 | 280 | 279 | 227 | 186 | 147 |
| 350 | 784 | 465 | 280 | 454 | 284 | 560 | 465 | 284 | 350 | 186 | 151 |
| 0 | 1.2 | 0 | 0 | 3.5 | 1.4 | 2.9 | 0 | 0 | 0 | 0 | 1.2 |
| 0 | .3 | 0 | 0 | .4 | .1 | .2 | 0 | 0 | 0 | 0 | .6 |

TABLE 10.03.--BOILED POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 12.6 | 3.8 | 9.9 | 10.3 | 11.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 10.5 | 2.9 | 8.7 | 9.1 | 10.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.9 | .9 | 1.0 | 1.2 | 1.3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | .1 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.2 | 1.3 | 1.2 | 1.1 | 1.1 |
| STANDARD DEVIATION.....%.....DO..... | .5 | .5 | .5 | .3 | .4 |
| MAXIMUM.....DO..... | 6.0 | 3.0 | 3.0 | 2.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 63 | 37 | 35 | 42 | 47 |
| STANDARD DEVIATION.....GM..... | 49 | 31 | 53 | 27 | 31 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 15 | | 7 | 7 | 15 |
| 25TH.....GM..... | 31 | | 15 | 30 | 30 |
| 50TH.....GM..... | 47 | | 30 | 41 | 41 |
| 75TH.....GM..... | 81 | | 35 | 60 | 60 |
| 90TH.....GM..... | 122 | | 60 | 77 | 81 |
| 95TH.....GM..... | 153 | | 61 | 85 | 102 |
| 99TH.....GM..... | 245 | | 328 | 129 | 169 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 156 | 95 | 81 | 113 | 123 |
| STANDARD DEVIATION.....GM..... | 104 | 95 | 46 | 69 | 71 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 160 | 83 | 165 | 109 | 141 |
| STANDARD DEVIATION.....GM..... | 94 | 29 | 203 | 47 | 56 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 177 | 0 | 11 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 93 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,164 | 368 | 1,164 | 508 | 508 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,701 | 368 | 1,522 | 508 | 627 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 153 | 88 | 87 | 112 | 124 |
| STANDARD DEVIATION.....GM..... | 94 | 77 | 81 | 66 | 67 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 45 | | 11 | 22 | 45 |
| 25TH.....GM..... | 92 | | 45 | 61 | 90 |
| 50TH.....GM..... | 122 | | 90 | 92 | 122 |
| 75TH.....GM..... | 184 | | 92 | 127 | 179 |
| 90TH.....GM..... | 254 | | 127 | 184 | 184 |
| 95TH.....GM..... | 358 | | 184 | 194 | 254 |
| 99TH.....GM..... | 488 | | 537 | 276 | 358 |
| MAXIMUM.....GM..... | 920 | 368 | 627 | 508 | 508 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 31.0 | 0 | .8 | 2.6 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .5 | 0 | .6 | 1.3 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 10.5 | 10.5 | 11.0 | 11.3 | 9.1 | 9.2 | 14.0 | 14.6 | 19.4 | 20.0 | 25.1 | 23.0 |
| 8.9 | 9.0 | 9.3 | 10.0 | 8.1 | 8.2 | 11.1 | 12.2 | 15.4 | 15.4 | 16.7 | 15.3 |
| 1.6 | 1.4 | 1.4 | 1.3 | .9 | 1.0 | 2.4 | 2.2 | 3.3 | 4.0 | 6.8 | 5.1 |
| 0 | .1 | .2 | 0 | .1 | (2) | .5 | .2 | .7 | .6 | 1.6 | 2.6 |
| 1.2 | 1.2 | 1.2 | 1.1 | 1.1 | 1.1 | 1.3 | 1.2 | 1.3 | 1.4 | 1.5 | 1.5 |
| .4 | .4 | .5 | .3 | .4 | .3 | .6 | .5 | .7 | .6 | .8 | .8 |
| 3.0 | 3.0 | 4.0 | 2.0 | 3.0 | 3.0 | 6.0 | 5.0 | 5.0 | 3.0 | 6.0 | 4.0 |
| 62 | 58 | 79 | 57 | 77 | 57 | 79 | 54 | 78 | 61 | 80 | 68 |
| 50 | 40 | 61 | 42 | 49 | 35 | 64 | 37 | 74 | 38 | 62 | 48 |
| 15 | 10 | 9 | 15 | 30 | 15 | 22 | 15 | 30 | 23 | 20 | 30 |
| 31 | 30 | 41 | 31 | 42 | 31 | 41 | 31 | 41 | 37 | 41 | 41 |
| 60 | 42 | 61 | 42 | 61 | 42 | 60 | 41 | 60 | 46 | 60 | 49 |
| 81 | 71 | 90 | 61 | 90 | 81 | 92 | 61 | 81 | 81 | 111 | 90 |
| 121 | 119 | 149 | 119 | 143 | 119 | 163 | 100 | 123 | 119 | 163 | 123 |
| 142 | 153 | 203 | 123 | 163 | 123 | 209 | 122 | 245 | 123 | 207 | 143 |
| 215 | 179 | 298 | 259 | 244 | 171 | 345 | 184 | 469 | 203 | 325 | 285 |
| 155 | 148 | 200 | 151 | 211 | 155 | 184 | 140 | 164 | 154 | 159 | 146 |
| 99 | 102 | 136 | 93 | 124 | 91 | 123 | 93 | 92 | 83 | 109 | 91 |
| 176 | 155 | 207 | 160 | 177 | 150 | 205 | 137 | 188 | 122 | 185 | 151 |
| 121 | 55 | 116 | 102 | 83 | 55 | 127 | 61 | 98 | 53 | 98 | 80 |
| 0 | 152 | 257 | 0 | 276 | 171 | 208 | 121 | 340 | 162 | 170 | 122 |
| 0 | 42 | 25 | 0 | 103 | 0 | 74 | 43 | 118 | 65 | 68 | 42 |
| 985 | 582 | 895 | 776 | 732 | 552 | 1,089 | 1,074 | 732 | 488 | 736 | 854 |
| 1,701 | 582 | 1,196 | 1,074 | 1,074 | 644 | 1,611 | 1,074 | 1,406 | 988 | 1,094 | 976 |
| 157 | 146 | 198 | 150 | 202 | 152 | 185 | 135 | 176 | 135 | 157 | 133 |
| 97 | 87 | 112 | 97 | 112 | 84 | 113 | 77 | 104 | 66 | 92 | 66 |
| 45 | 45 | 45 | 46 | 90 | 45 | 61 | 45 | 90 | 46 | 61 | 47 |
| 92 | 90 | 122 | 90 | 122 | 92 | 113 | 90 | 93 | 92 | 95 | 92 |
| 122 | 122 | 179 | 122 | 179 | 122 | 179 | 122 | 179 | 122 | 127 | 122 |
| 184 | 183 | 244 | 184 | 244 | 184 | 244 | 179 | 184 | 179 | 184 | 142 |
| 269 | 244 | 358 | 244 | 358 | 254 | 358 | 194 | 358 | 194 | 244 | 184 |
| 366 | 358 | 366 | 358 | 448 | 358 | 368 | 276 | 368 | 269 | 269 | 244 |
| 488 | 488 | 626 | 538 | 552 | 448 | 553 | 460 | 552 | 368 | 488 | 368 |
| 716 | 582 | 776 | 776 | 732 | 552 | 920 | 537 | 732 | 426 | 736 | 488 |
| 1.1 | 4.0 | 1.6 | 1.7 | 3.1 | .8 | 3.6 | 10.4 | 1.2 | 0 | 0 | 0 |
| .4 | 1.5 | .9 | .9 | .8 | .1 | .4 | .8 | .4 | 0 | 0 | 0 |

TABLE 10.04.--CREAMED, SCALLOPED, AND AU GRATIN POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.4 | 2.1 | 5.6 | 4.7 | 5.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.8 | 1.6 | 4.8 | 4.1 | 4.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .6 | .1 | .7 | .4 | .6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | .3 | .1 | .1 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 1.6 | 1.2 | 1.1 | 1.2 |
| STANDARD DEVIATION.....DO..... | .4 | 1.3 | .5 | .4 | .4 |
| MAXIMUM.....DO..... | 5.0 | 5.0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 69 | 44 | 35 | 46 | 55 |
| STANDARD DEVIATION.....GM..... | 46 | 75 | 33 | 31 | 37 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 10 | | 5 | 5 | 5 |
| 25TH.....GM..... | 39 | | 10 | 20 | 26 |
| 50TH.....GM..... | 61 | | 20 | 41 | 41 |
| 75TH.....GM..... | 82 | | 41 | 78 | 78 |
| 90TH.....GM..... | 124 | | 82 | 82 | 82 |
| 95TH.....GM..... | 163 | | 98 | 82 | 118 |
| 99TH.....GM..... | 235 | | 157 | 163 | 196 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 189 | 32 | 80 | 127 | 152 |
| STANDARD DEVIATION.....GM..... | 121 | 32 | 59 | 90 | 107 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 179 | 29 | 120 | 113 | 125 |
| STANDARD DEVIATION.....GM..... | 85 | 0 | 74 | 45 | 51 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 113 | 209 | 98 | 52 | 68 |
| STANDARD DEVIATION.....GM..... | 77 | 29 | 0 | 7 | 10 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 980 | 470 | 245 | 490 | 588 |
| MAXIMUM OVER 3 DAYS.....GM..... | 980 | 705 | 470 | 490 | 588 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 179 | 80 | 87 | 119 | 141 |
| STANDARD DEVIATION.....GM..... | 112 | 70 | 65 | 84 | 98 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 29 | | 15 | 15 | 15 |
| 25TH.....GM..... | 118 | | 30 | 61 | 61 |
| 50TH.....GM..... | 123 | | 61 | 118 | 118 |
| 75TH.....GM..... | 245 | | 118 | 123 | 235 |
| 90TH.....GM..... | 245 | | 235 | 235 | 245 |
| 95TH.....GM..... | 372 | | 235 | 245 | 245 |
| 99TH.....GM..... | 490 | | 245 | 353 | 587 |
| MAXIMUM.....GM..... | 980 | 235 | 245 | 490 | 588 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 35.4 | 0 | .7 | 2.0 | 3.2 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.5 | 0 | .9 | 2.1 | 2.9 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 6.0 | 5.2 | 6.9 | 4.1 | 5.5 | 4.4 | 5.7 | 5.4 | 8.6 | 5.9 | 5.2 | 6.4 |
| 5.7 | 4.3 | 6.0 | 3.5 | 5.2 | 3.9 | 5.0 | 4.7 | 8.2 | 5.3 | 5.0 | 5.5 |
| .2 | .9 | 1.0 | .5 | .3 | .5 | .7 | .6 | .4 | .7 | 0 | .4 |
| .1 | 0 | 0 | .1 | (2) | (2) | 0 | (2) | .1 | 0 | .2 | .4 |
| 1.1 | 1.2 | 1.2 | 1.2 | 1.1 | 1.1 | 1.2 | 1.2 | 1.1 | 1.2 | 1.2 | 1.3 |
| .4 | .4 | .4 | .5 | .3 | .4 | .4 | .4 | .3 | .5 | .5 | .6 |
| 4.0 | 2.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| 63 | 67 | 89 | 60 | 87 | 69 | 80 | 67 | 68 | 63 | 79 | 62 |
| 48 | 49 | 52 | 54 | 54 | 41 | 48 | 42 | 43 | 36 | 38 | 39 |
| 15 | 18 | 20 | 9 | 20 | 14 | 20 | 15 | 10 | 20 | | 5 |
| 39 | 39 | 55 | 39 | 41 | 39 | 41 | 39 | 39 | 39 | | 39 |
| 41 | 44 | 78 | 41 | 78 | 78 | 78 | 59 | 61 | 59 | | 64 |
| 82 | 82 | 118 | 78 | 123 | 82 | 82 | 82 | 82 | 82 | | 82 |
| 82 | 127 | 163 | 123 | 163 | 123 | 157 | 123 | 141 | 102 | | 118 |
| 163 | 163 | 163 | 137 | 201 | 163 | 163 | 157 | 163 | 123 | | 163 |
| 313 | 245 | 235 | 313 | 245 | 163 | 245 | 235 | 186 | 212 | | 163 |
| 169 | 176 | 236 | 135 | 254 | 184 | 222 | 180 | 195 | 165 | 230 | 184 |
| 93 | 122 | 117 | 75 | 159 | 103 | 130 | 109 | 122 | 72 | 114 | 106 |
| 228 | 158 | 233 | 216 | 199 | 192 | 191 | 173 | 192 | 192 | 0 | 147 |
| 139 | 93 | 101 | 126 | 74 | 56 | 77 | 75 | 48 | 71 | 0 | 107 |
| 296 | 0 | 0 | 196 | 93 | 54 | 0 | 137 | 117 | 0 | 117 | 44 |
| 19 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 16 |
| 588 | 980 | 790 | 470 | 735 | 490 | 735 | 705 | 558 | 470 | 490 | 490 |
| 940 | 980 | 875 | 940 | 735 | 588 | 735 | 735 | 558 | 637 | 490 | 490 |
| 174 | 169 | 228 | 154 | 240 | 182 | 209 | 171 | 183 | 161 | 194 | 143 |
| 98 | 115 | 119 | 91 | 142 | 95 | 119 | 96 | 107 | 76 | 97 | 106 |
| 31 | 29 | 59 | 28 | 61 | 41 | 59 | 44 | 29 | 29 | | 15 |
| 118 | 118 | 123 | 118 | 123 | 118 | 118 | 118 | 118 | 118 | | 44 |
| 123 | 123 | 235 | 123 | 235 | 235 | 235 | 123 | 123 | 123 | | 118 |
| 235 | 245 | 245 | 235 | 245 | 245 | 245 | 235 | 235 | 235 | | 235 |
| 245 | 245 | 369 | 245 | 490 | 245 | 368 | 245 | 306 | 245 | | 245 |
| 368 | 353 | 470 | 245 | 490 | 368 | 490 | 368 | 372 | 245 | | 353 |
| 490 | 490 | 490 | 470 | 706 | 490 | 490 | 470 | 558 | 470 | | 490 |
| 490 | 980 | 705 | 470 | 735 | 490 | 735 | 490 | 558 | 470 | 490 | 490 |
| 3.0 | 3.6 | 2.8 | 1.5 | 1.0 | 7.7 | 2.0 | 2.6 | 3.6 | .8 | .9 | 0 |
| 2.2 | 2.7 | 2.5 | 2.2 | .4 | 2.9 | .6 | .6 | 3.3 | .7 | 2.7 | 0 |

TABLE 10.05.--MASHED POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 26.5 | 11.5 | 29.0 | 30.1 | 32.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 21.4 | 8.0 | 23.0 | 24.9 | 25.9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 4.5 | 2.1 | 5.1 | 4.4 | 6.1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .6 | 1.4 | .9 | .8 | .9 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 1.6 | 1.3 | 1.2 | 1.3 |
| STANDARD DEVIATION.....DO..... | .5 | 1.0 | .6 | .5 | .5 |
| MAXIMUM.....DO..... | 9.0 | 6.0 | 4.0 | 4.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 67 | 41 | 37 | 46 | 56 |
| STANDARD DEVIATION.....GM..... | 51 | 42 | 29 | 38 | 40 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 18 | 4 | 7 | 8 | 13 |
| 25TH.....GM..... | 35 | 18 | 18 | 18 | 35 |
| 50TH.....GM..... | 70 | 35 | 35 | 35 | 35 |
| 75TH.....GM..... | 70 | 48 | 41 | 70 | 70 |
| 90TH.....GM..... | 140 | 70 | 70 | 70 | 105 |
| 95TH.....GM..... | 140 | 144 | 96 | 123 | 140 |
| 99TH.....GM..... | 280 | 210 | 140 | 175 | 210 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 163 | 73 | 91 | 112 | 133 |
| STANDARD DEVIATION.....GM..... | 100 | 55 | 69 | 75 | 80 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 167 | 98 | 86 | 112 | 135 |
| STANDARD DEVIATION.....GM..... | 96 | 66 | 46 | 59 | 66 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 176 | 100 | 96 | 150 | 152 |
| STANDARD DEVIATION.....GM..... | 104 | 61 | 49 | 91 | 63 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,260 | 368 | 630 | 420 | 525 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,310 | 630 | 630 | 1,260 | 840 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 160 | 79 | 87 | 112 | 130 |
| STANDARD DEVIATION.....GM..... | 98 | 52 | 62 | 70 | 74 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 39 | 25 | 13 | 25 | 26 |
| 25TH.....GM..... | 105 | 39 | 53 | 53 | 105 |
| 50TH.....GM..... | 105 | 53 | 92 | 105 | 105 |
| 75TH.....GM..... | 210 | 105 | 105 | 146 | 210 |
| 90TH.....GM..... | 210 | 151 | 158 | 210 | 210 |
| 95TH.....GM..... | 368 | 210 | 210 | 210 | 210 |
| 99TH.....GM..... | 420 | 221 | 210 | 368 | 420 |
| MAXIMUM.....GM..... | 1,260 | 221 | 630 | 420 | 525 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 68.0 | 0 | 1.0 | 1.0 | 4.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .5 | 0 | .3 | .2 | .5 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 35.9 | 31.6 | 32.3 | 30.1 | 24.9 | 23.2 | 28.2 | 22.5 | 26.0 | 21.9 | 26.2 | 22.5 |
| 28.2 | 25.9 | 25.8 | 25.4 | 20.5 | 19.3 | 21.9 | 18.5 | 19.7 | 16.8 | 19.5 | 17.7 |
| 6.9 | 5.2 | 6.3 | 4.4 | 4.1 | 3.4 | 5.5 | 3.6 | 5.7 | 4.3 | 4.5 | 3.6 |
| .8 | .6 | .2 | .4 | .3 | .5 | .9 | .4 | .5 | .8 | 2.2 | 1.2 |
| 1.3 | 1.2 | 1.3 | 1.2 | 1.2 | 1.2 | 1.3 | 1.2 | 1.3 | 1.3 | 1.4 | 1.3 |
| .5 | .5 | .5 | .5 | .5 | .5 | .6 | .5 | .6 | .6 | .9 | .9 |
| 4.0 | 4.0 | 4.0 | 4.0 | 6.0 | 4.0 | 4.0 | 6.0 | 6.0 | 5.0 | 4.0 | 9.0 |
| 71 | 58 | 86 | 63 | 84 | 63 | 84 | 60 | 69 | 62 | 86 | 61 |
| 46 | 42 | 62 | 45 | 61 | 44 | 60 | 45 | 45 | 45 | 63 | 49 |
| 18 | 13 | 22 | 18 | 27 | 18 | 27 | 13 | 18 | 18 | 35 | 13 |
| 35 | 35 | 53 | 35 | 53 | 35 | 35 | 35 | 35 | 35 | 35 | 35 |
| 70 | 47 | 70 | 62 | 70 | 53 | 70 | 48 | 70 | 53 | 70 | 53 |
| 70 | 70 | 105 | 70 | 105 | 70 | 105 | 70 | 70 | 70 | 105 | 70 |
| 140 | 105 | 140 | 105 | 140 | 105 | 105 | 105 | 140 | 140 | 175 | 116 |
| 148 | 140 | 210 | 140 | 210 | 140 | 210 | 140 | 140 | 140 | 245 | 175 |
| 210 | 210 | 280 | 263 | 350 | 245 | 280 | 216 | 280 | 280 | 292 | 228 |
| 175 | 145 | 211 | 160 | 209 | 160 | 200 | 148 | 166 | 140 | 186 | 133 |
| 102 | 86 | 127 | 91 | 110 | 89 | 109 | 88 | 93 | 74 | 98 | 64 |
| 168 | 146 | 215 | 172 | 219 | 153 | 197 | 155 | 164 | 155 | 204 | 159 |
| 81 | 76 | 91 | 100 | 144 | 74 | 98 | 89 | 83 | 71 | 110 | 76 |
| 151 | 170 | 352 | 150 | 191 | 194 | 222 | 156 | 146 | 165 | 202 | 172 |
| 63 | 94 | 268 | 52 | 111 | 92 | 128 | 101 | 44 | 77 | 55 | 84 |
| 840 | 550 | 1,260 | 945 | 1,260 | 630 | 1,050 | 840 | 630 | 630 | 525 | 420 |
| 840 | 1,260 | 2,100 | 1,050 | 2,310 | 1,260 | 2,100 | 1,260 | 876 | 945 | 876 | 945 |
| 165 | 141 | 206 | 158 | 208 | 158 | 195 | 147 | 159 | 143 | 179 | 136 |
| 91 | 82 | 119 | 94 | 121 | 86 | 105 | 88 | 90 | 77 | 96 | 73 |
| 39 | 33 | 53 | 53 | 69 | 53 | 65 | 26 | 53 | 39 | 69 | 26 |
| 105 | 105 | 105 | 105 | 105 | 105 | 105 | 105 | 105 | 105 | 105 | 105 |
| 158 | 105 | 210 | 105 | 210 | 146 | 210 | 105 | 158 | 105 | 185 | 105 |
| 210 | 210 | 210 | 210 | 210 | 210 | 210 | 210 | 210 | 210 | 210 | 210 |
| 210 | 210 | 315 | 210 | 420 | 210 | 315 | 210 | 210 | 210 | 315 | 210 |
| 420 | 292 | 420 | 315 | 420 | 315 | 420 | 315 | 315 | 210 | 420 | 210 |
| 420 | 420 | 631 | 420 | 631 | 420 | 525 | 420 | 584 | 420 | 438 | 315 |
| 630 | 550 | 1,260 | 945 | 1,260 | 630 | 840 | 840 | 630 | 630 | 525 | 420 |
| 5.4 | 4.1 | 2.9 | 3.5 | 14.1 | 6.3 | 10.5 | 10.1 | 1.5 | 1.4 | 1.9 | 0 |
| .6 | .5 | .5 | .7 | 1.2 | .4 | .6 | .5 | .4 | .3 | .9 | 0 |

TABLE 10.06.--HOME-FRIED AND HASH-BROWNEO POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 12.1 | 1.4 | 13.4 | 11.8 | 12.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 10.3 | 1.4 | 11.0 | 10.1 | 11.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.5 | 0 | 1.9 | 1.5 | 1.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | .5 | .2 | .4 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 1.0 | 1.2 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .5 | .0 | .6 | .4 | .5 |
| MAXIMUM.....DO..... | 5.0 | 1.0 | 4.0 | 3.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 51 | 19 | 30 | 35 | 42 |
| STANDARD DEVIATION.....GM..... | 41 | 18 | 27 | 28 | 30 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 13 | | 7 | 7 | 13 |
| 25TH.....GM..... | 28 | | 14 | 19 | 26 |
| 50TH.....GM..... | 43 | | 28 | 28 | 28 |
| 75TH.....GM..... | 57 | | 28 | 43 | 57 |
| 90TH.....GM..... | 103 | | 57 | 57 | 80 |
| 95TH.....GM..... | 113 | | 61 | 85 | 113 |
| 99TH.....GM..... | 207 | | 198 | 113 | 142 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 130 | 58 | 71 | 85 | 108 |
| STANDARD DEVIATION.....GM..... | 85 | 54 | 42 | 53 | 65 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 130 | 0 | 91 | 96 | 129 |
| STANDARD DEVIATION.....GM..... | 78 | 0 | 68 | 49 | 71 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 151 | 0 | 52 | 125 | 103 |
| STANDARD DEVIATION.....GM..... | 97 | 0 | 57 | 87 | 36 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,020 | 170 | 340 | 340 | 340 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,530 | 170 | 680 | 723 | 680 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 128 | 58 | 72 | 88 | 108 |
| STANDARD DEVIATION.....GM..... | 81 | 54 | 51 | 57 | 65 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 39 | | 8 | 21 | 39 |
| 25TH.....GM..... | 78 | | 43 | 43 | 78 |
| 50TH.....GM..... | 85 | | 64 | 85 | 85 |
| 75TH.....GM..... | 170 | | 85 | 85 | 155 |
| 90TH.....GM..... | 227 | | 155 | 170 | 170 |
| 95TH.....GM..... | 340 | | 170 | 170 | 210 |
| 99TH.....GM..... | 340 | | 340 | 340 | 340 |
| MAXIMUM.....GM..... | 850 | 170 | 340 | 340 | 340 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 58.6 | 0 | .8 | 1.0 | .9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.1 | 0 | .5 | .4 | .3 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 12.2 | 11.2 | 12.1 | 10.8 | 15.1 | 10.0 | 15.1 | 11.2 | 17.8 | 8.8 | 10.1 | 9.4 |
| 10.6 | 9.9 | 9.6 | 9.1 | 12.5 | 8.8 | 12.9 | 9.6 | 14.8 | 7.9 | 7.9 | 8.2 |
| 1.4 | 1.3 | 2.0 | 1.6 | 2.1 | 1.1 | 1.9 | 1.5 | 2.5 | .7 | 1.7 | 1.2 |
| .2 | 0 | .4 | (2) | .5 | .2 | .4 | .1 | .4 | .2 | .5 | 0 |
| 1.2 | 1.1 | 1.3 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.1 | 1.3 | 1.2 |
| .5 | .4 | .6 | .4 | .5 | .5 | .5 | .5 | .5 | .4 | .8 | .4 |
| 5.0 | 3.0 | 4.0 | 3.0 | 4.0 | 5.0 | 5.0 | 5.0 | 5.0 | 3.0 | 5.0 | 2.0 |
| 50 | 47 | 66 | 55 | 66 | 48 | 58 | 44 | 57 | 41 | 66 | 42 |
| 40 | 34 | 43 | 45 | 50 | 33 | 43 | 33 | 49 | 24 | 72 | 24 |
| 14 | 13 | 23 | 13 | 22 | 12 | 14 | 7 | 14 | 14 | 17 | 13 |
| 26 | 26 | 28 | 28 | 28 | 28 | 28 | 26 | 28 | 26 | 35 | 26 |
| 43 | 28 | 57 | 52 | 57 | 43 | 57 | 37 | 52 | 39 | 57 | 28 |
| 57 | 57 | 85 | 57 | 78 | 57 | 57 | 57 | 57 | 57 | 57 | 57 |
| 95 | 85 | 113 | 113 | 113 | 85 | 113 | 85 | 103 | 57 | 108 | 63 |
| 113 | 113 | 155 | 113 | 170 | 113 | 120 | 113 | 140 | 85 | 170 | 103 |
| 255 | 170 | 227 | 283 | 255 | 174 | 227 | 155 | 227 | 113 | 433 | 113 |
| 128 | 122 | 166 | 141 | 164 | 126 | 151 | 117 | 137 | 112 | 134 | 113 |
| 87 | 72 | 89 | 109 | 105 | 83 | 91 | 73 | 78 | 55 | 61 | 59 |
| 154 | 140 | 142 | 134 | 152 | 117 | 142 | 109 | 159 | 117 | 143 | 106 |
| 94 | 82 | 72 | 64 | 75 | 59 | 78 | 75 | 117 | 61 | 55 | 44 |
| 86 | 0 | 175 | 340 | 201 | 122 | 150 | 143 | 187 | 62 | 308 | 0 |
| 59 | 0 | 54 | 0 | 84 | 42 | 117 | 50 | 90 | 21 | 167 | 0 |
| 850 | 340 | 510 | 850 | 1,020 | 816 | 680 | 1,020 | 850 | 340 | 510 | 340 |
| 850 | 680 | 808 | 1,020 | 1,083 | 816 | 1,445 | 1,530 | 1,445 | 510 | 1,300 | 340 |
| 129 | 124 | 157 | 140 | 160 | 121 | 143 | 112 | 142 | 110 | 146 | 107 |
| 82 | 76 | 86 | 102 | 95 | 77 | 83 | 69 | 82 | 58 | 83 | 56 |
| 35 | 39 | 59 | 35 | 53 | 34 | 43 | 21 | 43 | 39 | 68 | 21 |
| 78 | 78 | 85 | 78 | 85 | 78 | 85 | 70 | 85 | 70 | 85 | 78 |
| 85 | 85 | 155 | 128 | 155 | 85 | 128 | 85 | 140 | 85 | 140 | 85 |
| 170 | 170 | 170 | 170 | 170 | 170 | 170 | 170 | 170 | 170 | 170 | 170 |
| 233 | 174 | 340 | 255 | 340 | 170 | 255 | 170 | 255 | 170 | 170 | 170 |
| 310 | 340 | 340 | 340 | 340 | 255 | 340 | 255 | 310 | 170 | 340 | 170 |
| 510 | 340 | 340 | 340 | 510 | 340 | 340 | 340 | 425 | 340 | 465 | 255 |
| 510 | 340 | 510 | 850 | 680 | 816 | 510 | 510 | 510 | 340 | 465 | 255 |
| 5.3 | 5.2 | 2.4 | 3.9 | 11.3 | 2.7 | 8.4 | 12.8 | .8 | 1.4 | 0 | 1.9 |
| 1.8 | 1.9 | 1.1 | 2.1 | 1.5 | .4 | .9 | 1.4 | .3 | .8 | 0 | 1.8 |

TABLE 10.07.--FRENCH FRIED POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 25.3 | 2.6 | 27.7 | 33.2 | 34.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 20.7 | 2.3 | 24.7 | 28.3 | 28.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.9 | .2 | 2.7 | 4.6 | 5.2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .7 | .1 | .3 | .4 | .6 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 1.2 | 1.1 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .5 | .7 | .4 | .5 | .5 |
| MAXIMUM.....DO..... | 6.0 | 4.0 | 3.0 | 3.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 27 | 8 | 16 | 21 | 23 |
| STANDARD DEVIATION.....GM..... | 22 | 4 | 13 | 15 | 17 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 7 | | 5 | 5 | 5 |
| 25TH.....GM..... | 14 | | 10 | 10 | 10 |
| 50TH.....GM..... | 19 | | 10 | 19 | 19 |
| 75TH.....GM..... | 36 | | 19 | 23 | 28 |
| 90TH.....GM..... | 48 | | 29 | 38 | 45 |
| 95TH.....GM..... | 68 | | 45 | 45 | 57 |
| 99TH.....GM..... | 113 | | 87 | 80 | 100 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 66 | 21 | 42 | 53 | 57 |
| STANDARD DEVIATION.....GM..... | 49 | 8 | 34 | 36 | 37 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 67 | 28 | 47 | 53 | 57 |
| STANDARD DEVIATION.....GM..... | 40 | 0 | 29 | 28 | 31 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 73 | 13 | 32 | 49 | 77 |
| STANDARD DEVIATION.....GM..... | 38 | 0 | 5 | 9 | 29 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 816 | 34 | 285 | 272 | 300 |
| MAXIMUM OVER 3 DAYS.....GM..... | 816 | 57 | 285 | 329 | 402 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 64 | 20 | 42 | 51 | 56 |
| STANDARD DEVIATION.....GM..... | 45 | 9 | 32 | 32 | 34 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 14 | | 14 | 14 | 14 |
| 25TH.....GM..... | 29 | | 29 | 29 | 29 |
| 50TH.....GM..... | 57 | | 29 | 57 | 57 |
| 75TH.....GM..... | 70 | | 57 | 57 | 60 |
| 90TH.....GM..... | 114 | | 70 | 86 | 114 |
| 95TH.....GM..... | 136 | | 102 | 120 | 136 |
| 99TH.....GM..... | 268 | | 136 | 136 | 136 |
| MAXIMUM.....GM..... | 680 | 34 | 285 | 272 | 272 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 213.6 | 0 | 4.2 | 7.6 | 14.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.8 | 0 | 1.3 | 1.1 | 1.9 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 35.6 | 33.8 | 38.6 | 34.9 | 37.1 | 26.7 | 23.1 | 16.3 | 12.3 | 7.6 | 8.2 | 6.5 |
| 28.9 | 27.9 | 27.5 | 26.8 | 27.4 | 22.5 | 19.2 | 14.5 | 11.0 | 7.3 | 7.9 | 5.3 |
| 6.0 | 5.0 | 8.5 | 6.4 | 8.0 | 3.8 | 3.4 | 1.7 | 1.3 | .2 | .3 | 1.0 |
| .7 | 1.0 | 2.7 | 1.7 | 1.7 | .5 | .5 | .1 | 0 | .1 | 0 | .2 |
| 1.3 | 1.2 | 1.4 | 1.4 | 1.4 | 1.2 | 1.2 | 1.1 | 1.2 | 1.1 | 1.0 | 1.2 |
| .5 | .5 | .7 | .7 | .7 | .5 | .5 | .4 | .4 | .3 | .2 | .5 |
| 4.0 | 5.0 | 5.0 | 6.0 | 5.0 | 5.0 | 4.0 | 4.0 | 3.0 | 3.0 | 2.0 | 3.0 |
| 28 | 26 | 36 | 29 | 34 | 26 | 27 | 23 | 23 | 22 | 24 | 25 |
| 21 | 21 | 30 | 24 | 28 | 19 | 22 | 18 | 15 | 15 | 17 | 26 |
| 7 | 5 | 10 | 7 | 10 | 7 | 7 | 6 | 10 | 5 | | 5 |
| 17 | 12 | 19 | 17 | 19 | 16 | 17 | 11 | 14 | 10 | | 10 |
| 19 | 19 | 29 | 23 | 23 | 19 | 19 | 19 | 19 | 19 | | 19 |
| 38 | 34 | 44 | 38 | 38 | 34 | 38 | 23 | 29 | 23 | | 19 |
| 50 | 45 | 68 | 52 | 67 | 45 | 48 | 42 | 42 | 45 | | 69 |
| 67 | 62 | 91 | 76 | 91 | 60 | 68 | 48 | 56 | 61 | | 96 |
| 95 | 101 | 159 | 136 | 148 | 95 | 131 | 110 | 76 | 74 | | 108 |
| 71 | 64 | 85 | 68 | 80 | 66 | 69 | 59 | 65 | 62 | 68 | 45 |
| 53 | 45 | 81 | 47 | 59 | 43 | 53 | 35 | 42 | 40 | 50 | 18 |
| 61 | 62 | 78 | 66 | 71 | 68 | 69 | 67 | 57 | 75 | 82 | 100 |
| 33 | 39 | 41 | 39 | 49 | 33 | 41 | 43 | 20 | 25 | 0 | 36 |
| 73 | 72 | 69 | 75 | 81 | 61 | 69 | 80 | 0 | 74 | 0 | 108 |
| 27 | 26 | 31 | 43 | 41 | 39 | 29 | 69 | 0 | 0 | 0 | 0 |
| 544 | 303 | 816 | 408 | 680 | 544 | 680 | 408 | 272 | 204 | 285 | 152 |
| 544 | 544 | 816 | 565 | 734 | 710 | 680 | 587 | 272 | 222 | 285 | 325 |
| 65 | 62 | 75 | 64 | 74 | 64 | 68 | 59 | 61 | 62 | 69 | 63 |
| 45 | 43 | 57 | 44 | 54 | 40 | 50 | 39 | 38 | 39 | 50 | 37 |
| 22 | 15 | 22 | 14 | 22 | 22 | 22 | 14 | 22 | 14 | | 14 |
| 29 | 29 | 57 | 29 | 43 | 35 | 43 | 29 | 38 | 29 | | 29 |
| 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | | 57 |
| 70 | 70 | 86 | 70 | 86 | 70 | 70 | 68 | 68 | 70 | | 67 |
| 114 | 114 | 134 | 114 | 134 | 114 | 114 | 113 | 91 | 134 | | 136 |
| 136 | 136 | 150 | 136 | 170 | 136 | 136 | 136 | 136 | 136 | | 136 |
| 204 | 272 | 327 | 272 | 272 | 204 | 272 | 227 | 227 | 204 | | 152 |
| 544 | 303 | 544 | 340 | 680 | 544 | 680 | 408 | 272 | 204 | 285 | 152 |
| 25.5 | 15.4 | 12.1 | 15.7 | 28.3 | 20.2 | 32.6 | 17.5 | 5.2 | 11.0 | 1.2 | 2.2 |
| 2.7 | 1.7 | 1.6 | 2.2 | 1.4 | 1.2 | 2.4 | 1.3 | 3.3 | 7.8 | 2.6 | 2.9 |

TABLE 10.08.--POTATO SALAD
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 6.4 | .2 | 4.8 | 4.9 | 4.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.5 | 0 | 4.0 | 4.2 | 4.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .8 | .2 | .7 | .7 | .3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | .1 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.2 | 2.0 | 1.2 | 1.2 | 1.1 |
| STANDARD DEVIATION.....DO..... | .5 | .0 | .5 | .4 | .3 |
| MAXIMUM.....DO..... | 5.0 | 2.0 | 3.0 | 3.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 72 | 26 | 33 | 42 | 53 |
| STANDARD DEVIATION.....GM..... | 55 | 0 | 26 | 33 | 33 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 10 | | 3 | 5 | 10 |
| 25TH.....GM..... | 42 | | 21 | 21 | 28 |
| 50TH.....GM..... | 63 | | 28 | 42 | 42 |
| 75TH.....GM..... | 83 | | 42 | 42 | 83 |
| 90TH.....GM..... | 167 | | 63 | 83 | 83 |
| 95TH.....GM..... | 167 | | 104 | 83 | 83 |
| 99TH.....GM..... | 250 | | 125 | 167 | 167 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 187 | 0 | 81 | 111 | 153 |
| STANDARD DEVIATION.....GM..... | 125 | 0 | 50 | 88 | 99 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 203 | 39 | 91 | 111 | 128 |
| STANDARD DEVIATION.....GM..... | 127 | 0 | 54 | 56 | 6 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 134 | 0 | 111 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 41 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,250 | 63 | 250 | 500 | 500 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,000 | 78 | 375 | 500 | 500 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 177 | 39 | 81 | 106 | 146 |
| STANDARD DEVIATION.....GM..... | 110 | 23 | 46 | 80 | 88 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 31 | | 16 | 16 | 30 |
| 25TH.....GM..... | 125 | | 63 | 63 | 83 |
| 50TH.....GM..... | 125 | | 63 | 94 | 125 |
| 75TH.....GM..... | 250 | | 125 | 125 | 250 |
| 90TH.....GM..... | 250 | | 125 | 155 | 250 |
| 95TH.....GM..... | 375 | | 125 | 250 | 250 |
| 99TH.....GM..... | 500 | | 250 | 500 | 500 |
| MAXIMUM.....GM..... | 750 | 63 | 250 | 500 | 500 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 17.6 | 0 | 0 | 0 | 2.1 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | 0 | 0 | 0 | 2.2 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.8 | 5.0 | 6.2 | 5.5 | 7.5 | 6.6 | 8.0 | 7.6 | 5.9 | 5.6 | 3.6 | 3.9 |
| 5.5 | 4.6 | 5.9 | 5.0 | 6.5 | 5.6 | 6.7 | 6.6 | 5.0 | 4.3 | 3.4 | 3.6 |
| .3 | .4 | .4 | .5 | .8 | .9 | 1.2 | .9 | .8 | 1.0 | .2 | .3 |
| 0 | (2) | 0 | 0 | .1 | .1 | (2) | (2) | .2 | .2 | 0 | 0 |
| 1.1 | 1.2 | 1.2 | 1.1 | 1.2 | 1.2 | 1.2 | 1.2 | 1.3 | 1.4 | 1.2 | 1.3 |
| .4 | .5 | .4 | .4 | .6 | .5 | .5 | .5 | .7 | .7 | .4 | .5 |
| 3.0 | 4.0 | 3.0 | 3.0 | 4.0 | 4.0 | 3.0 | 4.0 | 4.0 | 5.0 | 2.0 | 2.0 |
| 62 | 68 | 93 | 63 | 90 | 71 | 85 | 66 | 68 | 63 | 87 | 49 |
| 45 | 45 | 56 | 44 | 72 | 47 | 60 | 53 | 46 | 41 | 65 | 33 |
| 10 | 21 | 32 | 10 | 21 | 10 | 21 | 10 | 5 | 10 | | |
| 42 | 42 | 42 | 42 | 42 | 42 | 42 | 42 | 42 | 41 | | |
| 42 | 42 | 83 | 42 | 83 | 63 | 83 | 42 | 42 | 42 | | |
| 83 | 83 | 125 | 83 | 83 | 83 | 96 | 83 | 83 | 83 | | |
| 104 | 125 | 167 | 89 | 167 | 125 | 167 | 146 | 167 | 125 | | |
| 167 | 167 | 167 | 167 | 250 | 167 | 167 | 167 | 167 | 167 | | |
| 208 | 229 | 250 | 250 | 292 | 250 | 313 | 292 | 167 | 167 | | |
| 177 | 184 | 266 | 164 | 230 | 185 | 218 | 169 | 171 | 159 | 237 | 141 |
| 126 | 114 | 157 | 91 | 145 | 111 | 125 | 118 | 109 | 102 | 165 | 99 |
| 193 | 200 | 226 | 228 | 281 | 181 | 231 | 209 | 191 | 143 | 375 | 112 |
| 75 | 90 | 114 | 97 | 192 | 92 | 134 | 120 | 75 | 64 | 0 | 19 |
| 0 | 132 | 0 | 0 | 163 | 144 | 167 | 57 | 138 | 107 | 0 | 0 |
| 0 | 0 | 0 | 0 | 24 | 13 | 0 | 0 | 0 | 55 | 0 | 0 |
| 750 | 688 | 750 | 500 | 1,250 | 625 | 1,000 | 875 | 500 | 500 | 500 | 500 |
| 750 | 750 | 750 | 750 | 2,000 | 750 | 1,500 | 1,000 | 500 | 500 | 750 | 500 |
| 165 | 171 | 240 | 168 | 218 | 172 | 210 | 167 | 161 | 140 | 218 | 112 |
| 101 | 88 | 125 | 87 | 134 | 97 | 120 | 99 | 86 | 91 | 124 | 71 |
| 30 | 63 | 95 | 31 | 63 | 31 | 63 | 30 | 16 | 16 | | |
| 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 63 | | |
| 125 | 125 | 250 | 125 | 250 | 125 | 250 | 125 | 125 | 125 | | |
| 250 | 250 | 250 | 250 | 250 | 250 | 250 | 250 | 250 | 250 | | |
| 250 | 250 | 375 | 250 | 375 | 250 | 375 | 250 | 250 | 250 | | |
| 375 | 250 | 500 | 250 | 500 | 250 | 500 | 375 | 250 | 250 | | |
| 500 | 375 | 750 | 375 | 750 | 500 | 500 | 500 | 500 | 500 | | |
| 500 | 625 | 750 | 500 | 750 | 500 | 750 | 500 | 500 | 500 | 500 | 250 |
| 1.4 | 0 | 1.1 | 0 | 1.5 | 3.4 | .9 | 4.6 | 0 | 2.4 | 0 | 0 |
| 1.1 | 0 | 1.1 | 0 | .4 | .8 | .2 | .7 | 0 | 1.8 | 0 | 0 |

TABLE 10.09.--CABBAGE SALAD AND COLESLAW
 CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 9.7 | 0 | 3.4 | 4.9 | 8.5 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 8.5 | 0 | 3.1 | 4.4 | 7.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.1 | 0 | .3 | .4 | .8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | 0 | 0 | (2) | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 0 | 1.1 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .5 | 0 | .3 | .4 | .3 |
| MAXIMUM.....DO..... | 6.0 | 0 | 2.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 27 | 0 | 12 | 15 | 19 |
| STANDARD DEVIATION.....GM..... | 20 | 0 | 7 | 13 | 12 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 5 | | | 3 | 3 |
| 25TH.....GM..... | 15 | | | 5 | 10 |
| 50TH.....GM..... | 20 | | | 10 | 20 |
| 75TH.....GM..... | 40 | | | 20 | 20 |
| 90TH.....GM..... | 40 | | | 40 | 40 |
| 95TH.....GM..... | 60 | | | 40 | 40 |
| 99TH.....GM..... | 90 | | | 60 | 60 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 70 | 0 | 36 | 41 | 53 |
| STANDARD DEVIATION.....GM..... | 45 | 0 | 22 | 29 | 31 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 70 | 0 | 17 | 51 | 49 |
| STANDARD DEVIATION.....GM..... | 45 | 0 | 9 | 34 | 29 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 80 | 0 | 0 | 2 | 0 |
| STANDARD DEVIATION.....GM..... | 47 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,080 | 0 | 68 | 120 | 120 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,140 | 0 | 68 | 240 | 240 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 68 | 0 | 33 | 41 | 51 |
| STANDARD DEVIATION.....GM..... | 45 | 0 | 22 | 31 | 31 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 15 | | | 5 | 8 |
| 25TH.....GM..... | 40 | | | 15 | 30 |
| 50TH.....GM..... | 60 | | | 30 | 60 |
| 75TH.....GM..... | 90 | | | 60 | 60 |
| 90TH.....GM..... | 120 | | | 68 | 120 |
| 95TH.....GM..... | 120 | | | 120 | 120 |
| 99TH.....GM..... | 240 | | | 120 | 120 |
| MAXIMUM.....GM..... | 1,020 | 0 | 68 | 120 | 120 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 73.7 | 0 | 0 | .8 | 2.6 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.7 | 0 | 0 | .8 | 1.5 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 9.6 | 9.3 | 9.8 | 9.5 | 9.7 | 8.5 | 11.8 | 12.1 | 10.6 | 10.7 | 8.1 | 10.7 |
| 8.4 | 8.2 | 8.3 | 8.5 | 8.4 | 7.7 | 9.9 | 10.5 | 8.6 | 9.2 | 7.9 | 9.1 |
| 1.1 | 1.0 | 1.4 | .9 | 1.1 | .8 | 1.6 | 1.5 | 1.9 | 1.5 | .3 | 1.3 |
| .1 | (2) | 0 | .1 | .1 | .1 | .2 | .1 | .1 | .1 | 0 | .3 |
| 1.2 | 1.2 | 1.2 | 1.1 | 1.2 | 1.1 | 1.2 | 1.2 | 1.2 | 1.2 | 1.1 | 1.2 |
| .5 | .4 | .5 | .4 | .5 | .4 | .6 | .4 | .5 | .5 | .3 | .5 |
| 5.0 | 3.0 | 3.0 | 3.0 | 4.0 | 4.0 | 6.0 | 4.0 | 3.0 | 4.0 | 2.0 | 3.0 |
| 24 | 24 | 31 | 25 | 31 | 26 | 32 | 26 | 29 | 26 | 24 | 28 |
| 15 | 17 | 22 | 18 | 22 | 26 | 22 | 17 | 22 | 21 | 11 | 21 |
| 5 | 5 | 10 | 8 | 8 | 5 | 10 | 5 | 5 | 5 | | 3 |
| 13 | 10 | 15 | 15 | 20 | 13 | 20 | 15 | 13 | 10 | | 15 |
| 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 | | 20 |
| 30 | 33 | 40 | 40 | 40 | 40 | 40 | 40 | 40 | 30 | | 40 |
| 40 | 40 | 60 | 40 | 60 | 40 | 60 | 43 | 60 | 40 | | 60 |
| 50 | 60 | 80 | 53 | 80 | 47 | 80 | 60 | 80 | 73 | | 80 |
| 80 | 80 | 120 | 120 | 100 | 80 | 100 | 80 | 120 | 140 | | 100 |
| 62 | 65 | 77 | 68 | 83 | 68 | 82 | 69 | 66 | 61 | 69 | 70 |
| 34 | 42 | 54 | 44 | 57 | 45 | 46 | 42 | 41 | 32 | 32 | 45 |
| 68 | 61 | 88 | 63 | 70 | 64 | 71 | 68 | 88 | 88 | 53 | 77 |
| 29 | 40 | 38 | 29 | 29 | 96 | 35 | 30 | 41 | 53 | 7 | 32 |
| 37 | 47 | 0 | 88 | 119 | 92 | 94 | 56 | 50 | 60 | 0 | 93 |
| 12 | 0 | 0 | 30 | 28 | 35 | 65 | 14 | 0 | 0 | 0 | 10 |
| 180 | 240 | 360 | 425 | 540 | 1,080 | 240 | 240 | 240 | 360 | 120 | 240 |
| 240 | 360 | 360 | 425 | 540 | 1,140 | 660 | 300 | 360 | 450 | 120 | 300 |
| 60 | 61 | 77 | 66 | 78 | 67 | 78 | 66 | 70 | 64 | 64 | 71 |
| 34 | 40 | 51 | 41 | 49 | 63 | 43 | 39 | 43 | 41 | 33 | 38 |
| 15 | 14 | 30 | 15 | 23 | 14 | 30 | 15 | 15 | 15 | | 23 |
| 30 | 30 | 40 | 40 | 60 | 30 | 60 | 40 | 40 | 30 | | 46 |
| 60 | 60 | 60 | 60 | 60 | 60 | 60 | 60 | 60 | 60 | | 60 |
| 60 | 60 | 120 | 80 | 120 | 60 | 120 | 90 | 120 | 60 | | 120 |
| 120 | 120 | 120 | 120 | 120 | 120 | 120 | 120 | 120 | 120 | | 120 |
| 120 | 120 | 120 | 120 | 180 | 120 | 120 | 120 | 120 | 120 | | 120 |
| 120 | 180 | 240 | 120 | 240 | 240 | 240 | 180 | 240 | 180 | | 180 |
| 180 | 240 | 360 | 425 | 360 | 1,020 | 240 | 240 | 240 | 360 | 120 | 180 |
| 2.4 | 4.8 | 5.5 | 0 | 9.5 | 16.2 | 17.0 | 12.1 | 0 | 2.0 | 0 | .9 |
| 1.0 | 2.0 | 3.4 | 0 | 2.1 | 3.1 | 2.4 | 1.2 | 0 | .9 | 0 | .7 |

TABLE 10.10.--RAW CARROTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.0 | .8 | 3.4 | 5.4 | 9.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.0 | .6 | 3.0 | 4.3 | 7.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .8 | .2 | .4 | .9 | 1.5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | 0 | .1 | .2 | .5 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 1.3 | 1.2 | 1.3 | 1.3 |
| STANDARD DEVIATION.....DO..... | .6 | .5 | .8 | .7 | .8 |
| MAXIMUM.....DO..... | 7.0 | 2.0 | 6.0 | 6.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 18 | 16 | 11 | 17 | 17 |
| STANDARD DEVIATION.....GM..... | 19 | 11 | 13 | 18 | 22 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 1 | | | 1 | 1 |
| 25TH.....GM..... | 5 | | | 4 | 4 |
| 50TH.....GM..... | 17 | | | 13 | 10 |
| 75TH.....GM..... | 22 | | | 20 | 20 |
| 90TH.....GM..... | 41 | | | 33 | 37 |
| 95TH.....GM..... | 51 | | | 50 | 51 |
| 99TH.....GM..... | 96 | | | 122 | 110 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 44 | 27 | 29 | 36 | 38 |
| STANDARD DEVIATION.....GM..... | 44 | 1 | 27 | 29 | 45 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 47 | 50 | 18 | 50 | 34 |
| STANDARD DEVIATION.....GM..... | 35 | 0 | 16 | 39 | 24 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 46 | 0 | 77 | 40 | 73 |
| STANDARD DEVIATION.....GM..... | 31 | 0 | 0 | 28 | 48 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 500 | 50 | 112 | 183 | 300 |
| MAXIMUM OVER 3 DAYS.....GM..... | 500 | 100 | 232 | 366 | 495 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 43 | 37 | 28 | 38 | 38 |
| STANDARD DEVIATION.....GM..... | 40 | 12 | 25 | 33 | 41 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 4 | | | 4 | 4 |
| 25TH.....GM..... | 13 | | | 12 | 10 |
| 50TH.....GM..... | 31 | | | 31 | 28 |
| 75TH.....GM..... | 55 | | | 54 | 50 |
| 90TH.....GM..... | 100 | | | 61 | 72 |
| 95TH.....GM..... | 122 | | | 84 | 110 |
| 99TH.....GM..... | 183 | | | 183 | 165 |
| MAXIMUM.....GM..... | 500 | 50 | 100 | 183 | 300 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 7.0 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 8.6 | 6.5 | 4.5 | 5.5 | 3.4 | 5.2 | 4.0 | 4.9 | 3.8 | 3.9 | .9 | 2.8 |
| 7.0 | 5.5 | 3.9 | 4.7 | 2.9 | 4.1 | 3.2 | 4.2 | 3.1 | 2.9 | .9 | 1.8 |
| 1.3 | .9 | .6 | .5 | .5 | 1.0 | .6 | .5 | .7 | .8 | 0 | .8 |
| .2 | .2 | .1 | .3 | .1 | .1 | .2 | .2 | 0 | .2 | 0 | .2 |
| 1.3 | 1.2 | 1.2 | 1.3 | 1.2 | 1.3 | 1.3 | 1.2 | 1.2 | 1.4 | 1.0 | 1.5 |
| .6 | .5 | .4 | .9 | .5 | .5 | .7 | .5 | .4 | .8 | .0 | .8 |
| 4.0 | 4.0 | 3.0 | 7.0 | 3.0 | 3.0 | 4.0 | 4.0 | 2.0 | 4.0 | 1.0 | 4.0 |
| 16 | 13 | 16 | 17 | 23 | 21 | 23 | 16 | 12 | 22 | 24 | 16 |
| 18 | 13 | 15 | 19 | 24 | 18 | 23 | 14 | 18 | 20 | 12 | 14 |
| 1 | 1 | 1 | 1 | 4 | 3 | 2 | 1 | | 2 | | |
| 3 | 4 | 5 | 4 | 9 | 9 | 8 | 5 | | 9 | | |
| 10 | 9 | 11 | 9 | 17 | 17 | 17 | 17 | | 17 | | |
| 22 | 18 | 20 | 19 | 33 | 30 | 33 | 20 | | 33 | | |
| 37 | 33 | 40 | 50 | 41 | 46 | 50 | 36 | | 51 | | |
| 48 | 37 | 50 | 58 | 81 | 51 | 74 | 41 | | 61 | | |
| 73 | 65 | 72 | 96 | 167 | 85 | 102 | 81 | | 87 | | |
| 38 | 33 | 43 | 33 | 61 | 52 | 57 | 41 | 29 | 48 | 72 | 37 |
| 40 | 31 | 42 | 29 | 64 | 45 | 54 | 35 | 35 | 43 | 36 | 42 |
| 45 | 40 | 35 | 77 | 59 | 53 | 52 | 42 | 36 | 57 | 0 | 34 |
| 39 | 28 | 19 | 38 | 50 | 29 | 35 | 30 | 46 | 37 | 0 | 17 |
| 49 | 17 | 71 | 51 | 38 | 42 | 53 | 32 | 0 | 51 | 0 | 21 |
| 16 | 10 | 0 | 24 | 27 | 5 | 37 | 16 | 0 | 0 | 0 | 0 |
| 300 | 200 | 216 | 150 | 500 | 300 | 305 | 200 | 165 | 183 | 150 | 110 |
| 400 | 200 | 216 | 288 | 500 | 300 | 324 | 268 | 300 | 261 | 150 | 110 |
| 39 | 33 | 42 | 39 | 57 | 51 | 53 | 39 | 30 | 48 | 72 | 33 |
| 36 | 31 | 39 | 35 | 60 | 40 | 48 | 32 | 33 | 37 | 36 | 32 |
| 4 | 4 | 4 | 4 | 7 | 8 | 7 | 4 | | 4 | | |
| 8 | 8 | 12 | 13 | 20 | 28 | 20 | 13 | | 25 | | |
| 28 | 25 | 30 | 25 | 50 | 50 | 50 | 28 | | 50 | | |
| 55 | 50 | 61 | 50 | 61 | 61 | 72 | 50 | | 55 | | |
| 100 | 72 | 72 | 72 | 122 | 100 | 100 | 72 | | 100 | | |
| 100 | 100 | 120 | 122 | 144 | 122 | 144 | 108 | | 100 | | |
| 150 | 144 | 216 | 150 | 244 | 200 | 300 | 150 | | 183 | | |
| 220 | 200 | 216 | 150 | 500 | 300 | 305 | 200 | 150 | 183 | 150 | 110 |
| .9 | .9 | 2.7 | 0 | 0 | 1.4 | 0 | 0 | 0 | 1.1 | 0 | 0 |
| .4 | .5 | 3.8 | 0 | 0 | .4 | 0 | 0 | 0 | 1.2 | 0 | 0 |

TABLE 10.11.--RAW CELERY
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.5 | .2 | 1.8 | 3.1 | 5.4 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.6 | .2 | 1.6 | 2.5 | 4.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .7 | 0 | .1 | .3 | .8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | 0 | 0 | .3 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.3 | 1.0 | 1.1 | 1.4 | 1.3 |
| STANDARD DEVIATION.....DO..... | .6 | .0 | .3 | .8 | .6 |
| MAXIMUM.....DO..... | 5.0 | 1.0 | 2.0 | 5.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 14 | 3 | 8 | 13 | 12 |
| STANDARD DEVIATION.....GM..... | 13 | 0 | 6 | 14 | 9 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 3 | | | 2 | 3 |
| 25TH.....GM..... | 7 | | | 6 | 6 |
| 50TH.....GM..... | 10 | | | 9 | 9 |
| 75TH.....GM..... | 19 | | | 13 | 13 |
| 90TH.....GM..... | 28 | | | 20 | 20 |
| 95TH.....GM..... | 40 | | | 37 | 28 |
| 99TH.....GM..... | 65 | | | 76 | 59 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 34 | 10 | 25 | 27 | 29 |
| STANDARD DEVIATION.....GM..... | 26 | 0 | 18 | 17 | 19 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 36 | 0 | 14 | 20 | 29 |
| STANDARD DEVIATION.....GM..... | 25 | 0 | 6 | 7 | 23 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 38 | 0 | 0 | 51 | 25 |
| STANDARD DEVIATION.....GM..... | 17 | 0 | 0 | 22 | 2 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 280 | 10 | 84 | 112 | 120 |
| MAXIMUM OVER 3 DAYS.....GM..... | 330 | 10 | 84 | 228 | 176 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 33 | 10 | 22 | 29 | 28 |
| STANDARD DEVIATION.....GM..... | 24 | 0 | 14 | 18 | 20 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 8 | | | 4 | 8 |
| 25TH.....GM..... | 17 | | | 15 | 14 |
| 50TH.....GM..... | 28 | | | 28 | 25 |
| 75TH.....GM..... | 40 | | | 40 | 30 |
| 90TH.....GM..... | 60 | | | 56 | 51 |
| 95TH.....GM..... | 80 | | | 60 | 60 |
| 99TH.....GM..... | 120 | | | 60 | 120 |
| MAXIMUM.....GM..... | 204 | 10 | 60 | 84 | 120 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 58.7 | 0 | 0 | 2.3 | 4.8 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.6 | 0 | 0 | 3.2 | 3.9 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.7 | 4.8 | 3.0 | 4.1 | 3.0 | 4.0 | 4.9 | 5.7 | 7.2 | 5.5 | 4.3 | 5.4 |
| 5.2 | 4.1 | 2.5 | 3.5 | 2.4 | 3.4 | 3.6 | 4.4 | 5.1 | 4.2 | 3.2 | 3.9 |
| .4 | .7 | .4 | .4 | .5 | .5 | .9 | 1.1 | 1.6 | 1.0 | 1.1 | 1.1 |
| .1 | (2) | .1 | .3 | .1 | (2) | .4 | .3 | .4 | .3 | 0 | .3 |
| 1.1 | 1.2 | 1.2 | 1.3 | 1.3 | 1.3 | 1.4 | 1.4 | 1.4 | 1.3 | 1.3 | 1.4 |
| .4 | .4 | .4 | .7 | .6 | .5 | .7 | .7 | .7 | .7 | .4 | .7 |
| 3.0 | 3.0 | 3.0 | 4.0 | 4.0 | 5.0 | 4.0 | 5.0 | 5.0 | 5.0 | 2.0 | 4.0 |
| 12 | 11 | 12 | 16 | 14 | 14 | 18 | 14 | 17 | 15 | 18 | 14 |
| 10 | 7 | 10 | 15 | 10 | 12 | 16 | 14 | 14 | 10 | 12 | 11 |
| 3 | 3 | | 2 | 3 | 3 | 3 | 3 | 3 | 5 | | 3 |
| 6 | 6 | | 6 | 7 | 7 | 7 | 6 | 9 | 9 | | 6 |
| 9 | 9 | | 13 | 13 | 10 | 13 | 10 | 11 | 11 | | 10 |
| 13 | 13 | | 19 | 19 | 19 | 23 | 19 | 19 | 19 | | 19 |
| 23 | 20 | | 33 | 28 | 27 | 39 | 30 | 37 | 28 | | 33 |
| 33 | 27 | | 40 | 37 | 43 | 45 | 40 | 51 | 40 | | 36 |
| 40 | 37 | | 110 | 40 | 65 | 100 | 77 | 67 | 41 | | 45 |
| 30 | 27 | 30 | 37 | 39 | 37 | 40 | 32 | 37 | 37 | 41 | 33 |
| 22 | 16 | 26 | 27 | 25 | 31 | 28 | 29 | 24 | 22 | 28 | 25 |
| 32 | 30 | 31 | 35 | 28 | 35 | 48 | 38 | 36 | 33 | 47 | 28 |
| 9 | 12 | 16 | 16 | 15 | 19 | 33 | 28 | 30 | 14 | 11 | 18 |
| 51 | 30 | 30 | 48 | 41 | 62 | 35 | 33 | 39 | 40 | 0 | 35 |
| 17 | 0 | 0 | 29 | 19 | 4 | 15 | 14 | 3 | 9 | 0 | 2 |
| 120 | 90 | 153 | 240 | 120 | 280 | 200 | 204 | 168 | 120 | 120 | 100 |
| 200 | 112 | 153 | 330 | 204 | 280 | 300 | 280 | 200 | 177 | 120 | 136 |
| 31 | 27 | 30 | 37 | 35 | 34 | 39 | 32 | 36 | 35 | 44 | 30 |
| 22 | 16 | 24 | 26 | 22 | 24 | 29 | 27 | 25 | 19 | 25 | 18 |
| 10 | 8 | | 10 | 8 | 8 | 8 | 8 | 7 | 14 | | 8 |
| 17 | 17 | | 20 | 17 | 17 | 20 | 15 | 20 | 20 | | 17 |
| 28 | 28 | | 30 | 28 | 28 | 30 | 28 | 28 | 30 | | 28 |
| 34 | 31 | | 50 | 50 | 40 | 56 | 34 | 40 | 40 | | 40 |
| 56 | 50 | | 60 | 56 | 60 | 68 | 60 | 60 | 56 | | 60 |
| 84 | 56 | | 100 | 70 | 80 | 85 | 84 | 80 | 60 | | 68 |
| 102 | 80 | | 120 | 112 | 136 | 136 | 168 | 120 | 120 | | 80 |
| 120 | 90 | 153 | 120 | 120 | 140 | 200 | 204 | 168 | 120 | 120 | 80 |
| 1.1 | 4.8 | 2.2 | .9 | 5.4 | 10.6 | 5.3 | 14.3 | 1.8 | 2.5 | 2.7 | 0 |
| .8 | 4.0 | 4.4 | 1.2 | 3.6 | 4.0 | 1.6 | 2.6 | 1.6 | 2.0 | 9.3 | 0 |

TABLE 10.12.--RAW CUCUMBERS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 5.6 | .6 | 1.6 | 3.5 | 4.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.1 | .4 | 1.3 | 2.4 | 3.1 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.2 | .3 | .3 | 1.2 | .8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | 0 | 0 | .2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.4 | 1.4 | 1.2 | 1.4 | 1.3 |
| STANDARD DEVIATION.....DO..... | .7 | .5 | .4 | .6 | .6 |
| MAXIMUM.....DO..... | 6.0 | 2.0 | 2.0 | 3.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 37 | 30 | 16 | 27 | 30 |
| STANDARD DEVIATION.....GM..... | 45 | 26 | 14 | 32 | 43 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 3 | | | 1 | 3 |
| 25TH.....GM..... | 12 | | | 7 | 5 |
| 50TH.....GM..... | 23 | | | 15 | 18 |
| 75TH.....GM..... | 47 | | | 37 | 37 |
| 90TH.....GM..... | 79 | | | 53 | 75 |
| 95TH.....GM..... | 105 | | | 105 | 105 |
| 99TH.....GM..... | 220 | | | 179 | 330 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 77 | 24 | 46 | 50 | 66 |
| STANDARD DEVIATION.....GM..... | 72 | 16 | 40 | 46 | 70 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 88 | 89 | 30 | 72 | 52 |
| STANDARD DEVIATION.....GM..... | 80 | 0 | 20 | 68 | 51 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 105 | 0 | 0 | 0 | 138 |
| STANDARD DEVIATION.....GM..... | 99 | 0 | 0 | 0 | 117 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 840 | 158 | 110 | 316 | 440 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,580 | 178 | 110 | 536 | 990 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 80 | 63 | 40 | 58 | 68 |
| STANDARD DEVIATION.....GM..... | 76 | 63 | 36 | 50 | 73 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 8 | | | 8 | 8 |
| 25TH.....GM..... | 24 | | | 17 | 18 |
| 50TH.....GM..... | 70 | | | 46 | 40 |
| 75TH.....GM..... | 110 | | | 79 | 79 |
| 90TH.....GM..... | 158 | | | 140 | 140 |
| 95TH.....GM..... | 220 | | | 158 | 220 |
| 99TH.....GM..... | 316 | | | 220 | 316 |
| MAXIMUM.....GM..... | 840 | 158 | 110 | 220 | 440 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 22.3 | 0 | 0 | 0 | 1.0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .8 | 0 | 0 | 0 | 1.0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.2 | 4.6 | 3.9 | 6.3 | 4.2 | 6.0 | 6.7 | 8.0 | 6.1 | 7.2 | 6.4 | 3.3 |
| 2.2 | 3.7 | 3.5 | 5.0 | 3.5 | 4.7 | 4.5 | 5.8 | 4.5 | 4.6 | 3.3 | 2.1 |
| .9 | .7 | .2 | .9 | .7 | 1.1 | 1.8 | 1.8 | 1.2 | 1.4 | 2.0 | 1.0 |
| .1 | .2 | .2 | .3 | .1 | .2 | .4 | .4 | .3 | 1.2 | 1.0 | .2 |
| 1.4 | 1.3 | 1.2 | 1.3 | 1.2 | 1.3 | 1.5 | 1.4 | 1.4 | 1.6 | 1.8 | 1.7 |
| .6 | .6 | .5 | .8 | .5 | .6 | .7 | .7 | .6 | 1.0 | .8 | 1.0 |
| 4.0 | 4.0 | 3.0 | 6.0 | 3.0 | 3.0 | 5.0 | 4.0 | 3.0 | 5.0 | 3.0 | 4.0 |
| 34 | 31 | 31 | 27 | 38 | 36 | 47 | 36 | 33 | 45 | 54 | 28 |
| 28 | 38 | 30 | 32 | 54 | 39 | 53 | 39 | 33 | 74 | 37 | 20 |
| 4 | 3 | 3 | 3 | 3 | 4 | 4 | 3 | 8 | 5 | | |
| 12 | 7 | 12 | 7 | 9 | 12 | 16 | 12 | 12 | 12 | | |
| 23 | 18 | 18 | 18 | 23 | 23 | 29 | 23 | 23 | 26 | | |
| 47 | 37 | 47 | 37 | 37 | 53 | 70 | 47 | 37 | 53 | | |
| 73 | 73 | 78 | 68 | 93 | 73 | 97 | 76 | 92 | 73 | | |
| 76 | 100 | 93 | 73 | 108 | 93 | 146 | 105 | 93 | 110 | | |
| 147 | 220 | 140 | 158 | 373 | 220 | 293 | 220 | 167 | 526 | | |
| 77 | 69 | 75 | 63 | 80 | 84 | 84 | 76 | 82 | 80 | 89 | 86 |
| 64 | 90 | 70 | 72 | 85 | 68 | 71 | 71 | 57 | 65 | 54 | 59 |
| 76 | 99 | 77 | 74 | 132 | 80 | 109 | 86 | 78 | 70 | 99 | 29 |
| 52 | 80 | 34 | 77 | 140 | 72 | 87 | 72 | 83 | 49 | 47 | 19 |
| 59 | 57 | 110 | 63 | 149 | 118 | 137 | 87 | 42 | 119 | 110 | 70 |
| 20 | 23 | 20 | 21 | 92 | 92 | 102 | 75 | 38 | 155 | 0 | 0 |
| 220 | 660 | 280 | 474 | 840 | 440 | 500 | 632 | 280 | 632 | 220 | 220 |
| 440 | 660 | 420 | 660 | 1,120 | 948 | 1,220 | 880 | 500 | 1,580 | 330 | 220 |
| 75 | 72 | 76 | 62 | 96 | 83 | 96 | 78 | 71 | 83 | 93 | 50 |
| 58 | 82 | 64 | 64 | 115 | 72 | 80 | 67 | 64 | 84 | 50 | 52 |
| 12 | 8 | 8 | 8 | 8 | 12 | 12 | 9 | 12 | 9 | | |
| 35 | 20 | 32 | 17 | 24 | 32 | 35 | 28 | 20 | 17 | | |
| 55 | 46 | 55 | 55 | 55 | 70 | 70 | 70 | 53 | 53 | | |
| 110 | 105 | 110 | 70 | 110 | 110 | 110 | 110 | 110 | 110 | | |
| 158 | 158 | 140 | 140 | 220 | 185 | 220 | 158 | 158 | 220 | | |
| 220 | 220 | 220 | 220 | 280 | 220 | 280 | 220 | 220 | 316 | | |
| 220 | 280 | 280 | 280 | 840 | 316 | 316 | 280 | 280 | 316 | | |
| 220 | 660 | 280 | 474 | 840 | 440 | 440 | 440 | 280 | 316 | 220 | 220 |
| 1.2 | .8 | 1.8 | 0 | 1.9 | 3.1 | 2.1 | 4.7 | 1.8 | 2.0 | 0 | 1.8 |
| 1.3 | .6 | 2.7 | 0 | 1.0 | .8 | .4 | .6 | 1.8 | 1.0 | 0 | 3.2 |

TABLE 10.13.--LETTUCE AND TOSSED SALADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 50.7 | 0 | 16.6 | 30.4 | 42.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 26.6 | 0 | 12.7 | 20.5 | 26.4 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 16.4 | 0 | 3.1 | 7.5 | 12.4 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 7.8 | 0 | .8 | 2.3 | 4.0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.8 | 0 | 1.4 | 1.5 | 1.6 |
| STANDARD DEVIATION.....DO..... | 1.0 | 0 | .8 | .8 | .9 |
| MAXIMUM.....DO..... | 8.0 | 0 | 5.0 | 5.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 40 | 0 | 14 | 17 | 23 |
| STANDARD DEVIATION.....GM..... | 39 | 0 | 15 | 15 | 21 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 3 | | 2 | 2 | 3 |
| 25TH.....GM..... | 13 | | 4 | 5 | 9 |
| 50TH.....GM..... | 31 | | 9 | 13 | 18 |
| 75TH.....GM..... | 54 | | 18 | 23 | 31 |
| 90TH.....GM..... | 90 | | 32 | 36 | 47 |
| 95TH.....GM..... | 113 | | 47 | 43 | 67 |
| 99TH.....GM..... | 186 | | 65 | 71 | 104 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 67 | 0 | 29 | 34 | 43 |
| STANDARD DEVIATION.....GM..... | 62 | 0 | 26 | 29 | 34 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 73 | 0 | 34 | 39 | 48 |
| STANDARD DEVIATION.....GM..... | 54 | 0 | 27 | 23 | 31 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 80 | 0 | 44 | 35 | 49 |
| STANDARD DEVIATION.....GM..... | 55 | 0 | 25 | 21 | 32 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,080 | 0 | 186 | 186 | 216 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,296 | 0 | 270 | 326 | 432 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 65 | 0 | 30 | 34 | 43 |
| STANDARD DEVIATION.....GM..... | 59 | 0 | 29 | 26 | 33 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 10 | | 5 | 7 | 10 |
| 25TH.....GM..... | 20 | | 10 | 14 | 14 |
| 50TH.....GM..... | 55 | | 23 | 28 | 30 |
| 75TH.....GM..... | 93 | | 47 | 47 | 55 |
| 90TH.....GM..... | 140 | | 55 | 67 | 93 |
| 95TH.....GM..... | 186 | | 93 | 93 | 108 |
| 99TH.....GM..... | 270 | | 135 | 108 | 140 |
| MAXIMUM.....GM..... | 1,080 | 0 | 186 | 156 | 216 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 569.9 | 0 | .7 | 4.4 | 19.4 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.6 | 0 | .3 | .6 | 1.5 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 45.8 | 47.5 | 47.7 | 49.0 | 54.2 | 56.5 | 59.3 | 60.9 | 49.5 | 49.4 | 33.1 | 40.7 |
| 28.1 | 27.7 | 25.9 | 27.4 | 29.9 | 30.3 | 27.7 | 28.3 | 23.8 | 24.8 | 15.8 | 19.1 |
| 13.6 | 15.5 | 15.5 | 15.2 | 16.0 | 18.9 | 20.4 | 21.1 | 15.4 | 16.0 | 10.1 | 12.4 |
| 4.1 | 4.4 | 6.3 | 6.4 | 8.3 | 7.3 | 11.1 | 11.4 | 10.3 | 8.6 | 7.3 | 9.2 |
| 1.6 | 1.7 | 1.8 | 1.7 | 1.8 | 1.8 | 2.0 | 2.0 | 1.9 | 1.9 | 1.9 | 1.9 |
| .8 | .9 | 1.0 | .9 | 1.0 | 1.0 | 1.1 | 1.1 | 1.0 | 1.1 | 1.0 | 1.0 |
| 5.0 | 5.0 | 6.0 | 5.0 | 6.0 | 6.0 | 8.0 | 6.0 | 6.0 | 6.0 | 6.0 | 6.0 |
| 29 | 28 | 37 | 33 | 41 | 43 | 47 | 46 | 41 | 39 | 39 | 37 |
| 28 | 26 | 34 | 31 | 41 | 44 | 42 | 43 | 40 | 37 | 36 | 36 |
| 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 |
| 9 | 9 | 13 | 9 | 12 | 14 | 18 | 17 | 13 | 15 | 14 | 12 |
| 20 | 21 | 31 | 25 | 31 | 31 | 36 | 36 | 31 | 31 | 31 | 27 |
| 37 | 37 | 50 | 45 | 60 | 55 | 64 | 62 | 56 | 49 | 55 | 49 |
| 65 | 62 | 79 | 72 | 93 | 98 | 101 | 95 | 90 | 78 | 86 | 90 |
| 82 | 75 | 95 | 93 | 119 | 136 | 126 | 126 | 126 | 108 | 109 | 102 |
| 124 | 121 | 160 | 144 | 196 | 216 | 193 | 216 | 183 | 198 | 186 | 145 |
| 55 | 53 | 66 | 60 | 75 | 75 | 80 | 73 | 66 | 70 | 59 | 54 |
| 49 | 45 | 55 | 52 | 68 | 71 | 70 | 63 | 60 | 55 | 42 | 41 |
| 61 | 59 | 75 | 59 | 79 | 79 | 82 | 77 | 70 | 72 | 64 | 69 |
| 43 | 43 | 59 | 39 | 59 | 64 | 54 | 52 | 45 | 58 | 40 | 45 |
| 63 | 56 | 65 | 73 | 78 | 95 | 84 | 88 | 79 | 68 | 78 | 64 |
| 40 | 36 | 37 | 45 | 58 | 60 | 54 | 58 | 58 | 48 | 47 | 50 |
| 349 | 486 | 558 | 372 | 595 | 755 | 1,080 | 594 | 450 | 432 | 229 | 632 |
| 660 | 702 | 882 | 666 | 972 | 1,082 | 1,296 | 1,116 | 837 | 807 | 558 | 744 |
| 54 | 51 | 61 | 57 | 68 | 74 | 72 | 70 | 65 | 62 | 61 | 58 |
| 47 | 43 | 56 | 49 | 65 | 68 | 63 | 58 | 54 | 51 | 47 | 53 |
| 10 | 10 | 10 | 9 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 |
| 20 | 15 | 20 | 15 | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 |
| 47 | 41 | 54 | 51 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 54 |
| 81 | 70 | 93 | 93 | 108 | 108 | 108 | 104 | 93 | 93 | 93 | 93 |
| 108 | 108 | 110 | 108 | 162 | 162 | 153 | 140 | 135 | 110 | 113 | 108 |
| 135 | 110 | 186 | 142 | 186 | 204 | 186 | 186 | 162 | 150 | 162 | 140 |
| 216 | 216 | 270 | 216 | 279 | 306 | 234 | 270 | 279 | 225 | 204 | 204 |
| 330 | 280 | 558 | 325 | 595 | 755 | 1,080 | 558 | 450 | 324 | 216 | 539 |
| 17.5 | 29.7 | 13.9 | 21.5 | 71.2 | 65.4 | 116.0 | 138.9 | 20.0 | 32.0 | 9.1 | 10.3 |
| 1.1 | 1.7 | 1.2 | 1.7 | 1.8 | 1.2 | 2.0 | 1.6 | 1.9 | 1.9 | 2.7 | 1.3 |

TABLE 10.14.--LETTUCE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 33.2 | 0 | 9.4 | 18.2 | 26.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 21.2 | 0 | 8.4 | 13.9 | 19.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 8.8 | 0 | .9 | 3.3 | 5.7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 3.1 | 0 | .1 | 1.0 | 1.6 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.6 | 0 | 1.2 | 1.3 | 1.4 |
| STANDARD DEVIATION.....DO..... | .9 | 0 | .5 | .7 | .7 |
| MAXIMUM.....DO..... | 8.0 | 0 | 4.0 | 5.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 19 | 0 | 8 | 9 | 11 |
| STANDARD DEVIATION.....GM..... | 22 | 0 | 10 | 9 | 10 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 3 | | 1 | 2 | 2 |
| 25TH.....GM..... | 5 | | 3 | 3 | 3 |
| 50TH.....GM..... | 11 | | 5 | 7 | 9 |
| 75TH.....GM..... | 23 | | 9 | 10 | 17 |
| 90TH.....GM..... | 43 | | 18 | 20 | 28 |
| 95TH.....GM..... | 60 | | 30 | 28 | 32 |
| 99TH.....GM..... | 110 | | 60 | 45 | 45 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 36 | 0 | 21 | 19 | 24 |
| STANDARD DEVIATION.....GM..... | 42 | 0 | 27 | 16 | 19 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 40 | 0 | 25 | 25 | 29 |
| STANDARD DEVIATION.....GM..... | 34 | 0 | 18 | 17 | 18 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 45 | 0 | 30 | 21 | 24 |
| STANDARD DEVIATION.....GM..... | 36 | 0 | 0 | 10 | 12 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 755 | 0 | 180 | 130 | 135 |
| MAXIMUM OVER 3 DAYS.....GM..... | 809 | 0 | 180 | 157 | 230 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 36 | 0 | 21 | 21 | 24 |
| STANDARD DEVIATION.....GM..... | 38 | 0 | 25 | 16 | 18 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 8 | | 4 | 5 | 5 |
| 25TH.....GM..... | 10 | | 10 | 10 | 10 |
| 50TH.....GM..... | 28 | | 14 | 14 | 20 |
| 75TH.....GM..... | 55 | | 28 | 28 | 28 |
| 90TH.....GM..... | 83 | | 41 | 40 | 55 |
| 95TH.....GM..... | 110 | | 55 | 55 | 55 |
| 99TH.....GM..... | 165 | | 180 | 72 | 67 |
| MAXIMUM.....GM..... | 755 | 0 | 180 | 110 | 135 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 225.2 | 0 | 0 | 3.4 | 7.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.1 | 0 | 0 | .8 | 1.1 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 29.4 | 29.8 | 32.7 | 32.8 | 37.7 | 36.0 | 38.1 | 40.1 | 34.4 | 32.7 | 20.9 | 29.3 |
| 21.3 | 20.6 | 21.4 | 20.6 | 23.7 | 23.8 | 22.8 | 24.6 | 20.4 | 19.6 | 12.0 | 16.9 |
| 6.6 | 7.1 | 8.5 | 9.9 | 9.9 | 10.2 | 10.8 | 11.5 | 8.4 | 8.4 | 5.7 | 7.7 |
| 1.6 | 2.1 | 2.7 | 2.3 | 4.1 | 2.0 | 4.5 | 4.0 | 5.6 | 4.7 | 3.2 | 4.7 |
| 1.4 | 1.5 | 1.6 | 1.6 | 1.6 | 1.5 | 1.7 | 1.6 | 1.7 | 1.7 | 1.8 | 1.7 |
| .7 | .8 | .9 | .8 | .9 | .8 | 1.0 | .9 | .9 | 1.0 | 1.0 | 1.0 |
| 5.0 | 5.0 | 6.0 | 5.0 | 6.0 | 6.0 | 8.0 | 6.0 | 5.0 | 6.0 | 6.0 | 6.0 |
| 14 | 15 | 18 | 17 | 19 | 20 | 21 | 21 | 25 | 20 | 19 | 22 |
| 16 | 16 | 19 | 19 | 21 | 26 | 23 | 24 | 27 | 23 | 14 | 25 |
| 2 | 2 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 2 | 3 |
| 5 | 5 | 7 | 5 | 7 | 5 | 7 | 7 | 7 | 6 | 9 | 7 |
| 9 | 9 | 10 | 10 | 13 | 13 | 14 | 13 | 16 | 13 | 17 | 17 |
| 18 | 18 | 23 | 20 | 26 | 25 | 28 | 28 | 30 | 28 | 27 | 28 |
| 31 | 33 | 43 | 37 | 45 | 45 | 48 | 47 | 48 | 45 | 37 | 45 |
| 40 | 41 | 55 | 55 | 62 | 63 | 64 | 68 | 90 | 57 | 43 | 57 |
| 89 | 67 | 90 | 90 | 95 | 136 | 110 | 110 | 135 | 150 | 55 | 180 |
| 30 | 29 | 36 | 31 | 39 | 42 | 38 | 38 | 46 | 35 | 39 | 48 |
| 29 | 27 | 36 | 32 | 40 | 58 | 34 | 42 | 53 | 34 | 33 | 75 |
| 35 | 36 | 38 | 36 | 40 | 42 | 43 | 43 | 43 | 42 | 35 | 41 |
| 30 | 33 | 27 | 27 | 32 | 38 | 36 | 35 | 33 | 39 | 20 | 27 |
| 38 | 29 | 43 | 46 | 39 | 58 | 49 | 49 | 53 | 43 | 32 | 33 |
| 27 | 17 | 30 | 38 | 35 | 45 | 35 | 39 | 40 | 32 | 12 | 27 |
| 280 | 280 | 275 | 270 | 290 | 755 | 360 | 360 | 450 | 270 | 180 | 539 |
| 550 | 550 | 380 | 549 | 559 | 755 | 590 | 809 | 450 | 539 | 180 | 539 |
| 31 | 30 | 35 | 33 | 36 | 40 | 38 | 38 | 44 | 36 | 32 | 38 |
| 29 | 29 | 33 | 32 | 36 | 50 | 36 | 38 | 42 | 36 | 20 | 49 |
| 7 | 7 | 10 | 7 | 10 | 7 | 10 | 10 | 10 | 10 | 7 | 8 |
| 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 14 | 10 | 18 | 14 |
| 24 | 20 | 20 | 20 | 20 | 28 | 28 | 28 | 28 | 28 | 28 | 28 |
| 41 | 40 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 55 |
| 55 | 55 | 67 | 65 | 83 | 96 | 83 | 83 | 110 | 67 | 55 | 65 |
| 83 | 83 | 110 | 90 | 110 | 120 | 110 | 110 | 135 | 110 | 55 | 110 |
| 135 | 135 | 135 | 135 | 165 | 220 | 136 | 165 | 138 | 165 | 90 | 165 |
| 270 | 275 | 275 | 270 | 270 | 755 | 270 | 280 | 450 | 270 | 110 | 539 |
| 5.5 | 13.5 | 5.0 | 9.7 | 23.8 | 27.0 | 48.5 | 40.0 | 8.7 | 15.4 | 9.1 | 8.2 |
| .6 | 1.4 | .7 | 1.3 | 1.0 | .9 | 1.5 | .9 | 1.3 | 1.6 | 4.6 | 1.6 |

TABLE 10.15.--TOSSED SALADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 26.2 | 0 | 9.1 | 15.4 | 21.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 17.3 | 0 | 7.4 | 11.6 | 15.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 6.8 | 0 | 1.3 | 3.3 | 5.0 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.1 | 0 | .3 | .5 | 1.2 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.5 | 0 | 1.3 | 1.3 | 1.4 |
| STANDARD DEVIATION.....DO..... | .8 | 0 | .7 | .6 | .7 |
| MAXIMUM.....DO..... | 6.0 | 0 | 5.0 | 4.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 50 | 0 | 16 | 22 | 30 |
| STANDARD DEVIATION.....GM..... | 39 | 0 | 15 | 15 | 23 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 11 | | 2 | 4 | 6 |
| 25TH.....GM..... | 27 | | 8 | 12 | 16 |
| 50TH.....GM..... | 36 | | 15 | 18 | 25 |
| 75TH.....GM..... | 62 | | 18 | 31 | 36 |
| 90TH.....GM..... | 94 | | 31 | 36 | 62 |
| 95TH.....GM..... | 124 | | 54 | 49 | 72 |
| 99TH.....GM..... | 201 | | 90 | 72 | 113 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 102 | 0 | 38 | 52 | 63 |
| STANDARD DEVIATION.....GM..... | 60 | 0 | 25 | 30 | 35 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 107 | 0 | 41 | 50 | 70 |
| STANDARD DEVIATION.....GM..... | 55 | 0 | 29 | 26 | 34 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 117 | 0 | 54 | 50 | 76 |
| STANDARD DEVIATION.....GM..... | 59 | 0 | 32 | 22 | 32 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,080 | 0 | 186 | 186 | 216 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,296 | 0 | 270 | 326 | 432 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 101 | 0 | 39 | 49 | 64 |
| STANDARD DEVIATION.....GM..... | 56 | 0 | 29 | 27 | 34 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 30 | | 6 | 13 | 16 |
| 25TH.....GM..... | 60 | | 15 | 28 | 41 |
| 50TH.....GM..... | 93 | | 33 | 47 | 54 |
| 75TH.....GM..... | 108 | | 54 | 54 | 93 |
| 90TH.....GM..... | 186 | | 70 | 93 | 108 |
| 95TH.....GM..... | 216 | | 93 | 108 | 108 |
| 99TH.....GM..... | 279 | | 108 | 108 | 162 |
| MAXIMUM.....GM..... | 1,080 | 0 | 186 | 147 | 216 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 339.0 | 0 | .7 | 1.0 | 11.9 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 2.3 | 0 | .6 | .3 | 2.1 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 23.0 | 24.9 | 23.3 | 23.5 | 25.0 | 29.5 | 33.0 | 33.2 | 23.1 | 24.8 | 15.8 | 16.7 |
| 16.8 | 17.7 | 15.4 | 16.8 | 17.4 | 20.0 | 20.4 | 20.5 | 15.1 | 16.2 | 8.7 | 9.6 |
| 5.3 | 6.4 | 6.8 | 5.4 | 5.9 | 7.0 | 9.6 | 9.3 | 5.9 | 6.7 | 4.6 | 4.6 |
| .9 | .8 | 1.2 | 1.3 | 1.8 | 2.5 | 3.0 | 3.4 | 2.1 | 1.9 | 2.5 | 2.4 |
| 1.4 | 1.4 | 1.5 | 1.4 | 1.4 | 1.5 | 1.5 | 1.6 | 1.5 | 1.5 | 1.6 | 1.6 |
| .6 | .6 | .7 | .7 | .7 | .8 | .8 | .8 | .8 | .8 | .8 | .8 |
| 5.0 | 5.0 | 5.0 | 4.0 | 5.0 | 6.0 | 6.0 | 6.0 | 5.0 | 5.0 | 3.0 | 4.0 |
| 38 | 36 | 48 | 43 | 57 | 54 | 57 | 56 | 52 | 49 | 54 | 50 |
| 27 | 26 | 34 | 30 | 41 | 42 | 41 | 42 | 44 | 37 | 37 | 32 |
| 9 | 9 | 16 | 11 | 16 | 13 | 16 | 16 | 13 | 16 | 11 | 16 |
| 18 | 18 | 31 | 23 | 31 | 31 | 31 | 31 | 25 | 31 | 31 | 31 |
| 31 | 31 | 36 | 36 | 47 | 36 | 47 | 44 | 36 | 36 | 43 | 43 |
| 50 | 43 | 62 | 54 | 72 | 67 | 72 | 72 | 62 | 62 | 71 | 67 |
| 72 | 67 | 87 | 78 | 108 | 108 | 109 | 103 | 95 | 93 | 98 | 93 |
| 83 | 83 | 121 | 101 | 124 | 144 | 134 | 134 | 148 | 108 | 117 | 111 |
| 144 | 134 | 180 | 165 | 216 | 216 | 196 | 220 | 279 | 201 | 186 | 124 |
| 85 | 79 | 105 | 97 | 123 | 107 | 118 | 108 | 97 | 100 | 94 | 89 |
| 51 | 47 | 48 | 51 | 67 | 61 | 70 | 56 | 51 | 46 | 39 | 37 |
| 89 | 85 | 102 | 95 | 121 | 122 | 114 | 111 | 99 | 107 | 99 | 112 |
| 46 | 45 | 65 | 43 | 58 | 63 | 49 | 52 | 41 | 67 | 32 | 34 |
| 84 | 76 | 99 | 100 | 136 | 121 | 120 | 123 | 152 | 106 | 113 | 85 |
| 40 | 33 | 42 | 49 | 54 | 60 | 62 | 61 | 61 | 41 | 44 | 35 |
| 330 | 486 | 558 | 372 | 558 | 558 | 1,080 | 594 | 366 | 432 | 216 | 366 |
| 660 | 702 | 882 | 666 | 960 | 915 | 1,296 | 1,116 | 837 | 804 | 558 | 744 |
| 83 | 78 | 98 | 92 | 120 | 109 | 112 | 106 | 101 | 97 | 100 | 94 |
| 48 | 42 | 57 | 46 | 61 | 62 | 60 | 52 | 54 | 47 | 43 | 38 |
| 27 | 23 | 33 | 33 | 47 | 30 | 38 | 41 | 33 | 33 | 30 | 47 |
| 49 | 47 | 55 | 55 | 83 | 65 | 75 | 70 | 65 | 65 | 75 | 65 |
| 81 | 70 | 93 | 93 | 108 | 93 | 98 | 93 | 93 | 93 | 93 | 93 |
| 108 | 98 | 108 | 108 | 162 | 140 | 140 | 113 | 108 | 108 | 108 | 108 |
| 130 | 108 | 186 | 162 | 216 | 186 | 186 | 186 | 186 | 150 | 186 | 140 |
| 186 | 162 | 216 | 186 | 216 | 216 | 216 | 216 | 206 | 186 | 186 | 186 |
| 216 | 216 | 324 | 216 | 324 | 324 | 279 | 279 | 279 | 270 | 216 | 225 |
| 330 | 279 | 558 | 325 | 558 | 558 | 1,080 | 558 | 324 | 324 | 216 | 270 |
| 12.0 | 16.3 | 8.9 | 10.8 | 47.4 | 36.4 | 64.8 | 99.0 | 11.3 | 16.5 | 0 | 2.1 |
| 1.8 | 2.2 | 1.9 | 2.2 | 3.4 | 1.6 | 2.6 | 2.7 | 2.9 | 2.5 | 0 | .8 |

TABLE 10.16.--RAW MATURE ONIONS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 8.5 | 0 | 1.4 | 3.1 | 3.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 7.1 | 0 | 1.1 | 2.6 | 3.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.2 | 0 | .3 | .6 | .7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 0 | 1.2 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .5 | 0 | .4 | .5 | .4 |
| MAXIMUM.....DO..... | 5.0 | 0 | 2.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 13 | 0 | 9 | 8 | 8 |
| STANDARD DEVIATION.....GM..... | 16 | 0 | 8 | 11 | 7 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 1 | | | (2) | (2) |
| 25TH.....GM..... | 6 | | | 3 | 3 |
| 50TH.....GM..... | 7 | | | 6 | 6 |
| 75TH.....GM..... | 15 | | | 6 | 9 |
| 90TH.....GM..... | 24 | | | 12 | 18 |
| 95TH.....GM..... | 37 | | | 28 | 23 |
| 99TH.....GM..... | 73 | | | 73 | 37 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 31 | 0 | 20 | 20 | 22 |
| STANDARD DEVIATION.....GM..... | 33 | 0 | 17 | 35 | 19 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 34 | 0 | 27 | 16 | 16 |
| STANDARD DEVIATION.....GM..... | 44 | 0 | 13 | 15 | 16 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 31 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 22 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 660 | 0 | 70 | 220 | 110 |
| MAXIMUM OVER 3 DAYS.....GM..... | 990 | 0 | 90 | 220 | 110 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 31 | 0 | 22 | 19 | 20 |
| STANDARD DEVIATION.....GM..... | 33 | 0 | 18 | 30 | 19 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 3 | | | 1 | 1 |
| 25TH.....GM..... | 17 | | | 4 | 9 |
| 50TH.....GM..... | 18 | | | 17 | 18 |
| 75TH.....GM..... | 36 | | | 18 | 20 |
| 90TH.....GM..... | 57 | | | 35 | 36 |
| 95TH.....GM..... | 72 | | | 36 | 70 |
| 99TH.....GM..... | 180 | | | 220 | 110 |
| MAXIMUM.....GM..... | 350 | 0 | 70 | 220 | 110 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 210.7 | 0 | 0 | 7.4 | 7.2 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 5.3 | 0 | 0 | 11.2 | 8.5 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 6.0 | 5.3 | 9.9 | 7.9 | 12.4 | 9.8 | 11.7 | 10.4 | 7.8 | 6.9 | 5.6 | 3.9 |
| 5.5 | 4.4 | 7.9 | 6.7 | 10.4 | 8.5 | 9.0 | 8.7 | 6.5 | 5.7 | 4.9 | 2.6 |
| .4 | .8 | 1.5 | 1.0 | 1.6 | 1.2 | 2.2 | 1.3 | 1.1 | 1.0 | .7 | 1.2 |
| (2) | .1 | .5 | .1 | .3 | .1 | .6 | .3 | .3 | .2 | 0 | 0 |
| 1.1 | 1.2 | 1.3 | 1.2 | 1.2 | 1.2 | 1.4 | 1.2 | 1.3 | 1.3 | 1.1 | 1.4 |
| .3 | .5 | .7 | .5 | .5 | .4 | .6 | .6 | .6 | .5 | .3 | .6 |
| 3.0 | 3.0 | 5.0 | 3.0 | 4.0 | 4.0 | 4.0 | 4.0 | 3.0 | 3.0 | 2.0 | 3.0 |
| 10 | 10 | 13 | 10 | 13 | 10 | 16 | 11 | 22 | 16 | 19 | 14 |
| 9 | 11 | 14 | 10 | 12 | 11 | 18 | 13 | 45 | 32 | 16 | 14 |
| 1 | 1 | 1 | 1 | 1 | 1 | 2 | 1 | 1 | 1 | | |
| 6 | 6 | 6 | 3 | 6 | 5 | 6 | 6 | 6 | 6 | | |
| 7 | 6 | 7 | 6 | 10 | 6 | 12 | 6 | 12 | 12 | | |
| 12 | 12 | 14 | 12 | 18 | 12 | 18 | 12 | 23 | 18 | | |
| 18 | 21 | 24 | 23 | 28 | 18 | 37 | 23 | 31 | 24 | | |
| 24 | 37 | 42 | 37 | 37 | 25 | 47 | 37 | 46 | 35 | | |
| 47 | 55 | 73 | 53 | 63 | 63 | 100 | 73 | 330 | 257 | | |
| 26 | 26 | 32 | 27 | 35 | 27 | 38 | 28 | 41 | 31 | 53 | 29 |
| 20 | 30 | 37 | 26 | 31 | 29 | 37 | 34 | 38 | 31 | 48 | 31 |
| 35 | 30 | 29 | 25 | 32 | 27 | 37 | 30 | 99 | 71 | 47 | 36 |
| 24 | 21 | 26 | 22 | 28 | 22 | 34 | 21 | 160 | 105 | 10 | 24 |
| 18 | 13 | 27 | 22 | 29 | 19 | 43 | 25 | 38 | 26 | 0 | 0 |
| 0 | 6 | 16 | 5 | 10 | 5 | 29 | 18 | 7 | 11 | 0 | 0 |
| 110 | 220 | 274 | 140 | 220 | 220 | 360 | 350 | 660 | 440 | 210 | 140 |
| 140 | 220 | 274 | 158 | 280 | 220 | 369 | 350 | 990 | 770 | 210 | 158 |
| 27 | 26 | 29 | 25 | 33 | 26 | 36 | 27 | 50 | 39 | 52 | 31 |
| 20 | 27 | 29 | 26 | 29 | 27 | 35 | 31 | 68 | 53 | 43 | 27 |
| 3 | 3 | 3 | 3 | 5 | 3 | 3 | 3 | 3 | 5 | | |
| 15 | 10 | 16 | 10 | 18 | 10 | 18 | 14 | 18 | 18 | | |
| 18 | 18 | 18 | 18 | 20 | 18 | 25 | 18 | 35 | 18 | | |
| 35 | 35 | 36 | 33 | 36 | 35 | 38 | 35 | 55 | 36 | | |
| 54 | 54 | 55 | 54 | 70 | 55 | 70 | 55 | 70 | 55 | | |
| 70 | 70 | 70 | 70 | 110 | 70 | 110 | 70 | 220 | 220 | | |
| 110 | 110 | 220 | 110 | 140 | 150 | 180 | 175 | 330 | 330 | | |
| 110 | 220 | 220 | 140 | 220 | 220 | 280 | 350 | 330 | 330 | 210 | 110 |
| 9.3 | 3.8 | 20.5 | 7.8 | 23.7 | 36.7 | 42.0 | 37.6 | 3.5 | 9.7 | 0 | 1.4 |
| 6.7 | 2.8 | 11.2 | 5.6 | 4.0 | 6.0 | 5.4 | 4.1 | 3.1 | 6.4 | 0 | 2.7 |

TABLE 10.17.--RAW TOMATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 27.8 | .3 | 10.6 | 15.7 | 18.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 17.4 | .3 | 8.5 | 11.6 | 14.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 7.4 | 0 | 1.8 | 3.3 | 3.3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | 2.9 | 0 | .3 | .8 | .8 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.6 | 1.0 | 1.3 | 1.4 | 1.3 |
| STANDARD DEVIATION.....DO..... | 1.0 | .0 | .6 | .7 | .7 |
| MAXIMUM.....DO..... | 8.0 | 1.0 | 4.0 | 7.0 | 6.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 44 | 7 | 20 | 24 | 24 |
| STANDARD DEVIATION.....GM..... | 42 | 2 | 17 | 23 | 20 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 10 | | 4 | 5 | 7 |
| 25TH.....GM..... | 20 | | 10 | 10 | 10 |
| 50TH.....GM..... | 30 | | 15 | 20 | 20 |
| 75TH.....GM..... | 60 | | 29 | 30 | 30 |
| 90TH.....GM..... | 91 | | 46 | 51 | 45 |
| 95TH.....GM..... | 123 | | 60 | 71 | 61 |
| 99TH.....GM..... | 205 | | 80 | 111 | 120 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 78 | 21 | 43 | 54 | 53 |
| STANDARD DEVIATION.....GM..... | 59 | 7 | 32 | 55 | 34 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 91 | 0 | 61 | 54 | 66 |
| STANDARD DEVIATION.....GM..... | 56 | 0 | 23 | 32 | 43 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 110 | 0 | 57 | 64 | 59 |
| STANDARD DEVIATION.....GM..... | 62 | 0 | 19 | 32 | 18 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 907 | 30 | 182 | 615 | 308 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,638 | 30 | 240 | 615 | 490 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 81 | 21 | 46 | 52 | 55 |
| STANDARD DEVIATION.....GM..... | 55 | 7 | 32 | 44 | 33 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 30 | | 11 | 15 | 15 |
| 25TH.....GM..... | 45 | | 30 | 30 | 30 |
| 50TH.....GM..... | 62 | | 31 | 31 | 56 |
| 75TH.....GM..... | 113 | | 60 | 62 | 62 |
| 90TH.....GM..... | 123 | | 91 | 91 | 91 |
| 95TH.....GM..... | 182 | | 120 | 123 | 123 |
| 99TH.....GM..... | 246 | | 182 | 182 | 180 |
| MAXIMUM.....GM..... | 728 | 30 | 182 | 615 | 246 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 68.6 | 0 | .8 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .4 | 0 | .6 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 20.1 | 21.0 | 24.4 | 24.3 | 29.6 | 29.5 | 32.4 | 35.9 | 30.9 | 35.8 | 27.9 | 25.0 |
| 14.9 | 15.6 | 17.4 | 16.4 | 20.3 | 19.3 | 18.2 | 21.2 | 15.1 | 18.9 | 14.4 | 12.6 |
| 4.2 | 4.1 | 5.5 | 6.3 | 7.0 | 8.1 | 9.7 | 10.5 | 9.5 | 10.3 | 7.2 | 8.3 |
| .9 | 1.3 | 1.5 | 1.6 | 2.3 | 2.2 | 4.4 | 4.2 | 6.3 | 6.6 | 6.2 | 4.1 |
| 1.4 | 1.4 | 1.5 | 1.5 | 1.5 | 1.5 | 1.8 | 1.7 | 1.9 | 1.9 | 2.0 | 1.9 |
| .8 | .8 | .8 | .8 | .8 | .8 | 1.1 | 1.0 | 1.1 | 1.1 | 1.3 | 1.1 |
| 5.0 | 6.0 | 5.0 | 5.0 | 7.0 | 7.0 | 8.0 | 6.0 | 6.0 | 6.0 | 8.0 | 6.0 |
| 35 | 33 | 37 | 33 | 41 | 39 | 55 | 48 | 60 | 54 | 60 | 56 |
| 36 | 30 | 37 | 29 | 33 | 35 | 53 | 42 | 56 | 43 | 56 | 44 |
| 8 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 |
| 15 | 15 | 15 | 15 | 20 | 20 | 21 | 20 | 30 | 21 | 21 | 21 |
| 21 | 21 | 21 | 21 | 30 | 30 | 40 | 36 | 41 | 41 | 40 | 41 |
| 41 | 41 | 41 | 41 | 58 | 50 | 66 | 61 | 81 | 71 | 81 | 81 |
| 71 | 62 | 80 | 62 | 82 | 81 | 121 | 99 | 123 | 122 | 121 | 112 |
| 100 | 82 | 120 | 101 | 102 | 102 | 164 | 132 | 143 | 140 | 143 | 143 |
| 212 | 133 | 205 | 150 | 171 | 164 | 262 | 205 | 204 | 205 | 273 | 241 |
| 73 | 73 | 73 | 66 | 83 | 75 | 87 | 83 | 92 | 85 | 76 | 89 |
| 59 | 54 | 61 | 47 | 67 | 58 | 57 | 60 | 55 | 58 | 36 | 56 |
| 84 | 75 | 83 | 72 | 92 | 85 | 104 | 95 | 100 | 92 | 104 | 96 |
| 59 | 57 | 54 | 37 | 44 | 60 | 67 | 56 | 51 | 40 | 45 | 38 |
| 110 | 86 | 107 | 94 | 93 | 95 | 129 | 110 | 121 | 116 | 129 | 118 |
| 75 | 38 | 64 | 42 | 42 | 42 | 78 | 58 | 82 | 45 | 70 | 59 |
| 735 | 519 | 615 | 364 | 675 | 907 | 735 | 677 | 546 | 615 | 455 | 362 |
| 1,014 | 822 | 733 | 546 | 675 | 1,361 | 1,599 | 1,107 | 1,638 | 738 | 818 | 800 |
| 74 | 71 | 75 | 66 | 81 | 75 | 93 | 84 | 93 | 86 | 92 | 89 |
| 58 | 49 | 56 | 44 | 54 | 49 | 63 | 57 | 65 | 44 | 43 | 48 |
| 23 | 30 | 30 | 28 | 30 | 28 | 30 | 30 | 30 | 30 | 30 | 30 |
| 30 | 31 | 30 | 30 | 45 | 34 | 60 | 56 | 60 | 60 | 60 | 60 |
| 60 | 60 | 60 | 60 | 62 | 62 | 90 | 62 | 90 | 90 | 90 | 90 |
| 90 | 90 | 90 | 90 | 120 | 91 | 123 | 120 | 123 | 123 | 120 | 123 |
| 123 | 123 | 123 | 123 | 123 | 123 | 180 | 123 | 180 | 123 | 180 | 135 |
| 180 | 180 | 180 | 180 | 182 | 180 | 182 | 182 | 182 | 153 | 182 | 182 |
| 364 | 246 | 270 | 182 | 246 | 246 | 360 | 273 | 270 | 182 | 182 | 246 |
| 675 | 369 | 492 | 364 | 675 | 546 | 728 | 675 | 546 | 615 | 273 | 270 |
| 4.0 | 4.1 | 0 | 2.6 | 9.3 | 12.5 | 13.8 | 12.7 | 1.8 | 5.0 | 0 | 2.0 |
| .7 | .6 | 0 | .5 | .5 | .5 | .5 | .3 | .3 | .4 | 0 | .4 |

TABLE 10.18.--COOKED BROCCOLI
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 6.2 | 1.0 | 5.7 | 3.8 | 5.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 5.7 | 1.0 | 5.3 | 3.6 | 5.3 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .5 | 0 | .5 | .2 | .3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.1 | 1.0 | 1.1 | 1.1 | 1.0 |
| STANDARD DEVIATION.....DO..... | .3 | .0 | .3 | .2 | .2 |
| MAXIMUM.....DO..... | 3.0 | 1.0 | 2.0 | 2.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 41 | 14 | 20 | 23 | 29 |
| STANDARD DEVIATION.....GM..... | 27 | 9 | 13 | 16 | 18 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 12 | | 4 | 3 | 7 |
| 25TH.....GM..... | 26 | | 13 | 14 | 15 |
| 50TH.....GM..... | 31 | | 15 | 22 | 26 |
| 75TH.....GM..... | 57 | | 28 | 29 | 31 |
| 90TH.....GM..... | 62 | | 32 | 31 | 59 |
| 95TH.....GM..... | 99 | | 50 | 58 | 62 |
| 99TH.....GM..... | 137 | | 62 | 93 | 85 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 114 | 42 | 56 | 65 | 82 |
| STANDARD DEVIATION.....GM..... | 69 | 27 | 36 | 43 | 52 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 108 | 0 | 55 | 61 | 88 |
| STANDARD DEVIATION.....GM..... | 63 | 0 | 27 | 44 | 2 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 100 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 57 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 680 | 93 | 185 | 278 | 255 |
| MAXIMUM OVER 3 DAYS.....GM..... | 930 | 93 | 185 | 278 | 255 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 112 | 42 | 55 | 65 | 83 |
| STANDARD DEVIATION.....GM..... | 68 | 27 | 33 | 43 | 50 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 30 | | 11 | 10 | 22 |
| 25TH.....GM..... | 78 | | 39 | 43 | 47 |
| 50TH.....GM..... | 90 | | 46 | 57 | 80 |
| 75TH.....GM..... | 155 | | 85 | 88 | 90 |
| 90TH.....GM..... | 185 | | 90 | 93 | 175 |
| 95TH.....GM..... | 190 | | 93 | 175 | 185 |
| 99TH.....GM..... | 350 | | 185 | 278 | 255 |
| MAXIMUM.....GM..... | 680 | 93 | 185 | 278 | 255 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 9.0 | 0 | 0 | 0 | 1.8 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .3 | 0 | 0 | 0 | 1.7 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 4.6 | 5.1 | 4.3 | 4.1 | 6.4 | 6.4 | 7.1 | 8.1 | 7.1 | 6.9 | 4.0 | 7.1 |
| 4.4 | 4.8 | 3.9 | 3.8 | 6.0 | 5.7 | 6.6 | 7.3 | 6.4 | 5.6 | 3.4 | 6.0 |
| .2 | .3 | .4 | .3 | .4 | .7 | .5 | .7 | .7 | 1.3 | .7 | 1.1 |
| 0 | 0 | 0 | 0 | 0 | (2) | (2) | (2) | 0 | 0 | 0 | 0 |
| 1.0 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.2 | 1.2 |
| .2 | .2 | .3 | .3 | .3 | .3 | .3 | .3 | .3 | .5 | .4 | .4 |
| 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 3.0 | 3.0 | 3.0 | 2.0 | 3.0 | 2.0 | 2.0 |
| 33 | 31 | 36 | 38 | 54 | 42 | 48 | 42 | 41 | 41 | 38 | 40 |
| 25 | 21 | 20 | 20 | 34 | 21 | 34 | 26 | 24 | 24 | 22 | 27 |
| 8 | 7 | 15 | 13 | 14 | 15 | 14 | 13 | 13 | 14 | | 12 |
| 16 | 16 | 22 | 27 | 29 | 28 | 29 | 28 | 27 | 27 | | 26 |
| 28 | 28 | 30 | 31 | 52 | 35 | 43 | 31 | 31 | 31 | | 29 |
| 44 | 32 | 52 | 53 | 62 | 58 | 58 | 57 | 52 | 57 | | 57 |
| 58 | 57 | 58 | 62 | 110 | 62 | 62 | 63 | 73 | 62 | | 62 |
| 62 | 62 | 62 | 77 | 113 | 74 | 110 | 103 | 103 | 85 | | 71 |
| 155 | 113 | 117 | 117 | 146 | 123 | 227 | 123 | 117 | 129 | | 183 |
| 97 | 87 | 101 | 107 | 151 | 117 | 132 | 118 | 114 | 104 | 96 | 103 |
| 74 | 55 | 46 | 57 | 96 | 56 | 76 | 63 | 64 | 52 | 46 | 51 |
| 76 | 97 | 94 | 96 | 155 | 93 | 152 | 102 | 109 | 103 | 100 | 104 |
| 32 | 49 | 55 | 21 | 40 | 44 | 95 | 62 | 35 | 46 | 41 | 68 |
| 0 | 0 | 0 | 0 | 0 | 66 | 55 | 133 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 7 | 57 | 0 | 0 | 0 | 0 |
| 465 | 370 | 185 | 350 | 620 | 340 | 680 | 465 | 320 | 255 | 185 | 285 |
| 465 | 370 | 350 | 350 | 620 | 475 | 930 | 620 | 350 | 388 | 340 | 548 |
| 96 | 88 | 100 | 106 | 150 | 112 | 134 | 115 | 110 | 100 | 97 | 98 |
| 72 | 55 | 48 | 55 | 89 | 56 | 81 | 64 | 51 | 48 | 45 | 51 |
| 19 | 21 | 30 | 39 | 44 | 30 | 44 | 35 | 39 | 39 | | 39 |
| 47 | 51 | 62 | 78 | 88 | 78 | 85 | 78 | 85 | 78 | | 70 |
| 85 | 85 | 88 | 93 | 140 | 93 | 116 | 93 | 93 | 85 | | 85 |
| 127 | 93 | 155 | 131 | 175 | 155 | 175 | 160 | 155 | 131 | | 106 |
| 175 | 170 | 175 | 175 | 278 | 185 | 185 | 185 | 170 | 175 | | 175 |
| 185 | 175 | 175 | 185 | 340 | 190 | 310 | 190 | 185 | 185 | | 185 |
| 465 | 310 | 185 | 350 | 425 | 320 | 465 | 350 | 310 | 255 | | 285 |
| 465 | 370 | 185 | 350 | 620 | 340 | 680 | 465 | 310 | 255 | 185 | 285 |
| 2.2 | 1.0 | 0 | 0 | 0 | 0 | 0 | 4.0 | 0 | 0 | 0 | 0 |
| 2.2 | .8 | 0 | 0 | 0 | 0 | 0 | .6 | 0 | 0 | 0 | 0 |

TABLE 10.19.--COOKED CABBAGE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1: | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.7 | .4 | 3.2 | 3.3 | 3.8 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.2 | .3 | 2.7 | 3.0 | 3.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .5 | .2 | .5 | .3 | .3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 1.4 | 1.2 | 1.2 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .5 | .4 | .5 | .4 |
| MAXIMUM.....DO..... | 4.0 | 2.0 | 3.0 | 4.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 50 | 35 | 23 | 31 | 35 |
| STANDARD DEVIATION.....GM..... | 39 | 32 | 19 | 21 | 22 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 13 | | | 5 | 6 |
| 25TH.....GM..... | 25 | | | 13 | 19 |
| 50TH.....GM..... | 50 | | | 25 | 25 |
| 75TH.....GM..... | 50 | | | 50 | 50 |
| 90TH.....GM..... | 100 | | | 50 | 50 |
| 95TH.....GM..... | 102 | | | 75 | 75 |
| 99TH.....GM..... | 200 | | | 100 | 100 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 133 | 35 | 61 | 91 | 95 |
| STANDARD DEVIATION.....GM..... | 84 | 32 | 52 | 62 | 59 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 137 | 112 | 53 | 52 | 96 |
| STANDARD DEVIATION.....GM..... | 83 | 0 | 36 | 33 | 38 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 206 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 139 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 750 | 150 | 218 | 300 | 300 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,220 | 225 | 228 | 300 | 300 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 128 | 77 | 57 | 77 | 92 |
| STANDARD DEVIATION.....GM..... | 83 | 52 | 48 | 51 | 54 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 28 | | | 9 | 18 |
| 25TH.....GM..... | 75 | | | 38 | 38 |
| 50TH.....GM..... | 145 | | | 75 | 75 |
| 75TH.....GM..... | 150 | | | 93 | 150 |
| 90TH.....GM..... | 225 | | | 150 | 150 |
| 95TH.....GM..... | 300 | | | 150 | 150 |
| 99TH.....GM..... | 450 | | | 150 | 225 |
| MAXIMUM.....GM..... | 610 | 150 | 218 | 300 | 300 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 32.4 | 0 | .8 | 1.4 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.6 | 0 | 2.0 | 2.0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.9 | 4.5 | 4.5 | 4.3 | 3.1 | 3.9 | 5.3 | 6.0 | 5.9 | 6.7 | 10.1 | 6.5 |
| 3.8 | 4.3 | 4.3 | 4.0 | 2.7 | 3.4 | 4.8 | 5.6 | 5.0 | 5.2 | 9.0 | 4.7 |
| .1 | .2 | .2 | .3 | .4 | .4 | .5 | .4 | .6 | 1.4 | 1.1 | 1.9 |
| 0 | 0 | 0 | 0 | 0 | (2) | (2) | .1 | .3 | .1 | 0 | 0 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.2 | 1.2 | 1.1 | 1.3 | 1.4 | 1.2 | 1.3 |
| .3 | .3 | .3 | .3 | .4 | .4 | .4 | .4 | .5 | .7 | .6 | .5 |
| 3.0 | 3.0 | 2.0 | 2.0 | 3.0 | 4.0 | 3.0 | 4.0 | 3.0 | 4.0 | 4.0 | 2.0 |
| 42 | 43 | 49 | 45 | 58 | 48 | 56 | 50 | 50 | 65 | 58 | 57 |
| 28 | 36 | 22 | 30 | 39 | 33 | 32 | 48 | 37 | 52 | 36 | 32 |
| 7 | 9 | 13 | 13 | 13 | 13 | 13 | 9 | 19 | 19 | 25 | 24 |
| 25 | 25 | 38 | 25 | 30 | 25 | 36 | 25 | 25 | 38 | 25 | 25 |
| 38 | 25 | 50 | 48 | 50 | 50 | 50 | 48 | 50 | 50 | 50 | 50 |
| 50 | 50 | 50 | 50 | 75 | 50 | 75 | 50 | 50 | 100 | 75 | 75 |
| 100 | 75 | 75 | 75 | 100 | 100 | 100 | 100 | 100 | 102 | 100 | 100 |
| 100 | 141 | 100 | 100 | 145 | 102 | 117 | 102 | 100 | 125 | 138 | 102 |
| 150 | 150 | 102 | 150 | 200 | 140 | 150 | 250 | 250 | 407 | 200 | 150 |
| 124 | 119 | 146 | 126 | 153 | 131 | 157 | 129 | 126 | 153 | 148 | 133 |
| 84 | 88 | 63 | 85 | 86 | 77 | 88 | 92 | 62 | 81 | 70 | 70 |
| 110 | 176 | 73 | 123 | 168 | 115 | 137 | 151 | 185 | 153 | 198 | 134 |
| 37 | 115 | 52 | 41 | 84 | 60 | 55 | 82 | 99 | 119 | 59 | 42 |
| 0 | 0 | 0 | 0 | 0 | 237 | 133 | 312 | 43 | 200 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 140 | 0 | 33 | 0 | 0 |
| 450 | 450 | 305 | 450 | 435 | 375 | 600 | 750 | 450 | 610 | 413 | 305 |
| 450 | 750 | 305 | 450 | 600 | 712 | 600 | 1,220 | 750 | 1,220 | 600 | 450 |
| 117 | 121 | 129 | 119 | 150 | 125 | 146 | 132 | 118 | 142 | 144 | 129 |
| 79 | 91 | 65 | 81 | 84 | 75 | 82 | 94 | 73 | 94 | 55 | 68 |
| 22 | 26 | 38 | 38 | 38 | 28 | 38 | 28 | 36 | 38 | 75 | 38 |
| 75 | 75 | 75 | 75 | 75 | 75 | 75 | 75 | 75 | 75 | 75 | 75 |
| 75 | 75 | 150 | 75 | 150 | 145 | 150 | 132 | 113 | 150 | 150 | 150 |
| 150 | 150 | 150 | 150 | 150 | 150 | 150 | 150 | 150 | 150 | 150 | 150 |
| 150 | 225 | 200 | 218 | 300 | 225 | 300 | 300 | 150 | 300 | 225 | 150 |
| 300 | 300 | 300 | 300 | 300 | 300 | 300 | 305 | 300 | 300 | 225 | 300 |
| 450 | 450 | 305 | 450 | 435 | 305 | 450 | 450 | 450 | 610 | 300 | 305 |
| 450 | 450 | 305 | 450 | 435 | 305 | 600 | 610 | 450 | 610 | 300 | 305 |
| 2.0 | 1.9 | 0 | 2.1 | 2.3 | 4.1 | 9.7 | 2.8 | 0 | 1.1 | 0 | 4.2 |
| 2.3 | 1.9 | 0 | 2.9 | 1.7 | 1.7 | 3.2 | .6 | 0 | .7 | 0 | 4.8 |

TABLE 10.20.--COOKED CARROTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 9.8 | 21.7 | 11.7 | 8.0 | 8.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 8.6 | 16.4 | 10.2 | 7.5 | 7.9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.0 | 4.1 | 1.4 | .5 | .7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | 1.2 | .2 | .1 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.2 | 1.4 | 1.2 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .7 | .4 | .3 | .4 |
| MAXIMUM.....DO..... | 6.0 | 4.0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 30 | 34 | 22 | 18 | 22 |
| STANDARD DEVIATION.....GM..... | 21 | 23 | 19 | 12 | 15 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 6 | 8 | 5 | 3 | 6 |
| 25TH.....GM..... | 15 | 19 | 12 | 13 | 13 |
| 50TH.....GM..... | 25 | 33 | 13 | 13 | 20 |
| 75TH.....GM..... | 39 | 43 | 26 | 25 | 26 |
| 90TH.....GM..... | 52 | 57 | 50 | 32 | 39 |
| 95TH.....GM..... | 71 | 85 | 68 | 46 | 50 |
| 99TH.....GM..... | 103 | 132 | 85 | 63 | 75 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 81 | 88 | 56 | 49 | 58 |
| STANDARD DEVIATION.....GM..... | 53 | 61 | 45 | 31 | 33 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 80 | 67 | 64 | 54 | 75 |
| STANDARD DEVIATION.....GM..... | 41 | 36 | 41 | 24 | 35 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 73 | 57 | 40 | 52 | 45 |
| STANDARD DEVIATION.....GM..... | 34 | 31 | 11 | 9 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 736 | 425 | 255 | 155 | 192 |
| MAXIMUM OVER 3 DAYS.....GM..... | 736 | 425 | 300 | 233 | 300 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 79 | 71 | 54 | 49 | 59 |
| STANDARD DEVIATION.....GM..... | 50 | 41 | 38 | 31 | 33 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 19 | 14 | 10 | 10 | 18 |
| 25TH.....GM..... | 46 | 46 | 38 | 38 | 38 |
| 50TH.....GM..... | 75 | 57 | 39 | 39 | 50 |
| 75TH.....GM..... | 92 | 98 | 71 | 59 | 75 |
| 90TH.....GM..... | 150 | 128 | 113 | 78 | 92 |
| 95TH.....GM..... | 155 | 128 | 150 | 113 | 144 |
| 99TH.....GM..... | 276 | 213 | 192 | 155 | 155 |
| MAXIMUM.....GM..... | 736 | 213 | 213 | 155 | 192 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 25.2 | 0 | 0 | 1.4 | 3.8 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | 0 | 0 | .9 | 2.2 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 8.5 | 8.8 | 8.5 | 7.0 | 7.9 | 7.9 | 10.8 | 10.7 | 11.6 | 12.7 | 12.9 | 13.5 |
| 7.8 | 8.0 | 7.9 | 6.5 | 7.1 | 7.4 | 9.5 | 9.7 | 9.8 | 9.5 | 11.4 | 11.0 |
| .7 | .8 | .6 | .5 | .6 | .5 | 1.3 | .9 | 1.5 | 2.6 | 1.2 | 1.5 |
| .1 | .1 | 0 | 0 | .1 | .1 | .1 | .1 | .4 | .6 | .2 | 1.1 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.3 | 1.2 | 1.3 |
| .5 | .5 | .3 | .2 | .4 | .3 | .4 | .3 | .5 | .6 | .5 | .6 |
| 6.0 | 5.0 | 2.0 | 2.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| 30 | 28 | 31 | 25 | 35 | 28 | 36 | 28 | 36 | 38 | 31 | 37 |
| 22 | 20 | 17 | 16 | 24 | 17 | 27 | 18 | 23 | 24 | 16 | 22 |
| 3 | 8 | 8 | 6 | 11 | 5 | 13 | 6 | 13 | 8 | 13 | 13 |
| 15 | 13 | 16 | 13 | 17 | 15 | 25 | 15 | 19 | 25 | 17 | 25 |
| 25 | 25 | 26 | 25 | 26 | 25 | 26 | 25 | 26 | 31 | 26 | 31 |
| 32 | 32 | 50 | 26 | 50 | 33 | 50 | 32 | 50 | 50 | 46 | 50 |
| 52 | 52 | 52 | 50 | 57 | 50 | 52 | 50 | 71 | 64 | 50 | 75 |
| 75 | 76 | 52 | 52 | 78 | 52 | 92 | 52 | 96 | 88 | 51 | 77 |
| 125 | 103 | 100 | 100 | 125 | 91 | 153 | 103 | 101 | 102 | 90 | 128 |
| 79 | 74 | 87 | 74 | 95 | 77 | 97 | 78 | 89 | 88 | 85 | 96 |
| 53 | 46 | 50 | 48 | 61 | 46 | 69 | 46 | 54 | 48 | 40 | 43 |
| 88 | 85 | 74 | 47 | 94 | 83 | 91 | 67 | 100 | 87 | 81 | 82 |
| 34 | 45 | 19 | 16 | 51 | 24 | 56 | 33 | 28 | 31 | 27 | 37 |
| 125 | 88 | 0 | 0 | 88 | 52 | 73 | 106 | 73 | 83 | 46 | 67 |
| 0 | 17 | 0 | 0 | 30 | 5 | 18 | 38 | 21 | 30 | 0 | 37 |
| 375 | 300 | 300 | 300 | 400 | 300 | 736 | 310 | 300 | 300 | 155 | 225 |
| 375 | 388 | 300 | 300 | 450 | 300 | 736 | 464 | 303 | 414 | 271 | 383 |
| 79 | 75 | 86 | 71 | 94 | 75 | 94 | 76 | 88 | 84 | 78 | 86 |
| 48 | 46 | 48 | 46 | 60 | 43 | 65 | 44 | 49 | 41 | 36 | 44 |
| 10 | 23 | 23 | 14 | 29 | 18 | 28 | 18 | 38 | 23 | 38 | 18 |
| 39 | 39 | 48 | 39 | 48 | 46 | 50 | 46 | 57 | 50 | 48 | 66 |
| 75 | 75 | 75 | 75 | 78 | 75 | 75 | 75 | 75 | 75 | 75 | 78 |
| 92 | 78 | 113 | 78 | 136 | 78 | 138 | 92 | 113 | 113 | 93 | 113 |
| 150 | 150 | 155 | 150 | 155 | 150 | 155 | 150 | 150 | 150 | 150 | 150 |
| 155 | 155 | 155 | 150 | 192 | 150 | 184 | 155 | 155 | 150 | 150 | 150 |
| 225 | 233 | 300 | 300 | 310 | 233 | 310 | 276 | 288 | 162 | 155 | 225 |
| 300 | 300 | 300 | 300 | 400 | 300 | 736 | 310 | 300 | 225 | 155 | 225 |
| 1.0 | 1.2 | 3.1 | 0 | 0 | 0 | 5.3 | 4.0 | 0 | 2.9 | 0 | 2.5 |
| .5 | .6 | 2.4 | 0 | 0 | 0 | .9 | .5 | 0 | 1.0 | 0 | 1.4 |

TABLE 10.21.--COOKED WHOLE KERNEL CORN
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 23.9 | 3.2 | 25.8 | 30.1 | 34.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 20.4 | 2.2 | 22.2 | 25.4 | 29.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.2 | 1.0 | 3.3 | 4.5 | 4.7 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | .3 | .3 | .7 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 1.4 | 1.2 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .5 | .6 | .5 | .4 | .5 |
| MAXIMUM.....DO..... | 6.0 | 3.0 | 5.0 | 4.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 38 | 10 | 22 | 27 | 32 |
| STANDARD DEVIATION.....GM..... | 27 | 6 | 16 | 19 | 22 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 7 | | 4 | 7 | 7 |
| 25TH.....GM..... | 26 | | 13 | 14 | 18 |
| 50TH.....GM..... | 28 | | 19 | 26 | 28 |
| 75TH.....GM..... | 51 | | 28 | 28 | 41 |
| 90TH.....GM..... | 57 | | 42 | 55 | 57 |
| 95TH.....GM..... | 85 | | 55 | 57 | 62 |
| 99TH.....GM..... | 133 | | 85 | 110 | 112 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 97 | 28 | 58 | 72 | 81 |
| STANDARD DEVIATION.....GM..... | 59 | 21 | 41 | 48 | 44 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 96 | 16 | 56 | 62 | 75 |
| STANDARD DEVIATION.....GM..... | 52 | 7 | 28 | 37 | 39 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 110 | 0 | 49 | 68 | 100 |
| STANDARD DEVIATION.....GM..... | 53 | 0 | 15 | 41 | 41 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 850 | 85 | 255 | 510 | 340 |
| MAXIMUM OVER 3 DAYS.....GM..... | 869 | 85 | 267 | 510 | 680 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 95 | 22 | 56 | 68 | 78 |
| STANDARD DEVIATION.....GM..... | 56 | 17 | 40 | 45 | 41 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 21 | | 10 | 11 | 20 |
| 25TH.....GM..... | 65 | | 31 | 41 | 43 |
| 50TH.....GM..... | 83 | | 43 | 77 | 83 |
| 75TH.....GM..... | 123 | | 83 | 85 | 85 |
| 90TH.....GM..... | 170 | | 85 | 123 | 154 |
| 95TH.....GM..... | 170 | | 154 | 165 | 170 |
| 99TH.....GM..... | 330 | | 170 | 170 | 170 |
| MAXIMUM.....GM..... | 850 | 85 | 255 | 510 | 340 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 49.3 | 0 | 2.7 | 0 | 5.2 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .5 | 0 | .9 | 0 | .7 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 32.0 | 31.0 | 28.8 | 24.5 | 26.7 | 24.3 | 23.5 | 18.9 | 17.4 | 15.0 | 16.6 | 13.1 |
| 26.3 | 26.1 | 23.5 | 21.5 | 23.2 | 21.4 | 19.9 | 16.1 | 14.5 | 12.5 | 14.6 | 11.2 |
| 5.0 | 4.5 | 4.9 | 2.8 | 3.3 | 2.6 | 3.2 | 2.6 | 2.3 | 2.3 | 1.5 | 1.7 |
| .7 | .5 | .4 | .2 | .2 | .3 | .4 | .3 | .6 | .1 | .5 | .1 |
| 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.1 | 1.2 | 1.2 | 1.3 | 1.2 | 1.3 | 1.2 |
| .5 | .5 | .5 | .4 | .4 | .4 | .5 | .5 | .5 | .5 | .8 | .4 |
| 5.0 | 4.0 | 4.0 | 4.0 | 4.0 | 3.0 | 5.0 | 4.0 | 3.0 | 3.0 | 6.0 | 3.0 |
| 39 | 33 | 48 | 36 | 46 | 35 | 44 | 37 | 41 | 38 | 43 | 35 |
| 30 | 22 | 33 | 28 | 30 | 22 | 30 | 26 | 27 | 25 | 31 | 19 |
| 11 | 7 | 14 | 10 | 14 | 10 | 14 | 7 | 14 | 7 | 7 | 14 |
| 26 | 19 | 28 | 26 | 28 | 26 | 28 | 26 | 26 | 26 | 26 | 26 |
| 28 | 28 | 41 | 28 | 41 | 28 | 28 | 28 | 28 | 28 | 28 | 28 |
| 55 | 43 | 57 | 51 | 57 | 51 | 57 | 51 | 55 | 51 | 55 | 51 |
| 77 | 57 | 84 | 57 | 83 | 57 | 82 | 57 | 83 | 57 | 83 | 57 |
| 99 | 77 | 113 | 95 | 110 | 77 | 104 | 84 | 89 | 85 | 128 | 69 |
| 161 | 113 | 170 | 113 | 154 | 113 | 154 | 128 | 154 | 117 | 145 | 103 |
| 98 | 86 | 122 | 95 | 120 | 94 | 110 | 92 | 105 | 96 | 115 | 93 |
| 66 | 50 | 73 | 61 | 66 | 50 | 58 | 54 | 60 | 58 | 76 | 43 |
| 96 | 84 | 111 | 96 | 122 | 90 | 118 | 97 | 95 | 95 | 79 | 95 |
| 56 | 43 | 58 | 54 | 59 | 38 | 56 | 50 | 37 | 46 | 48 | 30 |
| 113 | 77 | 150 | 142 | 128 | 121 | 122 | 109 | 111 | 108 | 139 | 39 |
| 47 | 44 | 80 | 81 | 56 | 29 | 52 | 55 | 37 | 0 | 8 | 0 |
| 850 | 340 | 510 | 454 | 495 | 340 | 492 | 369 | 369 | 350 | 425 | 206 |
| 850 | 510 | 863 | 869 | 825 | 510 | 680 | 738 | 462 | 462 | 434 | 308 |
| 95 | 83 | 116 | 94 | 118 | 93 | 110 | 92 | 99 | 93 | 98 | 89 |
| 62 | 47 | 70 | 59 | 66 | 48 | 58 | 52 | 54 | 51 | 54 | 39 |
| 21 | 21 | 41 | 39 | 41 | 22 | 41 | 21 | 28 | 21 | 21 | 39 |
| 77 | 43 | 83 | 63 | 83 | 77 | 77 | 63 | 77 | 77 | 77 | 77 |
| 83 | 83 | 85 | 83 | 85 | 83 | 85 | 83 | 85 | 83 | 83 | 83 |
| 85 | 85 | 165 | 85 | 165 | 123 | 154 | 115 | 145 | 123 | 128 | 85 |
| 170 | 165 | 170 | 170 | 170 | 165 | 170 | 165 | 165 | 165 | 170 | 165 |
| 170 | 170 | 246 | 170 | 246 | 170 | 231 | 170 | 170 | 170 | 231 | 170 |
| 340 | 248 | 340 | 340 | 340 | 231 | 308 | 252 | 330 | 340 | 255 | 170 |
| 850 | 340 | 510 | 454 | 495 | 330 | 492 | 369 | 369 | 350 | 255 | 170 |
| 5.0 | 5.2 | 0 | 0 | 4.1 | 12.0 | 9.8 | 1.9 | 1.2 | 2.0 | 0 | 0 |
| .6 | .6 | 0 | 0 | .3 | .8 | .7 | .1 | .5 | .6 | 0 | 0 |

TABLE 10.22.--CREAM-STYLE CORN
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.3 | 8.5 | 5.6 | 4.7 | 5.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.9 | 6.2 | 5.0 | 4.0 | 4.6 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | 1.0 | .6 | .7 | .5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 1.3 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.1 | 1.6 | 1.1 | 1.2 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .9 | .3 | .4 | .3 |
| MAXIMUM.....DO..... | 5.0 | 5.0 | 2.0 | 2.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 53 | 39 | 36 | 40 | 46 |
| STANDARD DEVIATION.....GM..... | 38 | 24 | 29 | 34 | 28 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 11 | | 2 | 5 | 11 |
| 25TH.....GM..... | 28 | | 16 | 21 | 21 |
| 50TH.....GM..... | 43 | | 21 | 32 | 43 |
| 75TH.....GM..... | 85 | | 43 | 43 | 43 |
| 90TH.....GM..... | 85 | | 85 | 85 | 85 |
| 95TH.....GM..... | 128 | | 85 | 128 | 85 |
| 99TH.....GM..... | 171 | | 128 | 171 | 149 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 144 | 99 | 86 | 98 | 128 |
| STANDARD DEVIATION.....GM..... | 88 | 50 | 62 | 68 | 72 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 151 | 58 | 137 | 131 | 115 |
| STANDARD DEVIATION.....GM..... | 91 | 13 | 34 | 74 | 60 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 117 | 70 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 88 | 36 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 768 | 213 | 256 | 256 | 384 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,024 | 383 | 384 | 512 | 448 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 141 | 75 | 95 | 104 | 123 |
| STANDARD DEVIATION.....GM..... | 86 | 46 | 65 | 70 | 71 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 31 | | 8 | 16 | 16 |
| 25TH.....GM..... | 71 | | 57 | 64 | 64 |
| 50TH.....GM..... | 128 | | 64 | 84 | 128 |
| 75TH.....GM..... | 192 | | 128 | 128 | 128 |
| 90TH.....GM..... | 256 | | 213 | 256 | 256 |
| 95TH.....GM..... | 256 | | 256 | 256 | 256 |
| 99TH.....GM..... | 512 | | 256 | 256 | 256 |
| MAXIMUM.....GM..... | 768 | 213 | 256 | 256 | 384 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 11.9 | .6 | 0 | 0 | 1.5 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | .9 | 0 | 0 | 1.5 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 4.6 | 4.5 | 4.2 | 3.9 | 4.4 | 4.0 | 4.6 | 3.9 | 4.7 | 4.7 | 1.9 | 2.7 |
| 4.3 | 4.2 | 4.0 | 3.8 | 4.2 | 3.7 | 4.1 | 3.5 | 3.9 | 4.1 | 1.4 | 2.3 |
| .2 | .3 | .1 | .1 | .2 | .4 | .5 | .3 | .8 | .6 | .5 | .4 |
| 0 | 0 | 0 | 0 | 0 | 0 | (2) | (2) | 0 | .1 | 0 | 0 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.2 | 1.6 | 1.2 |
| .3 | .3 | .2 | .2 | .2 | .3 | .4 | .4 | .5 | .5 | 1.1 | .6 |
| 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 3.0 | 4.0 | 3.0 | 4.0 | 4.0 | 3.0 |
| 50 | 54 | 60 | 46 | 63 | 47 | 63 | 51 | 62 | 56 | 123 | 56 |
| 38 | 33 | 42 | 33 | 41 | 31 | 41 | 35 | 32 | 32 | 105 | 44 |
| 11 | 10 | 11 | 4 | 13 | 7 | 21 | 10 | 21 | 21 | | |
| 28 | 43 | 43 | 21 | 43 | 27 | 43 | 28 | 43 | 43 | | |
| 43 | 43 | 43 | 43 | 43 | 43 | 43 | 43 | 43 | 43 | | |
| 64 | 85 | 85 | 43 | 85 | 43 | 85 | 85 | 85 | 85 | | |
| 85 | 85 | 128 | 85 | 85 | 85 | 128 | 85 | 128 | 85 | | |
| 85 | 107 | 171 | 85 | 128 | 85 | 171 | 107 | 128 | 96 | | |
| 171 | 171 | 213 | 161 | 256 | 213 | 213 | 171 | 149 | 213 | | |
| 136 | 154 | 179 | 128 | 180 | 129 | 166 | 139 | 166 | 150 | 210 | 132 |
| 76 | 87 | 126 | 79 | 108 | 68 | 95 | 80 | 74 | 71 | 103 | 90 |
| 210 | 148 | 117 | 241 | 188 | 129 | 182 | 134 | 142 | 126 | 424 | 190 |
| 143 | 74 | 72 | 0 | 99 | 91 | 72 | 76 | 65 | 48 | 122 | 64 |
| 0 | 0 | 0 | 0 | 0 | 0 | 167 | 175 | 0 | 213 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 96 | 117 | 0 | 0 | 0 | 0 |
| 512 | 512 | 640 | 384 | 768 | 512 | 512 | 512 | 448 | 289 | 512 | 512 |
| 1,024 | 512 | 640 | 483 | 768 | 640 | 768 | 896 | 448 | 640 | 1,024 | 512 |
| 140 | 148 | 172 | 131 | 179 | 129 | 165 | 137 | 150 | 136 | 234 | 138 |
| 86 | 80 | 111 | 75 | 109 | 80 | 90 | 77 | 71 | 62 | 74 | 81 |
| 32 | 16 | 32 | 32 | 38 | 21 | 64 | 32 | 48 | 49 | | |
| 64 | 84 | 128 | 64 | 128 | 64 | 128 | 64 | 128 | 128 | | |
| 128 | 128 | 128 | 128 | 128 | 128 | 128 | 128 | 128 | 128 | | |
| 128 | 192 | 256 | 128 | 256 | 128 | 256 | 192 | 192 | 128 | | |
| 256 | 256 | 256 | 256 | 256 | 256 | 256 | 256 | 256 | 256 | | |
| 256 | 256 | 384 | 256 | 287 | 256 | 384 | 256 | 256 | 256 | | |
| 512 | 256 | 512 | 256 | 513 | 512 | 512 | 279 | 256 | 256 | | |
| 512 | 512 | 512 | 256 | 768 | 512 | 512 | 384 | 256 | 256 | 384 | 512 |
| 0 | 1.2 | 0 | 0 | 0 | 1.3 | 1.8 | 4.6 | 0 | .9 | 0 | 0 |
| 0 | 1.1 | 0 | 0 | 0 | .5 | .7 | 1.5 | 0 | .9 | 0 | 0 |

TABLE 10.23.--COOKED LIMA BEANS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.8 | 1.0 | 2.4 | 1.9 | 1.9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.5 | 1.0 | 2.1 | 1.5 | 1.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .3 | 0 | .3 | .4 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.2 | 1.0 | 1.2 | 1.3 | 1.1 |
| STANDARD DEVIATION.....DD..... | .4 | .0 | .4 | .4 | .3 |
| MAXIMUM.....DO..... | 3.0 | 1.0 | 2.0 | 2.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 43 | 24 | 21 | 21 | 29 |
| STANDARD DEVIATION.....GM..... | 33 | 22 | 21 | 15 | 20 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 7 | | | | |
| 25TH.....GM..... | 28 | | | | |
| 50TH.....GM..... | 29 | | | | |
| 75TH.....GM..... | 58 | | | | |
| 90TH.....GM..... | 85 | | | | |
| 95TH.....GM..... | 113 | | | | |
| 99TH.....GM..... | 170 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 114 | 71 | 52 | 56 | 84 |
| STANDARD DEVIATION.....GM..... | 82 | 67 | 30 | 34 | 45 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 120 | 0 | 77 | 40 | 64 |
| STANDARD DEVIATION.....GM..... | 77 | 0 | 64 | 33 | 66 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 101 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 65 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 875 | 175 | 175 | 175 | 175 |
| MAXIMUM OVER 3 DAYS.....GM..... | 875 | 175 | 350 | 175 | 350 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 110 | 71 | 54 | 49 | 79 |
| STANDARD DEVIATION.....GM..... | 75 | 67 | 38 | 31 | 47 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 21 | | | | |
| 25TH.....GM..... | 67 | | | | |
| 50TH.....GM..... | 88 | | | | |
| 75TH.....GM..... | 170 | | | | |
| 90TH.....GM..... | 175 | | | | |
| 95TH.....GM..... | 219 | | | | |
| 99TH.....GM..... | 350 | | | | |
| MAXIMUM.....GM..... | 875 | 175 | 175 | 88 | 175 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 11.4 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .9 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.8 | 2.3 | 2.6 | 1.8 | 2.7 | 2.5 | 3.5 | 3.2 | 4.5 | 4.3 | 2.9 | 3.7 |
| 1.6 | 2.2 | 2.4 | 1.8 | 2.3 | 2.3 | 3.1 | 2.8 | 4.3 | 3.7 | 2.7 | 2.8 |
| .2 | .1 | .2 | .1 | .4 | .2 | .4 | .4 | .3 | .6 | .2 | .8 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | (2) | 0 | 0 | 0 | .1 |
| 1.1 | 1.1 | 1.2 | 1.0 | 1.2 | 1.1 | 1.1 | 1.2 | 1.1 | 1.2 | 1.3 | 1.3 |
| .3 | .3 | .4 | .2 | .4 | .3 | .4 | .4 | .3 | .4 | .5 | .5 |
| 2.0 | 3.0 | 3.0 | 2.0 | 3.0 | 2.0 | 3.0 | 3.0 | 2.0 | 2.0 | 2.0 | 3.0 |
| 43 | 30 | 54 | 31 | 54 | 39 | 49 | 41 | 52 | 40 | 49 | 52 |
| 50 | 15 | 45 | 26 | 35 | 24 | 37 | 30 | 41 | 31 | 37 | 38 |
| | 7 | | | 7 | 7 | 15 | 7 | 15 | 7 | | |
| | 15 | | | 29 | 28 | 29 | 28 | 29 | 22 | | |
| | 29 | | | 57 | 29 | 29 | 29 | 44 | 29 | | |
| | 29 | | | 58 | 58 | 58 | 58 | 58 | 58 | | |
| | 58 | | | 113 | 58 | 113 | 58 | 88 | 68 | | |
| | 58 | | | 117 | 85 | 117 | 88 | 131 | 88 | | |
| | 58 | | | 170 | 117 | 248 | 170 | 233 | 175 | | |
| 112 | 91 | 146 | 96 | 150 | 113 | 125 | 109 | 138 | 98 | 130 | 123 |
| 148 | 46 | 101 | 78 | 105 | 68 | 74 | 73 | 82 | 62 | 95 | 81 |
| 132 | 59 | 205 | 11 | 119 | 89 | 169 | 105 | 214 | 129 | 175 | 142 |
| 34 | 26 | 125 | 0 | 32 | 47 | 92 | 61 | 132 | 64 | 0 | 61 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 146 | 0 | 0 | 0 | 7 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 875 | 175 | 700 | 340 | 510 | 350 | 394 | 510 | 394 | 350 | 263 | 350 |
| 875 | 175 | 744 | 340 | 510 | 350 | 744 | 525 | 700 | 525 | 350 | 510 |
| 114 | 86 | 141 | 91 | 137 | 106 | 129 | 103 | 138 | 101 | 111 | 123 |
| 133 | 45 | 94 | 78 | 83 | 61 | 79 | 59 | 87 | 61 | 60 | 83 |
| | 22 | | | 43 | 21 | 44 | 22 | 44 | 21 | | |
| | 44 | | | 88 | 58 | 88 | 67 | 88 | 77 | | |
| | 88 | | | 131 | 88 | 88 | 88 | 88 | 88 | | |
| | 88 | | | 175 | 170 | 175 | 170 | 175 | 117 | | |
| | 175 | | | 175 | 175 | 196 | 175 | 350 | 175 | | |
| | 175 | | | 340 | 175 | 340 | 175 | 350 | 175 | | |
| | 175 | | | 454 | 350 | 350 | 263 | 350 | 350 | | |
| 875 | 175 | 350 | 340 | 454 | 350 | 350 | 340 | 350 | 350 | 175 | 350 |
| .7 | 0 | .9 | 1.0 | 1.5 | 1.4 | 3.2 | 2.7 | 0 | 0 | 0 | 0 |
| 1.7 | 0 | 2.2 | 3.5 | 1.2 | .9 | 1.6 | 1.0 | 0 | 0 | 0 | 0 |

TABLE 10.24.--COOKED MIXED VEGETABLES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.4 | 11.4 | 3.7 | 3.1 | 4.0 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.0 | 7.7 | 3.3 | 2.7 | 3.8 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | 3.0 | .3 | .3 | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .1 | .7 | .1 | 0 | (2) |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.2 | 1.5 | 1.3 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .5 | .9 | .7 | .3 | .3 |
| MAXIMUM.....DO..... | 4.0 | 4.0 | 4.0 | 2.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 46 | 42 | 38 | 25 | 29 |
| STANDARD DEVIATION.....GM..... | 33 | 26 | 47 | 17 | 16 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 11 | 9 | | 7 | 7 |
| 25TH.....GM..... | 30 | 21 | | 16 | 16 |
| 50TH.....GM..... | 31 | 38 | | 20 | 31 |
| 75TH.....GM..... | 62 | 45 | | 31 | 31 |
| 90TH.....GM..... | 78 | 81 | | 39 | 61 |
| 95TH.....GM..... | 123 | 104 | | 62 | 62 |
| 99TH.....GM..... | 187 | 115 | | 94 | 78 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 118 | 99 | 86 | 66 | 83 |
| STANDARD DEVIATION.....GM..... | 70 | 57 | 83 | 38 | 44 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 126 | 79 | 97 | 79 | 101 |
| STANDARD DEVIATION.....GM..... | 63 | 37 | 26 | 38 | 12 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 138 | 96 | 256 | 0 | 43 |
| STANDARD DEVIATION.....GM..... | 77 | 18 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 454 | 344 | 454 | 187 | 187 |
| MAXIMUM OVER 3 DAYS.....GM..... | 935 | 344 | 767 | 281 | 233 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 117 | 81 | 89 | 69 | 82 |
| STANDARD DEVIATION.....GM..... | 69 | 47 | 78 | 40 | 44 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 28 | 23 | | 22 | 22 |
| 25TH.....GM..... | 91 | 57 | | 47 | 47 |
| 50TH.....GM..... | 94 | 64 | | 60 | 94 |
| 75TH.....GM..... | 182 | 115 | | 94 | 94 |
| 90TH.....GM..... | 187 | 128 | | 94 | 140 |
| 95TH.....GM..... | 187 | 135 | | 182 | 187 |
| 99TH.....GM..... | 374 | 344 | | 187 | 187 |
| MAXIMUM.....GM..... | 454 | 344 | 454 | 187 | 187 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 11.3 | 0 | 0 | 0 | 1.7 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .7 | 0 | 0 | 0 | 2.2 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.7 | 3.4 | 2.7 | 1.8 | 2.7 | 2.9 | 3.1 | 3.3 | 4.7 | 4.5 | 4.6 | 6.4 |
| 3.5 | 3.3 | 2.7 | 1.7 | 2.5 | 2.6 | 2.8 | 2.7 | 4.2 | 3.7 | 2.9 | 5.3 |
| .2 | .1 | 0 | .1 | .2 | .3 | .2 | .5 | .4 | .6 | 1.6 | .7 |
| 0 | 0 | 0 | 0 | .1 | 0 | .1 | (2) | 0 | .2 | .1 | .3 |
| 1.0 | 1.0 | 1.0 | 1.1 | 1.1 | 1.1 | 1.2 | 1.2 | 1.2 | 1.3 | 1.4 | 1.3 |
| .2 | .2 | .0 | .2 | .4 | .3 | .5 | .5 | .4 | .6 | .5 | .5 |
| 2.0 | 2.0 | 1.0 | 2.0 | 3.0 | 2.0 | 4.0 | 3.0 | 2.0 | 3.0 | 3.0 | 3.0 |
| 41 | 35 | 36 | 44 | 48 | 44 | 57 | 49 | 57 | 55 | 62 | 52 |
| 32 | 17 | 20 | 26 | 34 | 26 | 46 | 32 | 31 | 38 | 40 | 28 |
| 15 | 8 | | | 15 | 12 | 16 | 16 | 16 | 8 | | 30 |
| 30 | 30 | | | 31 | 31 | 31 | 31 | 31 | 31 | | 31 |
| 31 | 31 | | | 31 | 31 | 47 | 31 | 62 | 38 | | 39 |
| 47 | 47 | | | 62 | 61 | 62 | 62 | 62 | 62 | | 62 |
| 62 | 62 | | | 78 | 62 | 125 | 94 | 123 | 94 | | 94 |
| 94 | 62 | | | 121 | 94 | 125 | 125 | 123 | 130 | | 125 |
| 249 | 62 | | | 125 | 151 | 312 | 184 | 125 | 187 | | 125 |
| 112 | 102 | 107 | 134 | 126 | 114 | 146 | 121 | 152 | 128 | 124 | 148 |
| 63 | 51 | 60 | 80 | 69 | 58 | 86 | 68 | 69 | 69 | 47 | 86 |
| 158 | 81 | 0 | 53 | 159 | 133 | 128 | 135 | 186 | 149 | 145 | 91 |
| 144 | 10 | 0 | 31 | 94 | 44 | 70 | 56 | 1 | 33 | 67 | 28 |
| 0 | 0 | 0 | 0 | 125 | 0 | 192 | 78 | 0 | 156 | 109 | 77 |
| 0 | 0 | 0 | 0 | 0 | 0 | 91 | 0 | 0 | 58 | 0 | 18 |
| 374 | 187 | 374 | 374 | 374 | 454 | 374 | 374 | 369 | 374 | 281 | 374 |
| 748 | 187 | 374 | 374 | 748 | 468 | 935 | 561 | 374 | 561 | 561 | 374 |
| 116 | 101 | 107 | 124 | 129 | 116 | 146 | 122 | 149 | 128 | 133 | 123 |
| 75 | 50 | 60 | 80 | 73 | 58 | 85 | 66 | 54 | 56 | 60 | 75 |
| 46 | 23 | | | 46 | 23 | 47 | 47 | 47 | 47 | | 47 |
| 91 | 91 | | | 94 | 94 | 94 | 91 | 94 | 94 | | 91 |
| 94 | 94 | | | 94 | 94 | 116 | 94 | 182 | 121 | | 94 |
| 137 | 94 | | | 187 | 182 | 187 | 187 | 187 | 187 | | 182 |
| 187 | 187 | | | 187 | 187 | 234 | 187 | 187 | 187 | | 187 |
| 281 | 187 | | | 281 | 187 | 374 | 187 | 187 | 187 | | 374 |
| 374 | 187 | | | 374 | 281 | 374 | 374 | 281 | 187 | | 374 |
| 374 | 187 | 374 | 374 | 374 | 454 | 374 | 374 | 281 | 281 | 281 | 374 |
| 3.4 | 2.6 | 2.6 | .9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.2 | 3.5 | 6.9 | 3.2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

TABLE 10.25.--COOKED COMPEAS, FIELD PEAS, AND BLACKEYE PEAS
 CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.9 | .5 | 2.1 | 2.5 | 2.7 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.5 | .5 | 1.8 | 2.2 | 2.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .3 | 0 | .2 | .3 | (2) |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | .1 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 1.0 | 1.3 | 1.2 | 1.0 |
| STANDARD DEVIATION.....DO..... | .5 | .0 | .9 | .4 | .2 |
| MAXIMUM.....DO..... | 5.0 | 1.0 | 5.0 | 2.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 53 | 42 | 28 | 32 | 34 |
| STANDARD DEVIATION.....GM..... | 42 | 21 | 25 | 22 | 21 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 11 | | | | 7 |
| 25TH.....GM..... | 29 | | | | 28 |
| 50TH.....GM..... | 44 | | | | 29 |
| 75TH.....GM..... | 58 | | | | 44 |
| 90TH.....GM..... | 117 | | | | 58 |
| 95TH.....GM..... | 117 | | | | 58 |
| 99TH.....GM..... | 233 | | | | 117 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 140 | 127 | 73 | 93 | 98 |
| STANDARD DEVIATION.....GM..... | 101 | 64 | 70 | 68 | 56 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 136 | 0 | 56 | 63 | 175 |
| STANDARD DEVIATION.....GM..... | 90 | 0 | 38 | 17 | 0 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 193 | 0 | 69 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 89 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,050 | 175 | 350 | 263 | 350 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,050 | 175 | 350 | 263 | 350 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 131 | 127 | 63 | 84 | 97 |
| STANDARD DEVIATION.....GM..... | 88 | 64 | 50 | 60 | 57 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 22 | | | | 22 |
| 25TH.....GM..... | 88 | | | | 85 |
| 50TH.....GM..... | 88 | | | | 88 |
| 75TH.....GM..... | 175 | | | | 131 |
| 90TH.....GM..... | 196 | | | | 175 |
| 95TH.....GM..... | 350 | | | | 175 |
| 99TH.....GM..... | 350 | | | | 350 |
| MAXIMUM.....GM..... | 700 | 175 | 175 | 263 | 350 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 10.7 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .8 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 2.7 | 2.3 | 3.2 | 2.4 | 2.7 | 2.5 | 3.7 | 3.0 | 3.7 | 3.8 | 3.4 | 2.2 |
| 2.5 | 2.3 | 3.0 | 2.1 | 2.3 | 2.3 | 3.2 | 2.7 | 2.7 | 3.3 | 2.6 | 1.3 |
| .1 | (2) | .2 | .3 | .4 | .2 | .5 | .3 | 1.0 | .5 | .9 | .9 |
| .1 | 0 | 0 | 0 | .1 | 0 | 0 | (2) | 0 | 0 | 0 | 0 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.1 | 1.2 | 1.2 | 1.5 | 1.3 | 1.5 | 1.5 |
| .4 | .2 | .3 | .3 | .5 | .4 | .4 | .5 | .6 | .5 | .6 | .5 |
| 3.0 | 2.0 | 2.0 | 2.0 | 4.0 | 3.0 | 2.0 | 3.0 | 3.0 | 3.0 | 3.0 | 2.0 |
| 41 | 34 | 57 | 62 | 66 | 48 | 62 | 55 | 57 | 51 | 56 | 72 |
| 31 | 25 | 28 | 45 | 55 | 32 | 51 | 46 | 31 | 37 | 34 | 34 |
| 7 | 4 | | | 14 | 7 | 15 | 14 | | 11 | | |
| 29 | 19 | | | 29 | 28 | 29 | 29 | | 29 | | |
| 29 | 29 | | | 58 | 31 | 57 | 43 | | 29 | | |
| 58 | 39 | | | 88 | 58 | 59 | 58 | | 58 | | |
| 58 | 58 | | | 117 | 88 | 117 | 117 | | 117 | | |
| 88 | 88 | | | 117 | 117 | 143 | 175 | | 117 | | |
| 233 | 117 | | | 292 | 146 | 292 | 233 | | 175 | | |
| 109 | 97 | 161 | 162 | 153 | 135 | 165 | 143 | 183 | 140 | 144 | 188 |
| 57 | 69 | 76 | 101 | 98 | 94 | 127 | 104 | 95 | 108 | 93 | 109 |
| 174 | 175 | 159 | 181 | 179 | 113 | 157 | 145 | 68 | 112 | 122 | 130 |
| 115 | 0 | 21 | 103 | 89 | 36 | 111 | 107 | 38 | 55 | 42 | 33 |
| 87 | 0 | 0 | 0 | 280 | 0 | 0 | 233 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 350 | 350 | 350 | 350 | 525 | 438 | 1,050 | 875 | 438 | 525 | 350 | 438 |
| 700 | 350 | 350 | 700 | 875 | 438 | 1,050 | 875 | 438 | 525 | 350 | 438 |
| 109 | 96 | 151 | 163 | 161 | 126 | 157 | 136 | 115 | 113 | 110 | 143 |
| 60 | 67 | 63 | 100 | 93 | 83 | 106 | 94 | 75 | 68 | 46 | 83 |
| 22 | 11 | | | 44 | 22 | 43 | 22 | | 22 | | |
| 88 | 58 | | | 88 | 85 | 88 | 88 | | 88 | | |
| 88 | 88 | | | 175 | 88 | 131 | 88 | | 88 | | |
| 175 | 88 | | | 175 | 175 | 175 | 175 | | 175 | | |
| 175 | 175 | | | 350 | 263 | 340 | 263 | | 175 | | |
| 175 | 175 | | | 350 | 340 | 350 | 350 | | 263 | | |
| 350 | 350 | | | 525 | 350 | 526 | 525 | | 350 | | |
| 350 | 350 | 350 | 350 | 525 | 350 | 700 | 525 | 350 | 350 | 175 | 438 |
| 0 | 0 | .6 | 0 | 1.4 | 3.1 | 4.3 | 1.4 | 0 | 0 | 0 | 0 |
| 0 | 0 | 1.1 | 0 | 1.1 | 2.0 | 2.0 | .5 | 0 | 0 | 0 | 0 |

TABLE 10.26.--COOKED GREEN PEAS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 18.3 | 16.0 | 21.8 | 20.9 | 22.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 15.7 | 10.5 | 18.5 | 17.6 | 19.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.4 | 5.0 | 3.1 | 3.1 | 2.6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | .5 | .2 | .1 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.2 | 1.5 | 1.2 | 1.2 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .7 | .4 | .4 | .4 |
| MAXIMUM.....DO..... | 6.0 | 4.0 | 3.0 | 3.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 35 | 30 | 21 | 24 | 27 |
| STANDARD DEVIATION.....GM..... | 26 | 23 | 17 | 20 | 20 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 7 | 5 | 3 | 3 | 4 |
| 25TH.....GM..... | 19 | 14 | 7 | 14 | 14 |
| 50TH.....GM..... | 28 | 28 | 14 | 19 | 28 |
| 75TH.....GM..... | 55 | 43 | 28 | 28 | 28 |
| 90TH.....GM..... | 57 | 57 | 43 | 55 | 57 |
| 95TH.....GM..... | 85 | 67 | 57 | 57 | 57 |
| 99TH.....GM..... | 113 | 142 | 85 | 110 | 113 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 92 | 80 | 51 | 58 | 71 |
| STANDARD DEVIATION.....GM..... | 60 | 68 | 35 | 37 | 48 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 94 | 56 | 60 | 73 | 81 |
| STANDARD DEVIATION.....GM..... | 55 | 35 | 39 | 45 | 41 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 96 | 32 | 50 | 75 | 65 |
| STANDARD DEVIATION.....GM..... | 52 | 20 | 4 | 11 | 22 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 850 | 425 | 255 | 340 | 340 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,190 | 425 | 340 | 425 | 425 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 90 | 61 | 53 | 61 | 72 |
| STANDARD DEVIATION.....GM..... | 57 | 45 | 36 | 42 | 46 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 20 | 7 | 10 | 10 | 11 |
| 25TH.....GM..... | 43 | 29 | 21 | 41 | 43 |
| 50TH.....GM..... | 85 | 57 | 43 | 43 | 83 |
| 75TH.....GM..... | 85 | 80 | 85 | 85 | 85 |
| 90TH.....GM..... | 170 | 128 | 85 | 85 | 165 |
| 95TH.....GM..... | 170 | 135 | 128 | 165 | 170 |
| 99TH.....GM..... | 330 | 213 | 170 | 170 | 170 |
| MAXIMUM.....GM..... | 680 | 255 | 170 | 330 | 340 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 41.2 | 0 | 2.1 | 1.6 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .5 | 0 | .8 | .4 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY, 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 20.9 | 19.4 | 18.1 | 16.9 | 17.6 | 15.7 | 19.9 | 17.6 | 18.9 | 17.5 | 15.0 | 15.9 |
| 18.0 | 16.7 | 15.2 | 14.8 | 15.5 | 14.1 | 17.1 | 15.2 | 15.6 | 14.4 | 11.1 | 12.9 |
| 2.7 | 2.6 | 2.8 | 2.1 | 2.1 | 1.6 | 2.5 | 2.3 | 2.8 | 2.7 | 3.4 | 2.9 |
| .2 | .1 | .1 | 0 | (2) | .1 | .3 | .1 | .5 | .3 | .5 | .1 |
| 1.2 | 1.2 | 1.2 | 1.1 | 1.1 | 1.1 | 1.2 | 1.2 | 1.3 | 1.2 | 1.4 | 1.2 |
| .4 | .4 | .4 | .4 | .4 | .3 | .4 | .4 | .6 | .5 | .9 | .5 |
| 3.0 | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 4.0 | 5.0 | 3.0 | 6.0 | 3.0 |
| 34 | 32 | 44 | 36 | 41 | 34 | 42 | 35 | 41 | 39 | 45 | 34 |
| 23 | 23 | 39 | 28 | 27 | 23 | 30 | 24 | 29 | 27 | 35 | 22 |
| 7 | 7 | 7 | 7 | 14 | 7 | 11 | 7 | 11 | 14 | 14 | 7 |
| 19 | 14 | 28 | 18 | 28 | 21 | 28 | 19 | 28 | 28 | 28 | 19 |
| 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 |
| 43 | 43 | 57 | 55 | 57 | 55 | 57 | 43 | 57 | 55 | 57 | 43 |
| 57 | 57 | 83 | 57 | 57 | 57 | 71 | 57 | 83 | 57 | 110 | 57 |
| 78 | 83 | 110 | 83 | 85 | 57 | 110 | 85 | 98 | 85 | 113 | 57 |
| 113 | 113 | 170 | 113 | 113 | 113 | 125 | 113 | 170 | 140 | 158 | 128 |
| 87 | 82 | 109 | 93 | 109 | 92 | 109 | 92 | 99 | 98 | 96 | 93 |
| 54 | 52 | 66 | 58 | 66 | 54 | 66 | 59 | 53 | 68 | 69 | 56 |
| 90 | 94 | 129 | 111 | 108 | 97 | 107 | 86 | 111 | 95 | 117 | 63 |
| 48 | 43 | 106 | 68 | 53 | 52 | 62 | 42 | 51 | 34 | 41 | 38 |
| 77 | 88 | 100 | 0 | 112 | 33 | 109 | 114 | 111 | 101 | 127 | 127 |
| 8 | 20 | 8 | 0 | 28 | 0 | 63 | 63 | 43 | 31 | 42 | 0 |
| 495 | 425 | 850 | 510 | 680 | 413 | 680 | 340 | 340 | 680 | 340 | 340 |
| 495 | 425 | 1,190 | 850 | 680 | 510 | 1,020 | 850 | 598 | 680 | 475 | 383 |
| 86 | 83 | 112 | 96 | 107 | 91 | 106 | 88 | 97 | 94 | 96 | 82 |
| 52 | 46 | 73 | 62 | 64 | 55 | 61 | 52 | 50 | 59 | 60 | 48 |
| 20 | 20 | 32 | 28 | 32 | 20 | 31 | 21 | 21 | 31 | 21 | 21 |
| 43 | 43 | 83 | 54 | 83 | 56 | 83 | 54 | 83 | 83 | 56 | 43 |
| 85 | 83 | 85 | 83 | 85 | 85 | 85 | 83 | 85 | 85 | 85 | 85 |
| 85 | 85 | 165 | 128 | 165 | 85 | 165 | 85 | 124 | 85 | 114 | 85 |
| 170 | 170 | 170 | 170 | 170 | 170 | 170 | 170 | 170 | 170 | 170 | 165 |
| 170 | 170 | 255 | 170 | 170 | 170 | 170 | 170 | 170 | 170 | 170 | 170 |
| 255 | 170 | 340 | 340 | 340 | 330 | 340 | 255 | 170 | 255 | 330 | 170 |
| 495 | 255 | 510 | 510 | 680 | 413 | 495 | 340 | 340 | 680 | 330 | 340 |
| 4.4 | 2.6 | 0 | .7 | 5.8 | 1.5 | 10.8 | 10.1 | 0 | 1.6 | 0 | 0 |
| .9 | .5 | 0 | .3 | .7 | .2 | .9 | .7 | 0 | .4 | 0 | 0 |

TABLE 10.27.--COOKED SPINACH
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.5 | .9 | 2.8 | 3.2 | 5.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 4.0 | .9 | 2.6 | 3.2 | 5.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | 0 | .1 | (2) | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | .1 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER.... | 1.1 | 1.1 | 1.1 | 1.0 | 1.0 |
| STANDARD DEVIATION.....DO..... | .3 | .3 | .4 | .2 | .2 |
| MAXIMUM.....DO..... | 3.0 | 2.0 | 3.0 | 2.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 45 | 10 | 21 | 25 | 33 |
| STANDARD DEVIATION.....GM..... | 29 | 6 | 18 | 18 | 21 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 8 | | | 1 | 8 |
| 25TH.....GM..... | 30 | | | 15 | 17 |
| 50TH.....GM..... | 34 | | | 30 | 32 |
| 75TH.....GM..... | 65 | | | 34 | 34 |
| 90TH.....GM..... | 68 | | | 39 | 65 |
| 95TH.....GM..... | 103 | | | 63 | 68 |
| 99TH.....GM..... | 137 | | | 98 | 95 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 122 | 30 | 58 | 76 | 92 |
| STANDARD DEVIATION.....GM..... | 73 | 18 | 51 | 55 | 56 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 122 | 0 | 81 | 51 | 119 |
| STANDARD DEVIATION.....GM..... | 59 | 0 | 0 | 0 | 28 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 119 | 0 | 43 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 33 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 454 | 51 | 205 | 293 | 308 |
| MAXIMUM OVER 3 DAYS.....GM..... | 570 | 51 | 205 | 293 | 308 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 121 | 26 | 58 | 73 | 93 |
| STANDARD DEVIATION.....GM..... | 70 | 19 | 48 | 53 | 56 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 24 | | | 4 | 20 |
| 25TH.....GM..... | 78 | | | 45 | 51 |
| 50TH.....GM..... | 103 | | | 68 | 95 |
| 75TH.....GM..... | 185 | | | 98 | 103 |
| 90TH.....GM..... | 205 | | | 103 | 190 |
| 95TH.....GM..... | 205 | | | 190 | 195 |
| 99TH.....GM..... | 380 | | | 293 | 285 |
| MAXIMUM.....GM..... | 454 | 51 | 205 | 293 | 308 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER.... | 14.5 | 0 | 0 | 0 | 1.7 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .8 | 0 | 0 | 0 | 1.7 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 5.2 | 3.6 | 4.5 | 3.0 | 4.0 | 4.0 | 4.7 | 4.9 | 6.3 | 6.3 | 5.5 | 5.5 |
| 4.9 | 3.4 | 4.0 | 2.8 | 3.9 | 3.7 | 4.1 | 4.5 | 5.5 | 4.8 | 3.8 | 4.2 |
| .3 | .1 | .5 | .2 | .1 | .4 | .5 | .5 | .8 | 1.5 | 1.6 | 1.3 |
| (2) | 0 | 0 | 0 | 0 | 0 | .1 | (2) | 0 | 0 | 0 | 0 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.0 | 1.1 | 1.2 | 1.1 | 1.2 | 1.3 | 1.3 | 1.2 |
| .3 | .2 | .3 | .2 | .2 | .3 | .4 | .3 | .4 | .5 | .5 | .4 |
| 3.0 | 2.0 | 2.0 | 2.0 | 2.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 2.0 |
| 38 | 36 | 47 | 38 | 52 | 45 | 51 | 47 | 44 | 55 | 57 | 48 |
| 24 | 22 | 28 | 26 | 27 | 28 | 33 | 30 | 20 | 33 | 42 | 30 |
| 8 | 8 | 16 | | 16 | 10 | 16 | 15 | 14 | 16 | | 4 |
| 24 | 17 | 32 | | 33 | 32 | 32 | 32 | 32 | 32 | | 33 |
| 33 | 33 | 34 | | 49 | 34 | 34 | 34 | 34 | 48 | | 43 |
| 51 | 49 | 65 | | 65 | 63 | 65 | 65 | 65 | 68 | | 60 |
| 65 | 65 | 86 | | 69 | 68 | 86 | 73 | 68 | 101 | | 90 |
| 87 | 68 | 98 | | 103 | 85 | 133 | 127 | 68 | 127 | | 120 |
| 114 | 110 | 137 | | 130 | 171 | 167 | 137 | 103 | 137 | | 151 |
| 103 | 104 | 137 | 105 | 152 | 121 | 129 | 130 | 132 | 138 | 121 | 132 |
| 61 | 62 | 81 | 64 | 80 | 67 | 64 | 78 | 62 | 77 | 86 | 88 |
| 130 | 94 | 91 | 129 | 121 | 134 | 148 | 126 | 64 | 125 | 143 | 93 |
| 24 | 62 | 46 | 56 | 37 | 64 | 71 | 51 | 19 | 57 | 63 | 46 |
| 86 | 0 | 0 | 0 | 0 | 0 | 126 | 152 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 9 | 0 | 0 | 0 | 0 | 0 |
| 410 | 308 | 410 | 308 | 410 | 385 | 390 | 425 | 308 | 390 | 309 | 454 |
| 410 | 330 | 410 | 412 | 410 | 512 | 513 | 458 | 308 | 570 | 475 | 454 |
| 105 | 102 | 127 | 108 | 150 | 122 | 131 | 129 | 112 | 128 | 128 | 117 |
| 59 | 62 | 80 | 64 | 79 | 64 | 63 | 75 | 62 | 64 | 74 | 77 |
| 23 | 19 | 23 | | 48 | 29 | 37 | 38 | 41 | 45 | | 23 |
| 71 | 52 | 90 | | 98 | 95 | 95 | 95 | 52 | 90 | | 74 |
| 98 | 98 | 103 | | 146 | 103 | 103 | 103 | 95 | 103 | | 103 |
| 130 | 140 | 190 | | 195 | 180 | 195 | 190 | 172 | 190 | | 180 |
| 195 | 195 | 205 | | 205 | 195 | 205 | 205 | 205 | 205 | | 190 |
| 195 | 205 | 205 | | 293 | 205 | 205 | 256 | 205 | 205 | | 270 |
| 205 | 308 | 410 | | 390 | 380 | 308 | 385 | 308 | 308 | | 454 |
| 410 | 308 | 410 | 308 | 410 | 385 | 390 | 425 | 308 | 380 | 309 | 454 |
| 1.4 | 1.7 | .8 | .9 | 3.1 | 2.6 | 0 | 2.3 | 0 | 0 | 0 | 0 |
| 1.1 | 2.1 | 1.2 | 1.9 | 1.9 | 1.1 | 0 | .6 | 0 | 0 | 0 | 0 |

TABLE 10.28.--COOKED STRING BEANS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 27.3 | 19.7 | 25.1 | 25.4 | 31.6 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 22.7 | 13.0 | 20.5 | 21.5 | 26.7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 4.1 | 5.7 | 4.0 | 3.8 | 4.3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .5 | 1.1 | .5 | .2 | .7 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 1.6 | 1.2 | 1.2 | 1.2 |
| STANDARD DEVIATION.....DO..... | .5 | .8 | .5 | .4 | .5 |
| MAXIMUM.....DO..... | 6.0 | 5.0 | 4.0 | 3.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 35 | 36 | 20 | 20 | 26 |
| STANDARD DEVIATION.....GM..... | 29 | 27 | 16 | 18 | 20 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 6 | 3 | 3 | 3 | 5 |
| 25TH.....GM..... | 23 | 17 | 11 | 11 | 12 |
| 50TH.....GM..... | 23 | 29 | 12 | 18 | 23 |
| 75TH.....GM..... | 47 | 45 | 23 | 23 | 28 |
| 90TH.....GM..... | 69 | 76 | 45 | 45 | 47 |
| 95TH.....GM..... | 93 | 77 | 47 | 47 | 70 |
| 99TH.....GM..... | 140 | 134 | 85 | 70 | 93 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 88 | 82 | 48 | 53 | 63 |
| STANDARD DEVIATION.....GM..... | 57 | 68 | 37 | 52 | 40 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 90 | 77 | 49 | 51 | 71 |
| STANDARD DEVIATION.....GM..... | 51 | 38 | 26 | 26 | 34 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 110 | 61 | 63 | 44 | 81 |
| STANDARD DEVIATION.....GM..... | 82 | 34 | 32 | 15 | 33 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 840 | 344 | 220 | 700 | 280 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,820 | 401 | 304 | 700 | 420 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 86 | 69 | 48 | 51 | 64 |
| STANDARD DEVIATION.....GM..... | 54 | 47 | 33 | 46 | 38 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 18 | 9 | 8 | 8 | 16 |
| 25TH.....GM..... | 67 | 35 | 34 | 34 | 35 |
| 50TH.....GM..... | 70 | 57 | 35 | 35 | 68 |
| 75TH.....GM..... | 135 | 107 | 68 | 70 | 70 |
| 90TH.....GM..... | 140 | 128 | 70 | 70 | 135 |
| 95TH.....GM..... | 140 | 135 | 101 | 135 | 140 |
| 99TH.....GM..... | 280 | 213 | 140 | 140 | 210 |
| MAXIMUM.....GM..... | 840 | 344 | 220 | 700 | 280 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 91.4 | 2.0 | 0 | 4.2 | 8.2 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .7 | 1.3 | 0 | .8 | 1.2 |

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 31.1 | 29.4 | 29.5 | 24.8 | 25.4 | 23.4 | 29.4 | 27.6 | 27.2 | 29.8 | 32.1 | 27.2 |
| 25.2 | 24.9 | 25.9 | 21.5 | 22.6 | 19.9 | 24.9 | 22.8 | 20.5 | 22.3 | 25.8 | 18.6 |
| 5.5 | 4.3 | 3.4 | 3.0 | 2.5 | 3.2 | 4.0 | 4.3 | 5.4 | 6.7 | 4.4 | 7.0 |
| .4 | .2 | .2 | .2 | .3 | .3 | .5 | .5 | 1.3 | .8 | 1.8 | 1.6 |
| 1.2 | 1.2 | 1.2 | 1.2 | 1.1 | 1.2 | 1.2 | 1.2 | 1.4 | 1.4 | 1.3 | 1.5 |
| .5 | .5 | .4 | .4 | .4 | .4 | .5 | .5 | .7 | .7 | .7 | .8 |
| 4.0 | 5.0 | 3.0 | 4.0 | 4.0 | 3.0 | 6.0 | 6.0 | 4.0 | 5.0 | 5.0 | 6.0 |
| 31 | 30 | 36 | 33 | 38 | 33 | 42 | 37 | 45 | 42 | 38 | 42 |
| 31 | 25 | 29 | 22 | 24 | 22 | 37 | 27 | 36 | 33 | 30 | 29 |
| 5 | 6 | 9 | 11 | 11 | 6 | 12 | 11 | 11 | 11 | 11 | 9 |
| 18 | 15 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 |
| 23 | 23 | 23 | 23 | 34 | 23 | 35 | 23 | 41 | 35 | 23 | 35 |
| 45 | 45 | 47 | 45 | 47 | 47 | 47 | 47 | 47 | 47 | 47 | 47 |
| 47 | 47 | 68 | 47 | 70 | 49 | 90 | 70 | 93 | 93 | 77 | 90 |
| 70 | 70 | 90 | 79 | 93 | 76 | 93 | 93 | 117 | 93 | 93 | 93 |
| 117 | 135 | 140 | 117 | 123 | 113 | 175 | 140 | 228 | 187 | 161 | 140 |
| 75 | 76 | 91 | 89 | 100 | 85 | 103 | 92 | 110 | 94 | 99 | 94 |
| 47 | 57 | 51 | 58 | 56 | 48 | 59 | 54 | 86 | 57 | 73 | 55 |
| 76 | 79 | 114 | 79 | 101 | 87 | 111 | 99 | 90 | 97 | 66 | 91 |
| 42 | 52 | 79 | 36 | 40 | 44 | 54 | 52 | 43 | 57 | 29 | 48 |
| 138 | 98 | 99 | 73 | 114 | 95 | 186 | 91 | 124 | 140 | 101 | 80 |
| 160 | 19 | 24 | 13 | 41 | 36 | 142 | 51 | 51 | 50 | 43 | 39 |
| 840 | 700 | 630 | 420 | 420 | 338 | 700 | 560 | 700 | 420 | 420 | 280 |
| 1,820 | 700 | 770 | 420 | 560 | 560 | 1,820 | 700 | 700 | 595 | 514 | 560 |
| 75 | 74 | 93 | 83 | 99 | 85 | 106 | 91 | 99 | 92 | 87 | 86 |
| 54 | 55 | 58 | 51 | 53 | 47 | 60 | 52 | 60 | 47 | 55 | 47 |
| 17 | 17 | 27 | 26 | 34 | 18 | 34 | 26 | 28 | 17 | 17 | 18 |
| 35 | 35 | 68 | 68 | 68 | 68 | 70 | 68 | 70 | 68 | 68 | 68 |
| 70 | 70 | 70 | 70 | 70 | 70 | 87 | 70 | 70 | 70 | 70 | 70 |
| 70 | 70 | 140 | 105 | 140 | 135 | 140 | 135 | 140 | 140 | 135 | 135 |
| 140 | 140 | 140 | 140 | 140 | 140 | 140 | 140 | 140 | 140 | 140 | 140 |
| 140 | 140 | 140 | 140 | 147 | 140 | 210 | 140 | 140 | 140 | 140 | 140 |
| 270 | 280 | 280 | 280 | 280 | 270 | 280 | 280 | 280 | 210 | 280 | 280 |
| 840 | 700 | 560 | 420 | 420 | 280 | 560 | 420 | 454 | 280 | 397 | 280 |
| 4.2 | 4.8 | 5.1 | 5.2 | 13.2 | 5.9 | 12.0 | 18.5 | 0 | 6.0 | 0 | 2.1 |
| .5 | .6 | 1.1 | 1.2 | 1.2 | .4 | .7 | .8 | 0 | .8 | 0 | .5 |

TABLE 10.29.--COOKED SUMMER SQUASH
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 2.8 | .7 | 1.3 | 1.4 | 1.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.4 | .7 | 1.0 | 1.4 | 1.0 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .3 | 0 | .2 | .1 | .1 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 0 | 0 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.2 | 1.0 | 1.3 | 1.0 | 1.1 |
| STANDARD DEVIATION.....DO..... | .4 | .0 | .4 | .2 | .3 |
| MAXIMUM.....DO..... | 4.0 | 1.0 | 2.0 | 2.0 | 2.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 56 | 9 | 41 | 34 | 50 |
| STANDARD DEVIATION.....GM..... | 46 | 6 | 25 | 47 | 63 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 11 | | | | |
| 25TH.....GM..... | 36 | | | | |
| 50TH.....GM..... | 36 | | | | |
| 75TH.....GM..... | 72 | | | | |
| 90TH.....GM..... | 108 | | | | |
| 95TH.....GM..... | 143 | | | | |
| 99TH.....GM..... | 287 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 143 | 26 | 98 | 73 | 100 |
| STANDARD DEVIATION.....GM..... | 95 | 19 | 59 | 49 | 61 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 169 | 0 | 115 | 352 | 291 |
| STANDARD DEVIATION.....GM..... | 98 | 0 | 3 | 0 | 162 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 192 | 0 | 0 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 108 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 860 | 54 | 215 | 352 | 529 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,075 | 54 | 237 | 704 | 881 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 145 | 26 | 96 | 97 | 136 |
| STANDARD DEVIATION.....GM..... | 98 | 19 | 63 | 91 | 121 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 27 | | | | |
| 25TH.....GM..... | 105 | | | | |
| 50TH.....GM..... | 108 | | | | |
| 75TH.....GM..... | 215 | | | | |
| 90TH.....GM..... | 215 | | | | |
| 95TH.....GM..... | 352 | | | | |
| 99TH.....GM..... | 430 | | | | |
| MAXIMUM.....GM..... | 860 | 54 | 215 | 352 | 529 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 12.3 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 1.0 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.2 | 1.7 | 2.1 | 1.2 | 2.3 | 2.6 | 3.6 | 3.4 | 5.5 | 5.8 | 4.4 | 4.3 |
| 1.0 | 1.6 | 1.9 | 1.2 | 2.1 | 2.4 | 3.2 | 3.0 | 4.3 | 5.2 | 3.7 | 3.3 |
| .1 | .1 | .2 | .1 | .2 | .2 | .3 | .3 | 1.3 | .5 | .6 | .7 |
| .1 | 0 | 0 | 0 | (2) | 0 | .1 | .1 | 0 | .1 | 0 | .3 |
| 1.3 | 1.1 | 1.1 | 1.1 | 1.2 | 1.1 | 1.1 | 1.2 | 1.2 | 1.1 | 1.1 | 1.4 |
| .6 | .2 | .3 | .2 | .5 | .3 | .4 | .5 | .4 | .4 | .4 | .7 |
| 3.0 | 2.0 | 2.0 | 2.0 | 3.0 | 3.0 | 4.0 | 4.0 | 2.0 | 3.0 | 2.0 | 4.0 |
| 43 | 36 | 55 | 43 | 60 | 64 | 61 | 52 | 61 | 60 | 45 | 69 |
| 30 | 19 | 30 | 29 | 46 | 53 | 51 | 42 | 31 | 47 | 22 | 65 |
| | | | | 9 | 17 | 13 | 9 | 18 | 18 | | |
| | | | | 36 | 36 | 36 | 36 | 36 | 36 | | |
| | | | | 47 | 54 | 44 | 36 | 59 | 36 | | |
| | | | | 72 | 72 | 72 | 72 | 72 | 72 | | |
| | | | | 143 | 108 | 108 | 108 | 90 | 117 | | |
| | | | | 167 | 176 | 143 | 143 | 142 | 143 | | |
| | | | | 179 | 287 | 358 | 287 | 143 | 189 | | |
| 100 | 107 | 155 | 119 | 152 | 175 | 156 | 130 | 161 | 151 | 124 | 137 |
| 60 | 56 | 79 | 82 | 98 | 144 | 92 | 79 | 64 | 90 | 66 | 67 |
| 139 | 60 | 154 | 134 | 215 | 180 | 162 | 193 | 127 | 194 | 97 | 131 |
| 26 | 40 | 53 | 0 | 88 | 107 | 101 | 105 | 65 | 86 | 9 | 55 |
| 107 | 0 | 0 | 0 | 167 | 0 | 263 | 89 | 0 | 324 | 0 | 288 |
| 0 | 0 | 0 | 0 | 0 | 0 | 122 | 33 | 0 | 0 | 0 | 0 |
| 215 | 215 | 430 | 430 | 529 | 860 | 430 | 430 | 352 | 704 | 215 | 352 |
| 323 | 215 | 430 | 430 | 744 | 860 | 1,075 | 860 | 430 | 973 | 215 | 865 |
| 103 | 102 | 155 | 121 | 154 | 173 | 158 | 132 | 146 | 160 | 117 | 143 |
| 50 | 56 | 76 | 78 | 92 | 138 | 97 | 85 | 69 | 106 | 59 | 87 |
| | | | | 26 | 52 | 40 | 26 | 53 | 54 | | |
| | | | | 108 | 108 | 108 | 95 | 108 | 108 | | |
| | | | | 134 | 108 | 108 | 108 | 161 | 108 | | |
| | | | | 215 | 215 | 215 | 161 | 215 | 215 | | |
| | | | | 215 | 352 | 323 | 215 | 215 | 352 | | |
| | | | | 287 | 430 | 430 | 323 | 215 | 352 | | |
| | | | | 454 | 860 | 430 | 430 | 352 | 430 | | |
| 215 | 215 | 430 | 430 | 529 | 860 | 430 | 430 | 352 | 704 | 215 | 352 |
| 0 | 0 | 0 | 0 | 2.7 | 0 | 3.1 | 5.1 | 0 | 1.3 | 0 | 0 |
| 0 | 0 | 0 | 0 | 2.5 | 0 | 1.5 | 1.8 | 0 | 1.1 | 0 | 0 |

TABLE 10.30.--COOKED SWEET POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 4.1 | 10.8 | 3.8 | 3.1 | 3.2 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 3.7 | 8.0 | 3.7 | 3.1 | 2.9 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .4 | 1.7 | .1 | 0 | .3 |
| ON ALL 3 OF 3 DAYS.....PCT..... | (2) | 1.1 | .1 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE ^{1/}NUMBER..... | 1.2 | 1.5 | 1.1 | 1.1 | 1.1 |
| STANDARD DEVIATION.....00..... | .4 | .7 | .4 | .2 | .4 |
| MAXIMUM.....00..... | 4.0 | 3.0 | 3.0 | 2.0 | 3.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY ^{1/}GM..... | 52 | 40 | 36 | 34 | 38 |
| STANDARD DEVIATION.....GM..... | 38 | 24 | 35 | 19 | 25 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 14 | 9 | | 7 | 8 |
| 25TH.....GM..... | 34 | 21 | | 20 | 19 |
| 50TH.....GM..... | 38 | 38 | | 34 | 34 |
| 75TH.....GM..... | 68 | 45 | | 38 | 40 |
| 90TH.....GM..... | 85 | 76 | | 68 | 75 |
| 95TH.....GM..... | 132 | 81 | | 70 | 100 |
| 99TH.....GM..... | 200 | 147 | | 113 | 113 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 141 | 103 | 94 | 102 | 103 |
| STANDARD DEVIATION.....GM..... | 92 | 48 | 77 | 57 | 61 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 145 | 86 | 78 | 0 | 102 |
| STANDARD DEVIATION.....GM..... | 77 | 60 | 0 | 0 | 57 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE ^{1/}GM..... | 123 | 55 | 193 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 81 | 19 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,020 | 229 | 450 | 338 | 300 |
| MAXIMUM OVER 3 DAYS.....GM..... | 1,020 | 442 | 579 | 338 | 339 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE ^{1/}GM..... | 136 | 82 | 97 | 96 | 99 |
| STANDARD DEVIATION.....GM..... | 87 | 47 | 70 | 50 | 62 |
| PERCENTILES: ^{1/} | | | | | |
| 5TH.....GM..... | 38 | 19 | | 18 | 26 |
| 25TH.....GM..... | 86 | 57 | | 60 | 56 |
| 50TH.....GM..... | 114 | 64 | | 103 | 86 |
| 75TH.....GM..... | 185 | 115 | | 113 | 115 |
| 90TH.....GM..... | 225 | 128 | | 150 | 225 |
| 95TH.....GM..... | 238 | 135 | | 205 | 225 |
| 99TH.....GM..... | 450 | 229 | | 225 | 225 |
| MAXIMUM.....GM..... | 1,020 | 229 | 338 | 225 | 300 |
| QUANTITIES NOT SPECIFIED BY USERS: ^{1/} | | | | | |
| EATING OCCASIONS.....NUMBER..... | 13.4 | 0 | 0 | 0 | 1.0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .7 | 0 | 0 | 0 | 1.5 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 3.4 | 2.1 | 3.2 | 3.3 | 2.9 | 3.3 | 3.6 | 4.9 | 7.8 | 7.2 | 8.5 | 8.1 |
| 3.1 | 1.9 | 2.9 | 3.0 | 2.6 | 3.2 | 3.4 | 4.4 | 6.4 | 6.0 | 7.5 | 6.7 |
| .3 | .2 | .4 | .3 | .3 | .1 | .2 | .4 | 1.4 | 1.2 | .9 | 1.3 |
| 0 | 0 | 0 | 0 | 0 | (2) | (2) | (2) | 0 | .1 | 0 | .2 |
| 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.3 | 1.2 | 1.3 |
| .3 | .3 | .4 | .3 | .4 | .3 | .4 | .4 | .4 | .5 | .6 | .6 |
| 2.0 | 2.0 | 3.0 | 2.0 | 2.0 | 4.0 | 4.0 | 3.0 | 2.0 | 4.0 | 3.0 | 3.0 |
| 52 | 50 | 57 | 61 | 70 | 51 | 61 | 49 | 62 | 48 | 59 | 47 |
| 31 | 34 | 32 | 33 | 50 | 42 | 38 | 33 | 48 | 39 | 38 | 28 |
| 17 | | | | 22 | 9 | 20 | 17 | 19 | 11 | | 17 |
| 35 | | | | 38 | 34 | 38 | 33 | 34 | 22 | | 34 |
| 38 | | | | 62 | 38 | 50 | 38 | 40 | 38 | | 38 |
| 70 | | | | 75 | 68 | 75 | 67 | 75 | 60 | | 56 |
| 77 | | | | 133 | 78 | 103 | 76 | 123 | 98 | | 77 |
| 119 | | | | 150 | 137 | 150 | 113 | 153 | 113 | | 114 |
| 150 | | | | 340 | 178 | 205 | 200 | 265 | 300 | | 160 |
| 142 | 139 | 153 | 170 | 188 | 140 | 177 | 132 | 146 | 118 | 161 | 119 |
| 82 | 98 | 72 | 88 | 137 | 92 | 112 | 76 | 89 | 73 | 108 | 51 |
| 157 | 113 | 153 | 160 | 198 | 194 | 147 | 142 | 188 | 135 | 158 | 113 |
| 37 | 45 | 69 | 46 | 71 | 65 | 46 | 71 | 97 | 93 | 24 | 59 |
| 0 | 0 | 0 | 0 | 0 | 333 | 103 | 146 | 0 | 98 | 0 | 114 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 47 | 0 | 0 | 0 | 0 |
| 510 | 510 | 420 | 476 | 1,020 | 700 | 800 | 630 | 615 | 450 | 456 | 304 |
| 510 | 510 | 453 | 476 | 1,020 | 1,000 | 800 | 630 | 795 | 900 | 456 | 481 |
| 144 | 134 | 150 | 166 | 183 | 144 | 168 | 130 | 157 | 114 | 145 | 113 |
| 79 | 92 | 75 | 84 | 129 | 92 | 100 | 71 | 104 | 57 | 89 | 49 |
| 38 | | | | 56 | 28 | 57 | 42 | 57 | 33 | | 46 |
| 105 | | | | 113 | 103 | 114 | 99 | 103 | 76 | | 100 |
| 114 | | | | 168 | 114 | 130 | 114 | 119 | 113 | | 114 |
| 205 | | | | 225 | 200 | 210 | 180 | 205 | 130 | | 119 |
| 225 | | | | 255 | 228 | 238 | 225 | 228 | 200 | | 191 |
| 225 | | | | 450 | 400 | 410 | 225 | 410 | 210 | | 225 |
| 450 | | | | 1,020 | 450 | 450 | 315 | 615 | 357 | | 255 |
| 510 | 510 | 420 | 476 | 1,020 | 510 | 800 | 630 | 615 | 357 | 456 | 281 |
| 2.1 | 0 | .8 | 0 | 0 | 0 | 2.0 | 1.2 | 1.3 | 2.3 | 0 | 2.8 |
| 2.7 | 0 | 1.5 | 0 | 0 | 0 | 1.0 | .3 | 1.2 | 1.4 | 0 | 2.7 |

TABLE 10.31.--TOMATO JUICES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 3.9 | 0 | .8 | .9 | .9 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 2.6 | 0 | .7 | .8 | .7 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | .8 | 0 | 0 | (2) | .2 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .4 | 0 | .1 | 0 | 0 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.5 | 0 | 1.6 | 1.2 | 1.3 |
| STANDARD DEVIATION.....DO..... | .8 | 0 | 1.1 | .5 | .9 |
| MAXIMUM.....DO..... | 7.0 | 0 | 4.0 | 3.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 94 | 0 | 77 | 60 | 59 |
| STANDARD DEVIATION.....GM..... | 72 | 0 | 71 | 47 | 43 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 30 | | | | |
| 25TH.....GM..... | 45 | | | | |
| 50TH.....GM..... | 81 | | | | |
| 75TH.....GM..... | 121 | | | | |
| 90TH.....GM..... | 182 | | | | |
| 95TH.....GM..... | 243 | | | | |
| 99TH.....GM..... | 354 | | | | |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 191 | 0 | 211 | 184 | 135 |
| STANDARD DEVIATION.....GM..... | 105 | 0 | 222 | 143 | 41 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 208 | 0 | 0 | 61 | 177 |
| STANDARD DEVIATION.....GM..... | 104 | 0 | 0 | 0 | 98 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 204 | 0 | 121 | 0 | 0 |
| STANDARD DEVIATION.....GM..... | 99 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 1,879 | 0 | 790 | 729 | 456 |
| MAXIMUM OVER 3 DAYS.....GM..... | 2,844 | 0 | 790 | 729 | 699 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 190 | 0 | 147 | 156 | 133 |
| STANDARD DEVIATION.....GM..... | 85 | 0 | 73 | 61 | 48 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 91 | | | | |
| 25TH.....GM..... | 122 | | | | |
| 50TH.....GM..... | 182 | | | | |
| 75TH.....GM..... | 243 | | | | |
| 90TH.....GM..... | 243 | | | | |
| 95TH.....GM..... | 363 | | | | |
| 99TH.....GM..... | 486 | | | | |
| MAXIMUM.....GM..... | 1,150 | 0 | 243 | 243 | 243 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 13.6 | 0 | 0 | 0 | 0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | .6 | 0 | 0 | 0 | 0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 1.2 | 1.0 | 2.1 | 2.2 | 3.7 | 3.5 | 6.3 | 5.7 | 7.4 | 5.0 | 4.7 | 7.2 |
| 1.1 | .9 | 1.4 | 1.6 | 2.7 | 2.5 | 3.8 | 3.8 | 5.2 | 3.8 | 2.8 | 5.1 |
| .1 | .1 | .5 | .2 | .5 | .8 | 1.8 | 1.3 | .9 | .8 | 1.8 | 1.3 |
| 0 | 0 | .3 | .4 | .5 | .3 | .7 | .6 | 1.3 | .4 | .2 | .9 |
| 1.1 | 1.2 | 1.5 | 1.5 | 1.5 | 1.4 | 1.6 | 1.5 | 1.6 | 1.3 | 1.5 | 1.4 |
| .2 | .4 | .7 | .8 | .8 | .7 | .9 | .8 | .9 | .7 | .7 | .7 |
| 2.0 | 2.0 | 3.0 | 3.0 | 4.0 | 3.0 | 7.0 | 6.0 | 4.0 | 4.0 | 4.0 | 3.0 |
| 57 | 73 | 93 | 99 | 108 | 99 | 107 | 96 | 80 | 69 | 99 | 68 |
| 20 | 44 | 67 | 67 | 84 | 64 | 86 | 73 | 48 | 37 | 106 | 33 |
| | | | | 40 | 40 | 35 | 31 | 10 | 20 | | 40 |
| | | | | 61 | 61 | 61 | 61 | 41 | 41 | | 41 |
| | | | | 81 | 81 | 81 | 81 | 61 | 61 | | 61 |
| | | | | 122 | 141 | 151 | 121 | 121 | 81 | | 81 |
| | | | | 243 | 182 | 182 | 182 | 162 | 121 | | 122 |
| | | | | 324 | 243 | 243 | 243 | 172 | 182 | | 122 |
| | | | | 405 | 303 | 384 | 354 | 182 | 182 | | 161 |
| 169 | 187 | 181 | 217 | 204 | 214 | 206 | 187 | 170 | 165 | 117 | 155 |
| 58 | 103 | 79 | 111 | 87 | 114 | 156 | 68 | 97 | 60 | 34 | 58 |
| 93 | 199 | 209 | 183 | 273 | 228 | 205 | 210 | 193 | 141 | 219 | 146 |
| 42 | 64 | 135 | 43 | 131 | 102 | 74 | 127 | 40 | 53 | 56 | 43 |
| 0 | 0 | 188 | 195 | 238 | 195 | 225 | 219 | 136 | 154 | 567 | 121 |
| 0 | 0 | 9 | 76 | 88 | 52 | 140 | 73 | 35 | 30 | 0 | (2) |
| 304 | 547 | 486 | 605 | 547 | 729 | 1,879 | 1,573 | 486 | 304 | 972 | 363 |
| 304 | 547 | 972 | 1,089 | 1,214 | 1,094 | 2,844 | 2,541 | 547 | 547 | 1,701 | 484 |
| 159 | 183 | 191 | 194 | 222 | 211 | 198 | 189 | 153 | 154 | 199 | 144 |
| 63 | 95 | 94 | 84 | 96 | 93 | 99 | 67 | 68 | 55 | 104 | 49 |
| | | | | 121 | 121 | 92 | 92 | 30 | 91 | | 121 |
| | | | | 182 | 151 | 122 | 122 | 121 | 122 | | 122 |
| | | | | 182 | 182 | 182 | 182 | 122 | 122 | | 122 |
| | | | | 243 | 243 | 243 | 243 | 182 | 182 | | 182 |
| | | | | 365 | 363 | 243 | 243 | 243 | 243 | | 182 |
| | | | | 484 | 365 | 365 | 273 | 243 | 243 | | 242 |
| | | | | 514 | 607 | 486 | 484 | 365 | 304 | | 363 |
| 304 | 547 | 486 | 484 | 545 | 608 | 1,150 | 486 | 365 | 304 | 486 | 363 |
| 0 | 0 | 0 | 1.6 | 0 | 0 | 4.4 | 1.3 | 2.2 | 0 | 0 | 4.0 |
| 0 | 0 | 0 | 3.2 | 0 | 0 | .9 | .2 | 1.7 | 0 | 0 | 3.9 |

TABLE 10.32.--CATSUP

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER.... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 13.7 | .4 | 13.4 | 21.8 | 21.3 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 11.0 | .2 | 11.2 | 16.8 | 16.5 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 2.4 | .2 | 2.2 | 4.6 | 4.6 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .3 | 0 | .1 | .4 | .3 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER.... | 1.3 | 1.5 | 1.2 | 1.3 | 1.3 |
| STANDARD DEVIATION.....DO..... | .6 | .5 | .5 | .6 | .6 |
| MAXIMUM.....DO..... | 5.0 | 2.0 | 3.0 | 4.0 | 5.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 9 | 5 | 5 | 8 | 7 |
| STANDARD DEVIATION.....GM..... | 11 | (2) | 7 | 10 | 7 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 2 | | 1 | 1 | 2 |
| 25TH.....GM..... | 2 | | 2 | 2 | 2 |
| 50TH.....GM..... | 5 | | 4 | 5 | 5 |
| 75TH.....GM..... | 10 | | 5 | 10 | 10 |
| 90TH.....GM..... | 20 | | 10 | 20 | 15 |
| 95TH.....GM..... | 25 | | 15 | 25 | 23 |
| 99TH.....GM..... | 51 | | 34 | 46 | 46 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 21 | 15 | 12 | 18 | 17 |
| STANDARD DEVIATION.....GM..... | 28 | 0 | 10 | 24 | 17 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 21 | 9 | 14 | 20 | 18 |
| STANDARD DEVIATION.....GM..... | 22 | 0 | 11 | 18 | 15 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 22 | 0 | 73 | 17 | 14 |
| STANDARD DEVIATION.....GM..... | 21 | 0 | 0 | 5 | 7 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 495 | 15 | 83 | 273 | 137 |
| MAXIMUM OVER 3 DAYS.....GM..... | 495 | 17 | 220 | 273 | 137 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 20 | 11 | 13 | 17 | 16 |
| STANDARD DEVIATION.....GM..... | 25 | 4 | 13 | 22 | 16 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 3 | | 3 | 3 | 5 |
| 25TH.....GM..... | 6 | | 6 | 6 | 6 |
| 50TH.....GM..... | 15 | | 11 | 15 | 15 |
| 75TH.....GM..... | 28 | | 15 | 15 | 15 |
| 90TH.....GM..... | 38 | | 30 | 30 | 30 |
| 95TH.....GM..... | 68 | | 33 | 68 | 45 |
| 99TH.....GM..... | 137 | | 68 | 68 | 68 |
| MAXIMUM.....GM..... | 495 | 15 | 83 | 273 | 137 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER.... | 706.3 | 0 | 16.5 | 57.1 | 51.7 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 10.6 | 0 | 9.5 | 11.6 | 10.0 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 22.5 | 20.4 | 21.1 | 18.5 | 19.4 | 14.5 | 11.2 | 7.3 | 5.5 | 5.5 | 4.3 | 3.5 |
| 17.3 | 16.1 | 16.0 | 13.3 | 15.8 | 12.1 | 8.8 | 6.4 | 4.1 | 4.8 | 3.8 | 3.0 |
| 4.1 | 3.5 | 4.5 | 4.4 | 3.1 | 2.2 | 2.2 | .8 | 1.3 | .7 | .3 | .3 |
| 1.0 | .7 | .6 | .8 | .5 | .3 | .2 | .1 | .1 | 0 | .2 | .2 |
| 1.3 | 1.3 | 1.4 | 1.4 | 1.3 | 1.2 | 1.3 | 1.2 | 1.3 | 1.1 | 1.2 | 1.2 |
| .6 | .6 | .7 | .7 | .6 | .5 | .5 | .4 | .7 | .4 | .5 | .5 |
| 4.0 | 5.0 | 5.0 | 5.0 | 5.0 | 4.0 | 4.0 | 3.0 | 5.0 | 2.0 | 3.0 | 3.0 |
| 10 | 8 | 12 | 9 | 11 | 7 | 9 | 7 | 7 | 6 | 6 | 5 |
| 15 | 11 | 15 | 10 | 12 | 8 | 11 | 13 | 7 | 6 | 4 | 5 |
| 2 | 2 | 2 | 2 | 2 | 1 | 2 | 1 | 1 | 1 | | |
| 4 | 3 | 5 | 4 | 4 | 2 | 4 | 2 | 3 | 2 | | |
| 5 | 5 | 9 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | | |
| 10 | 10 | 15 | 10 | 14 | 10 | 10 | 7 | 10 | 5 | | |
| 20 | 15 | 25 | 23 | 23 | 15 | 18 | 11 | 10 | 10 | | |
| 27 | 23 | 35 | 28 | 35 | 23 | 23 | 20 | 23 | 15 | | |
| 91 | 51 | 91 | 47 | 60 | 46 | 51 | 46 | 46 | 40 | | |
| 23 | 18 | 27 | 22 | 25 | 18 | 23 | 20 | 18 | 16 | 17 | 14 |
| 39 | 21 | 33 | 22 | 26 | 18 | 30 | 41 | 23 | 13 | 14 | 14 |
| 26 | 19 | 34 | 22 | 30 | 17 | 20 | 11 | 11 | 16 | 9 | 3 |
| 26 | 15 | 31 | 23 | 31 | 14 | 18 | 6 | 5 | 18 | 0 | 0 |
| 24 | 30 | 18 | 15 | 26 | 32 | 17 | 12 | 27 | 0 | 6 | 15 |
| 25 | 32 | 8 | 9 | 14 | 26 | 14 | 7 | 0 | 0 | 0 | 0 |
| 454 | 303 | 360 | 273 | 273 | 137 | 341 | 495 | 137 | 68 | 60 | 68 |
| 454 | 333 | 360 | 288 | 375 | 273 | 341 | 495 | 137 | 120 | 60 | 68 |
| 23 | 18 | 26 | 20 | 25 | 18 | 21 | 17 | 16 | 15 | 14 | 13 |
| 34 | 24 | 29 | 23 | 25 | 18 | 26 | 36 | 18 | 15 | 9 | 13 |
| 4 | 5 | 6 | 3 | 6 | 4 | 6 | 3 | 3 | 3 | | |
| 6 | 6 | 11 | 6 | 11 | 6 | 6 | 6 | 6 | 6 | | |
| 15 | 15 | 15 | 15 | 15 | 15 | 15 | 15 | 15 | 15 | | |
| 30 | 15 | 30 | 28 | 30 | 17 | 30 | 15 | 15 | 15 | | |
| 45 | 30 | 57 | 45 | 60 | 33 | 33 | 30 | 30 | 30 | | |
| 68 | 45 | 68 | 68 | 68 | 60 | 60 | 45 | 33 | 60 | | |
| 137 | 137 | 137 | 120 | 137 | 102 | 137 | 137 | 137 | 68 | | |
| 454 | 273 | 273 | 273 | 205 | 137 | 341 | 495 | 137 | 68 | 33 | 68 |
| 62.6 | 72.2 | 52.2 | 62.8 | 89.0 | 83.5 | 78.5 | 63.9 | 1.9 | 11.2 | 0 | 3.3 |
| 9.9 | 12.5 | 12.7 | 16.5 | 9.0 | 8.9 | 11.2 | 10.6 | 2.3 | 10.2 | 0 | 8.0 |

TABLE 10.33.--MUSTARD
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | TOTAL | AGE (YEARS) AND SEX OF INDIVIDUAL | | | |
|---|--------|-----------------------------------|-----------------|-----------------|-----------------|
| | | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| | | | | | |
| TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 14.6 | 0 | 6.1 | 10.6 | 14.4 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 10.6 | 0 | 5.2 | 8.0 | 10.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 3.1 | 0 | .3 | 2.2 | 3.5 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .9 | 0 | .6 | .4 | .7 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE <u>1/</u>NUMBER..... | 1.4 | 0 | 1.3 | 1.4 | 1.4 |
| STANDARD DEVIATION.....DO..... | .7 | 0 | .6 | .6 | .7 |
| MAXIMUM.....DO..... | 6.0 | 0 | 3.0 | 3.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY <u>1/</u>GM..... | 4 | 0 | 2 | 3 | 3 |
| STANDARD DEVIATION.....GM..... | 4 | 0 | 2 | 3 | 3 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 1 | | (2) | (2) | (2) |
| 25TH.....GM..... | 2 | | 1 | 2 | 2 |
| 50TH.....GM..... | 2 | | 2 | 2 | 2 |
| 75TH.....GM..... | 5 | | 2 | 3 | 3 |
| 90TH.....GM..... | 7 | | 3 | 7 | 5 |
| 95TH.....GM..... | 10 | | 5 | 8 | 7 |
| 99TH.....GM..... | 20 | | 10 | 10 | 12 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 8 | 0 | 5 | 7 | 6 |
| STANDARD DEVIATION.....GM..... | 7 | 0 | 4 | 7 | 5 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 8 | 0 | 3 | 6 | 7 |
| STANDARD DEVIATION.....GM..... | 8 | 0 | 1 | 4 | 6 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE <u>1/</u>GM..... | 8 | 0 | 5 | 6 | 6 |
| STANDARD DEVIATION.....GM..... | 7 | 0 | 2 | 3 | 2 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 120 | 0 | 30 | 60 | 65 |
| MAXIMUM OVER 3 DAYS.....GM..... | 165 | 0 | 30 | 60 | 70 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE <u>1/</u>GM..... | 8 | 0 | 5 | 6 | 6 |
| STANDARD DEVIATION.....GM..... | 7 | 0 | 3 | 6 | 5 |
| PERCENTILES: <u>1/</u> | | | | | |
| 5TH.....GM..... | 3 | | 1 | 1 | 1 |
| 25TH.....GM..... | 5 | | 3 | 3 | 5 |
| 50TH.....GM..... | 5 | | 5 | 5 | 5 |
| 75TH.....GM..... | 10 | | 5 | 5 | 5 |
| 90TH.....GM..... | 15 | | 5 | 15 | 10 |
| 95TH.....GM..... | 20 | | 15 | 15 | 15 |
| 99TH.....GM..... | 30 | | 15 | 30 | 28 |
| MAXIMUM.....GM..... | 120 | 0 | 15 | 60 | 60 |
| QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u> | | | | | |
| EATING OCCASIONS.....NUMBER..... | 811.0 | 0 | 6.8 | 24.3 | 39.0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 10.5 | 0 | 8.4 | 9.8 | 10.2 |

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 17.9 | 16.6 | 20.4 | 16.6 | 21.1 | 16.6 | 18.2 | 11.5 | 10.7 | 7.5 | 5.5 | 3.9 |
| 12.9 | 11.7 | 14.9 | 11.9 | 14.1 | 12.8 | 12.1 | 9.5 | 8.4 | 5.8 | 3.4 | 2.2 |
| 3.6 | 4.0 | 4.1 | 3.7 | 4.9 | 3.1 | 4.7 | 1.7 | 2.0 | 1.6 | 2.2 | 1.5 |
| 1.5 | .9 | 1.5 | 1.0 | 2.0 | .7 | 1.5 | .3 | .4 | .1 | 0 | .2 |
| 1.5 | 1.4 | 1.4 | 1.5 | 1.6 | 1.3 | 1.5 | 1.2 | 1.3 | 1.3 | 1.4 | 1.6 |
| .8 | .7 | .8 | .7 | .9 | .6 | .8 | .5 | .5 | .5 | .5 | .7 |
| 5.0 | 5.0 | 4.0 | 5.0 | 6.0 | 5.0 | 6.0 | 3.0 | 3.0 | 3.0 | 2.0 | 3.0 |
| 4 | 3 | 5 | 3 | 5 | 3 | 4 | 3 | 3 | 2 | 3 | 3 |
| 4 | 3 | 7 | 3 | 6 | 3 | 4 | 2 | 3 | 2 | 3 | 2 |
| 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | | |
| 2 | 2 | 3 | 2 | 3 | 2 | 3 | 2 | 2 | 2 | | |
| 5 | 3 | 7 | 3 | 5 | 3 | 5 | 3 | 3 | 3 | | |
| 8 | 6 | 10 | 7 | 10 | 5 | 10 | 5 | 5 | 5 | | |
| 12 | 10 | 17 | 8 | 15 | 9 | 15 | 7 | 10 | 5 | | |
| 19 | 15 | 40 | 13 | 28 | 15 | 20 | 10 | 18 | 10 | | |
| 8 | 7 | 11 | 8 | 9 | 7 | 9 | 7 | 7 | 5 | 4 | 9 |
| 8 | 6 | 11 | 6 | 8 | 6 | 9 | 6 | 6 | 3 | 1 | 9 |
| 7 | 7 | 13 | 7 | 11 | 8 | 9 | 7 | 6 | 6 | 9 | 3 |
| 5 | 6 | 11 | 4 | 11 | 7 | 8 | 4 | 4 | 3 | 3 | 1 |
| 9 | 6 | 13 | 6 | 11 | 8 | 8 | 5 | 12 | 5 | 0 | 1 |
| 5 | 3 | 13 | 5 | 9 | 7 | 4 | 2 | 6 | 0 | 0 | 0 |
| 65 | 60 | 120 | 45 | 90 | 120 | 75 | 57 | 45 | 15 | 15 | 29 |
| 65 | 90 | 150 | 75 | 165 | 125 | 90 | 57 | 55 | 30 | 30 | 29 |
| 7 | 7 | 11 | 7 | 9 | 7 | 8 | 6 | 7 | 6 | 7 | 5 |
| 7 | 6 | 11 | 5 | 9 | 7 | 8 | 5 | 7 | 3 | 4 | 6 |
| 1 | 3 | 3 | 3 | 3 | 3 | 3 | 1 | 1 | 3 | | |
| 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | | |
| 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | | |
| 10 | 5 | 15 | 5 | 15 | 10 | 10 | 5 | 5 | 5 | | |
| 15 | 15 | 29 | 15 | 15 | 15 | 15 | 15 | 15 | 10 | | |
| 15 | 15 | 30 | 15 | 30 | 15 | 30 | 15 | 15 | 15 | | |
| 30 | 30 | 60 | 30 | 45 | 30 | 30 | 30 | 30 | 15 | | |
| 60 | 60 | 60 | 45 | 60 | 120 | 60 | 57 | 45 | 15 | 15 | 29 |
| 50.8 | 52.7 | 43.3 | 40.1 | 119.0 | 107.4 | 175.5 | 112.1 | 13.6 | 23.1 | 3.4 | 0 |
| 9.3 | 10.4 | 10.6 | 11.2 | 9.3 | 9.2 | 12.9 | 11.1 | 9.0 | 14.0 | 8.2 | 0 |

TABLE 10.34.--CUCUMBER PICKLES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

| MEASURE AND UNIT | AGE (YEARS) AND SEX OF INDIVIDUAL | | | | |
|---|-----------------------------------|-----------------|-----------------|-----------------|-----------------|
| | TOTAL | UNDER 1 | 1-2 | 3-5 | 6-8 |
| | | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE | MALE AND FEMALE |
| TOTAL INDIVIDUALS $\frac{1}{2}$ (WEIGHTED).....NUMBER..... | 37,874 | 498 | 1,045 | 1,719 | 1,841 |
| TOTAL INDIVIDUALS USING FOOD: | | | | | |
| AT LEAST ONCE IN 3 DAYS.....PCT..... | 9.2 | .2 | 4.6 | 6.2 | 8.1 |
| ON ONLY 1 OF 3 DAYS.....PCT..... | 7.9 | .2 | 4.4 | 5.6 | 7.2 |
| ON ONLY 2 OF 3 DAYS.....PCT..... | 1.0 | 0 | .3 | .5 | .8 |
| ON ALL 3 OF 3 DAYS.....PCT..... | .2 | 0 | 0 | 0 | .1 |
| EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS: | | | | | |
| AVERAGE $\frac{1}{2}$NUMBER..... | 1.2 | 1.0 | 1.1 | 1.1 | 1.1 |
| STANDARD DEVIATION.....DO..... | .5 | .0 | .4 | .3 | .4 |
| MAXIMUM.....DO..... | 5.0 | 1.0 | 3.0 | 2.0 | 4.0 |
| QUANTITY CONSUMED BY USERS: | | | | | |
| USED AT LEAST ONCE IN 3 DAYS: | | | | | |
| AVERAGE PER DAY $\frac{1}{2}$GM..... | 18 | 2 | 12 | 14 | 17 |
| STANDARD DEVIATION.....GM..... | 20 | 0 | 9 | 14 | 20 |
| AVERAGE PER DAY AT SPECIFIED PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 2 | | | 2 | 2 |
| 25TH.....GM..... | 6 | | | 5 | 5 |
| 50TH.....GM..... | 11 | | | 10 | 10 |
| 75TH.....GM..... | 22 | | | 22 | 22 |
| 90TH.....GM..... | 43 | | | 29 | 43 |
| 95TH.....GM..... | 50 | | | 45 | 64 |
| 99TH.....GM..... | 90 | | | 72 | 108 |
| USED ON ONLY 1 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 45 | 6 | 33 | 41 | 42 |
| STANDARD DEVIATION.....GM..... | 47 | 0 | 28 | 42 | 47 |
| USED ON ONLY 2 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 48 | 0 | 30 | 30 | 60 |
| STANDARD DEVIATION.....GM..... | 44 | 0 | 14 | 16 | 49 |
| USED ON ALL 3 OF 3 DAYS: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 47 | 0 | 0 | 0 | 55 |
| STANDARD DEVIATION.....GM..... | 34 | 0 | 0 | 0 | 0 |
| MAXIMUM IN ANY 1 DAY.....GM..... | 810 | 7 | 130 | 215 | 270 |
| MAXIMUM OVER 3 DAYS.....GM..... | 945 | 7 | 130 | 215 | 370 |
| QUANTITY CONSUMED PER EATING OCCASION: | | | | | |
| AVERAGE $\frac{1}{2}$GM..... | 45 | 6 | 32 | 38 | 45 |
| STANDARD DEVIATION.....GM..... | 45 | 0 | 26 | 36 | 46 |
| PERCENTILES: $\frac{1}{2}$ | | | | | |
| 5TH.....GM..... | 7 | | | 7 | 7 |
| 25TH.....GM..... | 16 | | | 13 | 15 |
| 50TH.....GM..... | 30 | | | 26 | 30 |
| 75TH.....GM..... | 65 | | | 60 | 65 |
| 90TH.....GM..... | 90 | | | 74 | 130 |
| 95TH.....GM..... | 130 | | | 135 | 135 |
| 99TH.....GM..... | 222 | | | 155 | 222 |
| MAXIMUM.....GM..... | 455 | 7 | 130 | 195 | 260 |
| QUANTITIES NOT SPECIFIED BY USERS: $\frac{1}{2}$ | | | | | |
| EATING OCCASIONS.....NUMBER..... | 217.1 | 0 | .8 | 8.3 | 11.0 |
| PROPORTION OF EATING OCCASIONS.....PCT..... | 5.3 | 0 | 1.5 | 7.0 | 6.5 |

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

| 9-14 | | 15-18 | | 19-34 | | 35-64 | | 65-74 | | 75 AND OVER | |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------------|--------|
| MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE | MALE | FEMALE |
| 2,089 | 2,158 | 1,394 | 1,473 | 3,928 | 5,346 | 4,929 | 7,069 | 1,118 | 1,738 | 536 | 993 |
| 8.6 | 9.1 | 9.9 | 8.5 | 11.7 | 11.0 | 10.7 | 9.1 | 7.8 | 6.6 | 7.6 | 6.3 |
| 7.7 | 8.1 | 8.6 | 7.3 | 10.0 | 9.8 | 8.7 | 7.8 | 6.4 | 5.9 | 6.2 | 5.2 |
| .6 | .8 | .9 | 1.0 | 1.4 | 1.1 | 1.5 | 1.1 | 1.4 | .6 | .9 | .9 |
| .2 | .2 | .5 | .1 | .3 | .1 | .5 | .2 | 0 | .1 | .5 | .2 |
| 1.1 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.3 | 1.2 | 1.2 | 1.1 | 1.3 | 1.2 |
| .4 | .5 | .6 | .5 | .5 | .4 | .6 | .5 | .4 | .4 | .7 | .5 |
| 3.0 | 4.0 | 4.0 | 4.0 | 4.0 | 4.0 | 5.0 | 4.0 | 2.0 | 3.0 | 3.0 | 3.0 |
| 18 | 19 | 18 | 23 | 19 | 15 | 21 | 17 | 15 | 15 | 18 | 17 |
| 19 | 28 | 19 | 28 | 18 | 16 | 21 | 19 | 13 | 16 | 24 | 14 |
| 3 | 2 | 2 | 2 | 3 | 2 | 2 | 2 | 2 | 2 | | 5 |
| 7 | 5 | 7 | 7 | 7 | 6 | 8 | 6 | 4 | 5 | | 7 |
| 10 | 10 | 11 | 12 | 12 | 10 | 13 | 11 | 10 | 10 | | 12 |
| 22 | 22 | 22 | 25 | 22 | 22 | 24 | 22 | 22 | 19 | | 22 |
| 43 | 43 | 43 | 49 | 43 | 35 | 45 | 39 | 26 | 28 | | 43 |
| 65 | 65 | 65 | 87 | 51 | 45 | 60 | 45 | 43 | 50 | | 49 |
| 87 | 108 | 90 | 127 | 87 | 87 | 90 | 90 | 65 | 90 | | 49 |
| 49 | 46 | 45 | 61 | 47 | 43 | 52 | 42 | 36 | 36 | 47 | 40 |
| 51 | 51 | 49 | 76 | 43 | 46 | 50 | 42 | 29 | 34 | 62 | 32 |
| 48 | 77 | 65 | 54 | 54 | 37 | 49 | 45 | 42 | 60 | 16 | 55 |
| 38 | 99 | 39 | 51 | 38 | 31 | 40 | 40 | 24 | 48 | 6 | 19 |
| 29 | 48 | 32 | 74 | 48 | 27 | 52 | 62 | 0 | 14 | 63 | 19 |
| 34 | 28 | 17 | 51 | 17 | 17 | 34 | 48 | 0 | 0 | 32 | 8 |
| 378 | 810 | 270 | 455 | 270 | 430 | 540 | 405 | 135 | 195 | 325 | 135 |
| 378 | 945 | 270 | 455 | 405 | 430 | 540 | 518 | 195 | 270 | 325 | 148 |
| 47 | 50 | 45 | 58 | 48 | 40 | 50 | 42 | 37 | 38 | 42 | 42 |
| 50 | 59 | 46 | 71 | 40 | 41 | 45 | 43 | 29 | 34 | 53 | 36 |
| 7 | 7 | 7 | 7 | 10 | 7 | 7 | 7 | 7 | 6 | | 13 |
| 15 | 13 | 20 | 14 | 20 | 15 | 20 | 16 | 13 | 15 | | 16 |
| 30 | 30 | 30 | 30 | 34 | 30 | 34 | 30 | 30 | 26 | | 30 |
| 65 | 65 | 60 | 65 | 65 | 60 | 65 | 65 | 60 | 57 | | 60 |
| 130 | 130 | 85 | 135 | 111 | 74 | 130 | 74 | 70 | 85 | | 130 |
| 150 | 195 | 130 | 195 | 130 | 130 | 135 | 130 | 78 | 130 | | 135 |
| 195 | 325 | 270 | 325 | 195 | 200 | 195 | 195 | 135 | 135 | | 135 |
| 378 | 405 | 270 | 455 | 270 | 390 | 270 | 405 | 135 | 135 | 325 | 135 |
| 15.9 | 10.2 | 10.4 | 6.8 | 25.5 | 52.1 | 39.8 | 27.4 | 2.6 | 6.4 | 0 | 0 |
| 7.9 | 4.4 | 6.1 | 4.6 | 4.6 | 7.6 | 6.0 | 3.6 | 2.5 | 4.9 | 0 | 0 |

APPENDIX A

System for Classifying Foods Into Food Groups Based on the First Three Digits of Seven-digit Food Codes

| <u>Major food groups</u> | <u>Major food subgroups</u> | <u>Minor food subgroups</u> | | |
|--|-----------------------------|---|---|---|
| 1 MILK AND MILK PRODUCTS | 11 Milk and Milk Drinks | 111 Milk, fluid--pasteurized, filled, buttermilk, and dry reconstituted | | |
| | | 112 Milk, concentrated fluid (canned) | | |
| | | 113 Milk, imitation | | |
| | | 114 Yogurt | | |
| | | 115 Chocolate, malted, shakes, and other flavored milk drinks | | |
| | | 116 Meal replacements with milk | | |
| | | 117 Baby formulas | | |
| | | 118 Milk, dry and powdered mixtures with dry milk, not reconstituted | | |
| | | 12 Cream and Cream Substitutes | 121 Sweet dairy cream (fluid whipped and dry) | |
| | | | 122 Sweet cream and whipped cream substitutes | |
| | | | 123 Sour dairy cream | |
| | | 13 Milk Desserts | 131 Milk desserts, frozen | 132 Milk desserts, not frozen |
| | | | | 133 Baby foods with milk |
| | | | | 134 Milk sauces and gravies |
| 135 Other milk products | | | | |
| 14 Cheeses | 141 Natural cheese | | | 142 Cottage cheese |
| | | 143 Cream cheese | | |
| | | 144 Processed cheese and cheese spreads | | |
| | | 145 Imitation cheese | | |
| | | 146 Cheese mixtures | | |
| | | 147 Cheese soups | | |
| | | 2 MEAT, POULTRY, FISH, AND MIXTURES | 21 Beef | 210 Meat, NFS ¹ , and beef, NFS ¹ |
| | | | | 211 Beef steaks with bone |
| 212 Beef steaks without bone | | | | |
| 213 Beef cuts with bone, not steaks | | | | |
| 214 Beef slices and chunks | | | | |
| 215 Ground beef patties and meat balls | | | | |
| 216 Other beef items | | | | |
| 217 Baby and junior beef | | | | |
| 22 Pork | 220 Pork, NFS ¹ | | 221 Pork chops | |
| | | | 222 Pork steaks and cutlets | |
| | | | 223 Ham | |
| | | | 224 Pork roasts, not ham | |
| | | | 225 Canadian bacon | |
| | | | 226 Bacon, salt pork, and fat back | |

¹NFS - not further specified

Food Group Codes--Continued

| <u>Major food groups</u> | <u>Major food subgroups</u> | <u>Minor food subgroups</u> |
|--------------------------|--|---|
| | | 227 Miscellaneous pork cuts |
| | | 228 Baby and junior pork, ham, and meat sticks |
| | 23 Lamb, Veal, Game, and Other Carcass Meat | 230 Lamb, NFS ¹ |
| | | 231 Lamb |
| | | 232 Veal |
| | | 233 Game |
| | | 234 Baby veal and lamb |
| | 24 Poultry | 241 Chicken |
| | | 242 Turkey |
| | | 243 Duck |
| | | 244 Rock cornish game hen and other poultry |
| | | 247 Baby and junior poultry |
| | 25 Organ Meats, Sausages, Luncheon Meats, and Spreads | 251 Organ meats and mixtures |
| | | 252 Frankfurters, sausages, luncheon meats, and meat spreads |
| | 26 Fish and Shellfish | 261 Finfish |
| | | 262 Other seafood |
| | | 263 Shellfish |
| | 27 Meat, Poultry, or Fish Combined With Nonmeat Items | 271 Meat, poultry, or fish in gravy, sauce, or creamed |
| | | 272 Meat, poultry, or fish combined with starch |
| | | 273 Meat, poultry, or fish with starch and vegetable |
| | | 274 Meat, poultry, or fish with vegetable, excluding white potatoes |
| | | 275 Sandwiches (reported as a unit) with meat, poultry, or fish |
| | | 276 Baby foods, combination with meat, poultry, or fish |
| | 28 Plate Meals, Soups and Gravies with Meat, Poultry or Fish Base and Plain Gelatin Drinks | 281 Frozen plate meals and soups |
| | | 283 Soups, broths, and extracts from meat poultry, or fish base |
| | | 284 Gelatin drink, plain |
| | | 285 Gravies, meat or poultry base, made with water |

¹NFS - not further specified

Food Group Codes--Continued

| <u>Major food groups</u> | <u>Major food subgroups</u> | <u>Minor food subgroups</u> |
|--|------------------------------------|------------------------------------|
| 3 EGGS, MIXTURES, AND SUBSTITUTES | 31 Eggs | 311 Chicken eggs |
| | | 312 Other poultry eggs |
| | | |
| | 32 Egg Mixtures | 321 Egg dishes |
| | | 322 Egg sandwiches |
| | | 323 Egg soups |
| | | 324 Meringues |
| | 33 Egg Substitutes | 330 Substitutes, NFS ¹ |
| | | 331 Made from powdered mixtures |
| | | 332 Made from frozen mixtures |
| | | 333 Made from liquid mixtures |
| | 4 DRY LEGUMES, NUTS, AND SEEDS | 41 Legumes |
| 411 Cooked or canned dried beans | | |
| 412 Cooked or canned dried bean mixtures | | |
| 413 Cooked dried peas and lentils and mixtures | | |
| 414 Soybean dried products | | |
| 415 Frozen meals with dried beans or peas as main course | | |
| 416 Soups, mainly legumes | | |
| 417 Baby-food legumes | | |
| 418 Meat substitutes | | |
| 419 Meat substitute sandwiches | | |
| 42 Nuts and Nut Butters and Mixtures | | |
| | 422 Nut butters | |
| | 423 Nut butter sandwiches | |
| | 424 Coconut beverages and mixtures | |
| | 425 Nut mixtures | |
| 43 Seeds and Seed Mixtures | 431 Seeds and seed mixtures | |
| 44 Carob and Carob Products | 441 Carob powders | |
| | 442 Carob chips | |
| 5 GRAIN PRODUCTS | 50 Flour | 500 Flour and dry mixes |
| | | 510 Bread, rolls, NFS ¹ |
| | 51 Yeast Breads and Rolls | 511 White bread and rolls |
| | | 512 Whole wheat bread and rolls |
| | | 513 Cracked wheat bread and rolls |
| | 514 Rye bread and rolls | |
| | 515 Oatmeal bread | |

¹NFS - Not further specified

Food Group Codes--Continued

| <u>Major food groups</u> | <u>Major food subgroups</u> | <u>Minor food subgroups</u> |
|--------------------------|--|--|
| | | 516 Multigrain bread |
| | | 517 Cottonseed bread |
| | | 518 Other breads |
| | 52 Quick Breads | 521 Biscuits |
| | | 522 Cornbread and corn muffins |
| | | 523 Other muffins and popovers |
| | | 524 Quickbreads excluding cornbread and muffins |
| | 53 Cakes, Cookies, Pies, and Pastries | 531 Cakes |
| | | 532 Cookies |
| | | 533 Pies |
| | | 534 Cobblers, eclairs, turnovers, and other pastries |
| | | 535 Danish, breakfast pastries, bars, and doughnuts |
| | | 536 Coffee cakes, not yeast type |
| | 54 Crackers and Salty Snacks from Grain Products | 541 Sweet crackers |
| | | 542 Low sodium (dietary) crackers |
| | | 543 Nonsweet crackers |
| | | 544 Salty snack products from grain products |
| | 55 Pancakes, Waffles, and French Toast | 551 Pancakes |
| | | 552 Waffles |
| | | 553 French toast |
| | | 554 Crepes |
| | | 555 Flour-water patties |
| | | 556 Flour-milk patties |
| | | 557 Rice flour cakes |
| | 56 Cooked Pastes and Cooked Cereals | 561 Pastes |
| | | 562 Cooked cereals |
| | 57 Ready-to-eat Cereals | 571 Cereal, ready-to-eat, NFS ¹ |
| | | 571 Buckwheat cereals |
| | | 572 Bran cereals |
| | | 573 Corn cereals |
| | | 574 Oat cereals |
| | | 575 Rice cereals |
| | | 576 Wheat cereals |
| | | 577 Multigrain cereals |
| | | 578 Baby cereals |
| | | 579 Other cereals |

¹NFS - not further specified

Food Group Codes--Continued

| <u>Major food groups</u> | <u>Major food subgroups</u> | <u>Minor food subgroups</u> |
|--------------------------|---|---|
| | 58 Mixtures, Mostly Grain | 581 Mixtures with animal protein 582 Mixtures without animal protein 583 Frozen plate meals 584 Soups with grain product as main ingredient 585 Baby food mixtures with animal protein |
| 6 FRUITS | 61 Citrus Fruits, and Juices | 611 Citrus fruits 612 Citrus fruit juices |
| | 62 Dried Fruits | 621 Dried fruits |
| | 63 Fruits and Berries (exclude citrus, dried) | 631 Fruit excluding berries 632 Berries 633 Mixtures of two or more fruits 634 Mixtures of fruits, berries, and nonfruit items |
| | 64 Fruit Juices and Nectars (exclude citrus) | 641 Juices 642 Nectars |
| | 67 Baby or Strained and Junior Fruits and Juices | 671 Fruits and fruit mixtures 672 Juices 673 Fruits with cereal 674 Fruit desserts and puddings |
| 7 VEGETABLES | 71 White Potatoes | 710 White potatoes, NFS ¹ 711 Baked, boiled, and canned 712 Chips and sticks 713 Creamed, scalloped, and au gratin 714 Fried 715 Mashed, stuffed, and puffs 716 Salad 717 Special recipes 718 Soups 719 Puerto Rican starchy vegetables |
| | 72 Dark-green Vegetables | 721 Dark-green leafy vegetables 722 Dark-green nonleafy vegetables 723 Dark-green vegetable soups |
| | 73 Deep-yellow Vegetables | 731 Carrots 732 Pumpkin 733 Squash 734 Sweetpotatoes 735 Deep-yellow vegetable soups |

¹NFS - not further specified

Food Group Codes--Continued

| <u>Major food groups</u> | <u>Major food subgroups</u> | <u>Minor food subgroups</u> |
|-----------------------------------|---|---|
| | 74 Tomatoes | 741 Raw tomatoes 742 Cooked tomatoes 743 Tomato juice 744 Tomato sauces 745 Tomato mixtures 746 Tomato soups 747 Tomato sandwiches |
| | 75 Other Vegetables and Vegetable Mixtures | 751 Raw vegetables 752 Cooked or canned vegetables 753 Cooked vegetables, mixture of two or more vegetables (including nuts) 754 Cooked vegetables with sauces, batters, casseroles 755 Olives, pickles, and relishes (excluding tomatoes) 756 Vegetable soups |
| | 76 Baby or Strained and Junior Vegetables and Mixtures, Mostly Vegetables | 761 Dark-green vegetables 762 Deep-yellow vegetables 764 Vegetables other than dark green, deep-yellow, and tomato 765 Potato mixtures with meat 766 Vegetable mixtures and meat with cereal 767 Vegetable mixtures and liver with cereal |
| | 77 Vegetable Mixtures with Animal Protein | 771 White potato mixtures 772 Puerto Rican starchy vegetable (viandas) mixtures 773 Other vegetable mixtures 775 Puerto Rican stews and soups with starchy vegetables (viandas) |
| 8 FATS, OILS, AND SALAD DRESSINGS | 81 Fats | 811 Table fats 812 Cooking fats 813 Other fats |
| | 82 Oils | 821 Vegetable oils |
| | 83 Salad Dressings | 831 Regular-type salad dressings 832 Low-calorie-type salad dressing |
| 9 SUGAR, SWEETS, AND BEVERAGES | 91 Sugar and Sweets | 911 Sugars 912 Sugar replacements and substitutes 913 Sirups, honey, and molasses 914 Jellies, jams, and preserves 915 Gelatin desserts and salads 916 Ices and popsicles |

Food Group Codes--Continued

| <u>Major food groups</u> | <u>Major food subgroups</u> | <u>Minor food subgroups</u> |
|---|-----------------------------|--|
| | | 917 Candies |
| | | 918 Chewing gum and cough drops |
| | 92 Nonalcoholic Beverages | 921 Coffee |
| | | 922 Coffee substitutes |
| | | 923 Tea |
| | | 924 Soft drinks |
| | | 925 Fruit ades and drinks |
| | | 926 Nonfruit beverages |
| | | 929 Sugar concentrates with vitamin C, powdered, not reconstituted |
| | 93 Alcoholic Beverages | 931 Beers and ales |
| | | 932 Cordials and liqueurs |
| | | 933 Cocktails |
| | | 934 Wines |
| | | 935 Distilled liquors |
| 0 FOOD ITEMS FOR WHICH NO NUTRITIVE VALUES ARE ASSIGNED | 00 Blank | 001 Artificial sweeteners |
| | | 002 Extracts, flavors, and vinegar |
| | | 003 Seasonings, spices, and herbs |
| | | 004 Vitamins, minerals, and supplements |
| | | 009 Cooking oils, sprays, and sticks |

APPENDIX B

Table Notes

Data in tables were derived as follows:

1. Total individuals include only those individuals with 3-day diet records.
2. Average number of eating occasions at which food was consumed in 3 days by users =

$$\frac{\text{Total number of eating occasions at which food was consumed}}{\text{Number of individuals using food at least once in 3 days}}$$

3. Average quantity per day consumed by individuals who used food at least once in 3 days =

$$\frac{\text{Total grams consumed by users divided by 3}}{\text{Number of individuals using at least once in 3 days}}$$

4. Percentiles are omitted when fewer than 50 individuals in a sex-age group reported the food.

5. Average quantity per day consumed by individuals who used food on only 1 of 3 days =

$$\frac{\text{Total grams consumed by 1-day users}}{\text{Number of individuals using on only 1 of 3 days}}$$

6. Average quantity per day consumed by individuals who used food on only 2 of 3 days =

$$\frac{\text{Total grams consumed by 2-day users divided by 2}}{\text{Number of individuals using on only 2 of 3 days}}$$

7. Average quantity per day consumed by individuals who used food on all 3 of 3 days =

$$\frac{\text{Total grams consumed by 3-day users divided by 3}}{\text{Number of individuals using on all 3 of 3 days}}$$

8. Average quantity consumed per eating occasion =

$$\frac{\text{Total grams consumed}}{\text{Total number of eating occasions}}$$

9. See number 4 above.

10. Quantities not specified by users are included in "quantity consumed by users" but excluded in "quantity consumed per eating occasion."