



United States
Department of
Agriculture

Agricultural
Research
Service

January 2021



TOTAL USUAL NUTRIENT INTAKE

from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level

What We Eat in America, NHANES 2015-2018

Table TH

Thiamin (mg) <i>EAR</i>	page 1	Vitamin D (µg) <i>EAR, UL</i>	page 13
Riboflavin (mg) <i>EAR</i>	2	Vitamin K (µg) <i>AI</i>	14
Niacin (mg) <i>EAR</i>	3	Calcium (mg) <i>EAR, UL</i>	15
Vitamin B6 (mg) <i>EAR, UL</i>	4	Phosphorus (mg) <i>EAR, UL</i>	16
Folate (µg DFE) <i>EAR</i>	5	Magnesium (mg) <i>EAR</i>	17
Folic acid (µg) <i>UL</i>	6	Iron (mg) <i>EAR, UL</i>	18
Choline (mg) <i>AI, UL</i>	7	Zinc (mg) <i>EAR, UL</i>	19
Vitamin B12 (µg) <i>EAR</i>	8	Copper (mg) <i>EAR, UL</i>	20
Vitamin C (mg) <i>EAR, UL</i>	9	Selenium (µg) <i>EAR, UL</i>	21
Vitamin C <i>Adult Smokers</i> (mg) <i>EAR, UL</i>	10	Sodium (mg) <i>AI, CDRR</i>	22
Vitamin C <i>Adult Non-Smokers</i> (mg) <i>EAR, UL</i>	11	Potassium (mg) <i>AI</i>	23
Vitamin C <i>Adult Smokers and Non-Smokers</i> (mg) <i>EAR, UL</i>	12	SAMPLE SIZE	24
		TABLE NOTES	25

Compared to Dietary Reference Intakes: **EAR**=Estimated Average Requirement; **AI**=Adequate Intake; **UL**=Tolerable Upper Intake Level;
CDRR=Chronic Disease Risk Reduction Intake; **AMDR**=Acceptable Macronutrient Distribution Range

Table TH 1. Thiamin (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR		
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)	
Over 350% poverty												
Males and females:												
1-3.....	18 (3.4)	1.18 (0.05)	0.69* (0.02)	0.78* (0.03)	0.92 (0.03)	1.11 (0.03)	1.35 (0.06)	1.64* (0.16)	1.93* (0.22)		0.4	<3
4-8.....	18 (3.7)	1.55 (0.06)	0.94* (0.03)	1.03 (0.03)	1.21 (0.03)	1.45 (0.03)	1.75 (0.09)	2.22 (0.28)	2.66* (0.30)		0.5	<3
Males:												
9-13.....	8* (2.8)	1.93 (0.09)	1.10* (0.05)	1.24* (0.05)	1.48 (0.05)	1.80 (0.06)	2.17 (0.08)	2.62* (0.12)	2.97* (0.19)		0.7	<3
14-18.....	5* (1.9)	2.23 (0.24)	1.19* (0.06)	1.33* (0.06)	1.59 (0.06)	1.93 (0.08)	2.31 (0.10)	2.76* (0.15)	3.13* (0.36)		1	<3
19-50.....	24 (3.4)	4.43 (0.90)	1.14 (0.04)	1.29 (0.04)	1.60 (0.05)	2.04 (0.06)	2.69 (0.15)	3.94 (0.57)	18.77 (11.17)		1	<3
51+.....	39 (3.1)	5.16 (0.87)	1.13 (0.05)	1.30 (0.05)	1.65 (0.06)	2.25 (0.11)	3.32 (0.14)	5.67 (1.07)	17.83 (10.77)		1	<3
19+.....	31 (2.2)	4.77 (0.63)	1.13 (0.04)	1.29 (0.04)	1.62 (0.04)	2.12 (0.05)	2.99 (0.10)	4.56 (0.86)	18.64 (7.65)		1	<3
Females:												
9-13.....	6* (2.7)	1.70 (0.06)	0.99* (0.04)	1.11* (0.05)	1.35 (0.05)	1.64 (0.06)	1.99 (0.08)	2.36* (0.10)	2.62* (0.14)		0.7	<3
14-18.....	12* (5.0)	1.85 (0.23)	0.84* (0.05)	0.96* (0.05)	1.17 (0.05)	1.46 (0.05)	1.84 (0.09)	2.43* (0.45)	3.21* (0.64)		0.9	7* (2.2)
19-50.....	25 (2.2)	4.11 (0.86)	0.84 (0.03)	0.95 (0.03)	1.18 (0.03)	1.52 (0.04)	2.18 (0.12)	3.69 (0.38)	9.98 (5.14)		0.9	7 (1.5)
51+.....	33 (3.0)	5.27 (0.85)	0.80 (0.03)	0.93 (0.03)	1.16 (0.03)	1.53 (0.05)	2.50 (0.15)	4.21 (0.67)	11.86 (13.77)		0.9	9 (1.2)
19+.....	29 (2.0)	4.72 (0.71)	0.82 (0.03)	0.94 (0.03)	1.17 (0.02)	1.53 (0.03)	2.35 (0.12)	3.92 (0.36)	11.00 (5.29)		0.9	8 (1.2)
All:												
1+.....	27 (1.5)	4.22 (0.36)	0.91 (0.02)	1.04 (0.02)	1.32 (0.02)	1.77 (0.03)	2.55 (0.06)	3.84 (0.15)	8.15 (2.68)		5 (0.5)	

NOTES: [†] Percent reporting supplement containing thiamin. Sample size presented on page 24.
 EAR=Estimated Average Requirement
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 2. Riboflavin (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)
Over 350% poverty											
Males and females:											
1-3.....	17 (3.5)	1.66 (0.06)	1.01* (0.05)	1.13* (0.05)	1.33 (0.05)	1.59 (0.06)	1.89 (0.07)	2.25* (0.11)	2.57* (0.15)	0.4	<3
4-8.....	15 (4.8)	1.92 (0.08)	1.18* (0.05)	1.30 (0.05)	1.52 (0.05)	1.82 (0.06)	2.18 (0.10)	2.69 (0.25)	3.18* (0.30)	0.5	<3
Males:											
9-13.....	6* (2.4)	2.22 (0.10)	1.20* (0.07)	1.37* (0.07)	1.68 (0.09)	2.11 (0.10)	2.62 (0.13)	3.22* (0.19)	3.64* (0.24)	0.8	<3
14-18.....	5* (1.9)	2.59 (0.26)	1.29* (0.08)	1.46* (0.08)	1.81 (0.09)	2.27 (0.10)	2.80 (0.13)	3.42* (0.20)	3.91* (0.46)	1.1	<3
19-50.....	25 (3.3)	4.15 (0.36)	1.28 (0.05)	1.52 (0.05)	2.02 (0.06)	2.78 (0.09)	3.80 (0.16)	5.38 (0.42)	11.48 (6.73)	1.1	<3
51+.....	39 (3.0)	7.37* (2.34)	1.30 (0.05)	1.57 (0.05)	2.15 (0.07)	3.12 (0.11)	4.46 (0.17)	6.58 (0.69)	21.93 (7.31)	1.1	<3
19+.....	31 (2.1)	5.65 (1.11)	1.29 (0.05)	1.54 (0.05)	2.08 (0.05)	2.91 (0.08)	4.12 (0.13)	5.99 (0.35)	17.32 (5.83)	1.1	<3
Females:											
9-13.....	5* (2.6)	2.04 (0.07)	1.08* (0.05)	1.25* (0.06)	1.55 (0.06)	1.95 (0.07)	2.43 (0.09)	2.94* (0.12)	3.30* (0.14)	0.8	<3
14-18.....	12* (5.0)	2.06 (0.24)	0.89* (0.06)	1.03* (0.06)	1.31 (0.06)	1.69 (0.07)	2.20 (0.13)	2.94* (0.36)	3.59* (0.49)	0.9	5* (1.7)
19-50.....	26 (2.3)	3.70 (0.29)	1.08 (0.04)	1.25 (0.05)	1.57 (0.05)	2.07 (0.07)	2.95 (0.17)	4.75 (0.54)	7.99 (2.35)	0.9	<3
51+.....	34 (3.1)	3.83 (0.51)	1.08 (0.03)	1.25 (0.04)	1.59 (0.04)	2.14 (0.08)	3.27 (0.21)	5.16 (0.67)	12.29 (9.66)	0.9	<3
19+.....	30 (2.2)	3.77 (0.31)	1.08 (0.03)	1.25 (0.03)	1.58 (0.04)	2.11 (0.06)	3.12 (0.16)	4.97 (0.38)	9.19 (5.33)	0.9	<3
All:											
1+.....	27 (1.5)	4.28 (0.49)	1.13 (0.02)	1.32 (0.02)	1.70 (0.03)	2.33 (0.04)	3.44 (0.08)	5.13 (0.18)	8.21 (2.11)		<3

NOTES: [†] Percent reporting supplement containing riboflavin. Sample size presented on page 24.
 EAR=Estimated Average Requirement
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 3. Niacin (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)
Over 350% poverty											
Males and females:											
1-3.....	19 (4.6)	14.6 (0.7)	8.0* (0.4)	9.2* (0.4)	11.1 (0.4)	13.8 (0.5)	16.9 (0.7)	20.8* (1.7)	24.3* (2.2)	5	<3
4-8.....	14 (3.8)	20.2 (0.9)	11.9* (0.5)	13.2 (0.6)	15.8 (0.6)	19.1 (0.7)	23.3 (1.2)	28.8 (2.3)	32.8* (2.4)	6	<3
Males:											
9-13.....	12* (4.0)	25.5 (1.1)	14.7* (0.8)	16.6* (0.8)	20.1 (0.9)	24.6 (1.0)	29.9 (1.3)	35.8* (1.7)	39.7* (1.9)	9	<3
14-18.....	14* (3.1)	31.6 (1.2)	18.4* (1.1)	20.6* (1.0)	24.8 (1.0)	30.3 (1.2)	36.7 (1.4)	44.0* (2.1)	49.4* (3.5)	12	<3
19-50.....	27 (3.3)	41.4 (1.7)	19.5 (0.9)	22.4 (0.9)	28.3 (1.1)	36.6 (1.4)	47.2 (2.1)	59.8 (2.8)	69.6 (3.6)	12	<3
51+.....	41 (3.5)	50.5 (5.4)	17.2 (0.8)	20.2 (0.8)	26.6 (1.0)	37.0 (1.5)	49.7 (1.5)	63.0 (2.2)	75.2 (4.2)	12	<3
19+.....	34 (2.3)	45.6 (2.6)	18.3 (0.8)	21.3 (0.8)	27.6 (0.9)	36.8 (1.1)	48.4 (1.3)	61.3 (1.7)	72.2 (2.6)	12	<3
Females:											
9-13.....	13* (3.4)	24.9 (1.9)	13.3* (0.7)	15.1* (0.8)	18.4 (0.8)	22.7 (0.8)	28.0 (1.0)	33.5* (1.3)	37.3* (1.6)	9	<3
14-18.....	25 (5.8)	24.6 (1.1)	12.7* (0.7)	14.4* (0.7)	17.8 (0.8)	22.5 (0.8)	29.2 (1.6)	37.9* (3.8)	44.3* (5.1)	11	<3
19-50.....	30 (2.0)	32.6 (2.9)	13.8 (0.6)	15.8 (0.6)	19.7 (0.6)	25.3 (0.8)	34.4 (1.4)	46.1 (1.9)	57.9 (5.9)	11	<3
51+.....	38 (3.2)	30.4 (2.1)	12.5 (0.5)	14.4 (0.5)	18.2 (0.6)	24.2 (1.1)	35.8 (1.9)	46.0 (1.8)	53.6 (3.5)	11	<3
19+.....	34 (2.0)	31.4 (1.8)	13.0 (0.5)	15.0 (0.5)	18.9 (0.4)	24.8 (0.7)	35.1 (1.2)	46.1 (1.4)	55.3 (3.2)	11	<3
All:											
1+.....	31 (1.4)	36.1 (1.3)	13.6 (0.2)	15.9 (0.3)	20.7 (0.4)	28.7 (0.5)	40.4 (0.7)	53.3 (1.1)	63.8 (1.7)		<3

NOTES: [†] Percent reporting supplement containing niacin. Sample size presented on page 24.
 EAR=Estimated Average Requirement
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 4. Vitamin B6 (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR		>UL	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)	UL	% (SE)
Over 350% poverty													
Males and females:													
1-3.....	47 (5.7)	1.58 (0.07)	0.85* (0.05)	0.96* (0.05)	1.16 (0.05)	1.46 (0.08)	1.88 (0.11)	2.36* (0.14)	2.73* (0.19)	0.4	<3	30	<3
4-8.....	49 (4.5)	2.01 (0.10)	1.06* (0.05)	1.19 (0.05)	1.44 (0.06)	1.83 (0.10)	2.46 (0.17)	3.17 (0.14)	3.51* (0.12)	0.5	<3	40	<3
Males:													
9-13.....	29 (6.0)	2.81 (0.54)	1.05* (0.09)	1.22* (0.09)	1.56 (0.11)	2.07 (0.14)	2.84 (0.29)	3.94* (0.75)	5.82* (9.88)	0.8	<3	60	<3
14-18.....	18 (3.9)	2.85 (0.33)	1.19* (0.11)	1.37* (0.12)	1.75 (0.12)	2.28 (0.15)	3.02 (0.26)	4.54* (0.84)	6.29* (0.51)	1.1	3* (1.7)	80	<3
19-50.....	28 (3.2)	4.79 (0.46)	1.30 (0.08)	1.56 (0.08)	2.13 (0.09)	3.03 (0.13)	4.44 (0.22)	6.72 (0.58)	10.94 (6.32)	1.1	<3	100	<3
51+.....	42 (3.2)	6.99 (1.14)	1.20 (0.07)	1.47 (0.08)	2.08 (0.10)	3.31 (0.22)	5.62 (0.34)	8.84 (1.01)	19.02 (9.69)	1.4	9 (1.5)	100	<3
19+.....	34 (2.1)	5.81 (0.61)	1.25 (0.07)	1.52 (0.07)	2.11 (0.08)	3.13 (0.12)	4.95 (0.20)	7.94 (0.58)	16.17 (6.21)	5	(1.0)	100	<3
Females:													
9-13.....	33 (6.9)	2.30 (0.15)	0.99* (0.06)	1.16* (0.06)	1.49 (0.06)	2.03 (0.11)	2.79 (0.19)	3.64* (0.26)	4.26* (0.35)	0.8	<3	60	<3
14-18.....	31 (5.9)	2.84 (0.46)	0.87* (0.07)	1.01* (0.06)	1.33 (0.07)	1.85 (0.09)	2.90 (0.37)	5.39* (1.14)	7.10* (6.68)	1	9* (2.5)	80	<3
19-50.....	35 (2.5)	5.07 (0.72)	1.05 (0.06)	1.22 (0.06)	1.59 (0.06)	2.19 (0.09)	3.55 (0.19)	5.75 (0.39)	10.95 (7.94)	1.1	6 (1.6)	100	<3
51+.....	40 (3.5)	5.99 (1.28)	0.97 (0.05)	1.13 (0.05)	1.49 (0.06)	2.16 (0.16)	4.30 (0.33)	7.40 (0.81)	21.21 (20.88)	1.3	16 (2.1)	100	<3
19+.....	38 (2.2)	5.55 (0.80)	1.00 (0.04)	1.17 (0.04)	1.53 (0.04)	2.18 (0.09)	3.91 (0.21)	6.62 (0.40)	17.88 (8.21)	12	(1.5)	100	<3
All:													
1+.....	36 (1.4)	5.11 (0.45)	1.06 (0.03)	1.25 (0.03)	1.67 (0.04)	2.47 (0.06)	4.11 (0.12)	6.78 (0.32)	11.43 (3.80)	7	(0.8)		<3

NOTES: [†] Percent reporting supplement containing vitamin B6. Sample size presented on page 24.
 EAR=Estimated Average Requirement; UL=Tolerable Upper Intake Level
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 5. Folate (µg DFE): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)
Over 350% poverty											
Males and females:											
1-3.....	40 (5.2)	408 (16)	202* (11)	233* (11)	292 (13)	380 (15)	491 (20)	618* (31)	718* (45)	120	<3
4-8.....	47 (4.3)	596 (30)	296* (15)	337 (15)	421 (18)	546 (25)	718 (43)	944 (81)	1083* (69)	160	<3
Males:											
9-13.....	25 (5.8)	681 (62)	334* (20)	382* (20)	474 (25)	606 (34)	785 (62)	1016* (90)	1179* (117)	250	<3
14-18.....	20 (4.1)	700 (33)	351* (19)	400* (19)	500 (22)	633 (26)	825 (46)	1115* (88)	1310* (88)	330	3* (1.4)
19-50.....	30 (3.2)	761 (31)	324 (16)	379 (16)	496 (18)	668 (24)	926 (41)	1256 (56)	1469 (67)	320	5 (1.2)
51+.....	41 (3.3)	877 (31)	313 (15)	372 (16)	501 (22)	742 (47)	1159 (40)	1448 (38)	1651 (56)	320	6 (1.2)
19+.....	35 (2.1)	815 (24)	319 (15)	376 (16)	498 (17)	694 (25)	1038 (36)	1366 (34)	1569 (50)	320	5 (1.1)
Females:											
9-13.....	34 (7.0)	635 (21)	320* (11)	366* (13)	459 (17)	600 (20)	768 (32)	952* (47)	1088* (65)	250	<3
14-18.....	31 (5.9)	635 (42)	266* (16)	305* (15)	388 (16)	522 (23)	798 (128)	1167* (117)	1323* (94)	330	14* (2.4)
19-50.....	33 (2.3)	704 (37)	255 (13)	298 (12)	390 (12)	542 (21)	925 (73)	1299 (76)	1686 (238)	320	13 (1.8)
51+.....	39 (3.3)	731 (46)	232 (14)	275 (14)	363 (15)	524 (36)	1026 (65)	1350 (91)	1695 (160)	320	17 (2.3)
19+.....	36 (2.1)	718 (37)	242 (12)	285 (12)	375 (12)	533 (27)	986 (58)	1324 (76)	1691 (183)	320	15 (1.9)
All:											
1+.....	35 (1.4)	740 (17)	270 (10)	319 (10)	425 (11)	605 (16)	958 (26)	1300 (28)	1543 (58)		9 (1.1)

NOTES: [†]Percent reporting supplement containing folate. Sample size presented on page 24.
 DFE=dietary folate equivalents; EAR=Estimated Average Requirement
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 6. Folic acid (µg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							>UL	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	UL	% (SE)
Over 350% poverty											
Males and females:											
1-3.....	40 (5.2)	170 (12)	59* (7)	74* (7)	105 (8)	153 (10)	216 (16)	288* (26)	345* (34)	300	9* (3.1)
4-8.....	47 (4.3)	266 (20)	101* (11)	122 (11)	167 (11)	236 (17)	334 (28)	465 (52)	547* (43)	400	15 (4.6)
Males:											
9-13.....	25 (5.8)	298 (29)	116* (10)	139* (11)	186 (12)	254 (14)	348 (30)	476* (52)	569* (64)	600	4* (2.1)
14-18.....	20 (4.1)	295 (22)	117* (13)	140* (13)	188 (14)	256 (16)	355 (27)	528* (60)	635* (52)	800	<3
19-50.....	30 (3.2)	294 (16)	78 (7)	101 (7)	152 (8)	236 (11)	373 (25)	566 (32)	675 (39)	1000	<3
51+.....	41 (3.3)	362 (17)	73 (7)	96 (8)	152 (10)	272 (25)	524 (20)	669 (23)	778 (50)	1000	<3
19+.....	35 (2.1)	326 (13)	76 (7)	99 (7)	152 (8)	248 (12)	450 (24)	624 (20)	729 (38)	1000	<3
Females:											
9-13.....	34 (7.0)	271 (15)	109* (9)	132* (11)	179 (13)	251 (15)	339 (17)	439* (28)	513* (43)	600	<3
14-18.....	31 (5.9)	277 (36)	83* (7)	102* (8)	141 (9)	209 (17)	358 (89)	575* (94)	658* (82)	800	<3
19-50.....	33 (2.3)	288 (17)	55 (5)	72 (5)	113 (6)	188 (9)	411 (41)	611 (36)	865 (134)	1000	<3
51+.....	39 (3.3)	307 (23)	46 (5)	62 (5)	100 (6)	177 (15)	480 (25)	642 (42)	867 (85)	1000	<3
19+.....	36 (2.1)	298 (16)	50 (4)	67 (5)	106 (4)	182 (10)	457 (29)	628 (31)	866 (100)	1000	<3
All:											
1+.....	35 (1.4)	304 (9)	63 (4)	84 (4)	132 (5)	221 (8)	422 (16)	604 (14)	732 (42)		<3

NOTES: [†] Percent reporting supplement containing folic acid. Sample size presented on page 24.
 UL=Tolerable Upper Intake Level
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 7. Choline (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							>AI		>UL	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	AI	% (SE)	UL	% (SE)
Over 350% poverty													
Males and females:													
1-3.....	22 (4.5)	205 (6)	122* (5)	137* (5)	164 (6)	199 (6)	239 (7)	283* (8)	310* (9)	200	49 (4.1)	1000	<3
4-8.....	25 (4.0)	223 (8)	133* (6)	149 (6)	179 (7)	216 (7)	260 (9)	305 (11)	334* (13)	250	30 (4.5)	1000	<3
Males:													
9-13.....	9* (3.1)	281 (13)	161* (11)	182* (11)	220 (12)	271 (13)	331 (14)	392* (16)	436* (21)	375	13* (3.6)	2000	<3
14-18.....	8* (2.1)	319 (14)	185* (11)	208* (12)	253 (12)	310 (14)	374 (17)	439* (19)	483* (22)	550	<3	3000	<3
19-50.....	8 (1.6)	400 (10)	214 (12)	246 (12)	308 (11)	387 (11)	477 (12)	569 (15)	631 (18)	550	12 (1.8)	3500	<3
51+.....	5 (1.4)	409 (8)	219 (9)	251 (8)	315 (6)	395 (8)	489 (11)	585 (15)	646 (18)	550	14 (1.8)	3500	<3
19+.....	6 (1.0)	404 (9)	216 (10)	249 (10)	311 (8)	391 (8)	483 (11)	576 (14)	638 (17)	550	13 (1.6)	3500	<3
Females:													
9-13.....	17 (5.6)	262 (10)	146* (8)	166* (8)	204 (9)	253 (10)	310 (12)	368* (18)	408* (22)	375	9* (2.8)	2000	<3
14-18.....	17 (4.3)	241 (13)	133* (9)	151* (10)	186 (11)	231 (13)	285 (14)	342* (16)	382* (20)	400	4* (1.4)	3000	<3
19-50.....	10 (1.5)	298 (8)	164 (5)	188 (5)	231 (5)	285 (7)	348 (8)	414 (10)	455 (11)	425	8 (1.4)	3500	<3
51+.....	8 (1.6)	294 (9)	165 (5)	187 (5)	229 (6)	284 (8)	346 (10)	412 (15)	455 (18)	425	8 (1.9)	3500	<3
19+.....	9 (1.1)	296 (7)	164 (4)	187 (5)	230 (5)	284 (7)	347 (9)	413 (11)	455 (13)	425	8 (1.5)	3500	<3
All:													
1+.....	9 (0.9)	334 (6)	166 (3)	192 (4)	243 (4)	314 (6)	404 (9)	503 (12)	568 (14)		12 (1.3)		<3

NOTES: [†]Percent reporting supplement containing choline. Sample size presented on page 24.
 AI=Adequate Intake; UL=Tolerable Upper Intake Level
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 8. Vitamin B12 (µg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)
Over 350% poverty											
Males and females:											
1-3.....	49 (5.3)	21.61* (16.24)	2.35* (0.15)	2.69* (0.15)	3.41 (0.17)	4.47 (0.23)	6.07 (0.35)	8.54* (1.06)	11.27* (475.54)	0.7	<3
4-8.....	49 (4.5)	6.74 (1.01)	2.60* (0.17)	2.98 (0.18)	3.78 (0.21)	5.04 (0.33)	7.23 (0.67)	9.54 (0.44)	10.71* (0.71)	1	<3
Males:											
9-13.....	29 (5.0)	43.72* (27.86)	2.24* (0.23)	2.72* (0.25)	3.72 (0.33)	5.26 (0.45)	7.59 (0.76)	11.17* (26.13)	69.80* (468.00)	1.5	<3
14-18.....	18 (3.9)	10.93 (2.70)	2.54* (0.25)	3.06* (0.27)	4.18 (0.31)	5.79 (0.41)	8.26 (0.89)	15.43* (3.49)	21.62* (24.30)	2	<3
19-50.....	32 (3.7)	49.08 (13.16)	2.61 (0.18)	3.24 (0.19)	4.76 (0.26)	7.37 (0.53)	12.85 (1.61)	32.88 (17.18)	117.09 (104.05)	2	<3
51+.....	45 (3.1)	103.03 (21.46)	2.50 (0.19)	3.21 (0.21)	4.91 (0.29)	8.85 (0.76)	29.82 (2.03)	107.41 (98.75)	1002.45 (359.49)	2	<3
19+.....	38 (2.2)	74.21 (9.85)	2.56 (0.18)	3.22 (0.19)	4.82 (0.22)	7.88 (0.47)	19.49 (2.85)	85.10 (24.44)	387.50 (180.90)	2	<3
Females:											
9-13.....	33 (6.9)	5.76 (0.51)	1.90* (0.13)	2.34* (0.14)	3.25 (0.19)	4.80 (0.34)	7.05 (0.67)	9.76* (0.97)	12.03* (1.51)	1.5	<3
14-18.....	31 (5.9)	23.32* (9.91)	1.47* (0.16)	1.83* (0.17)	2.64 (0.19)	4.07 (0.27)	7.65 (1.70)	16.86* (19.04)	54.54* (29.63)	2	13* (3.0)
19-50.....	37 (2.9)	60.85 (16.06)	2.01 (0.13)	2.43 (0.13)	3.42 (0.16)	5.29 (0.34)	10.66 (1.34)	49.00 (29.18)	254.85 (119.78)	2	5 (1.2)
51+.....	46 (3.9)	240.41 (58.96)	2.02 (0.12)	2.49 (0.13)	3.57 (0.19)	6.29 (1.10)	27.21 (5.42)	671.97 (410.96)	1053.63 (931.14)	2	5 (1.1)
19+.....	42 (2.5)	155.23 (37.56)	2.01 (0.11)	2.46 (0.12)	3.49 (0.14)	5.69 (0.40)	15.33 (2.44)	203.60 (175.51)	1003.07 (241.96)	2	5 (1.0)
All:											
1+.....	39 (1.5)	96.53 (15.66)	2.19 (0.09)	2.71 (0.09)	3.91 (0.11)	6.35 (0.23)	13.56 (1.18)	63.30 (20.69)	503.88 (120.41)	3	(0.6)

NOTES: [†] Percent reporting supplement containing vitamin B12. Sample size presented on page 24.
 EAR=Estimated Average Requirement
 It is advised that individuals over 50 meet their B12 requirement with fortified foods or supplements
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 9. Vitamin C (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR		>UL	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)	UL	% (SE)
Over 350% poverty													
Males and females:													
1-3.....	51 (5.9)	84.0 (8.2)	29.2* (3.9)	37.1* (4.0)	53.3 (5.6)	76.5 (7.5)	106.4 (10.7)	140.4* (13.3)	164.5* (16.0)	13	<3	400	<3
4-8.....	55 (4.3)	96.5 (6.5)	32.0* (2.6)	40.4 (2.8)	58.7 (3.7)	85.1 (4.8)	117.6 (7.2)	156.9 (11.3)	189.5* (16.5)	22	<3	650	<3
Males:													
9-13.....	30 (6.5)	98.2 (11.1)	32.6* (5.5)	41.3* (6.4)	59.7 (8.7)	87.2 (9.9)	122.5 (12.3)	165.6* (16.7)	196.2* (22.0)	39	8* (4.5)	1200	<3
14-18.....	21 (4.2)	117.7 (10.1)	27.1* (3.8)	34.2* (4.1)	50.6 (5.1)	75.3 (7.1)	111.9 (10.7)	178.5* (43.5)	363.8* (148.6)	63	38 (6.8)	1800	<3
19-50.....	35 (3.8)	163.7 (17.8)	25.8 (2.1)	35.0 (2.4)	57.6 (3.5)	97.7 (5.8)	166.1 (10.1)	290.4 (49.6)	542.1 (97.6)	75	37 (2.8)	2000	<3
51+.....	46 (3.3)	224.8 (22.9)	28.4 (2.3)	38.8 (3.0)	66.4 (5.1)	118.7 (8.0)	202.4 (14.0)	543.8 (116.3)	848.7 (191.2)	75	30 (2.9)	2000	<3
19+.....	40 (2.6)	192.2 (11.4)	26.8 (2.0)	36.6 (2.3)	61.2 (3.4)	106.4 (5.2)	182.6 (8.6)	377.6 (52.7)	665.2 (55.6)	75	33 (2.4)	2000	<3
Females:													
9-13.....	36 (7.4)	108.4 (13.1)	33.7* (4.6)	42.4* (4.8)	60.6 (5.8)	89.3 (8.0)	126.7 (10.2)	168.7* (16.9)	206.6* (24.1)	39	8* (3.1)	1200	<3
14-18.....	31 (5.9)	103.6 (13.5)	24.6* (3.3)	31.6* (4.0)	47.4 (5.4)	73.9 (7.3)	114.7 (11.9)	167.9* (19.7)	226.5* (200.7)	56	33 (6.3)	1800	<3
19-50.....	38 (2.2)	173.6 (31.3)	28.0 (2.6)	36.7 (2.9)	56.8 (3.5)	91.7 (4.5)	146.0 (7.1)	250.8 (27.1)	559.8 (152.3)	60	28 (2.8)	2000	<3
51+.....	47 (3.1)	198.0 (14.5)	29.0 (2.7)	38.4 (3.2)	61.9 (4.6)	107.1 (7.5)	195.4 (14.7)	555.0 (64.8)	761.1 (109.0)	60	24 (2.9)	2000	<3
19+.....	43 (1.8)	186.5 (18.4)	28.5 (2.5)	37.6 (2.9)	59.1 (3.7)	98.7 (4.9)	168.5 (8.8)	368.5 (111.3)	667.4 (52.6)	60	26 (2.5)	2000	<3
All:													
1+.....	41 (1.6)	174.0 (8.9)	28.0 (1.3)	37.1 (1.5)	58.9 (2.1)	97.2 (3.2)	162.5 (5.4)	309.7 (29.3)	620.5 (30.0)	27	(1.6)		<3

NOTES: [†] Percent reporting supplement containing vitamin C. Sample size presented on page 24.
 EAR=Estimated Average Requirement; UL=Tolerable Upper Intake Level
 EAR used is for non-smokers regardless of smoking status
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 10. Vitamin C (mg) Adult Smokers: Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

Percent reporting [†] % (SE)	Mean (SE)	Percentiles								<EAR		>UL		
		5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)	UL	% (SE)		
Over 350% poverty														
Males:														
19-50..... 25 (7.2)	--	--	--	--	--	--	--	--	--	--	110	--	2000	--
51+..... 36* (7.8)	--	--	--	--	--	--	--	--	--	--	110	--	2000	--
19+..... 29 (5.8)	164.5 (32.2)	18.3* (2.4)	25.0* (3.1)	41.5 (4.6)	73.4 (8.6)	143.6 (32.0)	482.3* (215.9)	728.9* (186.4)		110	67 (5.5)	2000	<3	
Females:														
19-50..... 26* (5.8)	--	--	--	--	--	--	--	--	--	--	95	--	2000	--
51+..... 18* (8.8)	--	--	--	--	--	--	--	--	--	--	95	--	2000	--
19+..... 22* (6.1)	102.9* (32.6)	16.6* (2.4)	21.3* (2.9)	33.7 (3.6)	55.0 (6.1)	86.7 (10.6)	134.8* (17.7)	185.6* (237.5)		95	79* (4.8)	2000	<3	

NOTES: [†] Percent reporting supplement containing vitamin C. Sample size presented on page 24.
 Smoking status determined by self-reported cigarette use. Individuals without smoking status data were excluded.
 EAR=Estimated Average Requirement for smokers; UL=Tolerable Upper Intake Level
 * Estimate may be less reliable due to small sample size and/or large relative standard error
 -- Estimate not displayed due to small sample size and for consistency of presentation

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 11. Vitamin C (mg) Adult Non-Smokers: Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR		>UL	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)	UL	% (SE)
Over 350% poverty													
Males:													
19-50.....	37 (4.1)	165.1 (17.8)	27.8 (2.1)	37.5 (2.5)	61.0 (3.7)	101.8 (6.4)	169.2 (11.5)	281.3 (42.5)	516.3 (111.7)	75	35 (3.1)	2000	<3
51+.....	47 (3.6)	228.7 (23.8)	30.3 (2.6)	41.3 (3.4)	69.3 (5.2)	122.9 (8.5)	206.2 (14.4)	530.1 (120.6)	865.0 (212.1)	75	28 (2.9)	2000	<3
19+.....	42 (2.8)	195.5 (12.0)	29.0 (2.2)	39.1 (2.6)	64.4 (3.6)	110.7 (5.7)	186.6 (9.4)	361.6 (53.7)	647.1 (62.1)	75	31 (2.4)	2000	<3
Females:													
19-50.....	39 (2.5)	181.6 (35.6)	30.8 (2.9)	39.7 (3.0)	60.4 (3.1)	95.3 (5.0)	149.5 (9.6)	259.7 (38.6)	582.7 (167.6)	60	25 (2.4)	2000	<3
51+.....	49 (3.1)	203.2 (16.1)	32.9 (3.6)	43.0 (4.2)	67.3 (5.6)	113.1 (8.9)	200.6 (19.7)	572.6 (62.7)	785.8 (111.3)	60	20 (3.2)	2000	<3
19+.....	45 (1.9)	193.0 (20.3)	31.8 (2.9)	41.3 (3.1)	63.6 (3.3)	103.4 (5.0)	173.4 (10.6)	404.1 (128.3)	678.3 (62.7)	60	22 (2.3)	2000	<3

NOTES: [†]Percent reporting supplement containing vitamin C. Sample size presented on page 24.
Smoking status determined by self-reported cigarette use. Individuals without smoking status data were excluded.
EAR=Estimated Average Requirement for non-smokers; UL=Tolerable Upper Intake Level

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 12. Vitamin C (mg) Adult Smokers/Non-Smokers: Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR		>UL	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)	UL	% (SE)
Over 350% poverty													
Males:													
19-50.....	35 (3.8)	163.1 (17.6)	25.6 (2.3)	34.8 (2.6)	57.3 (3.3)	97.4 (5.6)	165.5 (10.3)	289.5 (49.1)	539.7 (97.9)	110/75	39 (2.8)	2000	<3
51+.....	46 (3.3)	225.0 (22.7)	28.6 (2.5)	39.3 (3.1)	66.6 (5.0)	119.2 (8.3)	203.9 (13.6)	550.1 (121.8)	849.4 (193.1)	110/75	31 (2.9)	2000	<3
19+.....	40 (2.6)	191.9 (11.5)	26.8 (2.2)	36.6 (2.5)	61.0 (3.4)	106.3 (5.1)	183.2 (9.0)	373.4 (54.9)	664.8 (51.9)	110/75	35 (2.3)	2000	<3
Females:													
19-50.....	38 (2.2)	172.8 (32.2)	28.5 (3.2)	37.2 (3.5)	57.5 (3.6)	91.6 (5.0)	144.5 (9.2)	245.7 (27.8)	564.5 (174.3)	95/60	29 (3.2)	2000	<3
51+.....	47 (3.1)	197.1 (16.6)	29.2 (3.9)	38.8 (4.5)	62.2 (5.9)	106.6 (9.3)	191.5 (20.3)	556.8 (68.6)	759.0 (112.0)	95/60	25 (3.7)	2000	<3
19+.....	43 (1.8)	185.6 (19.0)	28.9 (2.9)	37.9 (3.2)	59.8 (3.6)	98.4 (5.1)	165.8 (10.3)	353.4 (133.3)	670.3 (52.4)	95/60	27 (2.6)	2000	<3

NOTES: [†]Percent reporting supplement containing vitamin C. Sample size presented on page 24.
Smoking status determined by self-reported cigarette use. Individuals without smoking status data were excluded.
EAR=Estimated Average Requirement for smokers and non-smokers; UL=Tolerable Upper Intake Level

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 13. Vitamin D (µg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)		Percentiles						<EAR		>UL		
		Mean (SE)	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)	UL	% (SE)
Over 350% poverty														
Males and females:														
1-3.....	50 (6.2)	9.3 (0.8)		3.0* (0.3)	3.8* (0.3)	5.3 (0.4)	7.8 (0.7)	11.8 (1.4)	17.2* (2.0)	20.6* (1.8)	10	66 (6.1)	63	<3
4-8.....	53 (4.4)	12.2 (2.5)		2.6* (0.3)	3.2 (0.3)	4.8 (0.4)	7.6 (0.8)	14.2 (1.5)	19.7 (1.4)	23.5* (6.7)	10	62 (5.1)	75	<3
Males:														
9-13.....	26 (5.8)	8.5 (1.0)		1.8* (0.1)	2.4* (0.2)	3.7 (0.2)	5.8 (0.5)	10.0 (1.9)	18.8* (2.8)	22.8* (2.5)	10	75 (5.3)	100	<3
14-18.....	20 (4.2)	12.6* (4.3)		1.6* (0.2)	2.1* (0.3)	3.3 (0.3)	5.2 (0.5)	8.7 (1.2)	19.8* (4.0)	36.3* (10.3)	10	79 (4.4)	100	<3
19-50.....	34 (3.4)	14.9 (1.7)		1.5 (0.1)	2.0 (0.1)	3.2 (0.2)	5.6 (0.4)	13.7 (2.1)	30.8 (12.4)	56.0 (9.2)	10	69 (3.2)	100	<3
51+.....	51 (3.2)	33.5 (5.7)		2.0 (0.2)	2.6 (0.2)	4.5 (0.4)	11.9 (2.3)	31.2 (2.3)	67.6 (9.7)	131.4 (28.7)	10	47 (3.7)	100	7 (1.8)
19+.....	42 (2.3)	23.6 (2.6)		1.7 (0.1)	2.2 (0.2)	3.7 (0.2)	7.0 (0.6)	23.2 (2.4)	54.7 (1.8)	85.4 (21.8)	10	59 (2.2)	100	4 (0.9)
Females:														
9-13.....	35 (7.1)	8.8 (1.4)		1.6* (0.2)	2.1* (0.2)	3.4 (0.3)	5.9 (0.7)	10.9 (1.6)	17.2* (2.3)	21.9* (5.5)	10	72 (5.5)	100	<3
14-18.....	34 (5.5)	13.3 (3.7)		1.0* (0.1)	1.4* (0.1)	2.4 (0.2)	4.5 (0.5)	10.9 (4.9)	30.6* (31.3)	83.0* (48.0)	10	73 (6.3)	100	<3
19-50.....	42 (2.7)	22.9 (2.7)		1.4 (0.1)	1.9 (0.2)	3.0 (0.2)	5.7 (0.4)	25.1 (2.3)	54.7 (14.9)	106.2 (23.6)	10	62 (2.6)	100	6 (1.8)
51+.....	59 (3.1)	35.3 (3.8)		1.6 (0.2)	2.2 (0.2)	3.9 (0.4)	18.7 (4.5)	50.1 (6.3)	82.4 (18.0)	126.3 (13.1)	10	44 (3.6)	100	9 (2.0)
19+.....	51 (2.1)	29.4 (2.8)		1.5 (0.1)	2.0 (0.2)	3.4 (0.2)	8.5 (2.0)	33.2 (2.5)	73.7 (11.6)	118.1 (12.8)	10	52 (2.6)	100	7 (1.7)
All:														
1+.....	45 (1.6)	23.7 (1.8)		1.6 (0.1)	2.1 (0.1)	3.6 (0.2)	7.1 (0.4)	25.5 (1.7)	55.2 (3.0)	95.8 (16.8)	10	58 (1.5)		5 (1.0)

NOTES: [†] Percent reporting supplement containing vitamin D. Sample size presented on page 24.
 EAR=Estimated Average Requirement; UL=Tolerable Upper Intake Level
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 14. Vitamin K (µg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							>AI	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	AI	% (SE)
Over 350% poverty											
Males:											
19-50.....	18 (2.7)	137.3 (7.4)	52.0 (3.5)	63.1 (3.9)	87.3 (5.0)	123.7 (6.6)	171.3 (9.2)	228.9 (12.8)	270.0 (15.3)	120	52 (4.2)
51+.....	32 (2.8)	152.0 (8.4)	58.0 (4.2)	70.7 (4.6)	98.2 (5.9)	137.7 (7.8)	191.0 (10.7)	250.3 (14.3)	292.0 (16.8)	120	61 (4.5)
19+.....	24 (1.9)	144.1 (7.3)	54.3 (3.3)	66.1 (3.7)	92.1 (5.0)	130.3 (6.4)	180.9 (9.3)	239.7 (12.7)	281.6 (15.2)	120	56 (3.9)
Females:											
19-50.....	18 (1.6)	149.8 (9.2)	54.1 (5.1)	66.4 (5.5)	92.7 (6.8)	132.9 (9.0)	188.5 (11.4)	253.8 (14.3)	301.7 (16.4)	90	77 (4.6)
51+.....	28 (2.8)	162.6 (20.7)	53.7 (5.3)	65.9 (6.0)	91.8 (6.9)	131.2 (9.1)	184.1 (11.2)	247.6 (15.2)	294.2 (18.4)	90	76 (4.6)
19+.....	23 (1.7)	156.6 (13.5)	53.8 (4.8)	66.1 (5.5)	92.3 (6.6)	131.9 (8.6)	186.2 (10.8)	250.7 (13.8)	297.3 (15.8)	90	76 (4.5)

NOTES: [†]Percent reporting supplement containing vitamin K. Sample size presented on page 24.
 Estimates for children unavailable due to very low reporting of supplements containing vitamin K
 AI=Adequate Intake

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 15. Calcium (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR		>UL	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)	UL	% (SE)
Over 350% poverty													
Males and females:													
1-3.....	10* (4.3)	951 (25)	573* (26)	646* (24)	772 (24)	932 (25)	1104 (29)	1286* (37)	1399* (44)	500	<3	2500	<3
4-8.....	13 (2.6)	994 (29)	598* (30)	669 (29)	803 (27)	971 (29)	1159 (34)	1349 (41)	1467* (46)	800	25 (3.6)	2500	<3
Males:													
9-13.....	12* (3.8)	1116 (47)	614* (44)	701* (43)	862 (44)	1075 (46)	1321 (49)	1581* (69)	1769* (82)	1100	53 (5.5)	3000	<3
14-18.....	15 (5.5)	1161 (56)	642* (46)	731* (50)	907 (50)	1129 (57)	1379 (64)	1623* (73)	1794* (83)	1100	47 (6.3)	3000	<3
19-50.....	39 (3.2)	1233 (34)	615 (30)	715 (28)	913 (27)	1177 (30)	1477 (36)	1814 (52)	2042 (64)	800	16 (2.1)	2500	<3
51+.....	50 (3.1)	1204 (36)	573 (24)	673 (24)	873 (26)	1140 (33)	1466 (46)	1820 (65)	2055 (76)		23 (2.3)	2000	6 (1.3)
19+.....	44 (2.5)	1219 (32)	594 (25)	695 (24)	894 (24)	1160 (28)	1472 (38)	1816 (53)	2047 (63)		19 (1.9)		4 (0.8)
Females:													
9-13.....	12* (4.7)	1045 (34)	555* (34)	642* (31)	799 (33)	1004 (35)	1240 (45)	1494* (63)	1667* (77)	1100	61 (4.2)	3000	<3
14-18.....	23 (5.0)	918 (45)	466* (43)	541* (43)	683 (42)	874 (44)	1107 (55)	1358* (70)	1531* (88)	1100	74 (4.3)	3000	<3
19-50.....	44 (2.6)	1070 (37)	524 (24)	603 (25)	761 (27)	980 (31)	1263 (39)	1590 (53)	1835 (94)	800	30 (3.2)	2500	<3
51+.....	57 (3.4)	1123 (41)	498 (24)	581 (25)	753 (29)	1015 (42)	1400 (64)	1831 (85)	2113 (89)	1000	49 (3.7)	2000	7 (1.3)
19+.....	51 (2.0)	1098 (28)	509 (21)	591 (21)	757 (23)	997 (29)	1327 (40)	1717 (57)	2007 (69)		40 (2.7)		4 (0.8)
All:													
1+.....	42 (1.4)	1137 (17)	548 (16)	637 (14)	814 (15)	1062 (17)	1372 (23)	1722 (33)	1975 (44)		31 (1.6)		3 (0.5)

NOTES: [†] Percent reporting supplement containing calcium. Sample size presented on page 24.
 EAR=Estimated Average Requirement; UL=Tolerable Upper Intake Level
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 16. Phosphorus (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR		>UL	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)	UL	% (SE)
Over 350% poverty													
Males:													
19-50.....	11 (2.2)	1713 (39)	1005 (40)	1132 (39)	1378 (38)	1678 (39)	2006 (44)	2332 (51)	2545 (59)	580	<3	4000	<3
51+.....	23 (2.5)	1626 (37)	931 (35)	1056 (33)	1296 (33)	1589 (36)	1914 (43)	2242 (50)	2450 (54)	580	<3		<3
19+.....	17 (1.8)	1672 (33)	967 (36)	1095 (36)	1339 (32)	1636 (33)	1966 (38)	2292 (46)	2506 (50)	580	<3		<3
Females:													
19-50.....	7 (1.2)	1281 (27)	777 (25)	875 (26)	1049 (25)	1257 (28)	1490 (30)	1716 (32)	1857 (39)	580	<3	4000	<3
51+.....	20 (2.3)	1224 (30)	734 (20)	827 (23)	994 (26)	1202 (29)	1428 (36)	1652 (44)	1792 (52)	580	<3		<3
19+.....	14 (1.2)	1251 (24)	753 (21)	848 (21)	1020 (21)	1228 (24)	1458 (28)	1682 (35)	1824 (41)	580	<3		<3

NOTES: [†] Percent reporting supplement containing phosphorus. Sample size presented on page 24.
 Estimates for children unavailable due to very low reporting of supplements containing phosphorus
 EAR=Estimated Average Requirement; UL=Tolerable Upper Intake Level

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 17. Magnesium (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)
Over 350% poverty											
Males:											
19-50.....	25 (2.8)	389 (12)	204 (9)	234 (9)	293 (9)	372 (11)	463 (15)	567 (22)	635 (28)	350	41 (3.2)
51+.....	41 (2.6)	419 (17)	207 (9)	238 (9)	302 (11)	386 (14)	493 (19)	617 (28)	738 (57)		39 (3.8)
19+.....	33 (2.2)	403 (12)	206 (8)	236 (8)	297 (8)	378 (9)	476 (13)	589 (21)	676 (34)		40 (2.5)
Females:											
19-50.....	21 (1.9)	326 (10)	171 (6)	195 (6)	239 (6)	299 (7)	373 (9)	462 (16)	551 (43)	265	35 (2.9)
51+.....	34 (3.5)	341 (18)	169 (7)	193 (7)	240 (9)	303 (12)	387 (17)	511 (42)	690 (108)		35 (4.0)
19+.....	28 (2.4)	334 (13)	170 (6)	194 (6)	240 (7)	301 (9)	380 (12)	486 (25)	610 (64)		35 (3.3)

NOTES: [†] Percent reporting supplement containing magnesium. Sample size presented on page 24.
 Estimates for children unavailable due to very low reporting of supplements containing magnesium
 EAR=Estimated Average Requirement

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 18. Iron (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR		>UL	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)	UL	% (SE)
Over 350% poverty													
Males and females:													
1-3.....	11 (2.9)	10.2 (0.5)	5.7* (0.2)	6.4* (0.2)	7.7 (0.3)	9.5 (0.4)	11.7 (0.6)	14.5* (1.1)	17.1* (2.3)	3	<3	40	<3
4-8.....	6* (1.4)	13.5 (0.3)	7.9* (0.3)	8.8 (0.3)	10.5 (0.2)	12.7 (0.2)	15.4 (0.3)	18.5 (0.7)	21.7* (2.5)	4.1	<3	40	<3
Males:													
9-13.....	4* (2.0)	16.5 (0.6)	9.2* (0.4)	10.4* (0.4)	12.7 (0.5)	15.7 (0.5)	19.4 (0.7)	23.7* (1.0)	26.9* (1.4)	5.9	<3	40	<3
14-18.....	4* (1.5)	17.6 (0.8)	9.9* (0.5)	11.1* (0.5)	13.7 (0.6)	16.9 (0.7)	20.7 (0.9)	24.9* (1.2)	28.0* (1.5)	7.7	<3	45	<3
19-50.....	10 (2.3)	18.1 (0.6)	9.1 (0.4)	10.5 (0.4)	13.2 (0.4)	16.7 (0.5)	21.2 (0.6)	27.0 (1.2)	32.2 (1.8)	6	<3	45	<3
51+.....	11 (1.9)	18.7 (0.9)	9.0 (0.4)	10.4 (0.4)	13.0 (0.4)	16.5 (0.4)	21.2 (0.7)	27.8 (1.6)	34.0 (2.6)	6	<3	45	<3
19+.....	11 (1.4)	18.4 (0.6)	9.1 (0.4)	10.4 (0.4)	13.1 (0.4)	16.6 (0.4)	21.2 (0.5)	27.4 (0.9)	33.0 (1.4)	6	<3	45	<3
Females:													
9-13.....	<3	15.6 (0.5)	8.5* (0.4)	9.7* (0.4)	12.0 (0.4)	14.8 (0.4)	18.3 (0.6)	22.2* (0.9)	25.1* (1.2)	5.7	<3	40	<3
14-18.....	10* (3.2)	15.1 (1.0)	7.2* (0.4)	8.2* (0.4)	10.1 (0.4)	12.9 (0.4)	16.5 (0.5)	21.8* (2.0)	29.4* (7.5)	7.9	14* (1.9)	45	<3
19-50.....	26 (2.4)	17.9 (0.9)	7.3 (0.3)	8.4 (0.3)	10.5 (0.3)	13.7 (0.4)	19.4 (1.2)	31.6 (1.6)	40.1 (4.9)	8.1	14 (1.4)	45	4 (1.2)
51+.....	21 (2.2)	15.0 (0.5)	6.9 (0.3)	8.0 (0.3)	9.9 (0.3)	12.7 (0.4)	16.8 (0.7)	26.2 (2.0)	31.4 (1.1)	5	<3	45	<3
19+.....	23 (1.6)	16.4 (0.4)	7.1 (0.3)	8.2 (0.3)	10.2 (0.2)	13.2 (0.3)	17.9 (0.7)	29.1 (1.0)	34.6 (1.4)	8	(0.9)	45	<3
All:													
1+.....	15 (0.9)	16.9 (0.3)	7.7 (0.2)	8.9 (0.2)	11.2 (0.2)	14.7 (0.2)	19.5 (0.3)	27.0 (0.6)	32.7 (0.7)	4	(0.4)		<3

NOTES: [†] Percent reporting supplement containing iron. Sample size presented on page 24.
 EAR=Estimated Average Requirement; UL=Tolerable Upper Intake Level
 EAR percentages determined by probability method for groups
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 19. Zinc (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR		>UL	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)	UL	% (SE)
Over 350% poverty													
Males and females:													
1-3.....	37 (5.9)	7.9 (0.2)	4.8* (0.2)	5.4* (0.2)	6.3 (0.2)	7.6 (0.2)	9.1 (0.3)	10.7* (0.4)	11.9* (0.8)	2.5	<3	7	62 (3.5)
4-8.....	44 (4.6)	10.0 (0.3)	6.1* (0.2)	6.7 (0.2)	7.9 (0.2)	9.5 (0.2)	11.3 (0.2)	13.5 (0.5)	15.8* (1.5)	4	<3	12	18 (2.1)
Males:													
9-13.....	22 (5.1)	11.9 (0.6)	6.3* (0.3)	7.2* (0.3)	8.9 (0.4)	11.1 (0.4)	13.8 (0.6)	17.0* (0.8)	19.3* (1.4)	7	9* (2.0)	23	<3
14-18.....	9* (2.4)	13.1 (0.6)	7.1* (0.4)	8.0* (0.4)	9.9 (0.4)	12.4 (0.5)	15.5 (0.7)	18.9* (1.1)	21.7* (1.5)	8.5	13* (2.7)	34	<3
19-50.....	26 (3.1)	16.4 (0.6)	8.2 (0.4)	9.3 (0.4)	11.6 (0.4)	14.8 (0.5)	19.2 (0.7)	25.4 (1.3)	29.9 (1.7)	9.4	10 (2.1)	40	<3
51+.....	39 (2.9)	20.4 (0.8)	8.1 (0.3)	9.2 (0.3)	11.8 (0.3)	16.1 (0.6)	24.4 (1.0)	33.3 (1.7)	41.2 (3.6)	9.4	11 (1.6)	40	5 (1.1)
19+.....	32 (2.1)	18.3 (0.5)	8.1 (0.3)	9.3 (0.3)	11.7 (0.3)	15.2 (0.4)	21.3 (0.7)	29.1 (1.0)	35.7 (1.8)	9.4	11 (1.7)	40	3 (0.7)
Females:													
9-13.....	28 (6.4)	11.0 (0.4)	5.7* (0.2)	6.6* (0.2)	8.2 (0.3)	10.4 (0.4)	13.1 (0.5)	15.9* (0.7)	18.2* (1.1)	7	13* (1.9)	23	<3
14-18.....	22 (5.1)	10.5 (0.7)	4.8* (0.3)	5.6* (0.3)	7.0 (0.3)	9.1 (0.4)	12.0 (0.7)	15.9* (1.5)	19.5* (3.3)	7.3	28 (4.2)	34	<3
19-50.....	28 (2.2)	13.2 (0.5)	5.8 (0.2)	6.7 (0.2)	8.3 (0.2)	10.6 (0.3)	14.7 (0.6)	23.4 (0.8)	26.8 (0.7)	6.8	11 (1.3)	40	<3
51+.....	36 (3.0)	14.7 (0.6)	5.7 (0.2)	6.5 (0.2)	8.2 (0.2)	10.8 (0.4)	18.7 (1.6)	26.7 (1.1)	32.9 (1.5)	6.8	12 (1.7)	40	<3
19+.....	32 (1.7)	14.0 (0.4)	5.7 (0.2)	6.6 (0.2)	8.2 (0.1)	10.7 (0.2)	16.5 (1.0)	25.1 (0.6)	29.6 (0.9)	6.8	12 (1.2)	40	<3
All:													
1+.....	31 (1.3)	15.2 (0.3)	6.2 (0.1)	7.1 (0.1)	9.1 (0.1)	12.3 (0.2)	18.1 (0.4)	25.9 (0.6)	31.3 (0.8)	11 (0.9)			5 (0.4)

NOTES: [†] Percent reporting supplement containing zinc. Sample size presented on page 24.
 EAR=Estimated Average Requirement; UL=Tolerable Upper Intake Level
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 20. Copper (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR		>UL	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)	UL	% (SE)
Over 350% poverty													
Males:													
19-50.....	19 (2.9)	1.7 (0.06)	0.8 (0.03)	0.9 (0.03)	1.1 (0.03)	1.5 (0.04)	1.9 (0.07)	2.6 (0.14)	3.2 (0.23)	0.7	<3	10	<3
51+.....	36 (2.8)	1.8 (0.05)	0.8 (0.03)	0.9 (0.03)	1.2 (0.04)	1.6 (0.05)	2.2 (0.08)	3.2 (0.15)	3.7 (0.13)	0.7	<3	10	<3
19+.....	27 (1.8)	1.7 (0.04)	0.8 (0.03)	0.9 (0.03)	1.2 (0.03)	1.5 (0.04)	2.1 (0.05)	2.9 (0.12)	3.5 (0.12)	0.7	<3	10	<3
Females:													
19-50.....	18 (1.6)	1.5 (0.05)	0.7 (0.02)	0.8 (0.02)	1.0 (0.02)	1.2 (0.03)	1.7 (0.06)	2.6 (0.22)	3.3 (0.13)	0.7	6 (0.8)	10	<3
51+.....	32 (3.0)	1.5 (0.06)	0.7 (0.03)	0.8 (0.03)	1.0 (0.03)	1.3 (0.05)	1.8 (0.08)	2.8 (0.17)	3.3 (0.10)	0.7	6 (1.3)	10	<3
19+.....	25 (1.8)	1.5 (0.04)	0.7 (0.02)	0.8 (0.02)	1.0 (0.02)	1.3 (0.03)	1.8 (0.06)	2.7 (0.13)	3.3 (0.07)	0.7	6 (1.0)	10	<3

NOTES: [†]Percent reporting supplement containing copper. Sample size presented on page 24.
 Estimates for children unavailable due to very low reporting of supplements containing copper
 EAR=Estimated Average Requirement; UL=Tolerable Upper Intake Level

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 21. Selenium (µg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							<EAR		>UL	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR	% (SE)	UL	% (SE)
Over 350% poverty													
Males:													
19-50.....	20 (2.9)	157 (6)	88 (4)	99 (4)	121 (5)	149 (6)	183 (8)	224 (10)	255 (12)	45	<3	400	<3
51+.....	37 (2.8)	160 (5)	83 (3)	95 (3)	117 (3)	147 (3)	188 (5)	242 (13)	285 (23)	45	<3	400	<3
19+.....	28 (1.9)	158 (5)	86 (3)	97 (3)	119 (3)	148 (4)	185 (5)	231 (11)	268 (14)	45	<3	400	<3
Females:													
19-50.....	21 (2.3)	114 (3)	63 (2)	71 (2)	86 (2)	105 (3)	129 (4)	158 (7)	184 (10)	45	<3	400	<3
51+.....	29 (2.9)	107 (4)	59 (2)	67 (2)	81 (2)	101 (3)	124 (4)	151 (6)	172 (9)	45	<3	400	<3
19+.....	25 (2.0)	111 (2)	61 (2)	69 (2)	84 (2)	103 (2)	126 (3)	155 (5)	178 (8)	45	<3	400	<3

NOTES: [†] Percent reporting supplement containing selenium. Sample size presented on page 24.
 Estimates for children unavailable due to very low reporting of supplements containing selenium
 EAR=Estimated Average Requirement; UL=Tolerable Upper Intake Level

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 22. Sodium (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							>AI		>CDRR	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	AI	% (SE)	CDRR	% (SE)
Over 350% poverty													
Males and females:													
1-3.....	12 (4.1)	1928 (75)	1157* (51)	1300* (51)	1553 (63)	1875 (73)	2243 (85)	2629* (108)	2880* (116)	800	>97	1200	94* (1.6)
4-8.....	11 (2.6)	2650 (78)	1651* (55)	1828 (59)	2158 (63)	2585 (74)	3070 (93)	3550 (113)	3871* (125)	1000	>97	1500	>97
Males:													
9-13.....	8* (3.3)	3455 (132)	2226* (110)	2454* (108)	2862 (116)	3382 (135)	3968 (151)	4542* (167)	4946* (189)	1200	>97	1800	>97
14-18.....	12* (5.4)	3979 (140)	2604* (106)	2858* (116)	3341 (123)	3915 (138)	4555 (165)	5157* (190)	5568* (208)	1500	>97	2300	>97
19-50.....	14 (2.6)	4425 (103)	2758 (126)	3061 (118)	3634 (114)	4334 (104)	5119 (99)	5893 (107)	6407 (119)	1500	>97	2300	>97
51+.....	11 (2.3)	4107 (62)	2490 (93)	2775 (85)	3331 (73)	4015 (60)	4784 (80)	5546 (109)	6054 (126)	1500	>97	2300	>97
19+.....	13 (1.8)	4277 (74)	2609 (104)	2913 (97)	3484 (85)	4188 (76)	4970 (79)	5748 (99)	6255 (113)	1500	>97	2300	>97
Females:													
9-13.....	13* (5.2)	3196 (125)	2015* (97)	2231* (104)	2632 (114)	3125 (123)	3686 (141)	4240* (164)	4592* (184)	1200	>97	1800	>97
14-18.....	12* (4.0)	2999 (120)	1876* (92)	2081* (97)	2456 (101)	2927 (119)	3464 (139)	4011* (168)	4355* (189)	1500	>97	2300	82 (4.4)
19-50.....	14 (2.2)	3212 (64)	1916 (54)	2163 (56)	2601 (58)	3136 (64)	3748 (76)	4360 (89)	4734 (99)	1500	>97	2300	86 (1.7)
51+.....	16 (2.9)	2942 (63)	1735 (49)	1953 (51)	2360 (53)	2876 (61)	3443 (73)	4018 (90)	4380 (103)	1500	>97	2300	78 (2.3)
19+.....	15 (1.9)	3070 (56)	1809 (46)	2039 (48)	2468 (49)	3000 (56)	3594 (67)	4195 (82)	4572 (93)	1500	>97	2300	82 (1.8)
All:													
1+.....	13 (0.9)	3572 (43)	1893 (39)	2183 (40)	2726 (43)	3440 (47)	4286 (63)	5139 (76)	5675 (90)		>97		91 (1.0)

NOTES: [†] Percent reporting supplement containing sodium. Sample size presented on page 24.
 AI=Adequate Intake; CDRR=Chronic Disease Risk Reduction Intake
 * Estimate may be less reliable due to small sample size and/or large relative standard error

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

Table TH 23. Potassium (mg): Mean and Percentiles of Total Usual Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, in the United States, 2015-2018

	Percent reporting [†] % (SE)	Mean (SE)	Percentiles							>AI	
			5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	AI	% (SE)
Over 350% poverty											
Males:											
19-50.....	16 (2.3)	3080 (68)	1732 (64)	1977 (64)	2441 (64)	3013 (70)	3641 (74)	4265 (84)	4680 (95)	3400	33 (2.7)
51+.....	28 (2.9)	3219 (66)	1833 (60)	2079 (58)	2565 (60)	3143 (67)	3795 (77)	4441 (94)	4851 (106)	3400	39 (2.8)
19+.....	22 (2.0)	3145 (59)	1777 (57)	2024 (57)	2496 (56)	3072 (61)	3710 (67)	4354 (82)	4765 (90)	3400	36 (2.4)
Females:											
19-50.....	11 (1.8)	2444 (59)	1444 (44)	1633 (48)	1974 (50)	2391 (57)	2852 (65)	3311 (76)	3598 (84)	2600	38 (3.3)
51+.....	20 (2.2)	2512 (61)	1509 (50)	1698 (52)	2040 (53)	2465 (60)	2929 (69)	3388 (81)	3678 (92)	2600	42 (3.5)
19+.....	16 (1.3)	2480 (56)	1479 (45)	1664 (47)	2008 (49)	2430 (55)	2894 (62)	3353 (73)	3642 (84)	2600	40 (3.2)

NOTES: [†] Percent reporting supplement containing potassium. Sample size presented on page 24.
 Estimates for children unavailable due to very low reporting of supplements containing potassium
 AI=Adequate Intake

CITATION: USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018
 Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>

SAMPLE SIZE

	Sample Size	Sample Size by Adult Smoking Status †		
		Smokers <i>Table TH 10</i>	Non-Smokers <i>Table TH 11</i>	Smokers/Non-Smokers <i>Table TH 12</i>
Over 350% poverty				
Males and females:				
1-3.....	191			
4-8.....	265			
Males:				
9-13.....	132			
14-18.....	139			
19-50.....	629	89	540	629
51+.....	656	61	594	655
19+.....	1285	150	1134	1284
Females:				
9-13.....	134			
14-18.....	141			
19-50.....	606	44	562	606
51+.....	622	41	580	621
19+.....	1228	85	1142	1227
All:				
1+.....	3515			

†Excludes individuals without smoking status data.

TABLE NOTES

The statistics in this table were estimated from day 1 and day 2 dietary recall interviews conducted in What We Eat in America (WWEIA), National Health and Nutrition Examination Survey (NHANES) 2015-2018 and 30-day dietary supplement interviews also conducted in the National Health and Nutrition Examination Survey (NHANES) 2015-2018. Trained interviewers, using the USDA 5-step Automated Multiple-Pass Method, conducted the 24-hour dietary recalls. The day 1 recalls were conducted in-person; the day 2 recalls were conducted by telephone 3-10 days after the day 1 recall. Food intakes were coded and nutrient values determined using the USDA Food and Nutrient Database for Dietary Studies 2015-2016 and 2017-2018.

The table includes data from individuals 1 year and over, excluding breast-fed children and pregnant or lactating females. Breast-fed children are excluded because human milk is not quantified in dietary recall interviews. The sample of pregnant and lactating females is not large enough to allow for reliable estimates. Individuals with missing or incomplete 30-day dietary supplement data are also excluded.

Sample weights designed for dietary analysis were used to allow estimates representative of the U.S. population for the years of collection.

The assessments presented in this table cover nutrient intakes from food, beverages (including water) and, dietary supplements. They do not contain intakes from medications.

Usual Nutrient Intakes

Usual intakes (expressed on a per day basis) are an estimate of the usual or habitual intake over the long term, as opposed to what was reported on the days surveyed. The National Cancer Institute (NCI) Method <https://epi.grants.cancer.gov/diet/usualintakes/method.html> was used to produce group usual intake distributions from food and beverages. Daily intake from dietary supplements derived from the 30-day questionnaire and the NCHS Dietary Supplement Database was added to the estimates of usual intake from food and beverages to produce the distributions of total usual nutrient intake presented in this table set. Information on this distribution is presented as mean and percentiles of usual nutrient intake with standard errors for a specific gender/age/lifestyle group.

When applicable, the estimated total usual nutrient intakes are compared to age/gender specific Dietary Reference Intakes (DRIs) <http://nationalacademies.org/HMD/Activities/Nutrition/SummaryDRIs/DRI-Tables.aspx>. The DRIs, nutrient reference values developed by the Institute of Medicine of the National Academies, are used to assess and plan the diets of healthy people. Estimates of percentages less than, within range of, or greater than the DRI, plus the standard error are presented for each age/gender/lifestyle group.

DRI Definitions

Estimated Average Requirement (EAR): The average daily nutrient intake level that is estimated to meet the requirements of half of the healthy individuals in a particular life stage and gender group. The EAR is used to estimate the prevalence of inadequate intakes in a population group.

Adequate Intake (AI): The recommended average daily intake level based on observed or experimentally determined approximations or estimates of nutrient intake by a group (or groups) of apparently healthy people that are assumed to be adequate.

Tolerable Upper Intake Level (UL): The highest average daily nutrient intake level that is likely to pose no risk of adverse health effects to almost all individuals in the general population. As intake increases above the UL, the potential risk of adverse effects may increase.

Chronic Disease Risk Reduction Intake (CDRR): Reductions in intake that exceed the CDRR are expected to reduce chronic disease risk within the apparently healthy population. Established for sodium using evidence of the beneficial effect of reducing sodium intake on cardiovascular disease risk, hypertension risk, systolic blood pressure, and diastolic blood pressure.

Symbol Legend

* Indicates an estimate that may be less statistically reliable than estimates that are not flagged. The rules for flagging estimates are as follows:

Mean: An estimated mean is flagged when based on a sample size of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect, or when the relative standard error is greater than 30 percent. The VIF used in this report is 2.52.

Percentiles: Estimated 50th percentile values are flagged when n is less than 30 times the VIF, where n is the sample size. Estimated values at the other percentiles are flagged when np is less than 8 times the VIF, where n is the sample size and p is 0.25 at the 25th and 75th percentile, 0.10 at the 10th and 90th percentile, and 0.05 at the 5th and 95th percentile.

Percentage above or below a DRI value: An estimated percentage between 25 and 75 percent is flagged when based on a sample size of less than 30 times the VIF, or when the relative standard error is greater than 30 percent. An estimated percentage less than or equal to 25 or greater than or equal to 75 is flagged when the smaller of np and $n(1-p)$ is less than 8 times the VIF, where n is the sample size and p is the percentage expressed as a fraction.

-- Estimate not displayed due to small sample size and for consistency of presentation.

CITATION

USDA, Agricultural Research Service, 2021. Total Usual Nutrient Intake from Food, Beverages, and Dietary Supplements, among Individuals Over 350% of Poverty Level, What We Eat in America, NHANES 2015-2018

Available <http://www.ars.usda.gov/nea/bhnrc/fsrg>